

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/229442747>

# Recent trends and some applications of isothermal titration calorimetry in biotechnology

ARTICLE *in* BIOTECHNOLOGY JOURNAL · JANUARY 2010

Impact Factor: 3.49 · DOI: 10.1002/biot.200900092

CITATIONS

31

READS

196

## 5 AUTHORS, INCLUDING:



**L. Selva Roselin**

Periyar Maniammai University

38 PUBLICATIONS 560 CITATIONS

[SEE PROFILE](#)



**Ming-Shen Lin**

National Central University

14 PUBLICATIONS 227 CITATIONS

[SEE PROFILE](#)



**Yung Chang**

Chung Yuan Christian University

137 PUBLICATIONS 2,932 CITATIONS

[SEE PROFILE](#)



**Wen-Yih Chen**

National Central University

140 PUBLICATIONS 2,433 CITATIONS

[SEE PROFILE](#)

## Review

# Recent trends and some applications of isothermal titration calorimetry in biotechnology

L. Selva Roselin<sup>1</sup>, Ming-Shen Lin<sup>1</sup>, Po-Hsun Lin<sup>1</sup>, Yung Chang<sup>2</sup> and Wen-Yih Chen<sup>1</sup>

<sup>1</sup> Department of Chemical and Materials Engineering, National Central University, Jhong-Li, Taiwan

<sup>2</sup> R&D Center for Membrane Technology and Department of Chemical Engineering, Chung Yuan Christian University, Jhong-Li, Taoyuan, Taiwan

Isothermal titration calorimeters (ITCs) are thermodynamic instruments used for the determination of enthalpy changes in any physical/chemical reaction. This can be applied in various fields of biotechnology. This review explains ITC applications, especially in bioseparation, drug development and cell metabolism. In liquid chromatography, the separation/purification of specific proteins or polypeptides in a mixture is usually achieved by varying the adsorption affinities of the different proteins/polypeptides for the adsorbent under different mobile-phase conditions and temperatures. Using ITC analysis, the binding mechanism of proteins with adsorbent solid material is derived by elucidating enthalpy and entropy changes, which offer valuable guidelines for designing experimental conditions in chromatographic separation. The binding affinity of a drug with its target is studied by deriving binding enthalpy and binding entropy. To improve the binding affinity, suitable lead compounds for a drug can be identified and their affinity tested by ITC. Recently ITC has also been used in studying cell metabolism. The heat produced by animal cells in culture can be used as a primary indicator of the kinetics of cell metabolism, which provides key information for drug bioactivity and operation parameters for process cell culture.

Received 9 April 2009  
Revised 17 August 2009  
Accepted 19 August 2009

**Keywords:** Binding affinity · Biorecognition · Bioseparation · Isothermal titration calorimetry · Thermodynamics

## 1 Introduction

All physical, chemical and biological processes are accompanied by heat exchange. The enthalpy change of a reaction,  $\Delta H^\circ$ , is a fundamental thermodynamic quantity that describes the amount of heat released or absorbed during the course of the reaction. The  $\Delta H^\circ$  of a reaction can, in general, be determined in one of the two ways; it can be determined directly using calorimetry or indirectly by

van't Hoff method. For a reversible reaction the van't Hoff relationship is described by Eq. (1),

$$\Delta H^\circ = -R \left( \frac{\partial \ln K}{\partial 1/T} \right) \quad (1)$$

where  $R$  is the gas constant,  $T$  is the absolute temperature (at constant pressure). The instantaneous slope of a plot of  $\ln K$  vs.  $1/T$ , multiplied by  $-R$ , is used to determine  $\Delta H^\circ$  [1].

Differential scanning calorimetry (DSC) and isothermal titration calorimetry (ITC) are the only methods for the direct determination of  $\Delta H$ . DSC measures the heat capacity of a reaction as a function of temperature. The heat capacity of the system is then measured against temperature as the direct thermodynamic observable. Therefore, in a single DSC experiment, transition midpoint –  $T_m$ , enthalpy ( $\Delta H$ ) and heat capacity change ( $\Delta C_p$ ) can be determined. ITC is the direct method to measure

**Correspondence:** Professor Wen-Yih Chen, Department of Chemical and Materials Engineering, National Central University, Jhong-Li 320, Taiwan  
**E-mail:** wychen@cc.ncu.edu.tw  
**Fax:** +886-3-422-5258

**Abbreviations:** EDL, electrical double layer; HEWL, hen egg white lysozyme; HIC, hydrophobic interaction chromatography; IEC, ion-exchange chromatography; IMAC, immobilized metal ion affinity chromatography; ITC, isothermal titration calorimetry/calorimeter

the heat change in a reaction at constant temperature. In ITC analysis, the measurement of heat is made at a given temperature, which allows direct determination of binding constants ( $K_a$ ), reaction stoichiometry ( $n$ ), enthalpy change ( $\Delta H$ ), free energy change ( $\Delta G$ ) and entropy change ( $\Delta S$ ). In addition, varying the temperature of the experiment allows the determination of the heat capacity ( $\Delta C_p$ ) for the reaction, thereby providing a complete thermodynamic profile of the molecular interaction.

When the van't Hoff enthalpy values for a particular reaction are compared to enthalpy for the same reaction obtained directly by calorimetric methods, there is often a significant difference between the two values. Furthermore, the binding mechanism may differ with temperature, and the heat capacity of the molecules involved may also vary with operating temperature. For example, the binding enthalpy of cytidine 2'-monophosphate (2'-CMP) to ribonuclease A (RNase A) revealed that the enthalpy obtained from the van't Hoff equation ( $\Delta H_{vH}$ ) differed from those obtained from microcalorimetric measurement ( $\Delta H_{ITC}$ ) at various temperatures and binding amounts [2]. This indicates that the binding reaction is more complex than the simple one-to-one binding model. The direct measurement of  $\Delta H_{ITC}$  represents the total energy change in the whole adsorption process, which includes dilution heat, binding heat, the energy required for desolvation, and the energy change after protein rearrangement. In contrast,  $\Delta H_{vH}$  is calculated on the basis of the reversible equilibrium process, and the analysis does not consider the contribution of protein conformation change. Furthermore, the  $\Delta H_{vH}$  is calculated from the free energy change at different temperatures. Since protein conformation is sensitive to temperature,  $\Delta C_p$  varies with temperature, causing variation of  $\Delta H_{vH}$ . Therefore, major contributing factors to the discrepancy between the  $\Delta H_{vH}$  and  $\Delta H_{ITC}$  are difference in the mechanism of protein adsorption caused by the changes in protein conformation and the interaction with solvent molecules and the hydrophobic ligand [3]. Weber has discussed the reasons for the differences between van't Hoff and calorimetric enthalpy values in detail [4]. Chaires [5] and Horn *et al.* [6] have noted that the enthalpy observed by van't Hoff analysis is considerably lower than that for calorimetric enthalpy and this discrepancy is associated with the reduced precision of van't Hoff analysis. Therefore, ITC is considered an accurate method for estimating the total heat change in various biological reactions. ITC is mostly used to measure strengths of interaction between proteins and ligands: either small molecules like drugs, or large ones such as nucleic acids or

other proteins. The biotechnological issues to which microcalorimetry has been successfully applied includes antibody studies [7]; cell metabolism [8]; drug receptor interactions [9]; enzyme interactions, *e.g.*, with co-enzymes [10], inhibitors [11], substrates [12], and drugs [13]; antigen-antibody interactions [14]; DNA-drug interactions [15]; protein interactions, *e.g.*, with other proteins, lipids, carbohydrates, DNA, ligands, and nucleic acids [16–20]; and biopolymer interactions, *e.g.*, with other biopolymers, drugs, ligands, [21–23], and enzyme activity analysis [24].

This review article reports our experiences and other reported results on ITC in the investigation of binding interaction between protein and solid surfaces for its application to bioseparation by liquid chromatography. In addition, we give a brief discussion on the application of this technique in drug development and studying cell metabolism.

## 2 Instrumental methodology

The calorimetric method for determination of binding constants was first developed in the early 1960s for the evaluation of thermodynamics of proton ionization from weak acids and bases, and for metal-ligand binding reactions [25]. Design improvements were made in an adiabatic calorimeter with no heat exchange taking place between the calorimetric vessel and its surroundings. The amount of heat that is evolved or absorbed in an ideal adiabatic calorimeter is equal to the product of the measured temperature change and the heat capacity of the vessel, including its content. A wide range of calorimeters are available to meet different needs of the user [26–29]. A typical isothermal titration calorimeter is composed of two identical cells (a reference and a sample cell) made of highly efficient thermal conducting material surrounded by an adiabatic jacket. Sensitive thermopile/thermocouple circuits are used to detect temperature differences between the reference cell (filled with buffer or water) and the sample cell containing the macromolecule. Known concentrations of degassed samples are taken in the injector and sample cell. Prior to the injection of the titrant, a constant power (<1 mW) is applied to the reference cell. This signal directs the feedback circuit to activate the heater located on the sample cell. This represents the baseline signal. During the experiment, the titrant is added into the sample cell in precisely known aliquots, causing heat to be either taken up or evolved. For an exothermic reaction, the temperature in the sample cell will increase, and the feedback power will be deactivated to maintain

equal temperatures between the two cells. For endothermic reactions, the reverse will occur, *i.e.*, the feedback circuit will increase power to the sample cell to maintain the temperature.

ITC instruments should be routinely calibrated by applying specified electrical pulses of approximately 5–10  $\mu$ cal/s. The total heat as determined by the area under the pulse should be within 2% of the expected value (Omega ITC Manual, Microcal Inc.). Baranauskienė *et al.* [30] have reported several chemical reactions to validate ITC equipment. Joel Tellinghuisen [31] used the heat of dilution of NaCl (aq) to calibrate a Micro Cal model (VP-ITC) instrument. The performance of the calorimeter can also be tested by measuring the heat of a standard chemical reaction, such as the protonation of tris(hydroxymethyl)aminomethane (THAM) by HCl [32]. It is advisable to use both instrumental calibration and one chemical calibration method to calibrate the instrument.

Measurements consist of the time-dependent input of power required to maintain equal temperature between the sample and reference cells. Observations are plotted as the power needed to maintain the reference and the sample cell at an identical temperature. This power is given as a function of time in seconds. The raw data for an experiment consists of a series of heat flow peaks (power), with every peak corresponding to a ligand injection. These heat flow peaks are integrated with respect to time, giving the total heat effect per injection. Peak integration is made either in a manual peak-by-peak fashion or automatically using the ITC software package. Baseline selection is an important factor in ITC data analysis and user input in the automated integration routine is limited. Therefore, manual peak-by-peak integration in which the operator defines the baseline regions used in the integration is preferred. This heat released or absorbed is in direct proportion to the amount of binding that occurs. Additionally, a volume correction is also performed due to dilution in the sample cell due to each injection.

Interpretation of calorimetric data always involves a model, usually expressed as an equation or set of equations. In the case of a receptor with multiple binding sites of different affinity, statistical thermodynamic treatment of the data is required. The amount of heat evolved on addition of ligand can be represented by Eq. (2) [33]:

$$Q = V_0 \Delta H_b [M]_i K_a [L] / (1 + K_a [L]) \quad (2)$$

where  $V_0$  is the volume of the cell,  $\Delta H_b$  is the enthalpy of binding per mole of ligand,  $[M]_i$  is the total macromolecule concentration including bound

and free fractions,  $K_a$  is the binding constant, and  $[L]$  is the free ligand concentration. For a more general model of binding, the multiple independent sites model, the macromolecule contains multiple ligand binding sites that are noninteracting. The total heat of binding can be described by

$$Q = V_0 [M]_i \sum (n_i \Delta H_i K_{ai} [L]) / (1 + K_{ai} [L]) \quad (3)$$

A detail discussion of the derivation of the above equations is presented in [33]. The method of analysis will be specific to the system under investigation and it is advised to attempt fitting according to a general model before using more specific models. The pattern of these heat effects as a function of the molar ratio [ligand]/[macromolecule] can then be derived to give the thermodynamic parameters.

The thermodynamic parameters are derived as follows:

$$\Delta G = \Delta G^0 + RT \ln Ka \quad (4)$$

At equilibrium  $\Delta G = 0$

$$\therefore \Delta G^0 = -RT \ln Ka = \Delta H^0 - T \Delta S^0 \quad (5)$$

where  $\Delta G$ ,  $\Delta G^0$ ,  $\Delta H^0$ ,  $\Delta S^0$  and  $K_a$  are free energy change, standard free energy change, standard enthalpy change, standard entropy change and affinity constant, respectively. The affinity constant  $K_a$  is derived from isotherm study. As  $\Delta H$  is dependent on concentration and is not considered as standard state (at standard state [conc] = 1 M), the  $\Delta H$  value is directly measured by ITC analysis.

From  $\Delta G^0$  and  $\Delta H$ ,  $\Delta S^*$  is derived as follows [34].

At extreme dilution condition

$$\Delta S^* = \Delta S_{unitary} + \Delta S_{cratic} \quad (6)$$

$\Delta S_{unitary}$  is given by Eq. (7) [35]

$$\Delta S_{unitary} = \Delta S^0 - \Delta S_{cratic} = \Delta S^0 - R \ln X_M \quad (7)$$

Where,  $\Delta S_{unitary}$  is independent on concentration.  $\Delta S_{cratic}$  is dependent on concentration.  $\Delta S_{cratic}$  is given by Eq. (8)

$$\Delta S_{cratic} = R \ln X_M \quad (8)$$

$X_M$  is the concentration of water in dilute aqueous solution, which is given by a value 55.6 M [36] and so the  $\Delta S_{cratic}$  is calculated as  $-7.98 \text{ cal mol}^{-1} \text{K}^{-1}$ . From this,  $\Delta S_{cratic}$  was derived and  $\Delta G_{unitary}$  was obtained from Eq. (9).

$$\Delta G_{unitary} = \Delta H + T \Delta S_{unitary} \quad (9)$$

### 3 Examples of ITC applications in biotechnology

#### 3.1 ITC applications in bioseparation

The chromatographic separation process is based on the difference in the surface interactions of the analyte and eluent molecules. There are two basic approaches for thermodynamic description of the HPLC retention phenomena, one is based on the partitioning theory and another is based on adsorption. Partition is a concentration change in the system due to the distribution of the components between two (or more) phases and adsorption is the concentration change in the system in presence of interface with another phase and due to the surface forces. One of the main problems with chromatographic separation is the difficulty in the optimization of different parameters to design effective process for protein separation/purification, which is presently fixed by trial and error. Knowledge of the binding mechanism of proteins at the liquid-solid interface provides valuable insights into the separation and purification of proteins. The separation/purification of specific proteins or polypeptides in a mixture is usually achieved by varying the adsorption affinities of the different proteins/polypeptides for the adsorbent. The thermodynamic characteristics provide valuable information for understanding the binding mechanism [37].

The feasibility of a protein or polypeptide being adsorbed (at a constant pressure  $P$  and temperature  $T$ ) is determined by the change in the standard Gibbs free energy,  $\Delta G^\circ$ , of the system. The relationship between standard enthalpy ( $\Delta H^\circ$ ) and entropy ( $\Delta S^\circ$ ) changes, the equilibrium binding affinity constant, ( $K_a$ ) and the corresponding standard Gibbs free energy change can be expressed by Eq. (5). Proteins or polypeptides can be separated by varying the magnitudes of the respective enthalpy and/or entropy changes, resulting in different binding affinities and different free energy changes. Using microcalorimetry, the heat of protein adsorption onto solid surfaces can be measured. The heat measured can be further quantified by the amount of protein adsorbed, which is obtained from the equilibrium binding isotherms. The heat of protein adsorption onto solid surfaces can be measured by ITC [38–50]. In this experimental technique, the adsorbent is suspended in the equilibrium buffer solution and placed in the sample cell: it is then titrated by protein solution prepared in the equilibrium buffer solution. The output signals are collected as power ( $P$ ) vs. time ( $t$ ) and are integrated and quantified by the amount of bound protein to give the enthalpy change of adsorption.

The adsorption enthalpy change can be calculated as:

$$Q_{ads} = V \times q \times \Delta H_{ads} \quad (10)$$

where  $Q_{ads}$  denotes the net heat attributed to the interaction between proteins and adsorbents (after corrected by the dilution heat of proteins and adsorbents) [J],  $V$  represents the volume of the adsorbent in the ampoule (mL),  $q$  denotes the amount of bound protein (mol/mL), and  $\Delta H_{ads}$  is the enthalpy change (kJ/mol).

Using isothermal calorimetric technique, detailed thermodynamic characteristics have been reported on hydrophobic interaction chromatography (HIC), immobilized metal ion affinity chromatography (IMAC), and ion-exchange chromatography (IEC).

The binding mechanism can be explained by considering five sequential subprocesses: (a) exclusion of water molecules or ions surrounding the protein surface, *i.e.*, the dehydration or deionization (removal of the electrical double layer, EDL) subprocess of the protein; (b) exclusion of water molecules or ions surrounding the adsorbent, *i.e.*, the dehydration or deionization subprocess of the adsorbent; (c) the hydrophobic interactions between the protein and the hydrophobic HIC adsorbent (in HIC); coordination interaction and/or non-specific interaction formed between protein and the immobilized metal ion (in IMAC); electrostatic interactions and/or nonspecific interactions, *e.g.*, hydrophobic interactions between the protein and the ion exchange adsorbent (in IEC); (d) the structural rearrangement of the protein upon adsorption/partition; and (e) the structural rearrangement of the excluded water molecules or ions in the bulk solvent.

##### 3.1.1 Hydrophobic interaction chromatography

ITC analysis is helpful in choosing a suitable hydrophobic adsorbent, salt type and salt concentration for separation/purification of polypeptide or proteins by HIC [38–43]. In addition, the values of adsorption enthalpy ( $\Delta H$ ) for different proteins provide guidelines to separate these proteins by HIC. For example, ITC analysis revealed that adsorption enthalpies ( $\Delta H$ ) of  $\alpha$ -chymotrypsinogen A with two adsorbents (CM-butyl and CM-octyl Sepharose) were lower than those of trypsinogen, in which  $\alpha$ -chymotrypsinogen A had a larger exposed hydrophobic area than trypsinogen (Figs. 1 and 2) [39]. These can be result from a higher exothermic amount of hydrophobic interaction and reduction in the endothermic amount for structural rearrangement of  $\alpha$ -chymotrypsinogen A than




for trypsinogen. As a result,  $\alpha$ -chymotrypsinogen A has a longer retention time than trypsinogen and are separated by HIC. In another study, peptides consisting of glycine and different lengths of tryptophan such as, GWG, GWWG, GWWWG were investigated for its adsorption behavior with hydrophobic adsorbents such as, octyl Sepharose and CM-octyl Sepharose [43]. Tryptophan is an unusual amino acid and plays an important role in biological systems [51–53]. The binding mechanism of tryptophan adsorption on octyl Sepharose and CM-octyl Sepharose was elucidated by equilibrium binding analysis, dilution heat of peptide solution and adsorption enthalpy. The mechanisms of CM-octyl Sepharose adsorption with GWG and GWWG involve hydrophobic and electrostatic interactions. However, the binding of GWWWG with the adsorbent is dominated by the hydrophobic interaction. These findings suggest that tryptophan chain length or the number and hydrophobicity of amino acid residues of a peptide is critical for the peptide binding and solvation mechanism. The thermodynamic data for peptides containing different tryptophan chain length offer fundamental information for protein separation and peptide design application.


ITC analysis is helpful in choosing adsorbents and provides valuable guidelines to design experimental parameters for the specific adsorbent for separating the proteins or polypeptides. Hydrophobic interaction differs for adsorbents with different chain length, type and density. Alkyl-based and phenyl-based adsorbents differ in their adsorption affinity for proteins and this can be result in different interaction mechanisms. Compared to n-alkyl-based HIC adsorbents, the interaction between proteins and HIC adsorbents con-

taining aromatic groups such as phenyl or benzyl ligands follow a less predictable binding behavior, attributed to the participation of  $\pi$ - $\pi$  interactions [54]. Comparison of the equilibrium binding affinity for alkyl and phenyl adsorbents with hen egg white lysozyme (HEWL) showed that the adsorption affinity of the HEWL with the Toyopearl butyl adsorbents was larger than that found with the Toyopearl phenyl adsorbent [39]. ITC analysis illustrated that both adsorption enthalpy ( $\Delta H_{\text{ads}}$ ) and adsorption entropy ( $\Delta S_{\text{ads}}$ ) values for Toyopearl butyl adsorbent with HEWL were higher than that with the Toyopearl phenyl adsorbent. The endothermic amount of the heat associated with the dehydration and the reduction in the EDL (*i.e.*, the a and b subprocesses) was smaller when the Toyopearl butyl adsorbent was used in comparison to the Toyopearl phenyl adsorbent. The binding strength associated with the hydrophobic interactions of HEWL with the Toyopearl phenyl adsorbent was stronger than that with the Toyopearl butyl adsorbent, leading to an increase in the exothermic amount of heat (*i.e.*, the c subprocess). This result provides evidence to support the conclusion that chromatographic processes based on the hydrophobic interactions are mostly driven by the entropy gain. This suggests that under these conditions separation/purification of proteins or polypeptides can be performed by considering temperature as the key parameter.

In another study, n-alkyl-based HIC adsorbents, such as the butyl- and octyl-containing adsorbents were compared [39, 40, 42]. The microcalorimetric results confirmed that the adsorption of two proteins ( $\alpha$ -chymotrypsinogen A and trypsinogen) on CM-butyl Sepharose was exothermic, while its adsorption on CM-octyl Sepharose was endothermic.



**Figure 1.** Adsorption enthalpies ( $\Delta H$ ) of  $\alpha$ -chymotrypsinogen A (CHY) or trypsinogen (TRY) onto CM-butyl Sepharose at various ligand densities, 4.0 M NaCl, and pH 10.0 (the number in the key of the figure represents the ligand density of the adsorbent in mol/mL) [39].



**Figure 2.** Adsorption enthalpies ( $\Delta H$ ) of  $\alpha$ -chymotrypsinogen A (CHY) or trypsinogen (TRY) onto CM-octyl Sepharose at various ligand densities, 4.0 M NaCl, and pH 10.0 (the number in the key of the figure represents the ligand density of the adsorbent, in mol/mL) [39].

The adsorption per mole of these proteins onto CM-octyl Sepharose required more energy for dehydration than that with CM-butyl Sepharose, and higher entropy gain with CM-octyl Sepharose as compared with CM-butyl Sepharose. Therefore, the interaction mechanism of the protein with CM-octyl Sepharose is more like a partitioning event, while with the CM-butyl Sepharose the mechanism is an adsorption-dominated process. Therefore, when CM-octyl Sepharose is used as the adsorbent, temperature is a powerful parameter that can be adjusted for better separation/purification of proteins or polypeptides by HIC.

The adsorbents CM-octyl Sepharose and octyl Sepharose have similar ligand density and both have exposed OH groups on the surface. However, CM-octyl Sepharose has partially unreacted carboxyl groups that remain on the surface. The adsorption behavior and thermodynamic parameters with octyl Sepharose and CM-octyl Sepharose with the oligomers (GWG, GWWG, GWWWG) showed that the binding affinities of each peptide for the two adsorbents are similar. However, the mechanism of adsorption for peptides with these hydrophobic ligands is quite different with respect to the binding enthalpy between peptides and adsorbents [43]. The adsorption of the peptides on octyl Sepharose is an entropy-driven process for all the peptides and is governed by hydrophobic interaction. In contrast, with CM-octyl Sepharose hydrophobic and electrostatic interactions are both included in the binding operation for GWG and GWWG. However, the binding of GWWWG with the adsorbent is dominated by the hydrophobic interaction. This finding is useful for fixing the eluent gradient and temperature for HIC separation. When both hydrophobic and electrostatic forces influence the interaction between proteins and the adsorbent, two parameters, such as temperature and salt concentration, are taken into consideration and are adjusted based on the relative values of  $\Delta H_{\text{ads}}$  and  $\Delta S_{\text{ads}}$ . When hydrophobic force alone influences the interaction between proteins and the adsorbent, good separation/purification of protein/polypeptide can be achieved by selecting suitable temperature.

Temperature plays an important role in separation/purification of proteins by HIC [55]. At different temperatures the interaction may be dominated by partitioning, adsorption, or both. Partitioning implies that the protein molecule increases exposure of the hydrophobic residue of the inner protein core to aqueous solution and penetrates the solid surface, whereas adsorption implies that the protein molecule is simply in surface contact with the adsorbent. ITC analysis helps in determining


the binding mechanism and in choosing a suitable temperature for separation/purification of proteins by HIC. For example, the affinity and enthalpy differences between two proteins ( $\alpha$ -chymotrypsinogen A and trypsinogen) with two adsorbents (butyl Sepharose and octyl Sepharose) under varying temperatures showed that the adsorption quantity of the proteins with these Sepharose adsorbents increased when the temperature was increased from 298 to 310 K [38, 42]. But  $\Delta H_{\text{ads}}$  value of  $\alpha$ -chymotrypsinogen A with butyl Sepharose was increased, while the  $\Delta H_{\text{ads}}$  value of trypsinogen was reduced. Comparison of the value of adsorption entropy ( $\Delta S_{\text{ads}}$ ) showed that the  $\alpha$ -chymotrypsinogen A exhibited higher value than trypsinogen. The higher entropy gain was mainly from the dehydration process, which characterized the partitioning process. This observation implies that as the temperature is increased, the solution polarity is reduced, and the decreased stability of the protein structure and the increased exposure of the hydrophobic residue of the inner protein core to aqueous solution lead to interaction mechanism changes from an adsorption-dominated process to a partitioning process. In addition, for octyl Sepharose, the  $\Delta H$  value of  $\alpha$ -chymotrypsinogen A was positive, but decreased with temperature increment. However, the  $\Delta H$  value of trypsinogen was positive and increased with temperature. Therefore, as the temperature increases, the interaction mechanism of the proteins for octyl Sepharose is a partitioning-dominated process. These observations are useful for understanding the temperature and eluent gradients required for chromatographic separation/purification of proteins/polypeptides.

The retention of protein is achieved by varying the salt concentration in the aqueous mobile phase. Horvath and co-workers [56] demonstrated that, in the absence of special binding effects, an increase in the salt molality in the mobile phase or the use of a salt with a greater molal surface tension increment will result in the increased retention. The molal surface tension increment is a concentration-independent characteristic of the salt that relates to its water-structuring potential. Thus, the molal surface tension increments of salts have been reported to highly correlate with their position in the lyotropic series [57], which is an empirical relationship discovered by Hofmeister [58] linking the solution properties of salts and their effect on the solubility of proteins. It has been noted that certain salts enter into specific interaction with proteins. The effect of salt type on protein retention can be related to the molal surface tension increment of the salt [56]. ITC analysis provides a clear picture

of the retention characteristics of different salt types. Higher affinities were obtained in the presence of  $\text{Na}_2\text{SO}_4$  compared to  $(\text{NH}_4)_2\text{SO}_4$  in two adsorbents (Toyopearl butyl and Toyopearl phenyl) [39]. ITC analysis demonstrated that lower values of  $\Delta H_{\text{ads}}$  were obtained in the presence of  $\text{Na}_2\text{SO}_4$  when compared to  $(\text{NH}_4)_2\text{SO}_4$  due to the greater ability to decrease the endothermic component of the dehydration of the protein and HIC adsorbent and to reduce the EDL (*i.e.*, the a and b subprocesses). In addition, the exothermic amount of heat (*i.e.*, the c subprocess) with  $\text{Na}_2\text{SO}_4$  was higher than with  $(\text{NH}_4)_2\text{SO}_4$ . This resulted in enhanced interaction between protein and the HIC adsorbent when  $\text{Na}_2\text{SO}_4$  was used. Since  $\text{Na}_2\text{SO}_4$  has a higher molal surface tension increment, a more solvophobic environment will occur for HEWL (and the other proteins) in an aqueous solution containing  $\text{Na}_2\text{SO}_4$  rather than  $(\text{NH}_4)_2\text{SO}_4$ . Thus, salts with higher molal surface-tension increments produce increased retention at equal molal salt concentrations. The retention of proteins on hydrophobic columns can be controlled by adjustment of salt concentrations [59]. The adsorption affinity of the myoglobin with either butyl or octyl Sepharose was increased with the salt concentrations [40]. ITC analysis illustrated that the  $\Delta H_{\text{ads}}$  of myoglobin adsorption onto butyl and octyl Sepharose were all decreased with increase in  $(\text{NH}_4)_2\text{SO}_4$  concentration. This was due to the heat revealed by van der Waal's forces of the protein with the adsorbent (*i.e.*, the (c) process) increased with  $(\text{NH}_4)_2\text{SO}_4$  concentrations. The increase of  $\Delta H_{\text{ads}}$  value was compensated by the larger positive value of  $\Delta S_{\text{ads}}$ . For a better separation/purification of protein/polypeptide by HIC, concentration of salt and temperature can be adjusted by considering the  $\Delta H_{\text{ads}}$  and  $\Delta S_{\text{ads}}$  values.

### 3.1.2 Immobilized metal ion affinity chromatography

ITC measurement can also find application in IMAC. The interaction mechanism of a protein with an immobilized metal ion is affected by the topography of the protein surface through various parameters, such as type of metal ion, pH of the medium and ionic salt concentration. ITC analysis is helpful for choosing a suitable metal ion and to set up suitable conditions for the chosen metal ion in IMAC [45–49]. The  $\Delta H_{\text{ads}}$  values of lysozyme adsorption onto immobilized Fe(III) are higher than those of Cu(II) at various pH values (Fig. 3) [48]. The heat required for the dehydration of Cu(II) is lower than that for the dehydration of Fe(III), and the heat generated from the formation of the coordination with Cu(II) is higher than with Fe(III). The coordinate bond formation of nitrogen atoms with



**Figure 3.** Enthalpy of adsorption ( $\Delta H_{\text{ads}}$ ) of lysozyme onto CS-IDA-Fe(III) or CS-IDA-Cu(II) at various pH values, 0.1 M NaCl, and 20 mM phosphate [48].

Cu(II) is more competent than with Fe(III), implying that the coordination of the nitrogen atom with Cu(II) is more thermodynamically stable than with Fe(III). The adsorption affinities and retention of proteins fused with histidine tags towards immobilized metal ions increased with the number of exposed histidines. The  $\Delta H_{\text{ads}}$  values of SjGST/His adsorption onto the Ni-NTA were lower than that of SjGST [49], because SjGST/His has an additional fusion tag containing six adjacent histidines in comparison to SjGST, which coordinated with immobilized  $\text{Ni}^{2+}$  resulting in a reduction in  $\Delta H_{\text{ads}}$  values. As a result, SjGST/His was retained more strongly than SjGST. This suggests that the retention of each protein can be adjusted on the basis of histidine tags.

### 3.1.3 Ion-exchange chromatography

In IEC, the binding strength of the protein with the ion-exchange resin can be affected by numerous factors, including pH value, salt concentration, and temperature, *etc.*, and these effects have been widely investigated [60, 61]. The thermodynamic data revealed that at a pH nearer to the *pI* value of protein, both electrostatic and hydrophobic forces mutually affect the interaction behavior in IEC [44, 50]. Similar observations are also made by Dieterle *et al.* [62] in adsorption studies of two human monoclonal antibodies on two different strong cation exchange resins.

## 3.2 ITC applications in drug development

ITC has an important application in the characterization and optimization of lead compounds in drug development. A lead compound in drug discovery is



a chemical compound that has pharmacological or biological activity and whose chemical structure is used as an active site for chemical modifications to improve selectivity or pharmacokinetic parameters. Compounds identified by high-throughput screening usually have weaker affinities towards the target. The binding affinity and selectivity of these compounds are increased by suitable lead compounds [63]. One of the major challenges for ITC has been the identification of ligands with high binding affinities. ITC has been applied in drug design, where the binding affinities of the drug to the target were identified from thermodynamic parameters, such as free energy change ( $\Delta G$ ), enthalpy change ( $\Delta H$ ), entropy change ( $\Delta S$ ) and binding affinity constant ( $K_a$ ) values. These thermodynamic parameters are related to each other by Eq. (5). For spontaneous binding between the drug and the target,  $\Delta G$  is negative and is directly related to binding affinity.

High affinity between drug and target is achieved only when enthalpy and/or entropy contribute favorable energy. For a favorable binding between the drug and the target, the  $\Delta H$  value is negative and  $\Delta S$  value is positive. The favorable enthalpy is associated with the formation of hydrogen bonds and van der Waal's interactions; and the unfavorable enthalpy is associated with the desolvation of polar groups. The enthalpy associated with the desolvation of polar groups is  $\sim 8$  kcal/mol [64], which is one order of magnitude higher than that of non-polar groups. For favorable enthalpy, the drug has to interact with the target by overcoming the unfavorable enthalpy resulted by desolvation. Therefore, the  $\Delta H$  value represents the strength of the drug-target interaction relative to those with solvent. Ruben *et al.* [65] have reported the optimization of an antimalarial target, namely plasmeprin II inhibitors. In plasmeprin II inhibitors, a hydrogen bond donor is placed at different positions in a phenyl ring and the enthalpy change was studied by ITC. The hydroxyl group located at para position showed favorable enthalpy and this was considered the best position. On the other hand, by placing a methyl group at the same position results in an unfavorable binding enthalpy of 4.4 kcal/mol, which confirmed the best functional group at that position is a hydrogen bond donor. This shows that ITC is the powerful tool for identifying the most appropriate position for hydrogen bond donors or acceptors in the drug and target. Thermodynamics data for inhibitor binding to renin were derived by ITC [66]. The binding reaction between inhibitor and rennin was driven by favorable enthalpy changes ( $\Delta H = -9.27$  kcal mol<sup>-1</sup>) but it was counterbalanced by unfavorable entropy changes ( $T\Delta S$

$= -2.00$  kcal mol<sup>-1</sup>). When the entropy term was improved, there was a significant loss in the enthalpy term. Synthetic efforts were initiated by molecular alterations and finally the entropy term was improved by designing a compound by adding an additional electrostatic interaction, which had a favorable entropy term ( $T\Delta S = 0.43$  kcal mol<sup>-1</sup>) with an almost unchanged  $\Delta H$  value. As a result the affinity of the optimized inhibitor was increased to about 45-fold compared with the initial compound.

Even though enthalpic interactions are required for extremely high affinity, the optimization of the binding enthalpy has been more difficult than the optimization of the binding entropy, the reason being that the enthalpy of desolvation of polar groups is very large and unfavorable [67]. In the binding thermodynamics of seven clinical protease inhibitors (indinavir, saquinavir, nelfinavir, ritonavir, amprenavir, lopinavir and atazanavir), the inhibitors (indinavir, nelfinavir, saquinavir and ritonavir) are dominated by a strong favorable entropy change [68]. The binding enthalpy is unfavorable for indinavir, nelfinavir and saquinavir, and only slightly favorable for ritonavir. Experimental results indicate that in the inhibitors, indinavir, ritonavir, saquinavir, and nelfinavir, the inhibitor/protein interactions are not strong enough to overcome the unfavorable enthalpy associated with the desolvation of protein and inhibitor group and the unfavorable enthalpy associated with the change in conformation of the protease upon binding. In all cases, favorable entropy contributes to the Gibbs energy. Two main factors contribute to the extremely favorable entropy change: a large favorable solvation entropy change that originates from the release of water molecules associated with the burial of a significant hydrophobic surface upon binding and a small unfavorable conformational entropy change. Therefore, entropic optimization is achieved by increasing the hydrophobicity of drug candidates and by introducing conformational constraints [69]. This favorable solvation entropy resulted from burial of hydrophobic groups. Protasevich *et al.* [70] studied the inhibitor affinity of four 2-(2-hydroxyphenoxy)phenol inhibitors, wherein the 4-position substituent varied from H to n-propyl to determine the contribution of the aliphatic chain to the binding to the wild-type (wt) enoyl-ACP reductase from *Escherichia coli* (FabI) and a drug-resistant mutant, (F203L)FabI, in which phenylalanine 203 is mutated to leucine. The inhibitor affinity to wt FabI and (F203L)FabI were increased with increasing aliphatic chain length, although the related affinity for (F203L)FabI was lower, and also, it showed no detectable binding to the 4-H inhibitor. The binding enthalpy became

less favorable with increasing aliphatic chain length. Therefore, the increase in affinity was due to favorable changes in solvation entropy. Hydrogen bond donor/acceptor functionality attached to a large aromatic ring can force part of the ring to be exposed to the solvent. This brings about a drop in the favorable desolvation entropy that partially or completely neutralizes any enthalpic gains due to hydrogen bond. In addition, when hydrogen bond donor and acceptor groups are stronger in the compound, these are hydrogen-bonded to water before binding. Therefore, the amount and type of hydrogen bonding is highly indicative of the specificity of the reaction. However, too many hydrogen donor groups on the drug molecule will cause low bioavailability due to poor cell membrane permeability. Conversely, less hydrogen binding results in poor specificity. Lipinski *et al.* [71] suggested that four parameters should be globally associated with solubility and permeability; namely molecular weight; Log P; the number of H-bond donors and the number of H-bond acceptors. The 'Lipinski rule of 5' states that: poor absorption or permeation are more likely when: there are more than 5 H-bond donors; the molecular weight is over 500; the Log P is over 5; and there are more than 10 H-bond acceptors. He also added that if two parameters are out of range, a poor absorption or permeability is possible. However, Li *et al* [72] reported in an analysis of the data in PDBcal that a diverse set of ligands with only 38% of the studies was obeying the Lipinski's Rule of Five. ITC data provide a way to optimize the number of hydrogen bonds with a minimum number of hydrogen bond donors and acceptors. Ernesto Freire [73] explained that for a highly favorable binding enthalpy it is not the number of polar groups present in the drug molecule that counts, but the quality of the hydrogen bonds that are made with the target protein. For example, two protease inhibitors, saquinavir and TMC-126, both have 11 polar atoms; however, saquinavir binds with unfavorable enthalpy (+1.5 kcal/mol) whereas TMC-126 binds with a highly favorable one (−12 kcal/mol) [74].

Biological systems are subject to enthalpy-entropy compensation, where the enthalpic gain from changing the structure of a compound to increase bonding interactions is offset by an entropic penalty, thus reducing the magnitude of change in free energy [75, 76]. The thermodynamics of the drug inhibitors acyclovir, ganciclovir, and 9-benzylguanine binding to human purine nucleoside phosphorylase (hsPNP) as a function of the substrate phosphate ion (Pi) concentration [77] showed that the binding reaction changed from an enthalpically driven reaction in the absence of Pi to an en-

tropically driven reaction at 10 mM Pi. The dependencies of the driving-nature of the binding reactions on Pi concentration can be simulated by Pi binding to its catalytic site. However, the binding constants are unaffected by change in the bound Pi concentration because of enthalpy-entropy compensation. Burial of the apolar surface of small-molecule hydrocarbons is known to be accompanied by a favorable entropy change in aqueous solution due to the release of water molecules from the apolar surface into bulk. In contrast, desolvation of polar dipeptides is accompanied by the formation of intersolute hydrogen bonds or salt bridges, which is enthalpically favorable [78]. Extrapolating from this understanding of small-molecule thermodynamics, it seems that increasing polar surface area burial in a protein-ligand complex could be used as a strategy to provide a more favorable enthalpy of binding. However, in the context of protein-ligand interactions, the thermodynamics of the burial of a polar surface between proteins and ligands is affected by the influence of the detailed topography of the binding site and the balance between unfavorable enthalpy change due to dehydration and favorable enthalpy change due to the formation of protein-ligand hydrogen bonds [79].

ITC can also be applied to improve drug delivery systems. Chitosan has been extensively used as a biomaterial because of its immunostimulatory, antibacterial and antifungal activities [80, 81]. However, chitosan suffers from low solubility at a physiological pH of 7.4, which limits its use as an absorption enhancer in a neutral environment, *e.g.*, for nasal peroral delivery systems. The solubility of chitosan was improved by using the derivatives of chitosan. The binding affinity of carboxymethylchitosan (OCMCS) with BSA was tested by ITC analysis [82]. ITC demonstrated that BSA bound to OCMCS with a molar ratio of 4.5:1. The binding isotherms appeared exothermic, which was ascribed to the contributions of H-bond and hydrophobic interactions.

### 3.3 ITC applications in cell metabolism

Isothermal microcalorimetry is currently used in the study of microorganisms, human and animal cellular systems, small animals, plants materials, and mammalian cells and tissues. Studies carried out in these areas have been published in Special Issues of *Thermochimica Acta* [83–85]. One important problem in studying cells using microcalorimetry is that the cultures may become anoxic due to limitations in the supply of oxygen [86, 87]. This problem has been solved using two

approaches. The first is to make the bioreactor the calorimeter (also termed macrocalorimetric technique) [88] and the second uses a combination of a bioreactor with a flow microcalorimeter [89]. This has a device to circulate the cells in suspension towards a microcalorimeter, which has a flow-through vessel (termed flow calorimetric technique). The principles and schematic representation of these different techniques have been extensively described elsewhere [90, 91].

The first macrocalorimetric technique, *i.e.*, dynamic calorimetry with a working volume of 10 L, was introduced in 1968 [92]. The major disadvantage arises from the brief interruption (5–9 min) of the cooling system, which resulted in relatively poor resolution especially for weakly exothermic or endothermic bioprocesses. The resolution of this system was further improved so that it had complete energy balance in a standard laboratory bioreactor with a working volume of 1.5 L without the need for disconnection of the cooling power [93]. Heat flow is calculated by mathematical models, to yield precise values.

Continuous calorimetry was developed using a 14-L laboratory bioreactor provided with a constant cooling power [94]. The reactor temperature was set using an electrical heater inside the reactor. Based on this principle, two commercially available reaction calorimeters were developed, the Bergh of Fermentation Calorimeter (BFK) [95, 96] and the Mettler-Toledo Reaction Calorimeter (RC1) [97]. The BFK type 1 has a 2-L working volume glass reactor and is double jacketed. The reactor temperature is monitored by a high resolution thermistor. A constant temperature difference is maintained by a silicone oil circulate. An advantage of the BFK-1 is that no corrections for heat losses through the reactor wall are required; however, this system could not provide sufficient cooling capacity for all applications, the thermopile wall is extremely fragile and the reactor design is impractical. For these reasons, a BFK type 2 was developed in which silicone oil is passed through the jacket of the housing to maintain the temperature at precisely that of the reactor contents, thus acting as an adiabatic shield.

Stockar's group [97] adapted a 2-L Mettler-Toledo RC1 heat accumulation calorimeter, capable of operating in isothermal, adiabatic or isoperibol modes. The detection limit for the reaction calorimeter was only  $\sim 2 \times 10^6$  cells/cm<sup>3</sup>, generally an order of magnitude higher than the starting inoculum for animal cultures, which in a standard batch culture generally rise to a maximum concentration of approximately  $1 \times 10^6$  cells/cm<sup>3</sup>. A series of improvements to the Mettler-Toledo RC1 reaction calorimeter in terms of resolution in sterile op-

eration, mixing and agitation, mass flows, and delivering air resulted in the detection limit sufficiently low to follow the course of batch cultures of many different types of animal cells [98, 99]. Two techniques for delivering air to the cell culture, *i.e.*, under surface aeration and purging by air, are investigated to study the consumption of glucose and the production of lactate [100]. The results indicated that the sparged culture was more aerobic and this resulted in less lactate and gave a lower  $Y_{\text{lac}/\text{glc}}$  yield of 0.96 than those under surface aeration ( $Y_{\text{lac}/\text{glc}} = 1.24$ ).

Garcia-Payo *et al.* [101] reported changes to the instrument in terms of limiting non-biological heat flows and altering the geometry of the glass reactors. The problems associated with this are direction of flow, slow maximum flow rate, narrow bore tubing and small volume of the measuring vessel. The Kemp research group developed a solution for measuring the heat flow rate of animal cell growing under the controlled conditions of the bioreactor [102]. It allows the measurement of the heat produced by circulating the cell suspension from the bioreactor to the flow calorimeter and then returning it to the bioreactor. The same research group also developed a new twin flow module calorimeter that is optimized for the use with cell suspensions as well as cells growing on microcarrier beads. The various fabricated instruments for flow calorimetry have been summarized in [91, 98]. Cultured animal cells, particularly those derived from either insect or mammalian tissues, are used increasingly in pharmaceutical industry. The heat produced by animal cells in culture can be used as the primary indicator of the kinetics of their metabolism [103]. Studies reported on Chinese hamster ovary cells (CHO320) genetically engineered to produce interferon- $\gamma$  (IFN- $\gamma$ ) showed an improved formulation that indicates faster cell growth and greater specific rate (flux) of IFN- $\gamma$  constitutive secretion, while decreasing glucose utilization and excretion of lactate that is toxic to cells.

Microcalorimetry was used successfully as a bioassay for the detection of the very small metabolic activation following binding of growth hormone to its cell surface receptor [104]. Activated metabolism and cell growth were observed (positive slope in heat flow) on the addition of trace amounts of added growth hormone (less than  $25 \times 10^{-12}$  mol/L). Information on the triggered cell division following the addition was also derived. Microcalorimetry can thus be considered a technique that can be used to test a vast number of therapeutically important substances on cells and tissues. Following the similar methodology, cell death was monitored on addition of anticancer



drugs [105]. The action of two anti-neoplastic drugs, methotrexate (MTX) and 6-thioguanine (6-TG), on a human T lymphoma cell line, CCRF-CEM, has been studied. The slope of the power-time curve after drug injection relative to that obtained during unperturbed growth was used to construct dose-response curves. Backman *et al.* [106] studied the growth and energy metabolism of cell with time by microcalorimetric study. The growth rate of cultured T lymphoma cells decreased with time. The changes could be attributed to the decrease in pH of the medium. Following these principles of studying cell death and cell growth by ITC analysis, this technique finds application in various fields, such as identification of some diseases, drug formulation and factors that affect the growth of a cell.

## 4 Conclusions

In this review we have presented recent trends and applications of ITC in bioseparation, drug development and studying the living organism. In addition, the methodology and analysis of the thermodynamic parameters derived have been described. In bioseparation, the mechanism of interaction has been elucidated from the thermodynamic parameters. The thermodynamic approaches serve as a framework for chromatographic separation. ITC application to drug development illustrate that binding affinities of the drug to the target can be identified from thermodynamic parameters. Improvements in terms of binding affinity and selectivity can be made using suitable lead compounds. The drug lead compounds are made by molecular alterations, and studying its affinity with the target by thermodynamic analysis by reaching an enthalpy and/or entropy to contribute favorable energy. ITC application in living organism can be used as the primary indicator of the kinetics of their metabolism and can be utilized in various fields especially in identification of some diseases and drug formulation.

*The authors would like to thank the National Science Council of the Republic of China, Taiwan and the Southern Science Base Industrial Park-NSC for financially supporting this research under Contract Numbers. NSC95-2221-E-008-172-MY3 and NSC97-2120-M-008-002*

*The authors have declared no conflict of interest.*

## 5 References

- [1] Sigurskjold, B. W., Bundle, D. R., Thermodynamics of oligosaccharide binding to a monoclonal antibody specific for a *Salmonella* O-antigen point to hydrophobic interactions in the binding site. *J. Biol. Chem.* 1992, 267, 8371–8376.
- [2] Naghibi, H., Tamura, A., Sturtevant, J. M., Significant discrepancies between van't Hoff and calorimetric enthalpies. *Proc. Natl. Acad. Sci. USA* 1995, 92, 5597–5599.
- [3] Finette, G. M., Mao, Q. M., Hearn, M. T., Comparative studies on the isothermal characteristics of proteins adsorbed under batch equilibrium conditions to ion-exchange, immobilised metal ion affinity and dye affinity matrices with different ionic strength and temperature conditions. *J. Chromatogr. A* 1997, 763, 71–90.
- [4] Weber, G., van't Hoff revised: Enthalpy of association of protein subunits. *J. Phys. Chem.* 1995, 99, 1052–1059.
- [5] Chaires, J. B., Possible origin of differences between van't Hoff and calorimetric enthalpy estimates. *Biophys. Chem.* 1997, 64, 15–23.
- [6] Horn, J. R., Russell, D., Lewis, E. A., Murphy, K. P., Van't Hoff and calorimetric enthalpies from isothermal titration calorimetry: Are there significant discrepancies? *Biochemistry* 2001, 40, 1774–1778.
- [7] Sottriffer, C. A., Flader, W., Cooper, A., Rode, B. M. *et al.*, Ligand binding by antibody IgE Lb4: Assessment of binding site preferences using microcalorimetry, docking, and free energy simulations. *Biophys. J.* 1999, 76, 2966–2977.
- [8] Guan, Y., Kemp, R. B., Medium design with the aid of heat flux measurement in mammalian cell culture, in: Westerhoff, H. V., Snoep, J. L., Sluse, F. E., Wijker, J. E., Kholodenko, B. N. (Eds.), *Biothermokinetics of the Living Cell*, BioThermoKinetics Press, Amsterdam 1996, pp. 387–397.
- [9] Bertini, I., Calderone, V., Fragai, M., Giachetti, A. *et al.*, Exploring the subtleties of drug-receptor interactions: The case of matrix metalloproteinases. *J. Am. Chem. Soc.* 2007, 129, 2466–2475.
- [10] Wolthers, K. R., Lou, X., Toogood, H. S., Leys, D., Scrutton, N. S., Mechanism of coenzyme binding to human methionine synthase reductase revealed through the crystal structure of the FNR-like module and isothermal titration calorimetry. *Biochemistry* 2007, 46, 11833–11844.
- [11] Andujar-Sanchez, M., Jara-Perez, V., Camara-Artigas, A., Thermodynamic determination of the binding constants of angiotensin-converting enzyme inhibitors by a displacement method. *FEBS Lett.* 2007, 581, 3449–3454.
- [12] Mudhivarthi, V. K., Bhambhani, A., Kumar, C. V., Novel enzyme/DNA/inorganic nanomaterials: A new generation of biocatalysts. *Dalton Trans.* 2007, 5483–5497.
- [13] Morgunova, E., Saller, S., Haase, I., Cushman, M. *et al.*, Lumazine synthase from *Candida albicans* as an anti-fungal target enzyme: Structural and biochemical basis for drug design. *J. Biol. Chem.* 2007, 282, 17231–17241.
- [14] Dormitzer, P. R., Ulmer, J. B., Rappuoli, R., Structure-based antigen design: A strategy for next generation vaccines. *Trends Biotechnol.* 2008, 26, 659–667.
- [15] Ren, J., Jenkins, T. C., Chaires, J. B., Energetics of DNA intercalation reactions. *Biochemistry* 2000, 39, 8439–8447.
- [16] Houtman, J. C., Brown, P. H., Bowden, B., Yamaguchi, H. *et al.*, Studying multisite binary and ternary protein interactions by global analysis of isothermal titration calorimetry data in SEDPHAT: Application to adaptor protein complexes in cell signaling. *Protein Sci.* 2007, 16, 30–42.



- [17] Meier-Andrejszki, L., Bjelic, S., Naud, J. F., Lavigne, P. *et al.*, Thermodynamics of b-HLH-LZ protein binding to DNA: The energetic importance of protein-DNA contacts in site-specific E-box recognition by the complete gene product of the Max p21 transcription factor. *Biochemistry* 2007, 46, 12427–12440.
- [18] Mosbahi, K., Walker, D., James, R., Moore, G. R., Kleanthous, C., Global structural rearrangement of the cell penetrating ribonuclease colicin E3 on interaction with phospholipid membranes. *Protein Sci.* 2006, 15, 620–627.
- [19] Fisher, R. J., Fivash, M. J., Stephen, A. G., Hagan, N. A. *et al.*, Complex interactions of HIV-1 nucleocapsid protein with oligonucleotides. *Nucleic Acids Res.* 2006, 34, 472–484.
- [20] Ciulli, A., Williams, G., Smith, A. G., Blundell, T. L., Abell, C., Probing hot spots at protein-ligand binding sites: A fragment-based approach using biophysical methods. *J. Med. Chem.* 2006, 49, 4992–5000.
- [21] Haq, I., Thermodynamics of drug-DNA interactions. *Arch. Biochem. Biophys.* 2002, 403, 1–15.
- [22] Chatterjee, A., Moulik, S. P., Majhi, P. R., Sanyal, S. K., Studies on surfactant-biopolymer interaction. I. Microcalorimetric investigation on the interaction of cetyltrimethylammonium bromide (CTAB) and sodium dodecylsulfate (SDS) with gelatin (Gn), lysozyme (Lz) and deoxyribonucleic acid (DNA). *Biophys. Chem.* 2002, 98, 313–327.
- [23] De, S., Robinson, D., Polymer relationships during preparation of chitosan-alginate and poly-L-lysine-alginate nanospheres. *J. Control Release* 2003, 89, 101–112.
- [24] Olsen, S., Applications of isothermal titration calorimetry to measure enzyme kinetics and activity in complex solutions. *Thermochim. Acta.* 2006, 448, 12–18.
- [25] Hansen, L. D., Christensen, J. J., Izatt, R. M., Entropy titration. A calorimetric method for the determination of  $\Delta G$ ,  $\Delta H$ , and  $\Delta S$ . *J. Chem. Soc. Chem. Commun.* 1965, 3, 36–37.
- [26] Eatough, D. J., Lewis, E. A., Hansen, L. D., Solution calorimetry: Determination of  $\Delta H$  and  $K_{eq}$  values, in: Grime, K. (Ed.), *Analytical Solution Calorimetry*, Wiley, New York 1985, pp. 137–161.
- [27] Wadsö, I., Isothermal microcalorimetry: Current problems and prospects. *J. Therm. Anal. Cal.* 2001, 64, 75–84.
- [28] Stanislaw, L. R., Recent developments in calorimetry. *Annu. Rep. Prog. Chem. Sect. C* 2002, 98, 157–217.
- [29] Ingemar, W., Trends in isothermal microcalorimetry. *Chem. Soc. Rev.* 1997, 26, 79–86.
- [30] Baranauskienė, L., Petrikaitė, V., Matulienė, J., Matulis, D., Titration calorimetry standards and the precision of isothermal titration calorimetry data. *Int. J. Mol. Sci.* 2009, 10, 2752–2762.
- [31] Tellinghuisen, J., Calibration in isothermal titration calorimetry: Heat and cell volume from heat of dilution of NaCl(aq). *Anal. Biochem.* 2007, 360, 47–55.
- [32] Hopkins, H. P. Jr., Fumero, J., Wilson, W. D., Temperature dependence of enthalpy changes for ethidium and propidium binding to DNA: Effect of alkylamine chains. *Biopolymers* 1990, 29, 449–459.
- [33] Indyk, L., Fisher, H. F., Theoretical aspects of isothermal titration calorimetry. *Methods Enzymol.* 1998, 295, 350–364.
- [34] Gurney, R. W., *Ionic Processes in Solution*. McGraw-Hill, New York 1953, p.89–95.
- [35] Kauzmann, W., Some factors in the interpretation of protein denaturation. *Adv. Protein Chem.* 1959, 14, 1–63.
- [36] Torigoe, H., Nakayama, T., Imazato, M., Shimada, I., Arata Y., Sarai, A., The affinity maturation of anti-4-hydroxy-3-nitrophenylacetyl mouse monoclonal antibody. *Biol. Chem.*, 1995, 270, 22218–22222.



**Wen-Yih Chen** was born in Kaohsiung, Taiwan. He finished his BSc degree in Chemical Engineering in National Central University, Taiwan. Then he pursued his Master and Ph.D. degree in Chemical Engineering in Rose-Hulman Institute of Technology and Oklahoma State University, USA, respectively. Before starting his academic career as Professor, he did his postdoctoral training in

the Life Science Division, Stanford Institute Research, International, USA. He is currently a Professor in the Department of Chemical and Materials Engineering, National Central University, Taiwan. He has spent almost 20 years building up thermodynamics approaches in biorecognition processes and one of the successful applications is in biochromatography. He studied the interaction mechanism by measuring interaction enthalpy between biomolecules by isothermal titration calorimetry (ITC). Other applications of ITC in his research group include protein conformation diseases, nonfouling biomaterials, nanostructure of biomaterials for stem cell purification and cultivation, *etc.*

- [37] Lin, F. Y., Chen, W. Y., Hearn, M. T., Thermodynamic analysis of the interaction between proteins and solid surfaces: Application to liquid chromatography. *J. Mol. Recognit.* 2002, 15, 55–93.
- [38] Huang, H. M., Lin, F. Y., Chen, W. Y., Ruaan, R. C., Isothermal titration microcalorimetric studies of the effect of temperature on hydrophobic interaction between proteins and hydrophobic adsorbents. *J. Colloid Interface Sci.* 2000, 229, 600–606.
- [39] Lin, F. Y., Chen, W. Y., Ruaan, R. C., Huang, H. M., Microcalorimetric studies of interactions between proteins and hydrophobic ligands in hydrophobic interaction chromatography: Effects of ligand chain length, density and the amount of bound protein. *J. Chromatogr. A* 2000, 872, 37–47.
- [40] Lin, F. Y., Chen, W. Y., Hearn, M. T., Microcalorimetric studies on the interaction mechanism between proteins and hydrophobic solid surfaces in hydrophobic interaction chromatography: Effects of salts, hydrophobicity of the sorbent, and structure of the protein. *Anal. Chem.* 2001, 73, 3875–3883.
- [41] Tsai, Y. S., Lin, F. Y., Chen, W. Y., Lin, C. C., Isothermal titration microcalorimetric studies of the effect of salt concentrations in the interaction between proteins and hydrophobic adsorbents. *Colloids Surf. A Physicochem. Eng. Aspects* 2002, 197, 111–118.
- [42] Chen, W. Y., Huang, H. M., Lin, C. C., Lin, F. Y., Chen, Y. C., Effect of temperature on hydrophobic interaction between proteins and hydrophobic adsorbents: Studies by isothermal titration calorimetry and the van't Hoff equation. *Langmuir* 2003, 19, 9395–9403.
- [43] Huang, H.-M., Chen, W.-Y., Ruaan, R.-C., Microcalorimetric studies of the interaction mechanism between designed peptides and hydrophobic adsorbents. *J. Colloid Interface Sci.* 2003, 26, 23–28.
- [44] Chen, W.-Y., Liu, Z.-C., Lin, P.-H., Fang, C.-I. *et al.*, The hydrophobic interactions of the ion-exchanger resin ligands with proteins at high salt concentrations by adsorption iso-

- therms and isothermal titration calorimetry. *Sep. Purif. Technol.* 2007, 54, 212–219.
- [45] Chen, W.-Y. Wu, C.-F., Liu, C.-C., Interactions of imidazole and proteins with immobilized Cu(II) ions: Effects of structure, salt concentrations and pH in affinity and binding capacity. *J. Colloid Interface Sci.* 1996, 180, 135–143.
- [46] Wu, C.-F., Chen, W.-Y., Lee, J. F., Microcalorimetric studies of the interactions of imidazole with immobilized Cu(II): Effects of pH value and salt concentration. *J. Colloid Interface Sci.* 1996, 183, 236–242.
- [47] Chen, W.-Y., Lee, J.-F., Wu, C.-F., Microcalorimetric studies of the interactions of lysozyme with immobilized Cu(II): Effects of pH value and salt concentration. *J. Colloid Interface Sci.* 1997, 190, 49–54.
- [48] Lin, F.-Y., Chen, W.-Y., Sang, L.-C. Microcalorimetric studies of the interactions of lysozyme with immobilized metal ions: Effects of ion, pH value and salt concentration. *J. Colloid Interface Sci.* 1999, 214, 373–379.
- [49] Lin, F. Y., Chen, W. Y., Chen, H.-M., Microcalorimetric study of the effect of hexa-histidine tag and denaturant on the interaction mechanism between protein and metal-chelating gel, *J. Colloid Interface Sci.* 2001, 238, 333–339.
- [50] Lin, F. Y., Chen, C. S., Chen, W. Y., Yamamoto, S., Microcalorimetric studies of the interaction mechanisms between proteins and Q-sepharose at pH near the isoelectric point (pI) effects of NaCl concentration, pH value, and temperature. *J. Chromatogr. A* 2001, 912, 281–289.
- [51] Ma, J. C., Dougherty, D. A., The Cation- $\pi$  Interaction. *Chem. Rev.* 1997, 97, 1303–1324.
- [52] Kyte, J., Doolittle, R., A simple method for displaying the hydrophobic character of a protein. *J. Mol. Biol.* 1982, 157, 105–132.
- [53] Hopp, T. P., Woods, K. R., Prediction of protein antigenic determinants from amino acid sequences. *Proc. Natl. Acad. Sci. USA* 1981, 78, 3824–3428.
- [54] Reubsaet, J. L. E., Vieskar, R. Characterisation of  $\pi$ - $\pi$  interactions which determine retention of aromatic compounds in reversed-phase liquid chromatography. *J. Chromatogr. A* 1999, 841, 147–154.
- [55] Haidacher, D., Vailaya, A., Horvath, C, Temperature effects in hydrophobic interaction chromatography. *Proc. Natl. Acad. Sci. USA*, 1996, 93, 2290–2295.
- [56] Melander, W. R. Corradini, D., Horvath, C., Salt-mediated retention of proteins in hydrophobic-interaction chromatography: Application of solvophobic theory. *J. Chromatogr. A* 1984, 317, 67–85.
- [57] Melander, W., Horvath, C., Salt effects on hydrophobic interaction in precipitation and chromatography of proteins: An interpretation of lyotropic series. *Arch. Biochem. Biophys.* 1977, 183, 200–215.
- [58] Hofmeister, F., Zur Lehre von der Wirkung der Salze – Zweite Mitteilung. *Arch. Exp. Pathol. Pharmacol.* 1888, 24, 247–260.
- [59] Szepeszy, L., Horvath, C., Specific salt effects in hydrophobic interaction chromatography of proteins. *Chromatographia* 1988, 26, 13–18.
- [60] Hallgren, E., Prediction of protein retention at gradient elution conditions in ion-exchange chromatography. *J. Chromatogr. A* 1999, 852, 351–359.
- [61] Yamamoto, S., Nomura, M., Sano, Y., Preparative chromatography of proteins: Design calculation procedure for gradient and stepwise elution. *J. Chromatogr. A* 1990, 512, 89–100.
- [62] Dieterle, M., Blaschke, T., Hasse, H., Microcalorimetric study of adsorption of human monoclonal antibodies on cation exchange chromatographic materials. *J. Chromatogr. A* 2008, 1205, 1–9.
- [63] Burnett, J. C., Ruthel, G., Stegmann, C. M., Panchal, R. G. *et al.*, Inhibition of metalloprotease botulinum serotype A from a pseudo-peptide binding mode to a small molecule that is active in primary neurons. *J. Biol. Chem.* 2007, 282, 5004–5014.
- [64] Sergio, C., Gianni, P., Mollica, V., Lepori, L., Group contributions to the thermodynamic properties of non-ionic organic solutes in dilute aqueous solution. *J. Solution Chem.* 1981, 10, 563–595.
- [65] Ruben, A. J. Kiso, Y., Freire, E., Overcoming roadblocks in lead optimization: A thermodynamic perspective. *Chem. Biol. Drug Des.* 2006, 67, 2–4.
- [66] Sarver, R. W., Peevers, J., Cody, W. L., Ciske, F. L. *et al.*, Binding thermodynamics of substituted diaminopyrimidine renin inhibitors. *Anal. Biochem.* 2007, 360, 30–40.
- [67] Velazquez-Campoy, A., Todd, M. J., Freire, E., HIV-1 protease inhibitors: Enthalpic versus entropic optimization of the binding affinity. *Biochemistry* 2000, 39, 2201–2207.
- [68] Todd, M. J., Luque, I., Velazquez-Campoy, A., Freire, E., Thermodynamic basis of resistance to HIV-1 protease inhibition: Calorimetric analysis of the V82F/I84V active site resistant mutant. *Biochemistry* 2000, 39, 11876–11883.
- [69] Freire, E., Isothermal titration calorimetry: Controlling binding forces in lead optimization. *Drug Discov. Today Technol.* 2004, 1, 295–296.
- [70] Protasevich, I., Brouillette, C. G., Snow, M. E., Dunham, S. *et al.*, Role of inhibitor aliphatic chain in the thermodynamics of inhibitor binding to *Escherichia coli* enoyl-ACP reductase and the Phe203Leu mutant: A proposed mechanism for drug resistance. *Biochemistry* 2004, 43, 13380–13389.
- [71] Lipinski, C. A., Lombardo, F., Dominy, B. W., Feeney, P. J., Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. *Adv. Drug Deliv. Rev.* 2001, 46, 3–26.
- [72] Li, L. D., Justin, J., Nowacki, J., O'Callaghan, B. J. *et al.*, A comprehensive dataset for receptor-ligand interactions with three-dimensional structures and binding thermodynamics from isothermal titration calorimetry. *Chem. Biol. Drug Des.* 2008, 71, 529–532.
- [73] Freire, E., Do enthalpy and entropy distinguish first in class from best in class? *Drug Discov. Today* 2008, 13, 869–874.
- [74] Ohtaka, H., Freire, E., Adaptive inhibitors of the HIV-1 protease. *Prog. Biophys. Mol. Biol.* 2005, 88, 193–208.
- [75] Ward, W. H., Holdgate, G. A., Isothermal titration calorimetry in drug discovery. *Prog. Med. Chem.* 2001, 38, 309–376.
- [76] Holdgate, G., The application of isothermal titration calorimetry to drug discovery, in Ladbury, J. E. (Ed.) *Bicalorimetry 2: Applications of calorimetry in the biological sciences*, Wiley, New York 2004, pp. 59–79.
- [77] Velazquez-Campoy, A., Muzammil, S., Ohtaka, H., Schon, A. *et al.*, Structural and thermodynamic basis of resistance to HIV-1 protease inhibition: Implications for inhibitor design. *Curr. Drug Targets Infect. Disord.* 2003, 3, 311–328.
- [78] Habermann, S. M., Murphy, K. P., Energetics of hydrogen bonding in proteins: A model compound study. *Protein Sci.* 1996, 5, 1229–1239.
- [79] Connelly, P. R., Aldape, R. A., Bruzzese, F. J., Chambers, S. P. *et al.*, Enthalpy of hydrogen bond formation in a protein-ligand binding reaction. *Proc. Natl. Acad. Sci. USA* 1994, 91, 1964–1968.
- [80] Jin, J., Song, M., Hourston, D. J., Novel chitosan-based films cross-linked by genipin with improved physical properties. *Biomacromolecules* 2004, 5, 162–168.

- [81] Iwasaki, N., Yamane, S. T., Majima, T., Kasahara, Y. *et al.*, Feasibility of polysaccharide hybrid materials for scaffolds in cartilage tissue engineering: Evaluation of chondrocyte adhesion to polyion complex fibers prepared from alginate and chitosan. *Biomacromolecules* 2004, 5, 828–833.
- [82] Wu, H., Zhu, A., Yuan, L., Interactions between O-carboxymethylchitosan and bovine serum albumin. *Mater. Chem. Phys.* 2008, 112, 41–46.
- [83] Kemp, R. B. (Ed.), *Special Issue Thermochim. Acta* 1990, 172.
- [84] Lamprecht, I., Hemminger, W., Hohne, G. W. H., Calorimetry in the biological sciences. *Special Issue Thermochim. Acta* 1991, 193, 452.
- [85] Kemp, R. B., Schaarschmidt, B., *Special Issue Thermochim. Acta* 1995, 250.
- [86] Kemp, R. B., Guan, Y., Heat flux and the calorimetric-respirometric ratio as measures of catabolic flux in mammalian cells. *Thermochim. Acta* 1997, 300, 199–222.
- [87] Backman, P., Bastos, M., Hallen, D., Lonnbro, P., Wadso, I., Heat conduction calorimeters: Time constants, sensitivity and fast titration experiments. *J. Biochem. Biophys. Methods* 1994, 28, 85–100.
- [88] Garcia-Payoa, M. C., Ampuero, S., Liub, J. S., Marison, I. W., von Stockar, U., The development and characterization of a high resolution bio-reaction calorimeter for weakly exothermic cultures. *Thermochim. Acta* 2002, 391, 29–39.
- [89] Kemp, R. B., Evans, P. M., Guan, Y., An enthalpy balance approach to studies of metabolic activity in mammalian cells. *J. Therm. Anal. Cal.* 1997, 49, 755–770.
- [90] Manson, I., Liu, J.-S., Ampuero, S., Von Stockar, U., Schenker, B., Biological reaction calorimetry: Development of high sensitivity bio-calorimeters. *Thermochim. Acta* 1998, 309, 157–173.
- [91] Kemp, R. B., Thermobiochemical Studies of Animal Cell systems in vitro: Evidence of their nature from Bioreactor Studies, in: Lorinczy, D. (Ed.): *The Nature of Biological Systems as Revealed by Thermal Methods*, Kluwer Academic, Dordrecht 2004, pp. 217–251.
- [92] Cooney, C. L., Wang, D. I. C., Mateles, R. I., Measurement of heat evolution and correlation with oxygen consumption during microbial growth. *Biotechnol. Bioeng.* 1969, 11, 269.
- [93] van Kleeff, B. H. A., Kuenen, J. G., Heijnen, J. J., Continuous measurement of microbial heat production in laboratory fermentors. *Biotechnol. Bioeng.* 1993, 41, 541–549.
- [94] Luong, J. H. T., Volesky, B., Determination of the heat of some aerobic fermentations. *Can. J. Chem. Eng.* 1980, 58, 497–504.
- [95] Meier-Schneiders, M., Weikmann, W., Grosshans, U., Busch, C., Steinbüchel, A., On-line-analysis of *Alcaligenes eutrophus* fermentations. *Can. J. Microbiol.* 1995, 41, 267.
- [96] Meier-Schneiders, M., Grosshans, U., Busch, C., Eigenberger, G., Biocalorimetry-supported analysis of fermentation processes. *J. Appl. Microbiol. Biotechnol.* 1995, 43, 431–439.
- [97] Cooney, M. J., Marison, I. W., van Gulik, W. M., von Stockar, U., Calorimetric and stoichiometric analysis of growth of *K. fragilis* in continuous culture: Nitrogen limitation imposed upon carbon limited growth. *J. Appl. Microbiol. Biotechnol.* 1996, 44, 643–653.
- [98] Duboc, P. M., I., von Stockar, U., Quantitative calorimetry and biochemical engineering, in: Kemp, R. B. (Ed.), *Handbook of Thermal Analysis and Calorimetry*, Vol. 4, *From Macromolecules to Man*, Elsevier, Amsterdam 1999, pp. 267–365.
- [99] Marison, I. L., J.-S. Ampuero, S. von Stockar, U., Schenker, B., Biological reaction calorimetry: Development of high sensitivity biocalorimeters. *Thermochim. Acta* 1998, 309, 147–137.
- [100] Gnaiger, E., Kemp, R. B., Anaerobic metabolism in aerobic mammalian-cells – Information from the ratio of calorimetric heat-flux and respirometric oxygen flux. *Biochim Biophys Acta* 1990, 1016, 328–332.
- [101] Garcia-Payoa, M. C., Ampuero, S., Liu, J. S., Marison, I. W., von Stockar, U., The development and characterization of a high resolution bioreaction calorimeter for weakly exothermic cultures. *Thermochim. Acta* 2002, 391, 25–39.
- [102] Guan, Y., Evans, P. M., Kemp, R. B., Specific heat flow rate: An on-line monitor and potential control variable of specific metabolic rate in animal cell culture that combines microcalorimetry with dielectric spectroscopy. *Biotechnol. Bioeng.* 1998, 58, 464–477.
- [103] Guan, Y. H., Lloyd, P. C., Kemp, R. B., A calorimetric flow vessel optimised for measuring the metabolic activity of animal cells. *Thermochim. Acta* 1999, 332, 211–220.
- [104] Schön, A., Walum, E., 37<sup>th</sup> Annual Meeting of the American Society for Cell Biology, Washington D.C. 1997.
- [105] Bermudez, J., Backman, P., Schon, A., Microcalorimetric evaluation of the effects of methotrexate and 6-thioguanine on sensitive T-lymphoma cells and on a methotrexate-resistant subline. *Cell Biophys.* 1992, 20, 111–123.
- [106] Backman, P., Kimura, T., Schon, A., Wadso, I., Effects of pH-variations on the kinetics of growth and energy metabolism in cultured T-lymphoma cells: A microcalorimetric study. *J. Cell. Physiol.* 1992, 150, 99–103.