

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/5233368>

Kinetic of leucrose formation from sucrose by dextransucrase

ARTICLE *in* BIOTECHNOLOGY AND BIOENGINEERING · AUGUST 1994

Impact Factor: 4.13 · DOI: 10.1002/bit.260440319 · Source: PubMed

CITATIONS

4

READS

19

3 AUTHORS, INCLUDING:



[Hans-Joachim Jöerdening](#)

Technische Universität Braunschweig

32 PUBLICATIONS 365 CITATIONS

SEE PROFILE

Errata

Kinetics of Leucrose Formation from Sucrose by Dextranucrase

Mathias Böker, Hans-Joachim Jördening, and Klaus Buchholz*

Sugar Institute, Institut für Technologie der Kohlenhydrate at the Technical University of Braunschweig, Langer Kamp 5, D-38106 Braunschweig, Germany

[Article in *Biotechnol. Bioenging.* 43(9) 856–864 (1994)]




Figure 6. Effect of fructose concentration on V_{Leu} at the following sucrose concentrations: (□) 584 mmol/L (immobilized dextranucrase), (▽) 183 mmol/L (immobilized DS), (○) 61.4 mmol/L (immobilized DS), and (▼) 183 mmol/L (free DS).




Figure 7. Effect of sucrose and fructose concentration on leucrose yield. Sucrose concentrations are (□) 584 mmol/L (immobilized DS), (▽) 183 mmol/L (immobilized DS), (▼) 183 mmol/L (free DS), and (○) 61.4 mmol/L (immobilized DS).

* To whom all correspondence should be addressed.




Figure 8. Reaction mechanism of DS with acceptor site.




Figure 9. Effect of temperature on the reaction rate and the distribution of product and byproducts. $F = 2.22 \text{ mol/L}$ and $s = 0.58 \text{ mol/L}$. Temperature = 5°, 15°, 25° C, (○) sucrose, (●) leucrose, (Δ) isomaltulose, and (▼) trehalulose.




Figure 10. (A) V_{leu} (○) and V_{ov} (Δ) as function of temperature. (B) Yield of leucrose (○), isomaltulose (Δ), and trehalulose (▲) as function of temperature.

