

Natural Product Synthesis

Sceptrin as a Potential Biosynthetic Precursor to Complex Pyrrole–Imidazole Alkaloids: The Total Synthesis of Ageliferin**

Phil S. Baran, Daniel P. O'Malley, and
Alexandros L. Zografos*


The isolation and characterization in 1981 of the first dimeric pyrrole-imidazole alkaloid, sceptrin (**1**, Scheme 1) by Faulkner, Clardy, and co-workers^[1] was a milestone event in marine

[*] Prof. Dr. P. S. Baran, D. P. O'Malley, Dr. A. L. Zografos
Department of Chemistry
The Scripps Research Institute
10550 North Torrey Pines Road, La Jolla, CA 92037 (USA)
Fax: (+1) 858-784-7375
E-mail: pbaran@scripps.edu

[**] We thank Dr. D. H. Huang and Dr. L. Pasternak for NMR spectroscopic assistance, Dr. G. Siuzdak for mass spectrometric assistance, and Dr. A. Saghatelian of the Cravatt group for assistance with HPLC separations. We are grateful to Dr. M. Assmann and Dr. T. Proksch for samples of natural ageliferin, Biotage (formerly Personal Chemistry) for a generous donation of process vials used in these studies, as well as the Hertz Foundation and the Department of Defense for predoctoral Fellowships (D.P.O.). Financial support for this work was provided by The Scripps Research Institute and Eli Lilly & Co.



Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.



Scheme 1. Structures of ageliferin (**2**), sceptrin (**1**), and debromooroidin (**3**) as well as retrosynthetic analysis of **1** and **2**.

natural product research as these compounds have extraordinary biological activity and stunning molecular architectures.^[2] Indeed, they have inspired a flurry of research in chemistry.^[2] Recently, the long-standing synthetic challenge posed by sceptrin (**1**) was solved with a concise sequence that proceeds in approximately 24% overall yield, can be conducted on a preparative scale, and does not necessitate chromatography.^[3]

Ageliferin (**2**),^[4] isolated in 1986 from *Agelas conifera* by Rinehart, is an antiviral agent^[4b] and may be a useful chemical tool for mechanistic studies of actin–myosin contractile systems.^[4c] It has been the subject of numerous synthetic efforts,^[5] all of which are based upon a widely accepted biosynthetic hypothesis^[6] wherein **2** is derived from two molecules of hymenidin (debromooroidin, **3**) by an enzymatic “Diels–Alderase” (Scheme 1). We were compelled to question this proposal upon noticing that in every instance in which **2** was isolated, **1** was by far the major constituent (see Figure 1 for an example).^[7] We reasoned that if **1** and **2** were derived from **3** by a divergent pathway (as is proposed^[6]), then the observed ratio of **1** and **2** after isolation should be reversed, solely on thermodynamic grounds. Thus, to explain this apparent discrepancy, we





Figure 1. HPLC chromatogram from the extracts of *Agelas conifera*. Reproduced from Ref. [7].

envisioned an alternative scenario wherein **1** rearranges to form **2**. Although such a rearrangement should not proceed thermally in a concerted fashion (see below), it would constitute an “allowed” event if the reaction proceeded in a stepwise fashion (radical or ionic processes) or through photochemical means.^[8,9]

Herein, we report the remarkable thermal conversion of sceptrin (**1**) into ageliferin (**2**). We also present an alternative biogenetic hypothesis commencing from **1** rather than **3** for other complex dimeric pyrrole-imidazole alkaloids, including the axinellamines^[10] and palau’amines.^[11]

Our explorations began with several unsuccessful attempts to effect photochemical [1,3] sigmatropic rearrangement of **1**. Thus, exposure of **1** to a 450-W Hanovia lamp (quartz or pyrex filter) for several days led only to gradual decomposition. Similarly, decomposition was observed when sceptrin (**1**) was heated in methanol at temperatures as high as 150 °C in a microwave. We also found the free base of **1** to be extremely unstable and to decompose into a variety of products which have not been fully characterized. However, when **1** was dissolved in water and heated to 195 °C for 1 minute by using microwave irradiation (Scheme 2), we obtained ageliferin ((±)-**2**) in 40% yield and identical in all respects to a natural sample, along with recovered (±)-**1** (52%). An NMR spectroscopic comparison of synthetic **1**, synthetic **2**, natural **2**, and the products of this reaction is



Scheme 2. Remarkable conversion of sceptrin (**1**) into ageliferin (**2**) and mechanistic analysis.

shown in Figure 2. Given the thermal instability of sceptrin^[3] and ageliferin, it is quite surprising that this reaction proceeds cleanly and reproducibly. In fact, decomposition begins after only 90 seconds at 195°C. If the reaction is performed at 195°C without a microwave, only sceptrin and decomposition products are observed.




Figure 2. Comparison of the ^1H NMR spectra (600 MHz, D_2O) of a) synthetic sceptrin, b) sceptrin after 1 minute at 195 °C, c) synthetic ageliferin (trifluoroacetic acid salt), and d) natural ageliferin (HCl salt).


This conversion constitutes, to the best of our knowledge, the first vinyl cyclobutane rearrangement of a natural product. However, perhaps more importantly, it also raises the intriguing question as to whether nature employs an enzyme to catalyze a similar process. It is equally possible that we have unearthed a completely abiotic route to **2**; thermolysis of **1** may just lead to an intermediate that can either return to **1** or irreversibly transform into **2**.

Three selected mechanistic portraits are depicted in Scheme 2. In the first scenario (path A), the cyclobutane is ruptured in a homolytic fashion and this is followed by a 6-*endo-trig* closure. An ionic process, reminiscent of the currently proposed biosynthesis of **2** from **3**,^[6] may also be invoked, as shown in path B. A series of concerted bond shifts, as depicted in path C, can also account for this structural reorganization. Thus, tautomerization to intermediate **4** followed by a thermal [1,2] shift (4→5-membered ring), a hydride shift, another [1,2] shift (5→6-

membered ring), and a final tautomerization could lead to **2**. All three pathways could conceivably lead to the stereochemistry of **2** since the bromopyrrole-bearing carbon centers are incapable of epimerization and could therefore direct ring closure in paths A and B. It follows then that the conversion **1**→**2** should be the same whether racemic or optically pure **1** is employed. Notably, the “allowed” thermal concerted [1,3] sigmatropic pathway would require inversion (suprafacial) of configuration at the migrating carbon atom, while we observe retention.^[8]


Although theoretical inquiries into the sigmatropic rearrangement of simple monocyclic vinyl cyclobutanes to cyclohexenes point to a radical mechanism,^[9] we believe an ionic process (perhaps similar to paths B or C) is operative here. The ambivalent reactivity of the 2-aminoimidazole C4–C5 double bond contrasts sharply with that of previously studied olefins in this rearrangement.^[12] For example, simply heating **1** at 80 °C in CD_3OD for 5 minutes with microwave irradiation led exclusively to $[\text{D}_2]$ sceptrin ($[\text{D}_2]$ -**1**) in quantitative yield (Scheme 3). Incidentally, this appears to be the first report of deuterium exchange at a carbon center on a 2-aminoimidazole under neutral conditions. It is postulated that the tautomeric form of sceptrin (**1**), structure **4**, is involved in this proton-addition/proton-loss sequence. In principle, a [1,2] rearrangement could then occur (see dotted arrows, Scheme 3), thus leading to the events shown in path C (Scheme 2) with eventual arrival at ageliferin (**2**). There is ample precedent for this type of ring expansion of cyclobutanes.^[13] Compelling evidence for the requirement of the 2-aminoimidazole subunit in this rearrangement was also garnered by submitting the known cyclobutane **5**^[14] to extended microwave irradiation (200 °C, 5 min) only to obtain the diacid **6** in quantitative yield (Scheme 4).

Based on these findings, it would not be surprising if other complex dimeric pyrrole-imidazole alkaloids such as the



Scheme 3. Facile synthesis of $[\text{D}_2]$ sceptrin ($[\text{D}_2]$ -**1**) points to the potential viability of path C (see Scheme 2).

palau'amines and axinellamines arise from sceptrin-type intermediates. Indeed, palau'amine (**9**), like ageliferin (**2**), was isolated along with sceptrin (**1**).^[11] For the purpose of inspiring new synthetic approaches to these alkaloids, we propose an alternative biogenic hypothesis for the generation



Scheme 4. Even after 5 minutes at 200 °C, only hydrolysis (no rearrangement) is observed with the known cyclobutane **5**.


of these compounds: We have chosen to depict the proposed transformation of the known natural product dibromosceptrin (**7**) into axinellamine A (**8**) as an example (Scheme 5); a ring-expansion pathway from a sceptrin-type intermediate could also be drawn for palau'amine (**9**).^[15] A ring-contraction pathway from an ageliferin-type structure could also lead to these alkaloids.^[16]

In summary, we have completed the first total synthesis of ageliferin (**2**, Scheme 1), identified the first vinyl cyclobutane rearrangement of a natural product, and advanced an alternative biosynthetic hypothesis for the formation of

complex pyrrole-imidazole alkaloids. The synthesis adds merely one additional step to the concise, practical, and analogue-friendly sceptrin synthesis developed in our laboratory,^[3] and it can thus easily be scaled up to allow for a full evaluation of the biological potential and structure–activity relationships of these intriguing alkaloids. Efforts are underway to harness the innate symmetry and reactivity of **1** and related structures to effect conversion into other naturally occurring pyrrole-imidazole alkaloids.

Received: February 4, 2004 [Z53937]

Keywords: alkaloids · biosynthesis · microwave reactions · natural products · total synthesis



Scheme 5. Proposed conversion of dibromosceptrin (**7**) into axinellamine A (**8**) in nature and the related structure of palau'amine (**9**).

- [1] R. P. Walker, D. J. Faulkner, D. Van Engen, J. Clardy, *J. Am. Chem. Soc.* **1981**, *103*, 6772–6773.
- [2] For an extensive review of the chemistry and biology of pyrrole-imidazole alkaloids, see: H. Hoffmann, T. Lindel, *Synthesis* **2003**, 1753–1783.
- [3] P. S. Baran, A. L. Zografos, D. P. O'Malley, *J. Am. Chem. Soc.* **2004**, *126*, 3726–3727.
- [4] a) P. A. Keifer, M. E. S. Koker, R. E. Schwartz, R. G. Hughes, Jr., D. Rittschof, K. L. Rinehart, *26th Interscience Conference on Antimicrobial Agents and Chemotherapy*, New Orleans, Sep. 28–Oct. 1, **1986**, No. 1281; b) K. L. Rinehart, *Pure Appl. Chem.* **1989**, *61*, 525–528; c) J. Kobayashi, M. Tsuda, T. Murayama, H. Nakamura, Y. Ohizumi, M. Ishibashi, M. Iwamura, T. Ohta, S. Nozoe, *Tetrahedron* **1990**, *46*, 5579–5586; d) P. A. Keifer, R. E. Schwartz, M. E. S. Koker, R. G. Hughes, D. Rittschof, K. L. Rinehart, *J. Org. Chem.* **1991**, *56*, 2965–2975; e) D. H. Williams, D. J. Faulkner, *Tetrahedron* **1996**, *52*, 5381–5390.
- [5] See Ref. [2] and a) Y. He, Y. Chen, H. Wu, C. J. Lovely, *Org. Lett.* **2003**, *5*, 3623–3626; b) I. Kawasaki, N. Sakaguchi, N. Fukushima, N. Fujioka, F. Nikaido, M. Yamashita, S. Ohta, *Tetrahedron Lett.* **2002**, *43*, 4377–4380.
- [6] All isolation reports such as those in Ref. [4] put forth this proposal. For a further elaboration of this hypothesis, see A. Al Mourabit, P. Potier, *Eur. J. Org. Chem.* **2001**, 237–243.
- [7] a) M. Assmann, M. Köck, *Z. Naturforsch. C* **2002**, *57*, 157–160; b) M. Assmann, E. Lichte, M. Köck, *Proceedings of the 6th International Sponge Symposium*, **2004**, in press; c) confirmed with Dr. Michael Assmann, personal communication.
- [8] I. Fleming, *Pericyclic Reactions*, Oxford Press, Oxford, **1999**, p. 75.
- [9] P. A. Leber, J. E. Baldwin, *Acc. Chem. Res.* **2002**, *35*, 279–287.
- [10] S. Urban, P. Leone, A. R. Carroll, G. A. Fechner, J. Smith, J. N. A. Hooper, R. J. Quinn, *J. Org. Chem.* **1999**, *64*, 731–735.
- [11] R. B. Kinnel, H.-P. Gehrken, R. Swali, G. Skoropowski, P. J. Scheuer, *J. Org. Chem.* **1998**, *63*, 3281–3286.
- [12] For studies that shed light on the chemistry of 2-aminoimidazoles, see Ref. [2] and Y.-Z. Xu, K. Yakushijin, D. A. Horne, *Tetrahedron Lett.* **1992**, *33*, 4385–4388; A. Olofson, K. Yakushijin, D. A. Horne, *J. Org. Chem.* **1998**, *63*, 1248–1253; Z.-K. Wan, G. H. C. Woo, J. K. Snyder, *Tetrahedron* **2001**, *57*, 5497–5507.
- [13] J. C. Namyslo, D. E. Kaufmann, *Chem. Rev.* **2003**, *103*, 1485–1537.
- [14] M. D'Auria, R. Racioppi, *Photochem. Photobiol.* **1998**, *112*, 145–148.
- [15] A natural product with the palau'amine skeleton and both pyrrole subunits intact is known (konbu'acidin A): J. Kobayashi, M. Suzuki, M. Tsuda, *Tetrahedron* **1997**, *53*, 15681–15684.
- [16] See Ref. [11] and N. Poje, M. Poje, *Org. Lett.* **2003**, *5*, 4265–4268.