

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/229015226>

Total Synthesis of (+/-)-Parvineostemonine

ARTICLE in CHEMISTRY - AN ASIAN JOURNAL · OCTOBER 2012

Impact Factor: 4.59 · DOI: 10.1002/asia.201200493 · Source: PubMed

CITATION

1

READS

44

4 AUTHORS, INCLUDING:



Tian Jinmiao

Lanzhou University

5 PUBLICATIONS 74 CITATIONS

SEE PROFILE



Yong Qiang Tu

Lanzhou University

151 PUBLICATIONS 3,015 CITATIONS

SEE PROFILE

Total Synthesis of (±)-Parvineostemonine

Zhi-Hua Chen, Jin-Miao Tian, Zhi-Min Chen, and Yong-Qiang Tu^{*[a]}

Roots and rhizomes of the stemonaceous plant have long been used for the treatment of respiratory diseases and as domestic insecticides in traditional folk medicine of East Asia.^[1] So far, over 130 structurally diverse *Stemona* alkaloids have been isolated from the monocotyledonous family Stemonaceae.^[2] These *Stemona* alkaloids have fascinating architectures and potential biological activities, which have provoked broad interests in their total synthesis.^[3–5] Among the *Stemona* alkaloids, parvineostemonine (**1a**), isolated from *Stemona parviflora* by Ye and co-workers in 2003,^[6] is a unique molecule which differs from the typical *Stemona* alkaloids such as **1b–1e** (Figure 1). Parvineostemonine fea-

cycloaddition.^[7] However, the total synthesis of parvineostemonine has not been achieved to date. Herein, we report an approach for the first total synthesis of (±)-parvineostemonine.

From a structural point of view, we found that **1a** has the same aza-bicyclo[3.2.1]octane unit (ring B and C) as tropinone. Thus, our synthesis of **1a** would focus on the formation of ring A and D from this commercially available material. A retrosynthetic analysis of **1a** is shown in Scheme 1.





Figure 1. Parvineostemonine (**1a**) and some selected examples of typical *Stemona* alkaloids.

tures a spiro-tetracyclic structure with a two-carbon bridge between C-3 and C-9 of the common pyrrolo[1,2-*a*]azepine (ring A and B) of the *Stemona* alkaloids. The piperidine moiety (ring C) contains five contiguous stereogenic carbons, wherein the oxa-quaternary carbon (C-11) is part of a spirobutenolide ring (ring D). Recently, Hsung and co-workers described a skeletal synthetic study toward parvineostemonine that features a highly stereoselective [4+3]




Scheme 1. Retrosynthetic analysis of (±)-parvineostemonine.

We envisioned that **1a** could be obtained by isomerization of an exocyclic double bond and reduction of the lactam carbonyl of **2**, the lactone ring D of which could be conveniently constructed by a stereoselective carbonyl allylation/lactonization of tricyclic compound **3**. We further expected that ring A in compound **3** might be formed by a key radical-mediated 7-*exo*-trig cyclization from halide precursor **4**, which could be derived from **5** through several transformations, including oxidation and cross-coupling reactions. We anticipated that amide **5** could be easily obtained in large-scale through a one-pot demethylation and acylation manipulations of inexpensive tropinone **6**.

According to the above-described plan, we commenced our synthesis from the preparation of the known ammonium salt **7**,^[8] as depicted in Scheme 2. Demethylation of tropinone **6** with 1-chloroethyl chloroformate afforded the crude ammonium salt **7**, which was directly treated with 4-chlorobutyl chloride to afford amide **5** in 96% yield. Initial attempts to obtain the α -diketone **9** directly from **5** with SeO₂ in refluxing 1,4-dioxane^[9] led to the decomposition of starting material. Thus, an alternative approach involving hydroxylation of **5** and subsequent oxidation was then employed. Treatment of **5** with KOH/PhI(OAc)₂^[10] followed by


[a] Z.-H. Chen, J.-M. Tian, Z.-M. Chen, Prof. Dr. Y.-Q. Tu
State Key Laboratory of Applied Organic Chemistry and Department of Chemistry
Lanzhou University
Lanzhou 730000 (P. R. China)
Fax: (+86) 931-8912582
E-mail: tuyq@lzu.edu.cn

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/asia.201200493>.



a one-pot hydrolyzation with hydrochloric acid provided α -hydroxy ketone **8** as a single diastereoisomer. As the hydroxy moiety in **8** would be oxidized to form a ketone, we did not determine the relative configuration of the newly formed stereocenter at the C-11 position. Swern oxidation^[11] of **8** did not afford α -diketone **9** and instead afforded a complex mixture of products. The successful oxidation of **8** was carried out by using Dess–Martin periodinane (DMP),^[12] thus furnishing α -diketone **9**, which was labile on silica-gel column chromatography. Therefore, the crude product **9** was immediately treated with triflic anhydride in CH_2Cl_2 at -78°C to afford enone **10** in 59 % yield over two steps. The introduction of an ethyl moiety was accomplished by using a Suzuki cross-coupling reaction^[13] of **10**, which was treated with Et_3B in the presence of $[\text{PdCl}_2(\text{dppf})]$ (dppf = diphenylphosphinoferrocene) to give radical cyclization precursor **11** in 82 % yield. Given that the corresponding iodo precursor would probably have a higher reactivity in the radical reaction, we also prepared iodide **12** from **11** by a Finkelstein reaction with NaI in refluxing acetone.^[14]

Next, the key radical-mediated 7-*exo*-trig cyclization to access tricyclic skeleton **3** was investigated (Scheme 3).



Scheme 3. Key radical-mediated 7-*exo*-trig cyclization.

Treatment of **11** or **12** with $\text{SmI}_2^{[15]}$ in tetrahydrofuran (THF) at 0°C to 66°C did not produce the cyclization product. By examining the general azobisisobutyronitrile (AIBN)/ $n\text{Bu}_3\text{SnH}$ reaction conditions,^[16] we found that the use of chlorinate **11** did not result in radical cyclization. However, iodide **12** could be converted into the desired tricyclic skeleton **3** along with its epimer **13** and the reduced product **14** under the same reaction conditions. Further studies showed that both the initiator and reaction tempera-







Figure 2. X-ray structure of the tricyclic compound **13**. Ellipsoids set at 30 % probability.^[18]

C-10 position was not in agreement with that of parvinecostemonine (**1a**). To convert epimer **13** into the requisite compound **3**, we investigated the adjustment of the stereochemistry at the C-10 position in **13** by an enolization/protonation sequence^[19] from the β face. A variety of bases such as lithium hexamethyldisilazane (LHMDS), NaHMDS, KHMDs, lithium diisopropylamide (LDA), and *t*BuOK and proton sources such as H₂O, phenol, and 2,6-di-*tert*-butyl-4-methylphenol were employed. However, in all cases a mixture of **3** and **13** were obtained in an approximate ratio of 1:3 (Scheme 4). Ultimately, tricycle **3** was obtained in 40% yield after subjecting **13** to epimerization for two times.

Scheme 4. Epimerization of **13**.

Having successfully established the cagelike skeleton **3**, we turned our attention to the stereoselective construction of lactone ring D. As shown in Scheme 5, the key carbonyl allylation/lactonization^[20] was achieved by treatment of **3** with ethyl (2-bromomethyl)acrylate^[21] and zinc powder in refluxing THF, and this provided **2** as a single diastereoisomer in 78% yield. On the basis of molecular model studies, we assumed that the zinc nucleophile formed in situ would attack the carbonyl carbon (C-11) in **3** from the less sterical-



Scheme 5. Synthesis of (±)-parvineostemonine.

ly hindered α face, which might account for the observed high diastereoselectivity. Then, RhCl_3 -mediated isomerization of the exocyclic double bond^[22] of lactone **2** afforded **15** in high yield. With **15** in hand, we attempted to selectively reduce the lactam carbonyl without affecting the unsaturated lactone moiety in **15**. Unfortunately, direct reduction of the lactam using our previously reported one-pot protocol ($\text{MeOTf}/\text{CH}_2\text{Cl}_2$; $\text{NaBH}_3\text{CN}/\text{EtOH}$)^[5c,h] resulted in recovery of **15**. Therefore, an alternative method was performed that involved formation of thioamide **16**, which was then subsequently reduced. The reaction of **15** with $\text{P}_4\text{S}_{10}/\text{HMDO}$ ^[23] (hexamethyldisiloxane) in refluxing toluene gave an undesired dithiocarbonyl byproduct.^[24] Replacement of P_4S_{10} with 2 equivalents of Lawesson's reagent^[25] in refluxing toluene only resulted in a trace amount of thioamide **16**. However, to our delight, by increasing the amount of Lawesson's reagent to 8 equivalents and keeping the reaction time to 1 hour enhanced the yield of thioamide **16** to 56% along with recovery of starting material **15** in 31% yield. It should be noted that prolonging the reaction time led to complete consumption of **15**, whilst affording a significant amount of dithiocarbonyl byproduct. Finally, reduction of **16** using W-2 Raney Ni^[26] in EtOH gave (±)-parvineostemonine (**1a**), the NMR spectra of which were identical to those of the natural product.^[27]

In summary, the first total synthesis of (±)-parvineostemonine (**1a**) has been achieved in 11 steps with an overall yield of 9.1% from commercially available tropinone **6**. Key transformations include a radical-mediated 7-*exo*-trig cyclization to construct the tricyclic cage-like skeleton **3** and a stereoselective carbonyl allylation/lactonization to form the spirobutenolide ring D.

Experimental Section

Ketones (**3**)

A solution of $n\text{Bu}_3\text{SnH}$ (394 μL , 1.465 mmol) and ACN (47.6 mg, 0.195 mmol) in anhydrous toluene (8 mL) was added dropwise to a boiling solution of **12** (338.3 mg, 0.975 mmol) in anhydrous toluene (30 mL) under argon over 3.5 h by employing a syringe-pump technique, and the mixture was further heated at reflux for 1 h. After cooling the solution was evaporated and the residue was purified by silica-gel chromatography ($\text{EtOAc}/\text{petroleum ether}$, 7:13–1:1) to give **13** (79.1 mg, 37%) as a white solid and **3** (51.7 mg, 24%) as a light yellow oil. Compound **13**: m.p. 104–106°C; ^1H NMR (400 MHz, CDCl_3): δ = 4.65–4.63 (d, J =

6.4 Hz, 1H), 4.20–4.17 (d, J = 8.4 Hz, 1H), 2.83–2.76 (dt, J = 12.8, 2.4 Hz, 1H), 2.46–2.37 (m, 2H), 2.15–2.10 (dd, J = 11.2, 5.6 Hz, 1H), 1.97–1.80 (m, 6H), 1.71–1.68 (m, 1H), 1.62–1.55 (m, 2H), 1.51–1.43 (m, 1H), 0.89–0.86 ppm (t, J = 7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3): δ = 212.4, 175.3, 64.6, 58.2, 46.1, 43.3, 36.2, 32.9, 30.0, 28.0, 22.1, 19.2, 11.3 ppm; IR (neat): $\tilde{\nu}$ = 2920, 1718, 1653, 1460 cm^{-1} ; MS (EI): m/z : 221, 193, 165, 150, 123, 85, 69; HRMS (ESI): m/z : calcd for $\text{C}_{13}\text{H}_{20}\text{NO}_2$ [$M+\text{H}$] $^+$: 222.1489; found 222.1487. Compound **3**: ^1H NMR (400 MHz, CDCl_3): δ = 4.68–4.66 (d, J = 7.6 Hz, 1H), 4.30–4.28 (d, J = 7.2 Hz, 1H), 2.80–2.73 (ddd, J = 14.4, 7.2, 7.2 Hz, 1H), 2.48–2.36 (m, 4H), 2.24–2.14 (m, 1H), 2.03–1.84 (m, 3H), 1.78–1.60 (m, 3H), 1.46–1.38 (m, 1H), 1.26–1.16 (m, 1H), 0.90–0.86 ppm (t, J = 7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3): δ = 204.7, 173.5, 64.9, 57.6, 47.3, 44.3, 35.3, 27.9, 26.6, 26.3, 19.2, 17.2, 11.8 ppm; IR (neat): $\tilde{\nu}$ = 2925, 1723, 1660, 1460 cm^{-1} ; MS (EI): m/z : 221, 193, 164, 137, 113, 97, 67; HRMS (ESI): m/z : calcd for $\text{C}_{13}\text{H}_{20}\text{NO}_2$ [$M+\text{H}$] $^+$: 222.1489; found 222.1487.

Lactone (**2**)

Zinc powder (254.2 mg, 3.887 mmol) was added to a solution of ketone **3** (85.9 mg, 0.389 mmol) in anhydrous THF (5 mL) under argon. The solution was heated, and when reflux started, a solution of ethyl (2-bromomethyl)acrylate (112.5 mg, 0.583 mmol) in anhydrous THF (2 mL) was added dropwise over 5 min. After stirring for an additional 15 min, the resultant mixture was cooled to room temperature, quenched with water (100 μL), and concentrated under vacuum. The residue was purified by silica-gel chromatography ($\text{EtOAc}/\text{acetone}$, 4:1) to afford **2** (87.3 mg, 78%) as a white amorphous solid; ^1H NMR (400 MHz, CDCl_3): δ = 6.23–6.22 (t, J = 2.8 Hz, 1H), 5.62–5.61 (t, J = 2.4 Hz, 1H), 4.61–4.60 (d, J = 7.2 Hz, 1H), 4.28–4.26 (d, J = 7.6 Hz, 1H), 2.83–2.67 (m, 3H), 2.31–2.18 (m, 3H), 2.15–1.99 (m, 3H), 1.95–1.92 (m, 1H), 1.78–1.65 (m, 4H), 1.41–1.33 (m, 1H), 1.32–1.23 (m, 1H), 0.90–0.86 ppm (t, J = 7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3): δ = 174.9, 169.1, 134.4, 122.2, 86.9, 62.6, 57.6, 43.2, 37.9, 35.9, 35.8, 27.6, 25.5, 24.1, 19.4, 16.7, 11.5 ppm; IR (neat): $\tilde{\nu}$ = 2958, 2922, 1751, 1633, 1458 cm^{-1} ; MS (EI): m/z : 289, 253, 211, 169, 141, 113, 99, 71; HRMS (ESI): m/z : calcd for $\text{C}_{17}\text{H}_{24}\text{NO}_3$ [$M+\text{H}$] $^+$: 290.1751; found 290.1750.

Thioamide (**16**)

Lawesson's reagent (543.1 mg, 1.343 mmol) was added to a solution of **15** (48.5 mg, 0.168 mmol) in anhydrous toluene (10 mL) under argon and the solution was heated at reflux for 1 h. After cooling the solvent was evaporated and the residue was purified by silica-gel chromatography ($\text{MeOH}/\text{CH}_2\text{Cl}_2$, 1:200–2:98) to afford **16** (28.7 mg, 56%) as a light yellow amorphous solid and recovered starting material **15** (14.8 mg, 31%). Thioamide **16**: ^1H NMR (600 MHz, CDCl_3): δ = 6.84–6.83 (d, J = 1.8 Hz, 1H), 4.96–4.95 (d, J = 7.8 Hz, 1H), 4.53–4.52 (d, J = 7.2 Hz, 1H), 3.35–3.33 (m, 1H), 3.17–3.12 (m, 1H), 2.35–2.28 (m, 1H), 2.21–2.17 (m, 1H), 2.04–1.98 (m, 3H), 1.94–1.93 (d, J = 1.2 Hz, 3H), 1.92–1.90 (d, J = 9.0 Hz, 2H), 1.72–1.61 (m, 2H), 1.32–1.27 (m, 1H), 1.08–1.01 (m, 1H), 0.89–0.84 (m, 1H), 0.83–0.80 ppm (t, J = 7.2 Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3): δ = 202.2, 172.0, 148.0, 132.9, 92.4, 67.0, 60.4, 45.3, 39.3, 37.5, 26.8, 24.4, 23.7, 20.2, 16.1, 11.4, 10.8 ppm; IR (neat): $\tilde{\nu}$ = 2926, 1715, 1459, 1374, 1240, 1047 cm^{-1} ; MS (EI): m/z : 305, 265, 223, 181, 149, 113, 85; HRMS (ESI): m/z : calcd for $\text{C}_{17}\text{H}_{24}\text{NO}_2\text{S}$ [$M+\text{H}$] $^+$: 306.1522; found 306.1522.

Parvineostemonine (**1a**)

Raney Ni W-2 (excess) was added to a solution of **16** (28.7 mg, 0.094 mmol) in anhydrous EtOH (10 mL) under argon at room temperature. After stirring for 2 h, the solvent was evaporated and the residue was purified by silica-gel chromatography ($\text{MeOH}/\text{CH}_2\text{Cl}_2$, 1:19) to afford (±)-parvineostemonine (**1a**) (23.3 mg, 90%) as a white amorphous solid; ^1H NMR (400 MHz, CDCl_3): δ = 6.88–6.87 (d, J = 1.6 Hz, 1H), 3.73–3.72 (d, J = 6.8 Hz, 1H), 3.66–3.59 (dt, J = 12.8, 3.6 Hz, 1H), 3.15–3.11 (ddd, J = 12.0, 3.2, 3.2 Hz, 1H), 2.99–2.97 (d, J = 6.8 Hz, 1H), 2.10–1.94 (m, 4H), 1.92–1.91 (d, J = 1.2 Hz, 3H), 1.89–1.81 (m, 3H), 1.78–1.64 (m, 3H), 1.58–1.52 (m, 1H), 1.39–1.23 (m, 2H), 1.04–0.98 (m, 1H), 0.82–0.78 ppm (t, J = 7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3): δ = 174.0,

152.8, 130.6, 89.4, 66.0, 56.9, 46.5, 38.2, 37.9, 28.2, 28.1, 27.1, 26.9, 24.0, 17.1, 11.7, 10.6 ppm; IR (neat): $\tilde{\nu}$ = 2923, 1747, 997, 731 cm^{-1} ; MS (EI): m/z : 275, 246, 178, 137, 122, 109, 95, 69; HRMS (ESI): m/z : calcd for $\text{C}_{17}\text{H}_{26}\text{NO}_2$ $[M+H]^+$: 276.1958; found 276.1965.

Acknowledgements

The authors thank Prof. Y. Ye (Shanghai Institute of Materia Medica) for providing us with the NMR spectra of parvineostemonine. This work was supported by NSFC (Grant Nos. 21072085, 20921120404, 21102061, and 20972059), the National Basic Research Program of China ("973" Program, Grant No. 2010CB833203), "111" Program of MOE, and scholarship award for excellent doctoral student of MOE (86007).

Keywords: allylation • parvineostemonine • radicals • *stemona* alkaloids • total synthesis

- [1] a) H. Iizuka, H. Irie, N. Masaki, K. Osaki, S. Ueno, *J. Chem. Soc. Chem. Commun.* **1973**, 125–126; b) K. Sakata, K. Aoki, C.-F. Chang, A. Sakurai, S. Tamura, S. Murakoshi, *Agric. Biol. Chem.* **1978**, *42*, 457–463; c) Y. Ye, G.-W. Qin, R.-S. Xu, *Phytochemistry* **1994**, *37*, 1205–1208; d) X.-Z. Yang, J.-Y. Zhu, C.-P. Tang, C.-Q. Ke, G. Lin, T.-Y. Cheng, J. A. Rudd, Y. Ye, *Planta Med.* **2009**, *75*, 174–177.
- [2] For recent reviews, see: a) R. A. Pilli, M. C. F. de Oliveira, *Nat. Prod. Rep.* **2000**, *17*, 117–127; b) R. A. Pilli, G. B. Rosso, M. C. F. de Oliveira in *The Alkaloids*, Vol. 62 (Ed.: G. A. Cordell), Elsevier, New York, **2005**, pp. 77–173; c) H. Greger, *Planta Med.* **2006**, *72*, 99–113; d) R. A. Pilli, G. B. Rosso, M. C. F. de Oliveira, *Nat. Prod. Rep.* **2010**, *27*, 1908–1937; e) S. Kongiatpaiboon, J. Schinnerl, S. Felsinger, V. Keeratinijakal, S. Vajrodaya, W. Gritsanapan, L. Breckner, H. Greger, *J. Nat. Prod.* **2011**, *74*, 1931–1938.
- [3] For reviews on synthetic studies of *Stemona* alkaloids, see: R. Alibés, M. Figueredo, *Eur. J. Org. Chem.* **2009**, 2421–2435; ref. [2a, 2b, 2c].
- [4] For recent synthetic studies reported on *Stemona* alkaloids, see: a) T. Honda, T. Matsukawa, K. Takahashi, *Org. Biomol. Chem.* **2011**, *9*, 673–675; b) Y. Wang, L. Zhu, Y. Zhang, R. Hong, *Angew. Chem.* **2011**, *123*, 2839–2842; *Angew. Chem. Int. Ed.* **2011**, *50*, 2787–2790; c) A. T. Hoye, P. Wipf, *Org. Lett.* **2011**, *13*, 2634–2637; d) R. W. Bates, S. Sridhar, *J. Org. Chem.* **2011**, *76*, 5026–5035; e) S.-C. Tuo, J.-L. Ye, A.-E. Wang, S.-Y. Huang, P.-Q. Huang, *Org. Lett.* **2011**, *13*, 5270–5273; f) J. Chen, J. Chen, Y. Xie, H. Zhang, *Angew. Chem.* **2012**, *124*, 1048–1051; *Angew. Chem. Int. Ed.* **2012**, *51*, 1024–1027.
- [5] For previous studies on *Stemona* alkaloids from our laboratory, see: a) Y.-M. Zhao, P. Gu, Y.-Q. Tu, C.-A. Fan, Q. Zhang, *Org. Lett.* **2008**, *10*, 1763–1766; b) Y.-M. Zhao, P. Gu, H.-J. Zhang, Q.-W. Zhang, C.-A. Fan, Y.-Q. Tu, F.-M. Zhang, *J. Org. Chem.* **2009**, *74*, 3211–3213; c) M. Tang, C.-A. Fan, F.-M. Zhang, Y.-Q. Tu, *Tetrahedron* **2009**, *65*, 5716–5719; d) Z.-H. Chen, Y.-Q. Zhang, Z.-M. Chen, Y.-Q. Tu, F.-M. Zhang, *Chem. Commun.* **2011**, *47*, 1836–1838; e) Z.-H. Chen, Y.-Q. Tu, S.-Y. Zhang, F.-M. Zhang, *Org. Lett.* **2011**, *13*, 724–727; f) Z.-H. Chen, Y.-Q. Tu, S.-Y. Zhang, F.-M. Zhang, *Scientia Sinica Chimica* **2011**, *41*, 474–478; g) Z.-H. Chen, Z.-M. Chen, Y.-Q. Zhang, Y.-Q. Tu, F.-M. Zhang, *J. Org. Chem.* **2011**, *76*, 10173–10186. For previous studies on constructing the skeleton of *Stemona* alkaloids from our laboratory, see: h) P. Gu, Y.-M. Zhao, Y. Ma, Y.-Q. Tu, F.-M. Zhang, *Org. Lett.* **2006**, *8*, 5271–5273; i) M. Yang, Y.-M. Zhao, S.-Y. Zhang, Y.-Q. Tu, F.-M. Zhang, *Chem. Asian J.* **2011**, *6*, 1344–1347; j) Q.-W. Zhang, K. Xiang, Y.-Q. Tu, S.-Y. Zhang, X.-M. Zhang, Y.-M. Zhao, T.-C. Zhang, *Chem. Asian J.* **2012**, *7*, 894–898.
- [6] C. Q. Ke, Z. S. He, Y. P. Yang, Y. Ye, *Chin. Chem. Lett.* **2003**, *14*, 173–175.
- [7] J. E. Antoline, R. P. Hsung, J. Huang, Z. Song, G. Li, *Org. Lett.* **2007**, *9*, 1275–1278.
- [8] G. L. Ellis, R. Amewu, S. Sabbani, P. A. Stocks, A. Shone, D. Stanford, P. Gibbons, J. Davies, L. Vivas, S. Charnaud, E. Bongard, C. Hall, K. Rimmer, S. Lozanom, M. Jesus, D. Gargallo, S. A. Ward, P. M. O'Neill, *J. Med. Chem.* **2008**, *51*, 2170–2177.
- [9] G. Mehta, H. M. Shinde, *Tetrahedron Lett.* **2003**, *44*, 7049–7053.
- [10] R. M. Moriarty, H. Hu, S. C. Gupta, *Tetrahedron Lett.* **1981**, *22*, 1283–1286.
- [11] M. G. Nilson, R. L. Funk, *J. Am. Chem. Soc.* **2011**, *133*, 12451–12453.
- [12] D. B. Dess, J. C. Martin, *J. Org. Chem.* **1983**, *48*, 4155–4156.
- [13] F. Frébault, M. T. Oliveira, E. Wöstefeld, N. Maulide, *J. Org. Chem.* **2010**, *75*, 7962–7965.
- [14] A. K. Ghosh, J. Li, *Org. Lett.* **2011**, *13*, 66–69.
- [15] K. C. Nicolaou, S. P. Ellery, J. S. Chen, *Angew. Chem.* **2009**, *121*, 7276–7301; *Angew. Chem. Int. Ed.* **2009**, *48*, 7140–7165.
- [16] L. Yet, *Tetrahedron* **1999**, *55*, 9349–9403.
- [17] a) T. Taniguchi, G. Tanabe, O. Muraoka, H. Ishibashi, *Org. Lett.* **2008**, *10*, 197–199; b) T. Taniguchi, H. Ishibashi, *Tetrahedron* **2008**, *64*, 8773–8779.
- [18] CCDC 871427 (**13**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [19] M. D. Lainchbury, M. I. Medley, P. M. Taylor, P. Hirst, W. Dohle, K. I. Booker-Milburn, *J. Org. Chem.* **2008**, *73*, 6497–6505.
- [20] a) A. P. Rauter, J. Figueiredo, M. Ismael, T. Canda, J. Font, M. Figueredo, *Tetrahedron: Asymmetry* **2001**, *12*, 1131–1146; b) F. Sánchez-Izquierdo, P. Blanco, F. Busqué, R. Alibés, P. de March, M. Figueredo, J. Font, T. Parella, *Org. Lett.* **2007**, *9*, 1769–1772.
- [21] H.-S. Byun, K. C. Reddy, R. Bittman, *Tetrahedron Lett.* **1994**, *35*, 1371–1374.
- [22] P. Grieco, M. Nishizawa, N. Marinovic, W. J. Ehmann, *J. Am. Chem. Soc.* **1976**, *98*, 7102–7104.
- [23] T. J. Curphey, *J. Org. Chem.* **2002**, *67*, 6461–6473.
- [24] Reaction of **15** with P_4S_{10} /HMDO at lower temperatures only resulted in recovery of starting material.
- [25] T. Ozturk, E. Ertas, O. Mert, *Chem. Rev.* **2007**, *107*, 5210–5278.
- [26] R. J. Sundberg, J. D. Bloom, *J. Org. Chem.* **1980**, *45*, 3382–3387.
- [27] See the Supporting Information.

Received: June 3, 2012

Published online: July 6, 2012