

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/26765487>

Enantioselective Synthesis of (–)-β-Santalol by a Copper-Catalyzed Enynol Cyclization-Fragmentation Reaction

ARTICLE *in* ANGEWANDTE CHEMIE INTERNATIONAL EDITION · SEPTEMBER 2009

Impact Factor: 11.26 · DOI: 10.1002/anie.200903449 · Source: PubMed

CITATIONS

29

READS

29

5 AUTHORS, INCLUDING:



Iris Magpantay

Firmenich


10 PUBLICATIONS 186 CITATIONS

SEE PROFILE

Rearrangement Reactions

Enantioselective Synthesis of (–)-β-Santalol by a Copper-Catalyzed Enynol Cyclization–Fragmentation Reaction**


Charles Fehr,* Iris Magpantay, Jeremy Arpagaus, Xavier Marquet, and Magali Vuagnoux



We report herein the first enantioselective, direct synthesis of the highly prized natural sandalwood odorant (–)-β-santalol ((–)-1).^[1] The key step in the synthesis is an efficient copper-catalyzed rearrangement of an enynol.

The increasing scarcity of East Indian sandalwood oil^[2] has motivated research chemists over the last 40 years to develop a synthesis of the most valuable component, (–)-β-santalol ((–)-1), which makes up 20–25% of the composition of the oil, or its racemate.^[1] Despite extensive efforts, this excellent fragrance ingredient is still not commercially available.


Recently, we reported the cost-efficient copper-catalyzed cycloisomerization of 5-en-1-yn-3-ols (cyclopropanation/1,2-alkyl shift; Scheme 1, **A**→**E**)^[3] and related enynol esters.^[4]



Scheme 1. Enynol cycloisomerization pathways.

These cyclopropanation reactions,^[5] which lead selectively to complex polycyclic compounds, are generally catalyzed by platinum^[6] or gold.^[7] During further studies on enynols of type **A**, we discovered that a cyclization–fragmentation pathway (Scheme 1, **A**→**D**) could compete with the cyclopropanation, and that [Cu(CH₃CN)₄]BF₄ was the reagent of choice for promoting this rearrangement, as opposed to Ag,^[8] Au,^[7b] or Pt catalysts.^[9] These results prompted us to devise a

new synthesis of (–)-β-santalol ((–)-1). If enynol **2** could be induced to preferentially undergo the cyclization–fragmentation to afford **3** (*E* or *Z*) with an *exo*-oriented enal side chain, at the expense of the cyclopropanation product **4**, this transformation would open a new direct route to (–)-1 (Scheme 2).



Scheme 2. Projected cyclization–fragmentation reaction en route to (–)-1.


The successful realization of this synthetic plan began with a highly enantioselective and *exo*-selective Diels–Alder reaction between cyclopentadiene and crotonaldehyde (Scheme 3). Chiral amines have been employed as efficient catalysts for enantioselective Diels–Alder reactions through the reversible formation of a dienophilic iminium salt.^[10] The most impressive results in terms of enantioselectivity and *exo* selectivity for the formation of **6** were reported by the research groups of MacMillan,^[10a] Maruoka,^[10b] and Hayashi.^[10c] The system described by Hayashi et al.,^[10c] a concentrated two-phase system in water at 20°C, is by far the most practical. Hayashi et al. used catalyst **5** (5 mol %) and a threefold excess of cyclopentadiene and obtained **6** (99% *ee*) and **7** (99% *ee*) in a 72:28 ratio in 72% combined yield. When we carried out the reaction under the same conditions, we obtained **6** (96% *ee*) and **7** (77% *ee*) in a 72:28 ratio and 55% combined yield. We were able to optimize this reaction by using equimolar amounts of cyclopentadiene and crotonaldehyde and only 1.5 mol % of catalyst **5** (**6**+**7**: 61% yield; **6** (95% *ee*)/**7** (78% *ee*) 70:30).^[11] The use of just 1 mol % of **5** still led to the formation of **6** (94% *ee*) and **7** (80% *ee*) in a 69:31 ratio.

The cycloadducts **6** and **7** were hydrogenated as a mixture to give the corresponding saturated alcohols, which were converted into methyl carbonates and pyrolyzed to afford hydrocarbons **8** and **9** in high overall yield.^[12,13] Upon treatment with trimethylsilylpropynal,^[14] **8** underwent the expected ene reaction to afford **2** after desilylation. However,

[*] Dr. C. Fehr, I. Magpantay, J. Arpagaus, X. Marquet, Dr. M. Vuagnoux^[4]
Firmenich SA, Corporate R&D Division
P. O. Box 239, 1211 Geneva 8 (Switzerland)
Fax: (+41) 22-780-3334
E-mail: charles.fehr@firmenich.com

[†] Postdoctoral fellow at Firmenich SA (July 2007–June 2008).

[**] We thank Dr. S. Samanta and Prof. Y. Hayashi (Tokyo University of Science (Japan)) for helpful advice concerning the Diels–Alder reaction.



Scheme 3. Reagents and conditions: a) Raney Ni (2%), 20 bar, 80°C, 24 h (100%); b) ClCO₂Me (2 equiv), pyridine, toluene, 0°C→RT, 1 h (90%); c) pyrolysis (415°C; quartz tube, 3 m; N₂ stream; 90%); d) TMSCHCHO (1.0 equiv with respect to **8**), Me₂AlCl (1.0 equiv), CH₂Cl₂, -20°C (72%); e) K₂CO₃ (10 equiv), MeOH (89%); f) H₂, 5% Pd/CaCO₃ (5%), MeOH/H₂O (96:4), 4 h (85%); g) EtPPh₃I (1.1 equiv), BuLi (1.1 equiv), THF, -78°C; then BuLi (1.2 equiv), -78°C→0°C; then paraformaldehyde (6 equiv), 0°C→RT, 1 h (50%). [A] toluene, 70°C, 1 h; [B] CH₂Cl₂, -10°C, 2 h; [C] ClCH₂CH₂Cl, 50°C, 140 min; [D] THF, H₂O, 70°C, 8 h (87% conversion). TMS=trimethylsilyl.

9, which has a less accessible α hydrogen atom, underwent a “homoeene reaction” exclusively to afford **10** after desilylation.^[15] The formation of **10** could be prevented completely by adapting the amount of trimethylsilylpropynal used to that of **8**.

We tested Pt, Au, and Cu catalysis for the key cycloisomerization reaction of enynol **2**, which existed as one major and one minor diastereomer. PtCl₂ catalysis led exclusively to the undesired cyclopropane **4** in 88% yield. In the presence of catalytic amounts of [PPh₃AuCl] and AgSbF₆, the cycloisomerization leading to **4** was again favored (**4/3** 91:9). Evidently, the AgCl formed in situ does not influence the reaction course. In sharp contrast, [Cu(CH₃CN)₄]BF₄

(5 mol%) efficiently promoted the desired cyclization–fragmentation reaction to afford the desired aldehyde **3**^[16] in high yield with excellent chemoselectivity and face selectivity (**3/4** 98:2).^[17] The use of AgNO₃ (10 mol%) and KNO₃ (1 equiv)^[8] also led to a selective reaction (**3/4** 99:1); however, the reaction was slower (87% conversion after 8 h) and the yield lower (64%).

The *ee* value of **3** could be increased to 97% through one low-temperature crystallization (pentane). Finally, site-selective hydrogenation with Pd/CaCO₃, followed by a Wittig reaction under the modified conditions described by Corey and Yamamoto with hydroxyalkylation of the intermediate betaine^[18] afforded the target (–)-**1** (Z/E 97:3; 43% (2 steps)),^[19] the sandalwood odor of which was excellent.

In conclusion, we have developed an efficient enantioselective synthesis of (–)- β -santalol. The key step in the synthesis consists of a highly selective copper-catalyzed cyclization–fragmentation of an enynol. The scope and generality of this novel methodology will be reported in due course.

Experimental Section

6 and 7: A mixture of catalyst **5** (314 mg, 0.45 mmol), water (3.78 mL, 210 mmol), toluene (0.24 mL; internal standard), and crotonaldehyde (95% *trans*; distilled and stored in the freezer; 2.10 g (2.46 mL), 30.0 mmol) was stirred at 20°C for 5 min and then treated with freshly prepared cyclopentadiene monomer (stored in a dewar containing dry ice; 1.98 g (2.44 mL), 30.0 mmol). The two-phase system was stirred at 23°C for 6 h, then cooled to 10°C, treated with saturated aqueous NaHCO₃ solution (2.5 mL) and solid NaCl (100 mg), and stirred for 15 min. The two-phase system was separated, and the aqueous phase was washed with pentane (3 mL). Concentration (0°C, 50 mbar) and bulb-to-bulb distillation (oven temperature 100°C, 4 mbar) afforded a mixture of **6** and **7** (70:30; 2.07 g, 61%) with 88% purity. This material, which contained traces of toluene, dicyclopentadiene, and two unidentified products that may have originated from *cis*-crotonaldehyde, was used for the ensuing hydrogenation.

(–)-**3**: [Cu(CH₃CN)₄]BF₄ (0.294 g, 0.93 mmol) was added to a solution of **2** (3.24 g, 18.4 mmol) in 1,2-dichloroethane (100 mL) at room temperature under nitrogen, and the resulting mixture was stirred at 50°C for 140 min. The dark-gray mixture was then cooled to room temperature, filtered through a short pad of silica gel, and concentrated under vacuum to give an orange oil (3.87 g). Bulb-to-bulb distillation of this oil (oven temperature 100°C, 0.08 mbar) afforded (–)-**3** (3.12 g, 94%, 92% *ee*) with 98% purity (the impurity corresponding to the remaining 2% was **4**). Crystallization once from pentane at –78°C afforded (–)-**3** (2.79 g) with 97% *ee*. $[\alpha]_{\text{D}}^{20} = -267.4 \text{ deg cm}^3 \text{ g}^{-1} \text{ dm}^{-1}$ (CHCl₃, $c = 1.06 \times 10^{-2} \text{ g cm}^{-3}$).

¹H NMR (400 MHz, CDCl₃): δ = 9.52 (d, J = 7.8 Hz, 1H), 6.77 (d, J = 15.7 Hz, 1H), 6.09 (dd, J = 15.7, 7.8 Hz, 1H), 5.01 (s, 1H), 4.58 (s, 1H), 2.76 (br s, 1H), 2.18 (br s, 1H), 1.69–1.80 (m, 2H), 1.52–1.58 (m, 2H), 1.30–1.37 (m, 1H), 1.24–1.27 (m, 1H), 1.23 ppm (s, 3H).

¹³C NMR (100 MHz, CDCl₃): δ = 194.5 (d), 165.4 (d), 160.4 (s), 130.1 (d), 104.1 (t), 49.7 (s), 46.5 (d), 46.3 (d), 37.1 (t), 29.8 (t), 23.0 (t), 22.7 ppm (q).

MS: m/z (%): 176 [M^+] (23), 161 (71), 147 (63), 133 (75), 119 (62), 105 (100), 91 (99), 79 (58), 77 (58), 39 (34).

Received: June 25, 2009

Published online: ■ ■ ■ ■ ■, 2009

Keywords: copper · cyclopropanes · enantioselectivity · fragrances · rearrangement


- [1] a) For the synthesis of (–)- β -santalol (very strong sandalwood odor) and (+)- β -santalol (odorless) through the use of chiral auxiliaries see: A. Krotz, G. Helmchen, *Liebigs Ann. Chem.* **1994**, 601; A. Krotz, G. Helmchen, *Tetrahedron: Asymmetry* **1990**, *1*, 537; for related studies, see: W. Oppolzer, C. Chapuis, D. Dupuis, M. Guo, *Helv. Chim. Acta* **1985**, *68*, 2100; N. Baldovini, G. Solladié, *Tetrahedron: Asymmetry* **2002**, *13*, 885; b) for the synthesis of (\pm)- β -santalol, see: D. Solas, J. Wolinsky, *J. Org. Chem.* **1983**, *48*, 1988; H. Monti, C. Corriol, M. Bertrand, *Tetrahedron Lett.* **1982**, *23*, 5539; K. Sato, O. Miyamoto, S. Inoue, K. Honda, *Chem. Lett.* **1981**, 1183; P. A. Christenson, B. J. Willis, *J. Org. Chem.* **1979**, *44*, 2012; H. C. Kretschmar, W. F. Erman, *Tetrahedron Lett.* **1970**, *11*, 41.
- [2] The wood of *Santalum album* L. has been overharvested, and the Indian government now bans exportation of this threatened species (2004 IUCN Red List).
- [3] C. Fehr, I. Farris, H. Sommer, *Org. Lett.* **2006**, *8*, 1839.
- [4] C. Fehr, J. Galindo, *Angew. Chem.* **2006**, *118*, 2967; *Angew. Chem. Int. Ed.* **2006**, *45*, 2901.
- [5] For reviews on enyne cycloisomerization reactions, see: H. C. Shen, *Tetrahedron* **2008**, *64*, 7847; E. Jiménez-Núñez, A. M. Echavarren, *Chem. Rev.* **2008**, *108*, 3326; V. Michelet, P. Y. Toullec, J.-P. Genêt, *Angew. Chem.* **2008**, *120*, 4338; *Angew. Chem. Int. Ed.* **2008**, *47*, 4268; B. Crone, S. F. Kirsch, *Chem. Eur. J.* **2008**, *14*, 3514; A. Fürstner, P. W. Davies, *Angew. Chem.* **2007**, *119*, 3478; *Angew. Chem. Int. Ed.* **2007**, *46*, 3410; E. Jiménez-Núñez, A. M. Echavarren, *Chem. Commun.* **2007**, 333; D. J. Gorin, D. Toste, *Nature* **2007**, *446*, 395; N. Marion, S. P. Nolan, *Angew. Chem.* **2007**, *119*, 2806; *Angew. Chem. Int. Ed.* **2007**, *46*, 2750; J. Marco-Contelles, E. Soriano, *Chem. Eur. J.* **2007**, *13*, 1350; L. Zhang, J. Sun, S. A. Kozmin, *Adv. Synth. Catal.* **2006**, *348*, 2271; C. Nieto-Oberhuber, S. López, E. Jiménez-Núñez, A. M. Echavarren, *Chem. Eur. J.* **2006**, *12*, 5916; S. Ma, S. Yu, Z. Gu, *Angew. Chem.* **2006**, *118*, 206; *Angew. Chem. Int. Ed.* **2006**, *45*, 200; C. Bruneau, *Angew. Chem.* **2005**, *117*, 2380; *Angew. Chem. Int. Ed.* **2005**, *44*, 2328; A. M. Echavarren, C. Nevado, *Chem. Soc. Rev.* **2004**, *33*, 431; C. Aubert, O. Buisine, M. Malacria, *Chem. Rev.* **2002**, *102*, 813.
- [6] X. Moreau, J.-P. Goddard, M. Bernard, G. Lemièrre, J. M. López-Romero, E. Mainetti, N. Marion, V. Mouriès, S. Thorimbert, L. Fensterbank, M. Malacria, *Adv. Synth. Catal.* **2008**, *350*, 43; C. Blaszykowski, Y. Harrak, C. Brancour, K. Nakama, A.-L. Dhimane, L. Fensterbank, M. Malacria, *Synthesis* **2007**, 2037; E. Soriano, J. Marco-Contelles, *J. Org. Chem.* **2007**, *72*, 2651; J. Barluenga, L. Riesgo, R. Vicente, L. A. López, M. Tomás, *J. Am. Chem. Soc.* **2007**, *129*, 7772 (Pt and Cu); S. Anjum, J. Marco-Contelles, *Tetrahedron* **2005**, *61*, 4793; E. Soriano, P. Ballesteros, J. Marco-Contelles, *Organometallics* **2005**, *24*, 3182; E. Soriano, J. Marco-Contelles, *J. Org. Chem.* **2005**, *70*, 9345; C. Blaszykowski, Y. Harrak, M.-H. Gonçalves, J.-M. Cloarec, A.-L. Dhimane, L. Fensterbank, M. Malacria, *Org. Lett.* **2004**, *6*, 3771; Y. Harrak, C. Blaszykowski, M. Bernard, K. Cariou, E. Mainetti, V. Mouriès, A.-L. Dhimane, L. Fensterbank, M. Malacria, *J. Am. Chem. Soc.* **2004**, *126*, 8656; V. Mamane, T. Gress, H. Krause, A. Fürstner, *J. Am. Chem. Soc.* **2004**, *126*, 8654.
- [7] a) N. Marion, G. Lemièrre, A. Correa, C. Costabile, R. S. Ramon, X. Moreau, P. de Frémont, R. Dahmane, A. Hours, D. Lesage, J.-C. Tabet, J.-P. Goddard, V. Gandon, L. Cavallo, L. Fensterbank, M. Malacria, S. P. Nolan, *Chem. Eur. J.* **2009**, *15*, 3243; I. D. G. Watson, S. Ritter, F. D. Toste, *J. Am. Chem. Soc.* **2009**, *131*, 2056; X. Moreau, J.-P. Goddard, M. Bernard, G. Lemièrre, J. M. López-Romero, E. Mainetti, N. Marion, V. Mouriès, S. Thorimbert, L. Fensterbank, M. Malacria, *Adv. Synth. Catal.* **2008**, *350*, 43; A. Fürstner, P. Hannen, *Chem. Eur. J.* **2006**, *12*, 3006; G. Lemièrre, V. Gandon, K. Cariou, T. Fukuyama, A.-L. Dhimane, L. Fensterbank, M. Malacria, *Org. Lett.* **2007**, *9*, 2207; A. Buzas, F. Gagosz, *J. Am. Chem. Soc.* **2006**, *128*, 12614; E. J. Cho, M. Kim, D. Lee, *Org. Lett.* **2006**, *8*, 5413; M. J. Johansson, D. J. Gorin, S. T. Staben, F. D. Toste, *J. Am. Chem. Soc.* **2005**, *127*, 18002; M. R. Luzung, J. P. Markham, F. D. Toste, *J. Am. Chem. Soc.* **2004**, *126*, 10858; b) F. Gagosz, *Org. Lett.* **2005**, *7*, 4129; c) see also: C. M. Grisé, E. M. Rodrigue, L. Barriault, *Tetrahedron* **2008**, *64*, 797.
- [8] N. Bluthé, J. Goré, M. Malacria, *Tetrahedron* **1986**, *42*, 1333.
- [9] The copper reagent possibly produces an intermediate of type **B** with more-localized charges and thus favors the fragmentation pathway. As **B** and **C** are two extreme representations of the same intermediate, the ease of fragmentation may also be related to the unlikelihood that a 1,2-R shift would take place in this system (owing to the electrophilicity of the carbenoid).
- [10] a) D. W. C. MacMillan, PCT Int. Appl. 2003002491; K. A. Ahrendt, C. J. Borths, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2000**, *122*, 4243; b) T. Kano, Y. Tanaka, K. Maruoka, *Chem. Asian J.* **2007**, *2*, 1161; c) Y. Hayashi, S. Samanta, H. Gotoh, H. Ishikawa, *Angew. Chem.* **2008**, *120*, 6736; *Angew. Chem. Int. Ed.* **2008**, *47*, 6634; d) Q. Chu, W. Zhang, D. Curran, *Tetrahedron Lett.* **2006**, *47*, 9287; M. Lemay, L. Aumand, W. W. Ogilvie, *Adv. Synth. Catal.* **2007**, *349*, 441 (enantiomer); C. Biaggi, M. Benaglia, S. Rossi, S. Proto, R. Annunziata, *Tetrahedron Lett.* **2007**, *48*, 8521; C. S. Pecinovsky, G. D. Nicodemus, D. L. Gin, *Chem. Mater.* **2005**, *17*, 4889.
- [11] The *ee* values were determined by GC on a chiral stationary phase of the corresponding trifluoroacetates obtained by reduction (LiAlH₄, Et₂O) and esterification ((CF₃CO)₂O) according to: D. Seebach, R. Dahinden, R. E. Marti, A. K. Beck, D. A. Plattner, F. N. M. Kühnle, *J. Org. Chem.* **1995**, *60*, 1788.
- [12] We chose to prepare compounds **8** and **9** by this method based on a known procedure (G. Joachimsman-Dufresne, M. Blanchard, *Bull. Soc. Chim. Fr.* **1968**, 385), as the conditions are neutral (in contrast to acetate pyrolysis), industrially sound (as opposed to mesylate elimination), and high-yielding.
- [13] Surprisingly, the *ee* values of **8** and, to a larger extent, **9** were lower than those of **6** and **7**, although the diastereomeric ratio remained the same. Compounds **8** and **9** possibly undergo a thermal 1,2-H shift.
- [14] K. J. H. Kruithof, R. F. Schmitz, G. W. Klumpp, *Tetrahedron* **1983**, *39*, 3073.
- [15] For another example, see: R. L. Snowden, R. Näf, *Tetrahedron Lett.* **1984**, *25*, 2877.
- [16] For a long synthesis of (\pm)-**3** of unspecified configuration (8 steps, 13% yield), see: M. Baumann, W. Hoffmann (BASF), EP 0010213, **1980** [*Chem. Abstr.* **1980**, *93*, 185844]; spectral data are not reported.
- [17] The use of [Cu(CH₃CN)₄]BF₄ in toluene (70°C, 1 h) led to completely selective cyclization–fragmentation; however, the yield was inferior (55%).
- [18] E. J. Corey, H. Yamamoto, *J. Am. Chem. Soc.* **1970**, *92*, 226.
- [19] We are presently tackling the longstanding problem that no more general alternative exists for the synthesis of *Z* trisubstituted allylic alcohols.

Communications

Rearrangement Reactions

C. Fehr,* I. Magpantay, J. Arpagaus,
X. Marquet, M. Vuagnoux — ■■■■—■■■■

Enantioselective Synthesis of (–)-β-Santalol by a Copper-Catalyzed Enynol Cyclization–Fragmentation Reaction



The right cat for the desired odor: The key step in an enantioselective synthesis of the prized fragrance (–)-β-santalol was a highly selective copper-catalyzed cyclization–fragmentation reaction of an enynol (see scheme). When a platinum catalyst was used for the cyclization step, the desired fragmentation did not take place; instead, a product containing a cyclopropane ring was formed with 100% selectivity.