

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/262680983>

Ebselen, a promising antioxidant drug: Mechanisms of action and targets of biological pathways

ARTICLE *in* MOLECULAR BIOLOGY REPORTS · MAY 2014

Impact Factor: 2.02 · DOI: 10.1007/s11033-014-3417-x · Source: PubMed

CITATIONS

20

READS

316

2 AUTHORS, INCLUDING:



[Gajendra Kumar Azad](#)

Hebrew University of Jerusalem

18 PUBLICATIONS 148 CITATIONS

SEE PROFILE

Ebselen, a promising antioxidant drug: mechanisms of action and targets of biological pathways

Gajendra Kumar Azad · Raghuvir S. Tomar

© Springer Science+Business Media Dordrecht 2014

Abstract Ebselen, an organoselenium compound, mimics glutathione peroxidase activity. It is a multifunctional compound, which catalyzes several essential reactions for the protection of cellular components from oxidative and free radical damage. Based on a number of *in vitro* and *in vivo* studies, various mechanisms are proposed to understand the biomedical actions of ebselen in health and diseases. It modulates metallo-proteins, enzymatic cofactors, gene expression, epigenetics, antioxidant defenses and immune systems. Owing to these properties, ebselen is currently under clinical trials for the prevention and treatment of various disorders such as cardiovascular diseases, arthritis, stroke, atherosclerosis, and cancer. A few ebselen-based pharmaceutical agents are under extensive investigation. As ebselen has been shown to have significant cellular toxicity, appropriate studies are needed to redesign the ebselen-based therapy for clinical trials. This review summarizes current understanding of the biochemical and molecular properties, and pharmacological applications of ebselen and future directions in this area of research.

Keywords Ebselen · Organoselenium compounds · Redox biology · Antioxidant · Enzyme mimic

Introduction

Ebselen (2-phenyl-1,2-benzisoselenazol-3(2H)-one) is an organoselenium compound (Fig. 1), chemically is an

electrophile [1]. The general mechanism of action of ebselen is through reactions with specific cysteine thiol groups in proteins [2, 3]. Ebselen (EBS) catalyzes the reduction of reactive oxygen species (ROS) in a manner similar to glutathione peroxidase (GPx), which operates by a simple, three-step mechanism involving changes in the oxidation state of the active site (SeCys) residue (Fig. 2). ROS oxidize the resting state selenol (EBS-SeH) to selenenic acid (EBS-SeOH), which is reduced to the active selenol by glutathione (GSH) through a selenenyl sulfide intermediate (EBS-SeSG) [4, 5].

Ebselen also reacts with the thioredoxin (Trx) system [3]. Ebselen is an excellent substrate for the mammalian thioredoxin reductase (TrxR) and acts as a highly efficient oxidant of the reduced Trx, and catalyzes the hydrogen peroxide (H_2O_2) reductase activity of TrxR. Similarly, ebselen has also been shown to act as a dehydroascorbic acid (DHA) reductase mimetic [6]. A considerable extent of the pharmacological activity of ebselen is attributed to its antioxidant action. Recent animal studies have demonstrated the anti-inflammatory activities of ebselen in a variety of experimental animal models [7, 8]. Due to its antioxidant behavior, it exhibits anti-lipoperoxidative, anti-atherosclerotic, anti-thrombotic, and cytoprotective properties [9–15]. Ebselen is a lipid-soluble compound, and therefore readily enters the cell. Ebselen potently inhibits lipid peroxidation through a GPx-like action, as it has the ability to oxidize glutathione thiol (GSH) and reduce hydrogen peroxide to water [16–19]. Moreover, ebselen has a strong protective ability against cytotoxicity and DNA damage caused by the reactive oxygen species (ROS) generated during various cellular processes [20]. ROS, including H_2O_2 , superoxide radical ($O_2^{\bullet-}$), hydroxyl radical ($\bullet OH$), and the reactive nitrogen species (RNS) such as peroxynitrite ($ONOO^-$) have been shown to increase upon

G. K. Azad · R. S. Tomar (✉)
Laboratory of Chromatin Biology, Department of Biological Sciences, Indian Institute of Science Education and Research, Bhopal 462023, India
e-mail: rst@iiserb.ac.in




Fig. 1 Chemical structure of Ebselen (2-Phenyl-1,2-benzoselenazol-3-one)




Fig. 2 Catalytic mechanism of action of ebselen. Ebselen (*EBS*) catalyzes the reduction of reactive oxygen species (*ROS*), which operates by a simple, three-step mechanism involving changes in the oxidation state of the active site Se molecule. *ROS* oxidize the resting state selenol (*EBS-SeH*) to selenenic acid (*EBS-SeOH*), which is reduced to the active selenol by glutathione (*GSH*) through a selenenyl sulfide intermediate (*EBS-SeSG*)

ischemia and reperfusion damage [17]. Ebselen rapidly reacts with peroxynitrite and other free radicals generated during ischemia or reperfusion [21–23] and protects against the deleterious effects of these *ROS* and *RNS* [5, 8, 24–28]. Due to this wide range of effects, ebselen has been investigated for the treatment of various ailments such as arthritis, stroke, organ transplantation, anti-asthmatic, ozone toxicity, diabetes-related atherosclerosis and nephropathy, and as an inhibitor of the neuronal demyelination [11, 29–32].

Contrary to the beneficial effects shown by ebselen, it also causes cellular toxicity. Based on various studies it is known to cause deterioration of the mitochondrial function by stimulating Ca^{2+} release from the mitochondria through the NAD^+ hydrolysis-dependent mechanism [15, 33, 34]. Furthermore, high concentrations of ebselen induce the necrotic cell death of Sp2/0-Ag14 hybridoma cells [35]. It also causes genotoxicity, induction of DNA damage in V79 cells [36], and apoptosis in HepG2 cells [37]. Recently, it has been that ebselen can also induce DNA damage in the yeast (*Saccharomyces cerevisiae*) leading to activation of the checkpoint kinase proteins and derepression of DNA damage response genes [38]. Ebselen also inhibits the

DNA binding mechanisms of the cysteine- and zinc-containing transcription factors including TFIIIA, Sp1, and NF-kappa B [39]. Ebselen reacts with metallothionein (*MT*) and other biological zinc/sulfur coordination sites to release zinc. As a result, zinc finger motifs are highly reactive towards this compound, possibly affecting the gene expression, DNA repair and, therefore, genomic stability [40].

Taken together, ebselen targets a wide variety of proteins and modulates several biological processes, leading to both beneficial and harmful actions in the biological systems. Ebselen is being used in clinical trials for the treatment of various diseases. The outcome of these studies may provide additional insights into the mechanisms underlying the biological properties of ebselen. The focus of this review is to summarize the biological/biochemical and molecular pathways targeted by this antioxidant molecule.

Ebselen localization inside the cell

Experimentally, it has been demonstrated that upon administration into the bloodstream, extracellular ebselen is transported as an albumin complex [41] bound to the reactive cysteine residue in albumin [42]. It is difficult to track the intracellular movement of ebselen within the cell because it is not fluorescent. The attachment of tags or reporter groups may fundamentally change the physicochemical properties of ebselen. However, ebselen does contain one potentially useful spectroscopic marker, in that it integrates an organoselenium moiety within its structure. As a trace element, selenium (*Se*) is present in cultured cells in extremely low concentrations. Therefore, the *Se* content of the ebselen-treated cells would be expected to be much higher than that of cells with normal background *Se* levels. Based on this selective increase of *Se*, it is anticipated that the analysis of cellular *Se* distribution would provide information on the intracellular localization and concentration of ebselen. Recently, Aitken et al. [43] have adopted this approach to analyze the intracellular localization of ebselen by using synchrotron radiation-induced X-ray emission (*SRIXE*) spectroscopy. Based on their results ebselen uptake appears to peak after 2 h, following which the cellular *Se* content declines, probably due to metabolic complexation and drug efflux. Analysis of the elemental maps suggests that ebselen or its metabolites immediately localizes in the ER following uptake and remains predominantly partitioned within this organelle for the 4 h of the experimental duration [43].

Several reports indicate that the mitochondria are also potential targets of ebselen [44–46]. In rat liver, the effect of ebselen on the electron transport chain has been studied.

Table 1 List of the proteins whose activity is inhibited by ebselen

| S. No. | Enzyme/protein | Action of ebselen | References |
|--------|--|---|------------|
| 1 | JMJD2A | Disrupts Zn binding site | [63] |
| 2 | TFIIIA and xeroderma pigmentosum group A | Disrupts Zn finger motif | [39, 64] |
| 3 | Alcohol dehydrogenase and metallothioneine (MT) | Disrupts Zn finger motif | [40] |
| 4 | Fpg (formamidopyrimidine-DNA glycosylase) | Disrupts Zn finger motif | [64] |
| 5 | Lipoxygenases | Alteration of the iron ligand sphere | [50, 51] |
| 6 | NO synthases | Reacts with its critical thiol group | [52] |
| 7 | NADPH oxidases (Nox) | Inhibits assembly of the Nox2 regulatory subunits | [53] |
| 8 | Horseradish peroxidase | Interferes with sulfhydryl groups | [55] |
| 9 | H ⁺ /K ⁺ -ATPase | Interferes with sulfhydryl groups | [54] |
| 10 | Lactate dehydrogenase | Interacts with critical thiol groups | [56] |
| 11 | P50 subunit of NF-κB and Sp1 | Disrupts its Zn finger motif | [39] |
| 12 | Heme enzyme indoleamine 2,3-dioxygenase (IDO) | reacts with multiple cysteine residues and disrupts its active site | [58] |
| 13 | Na ⁺ , K ⁺ -ATPase | Interacts with critical thiol groups of this enzyme | [57] |
| 14 | Inositol Monophosphatase (IMPase) | Covalent and irreversible inhibition | [130] |
| 15 | <i>P. falciparum</i> Hexokinase (PfHK) | Non-covalent interaction with HK | [153] |
| 16 | <i>Pseudomonas aeruginosa</i> Diguanylate cyclases | Reacts with its critical thiol group | [150] |
| 17 | divalent metal transporter-1 (DMT1) | Noncovalent interaction | [102, 103] |

At low micromolar concentrations, ebselen inhibits the flow of reducing equivalents from NADPH-cytochrome P450 reductase to both of its electron acceptors, cytochrome P450 and cytochrome *c* [47]. In addition, ebselen has been shown to protect the mitochondrial membranes against peroxidative reactions in the liver mitochondria isolated from glutathione-depleted rats [48]. Furthermore, ebselen significantly prevents the mitochondrial depolarization in ischemic astrocytes [25]. Another report suggests that depending on the concentration used, ebselen exerts deleterious actions on astrocyte physiology by altering the permeability of the mitochondrial membrane as well as calcium homeostasis [49]. Taken together, these results indicate that the ER and mitochondria are the most preferred intracellular targets of ebselen and, presumably, the pharmacological activity of this compound might be attributed to its ability to modulate the function of these two organelles. Once inside the cell, ebselen covalently and non-covalently interacts with several biomolecules. In the next section, we have described some of the well-established molecular targets of ebselen.

Intracellular targets of ebselen

In the last few years, many analytical methods (e.g., X-ray diffraction, high performance liquid chromatography, nuclear magnetic resonance, infrared, circular dichroism, and ultraviolet–visible (UV/Vis) spectroscopies) have been utilized to characterize the drug-protein systems. These techniques have been used to show that ebselen reacts with

the thiols to form stable selenenosulfide bonds [50]. Moreover, ebselen appears to act on multiple pathways and protein targets, and its mode of action in cells is pleiotropic. The reactivity of ebselen with the protein thiols makes it a potent modulator for proteins that require cysteine for normal function. Ebselen inhibits a number of enzymes involved in various biological processes, such as lipoxygenases [51, 52], nitric oxide synthases (NOS) [53], NADPH oxidases [54], H⁺-K⁺-ATPase [55], horseradish peroxidase [56], glutamate dehydrogenase [57], lactate dehydrogenase [58], and many others listed in Table 1. In most cases, ebselen targets critical thiol residues of enzymes, and inhibition can be reversed by the addition of reducing agents such as dithiothreitol (DTT) [53, 55, 59–61].

The zinc-finger (ZF) domain is another ebselen target. ZF transcription factors contain cysteine and histidine residues (characteristically Cys4, Cys3His, or Cys2His2) that tetrahedrally coordinate Zn²⁺ ions to ensure appropriate folding of the ZF tertiary structure, which is required for the biological recognition [62, 63]. The Zn²⁺ ion coordinates to a ZF in its reduced state, in which all cysteine residues are in the thiolate form. Alterations in the oxidation state of cysteine cause the release of Zn²⁺, leading to the disruption of tertiary structure and ultimately the loss of ZF function. Ebselen inhibits the function of several ZF-containing proteins by a similar mechanism [64]. For example, JMJD2A, a 2-oxoglutarate-dependent N(epsilon)-methyl lysine histone demethylase, is inhibited by the disruption of its Zn-binding site [65]. Ebselen reacts with metallothionein (MT) and alcohol dehydrogenase to

release zinc [40]. Ebselen also inhibits the DNA binding mechanisms of several ZF-containing proteins, including TFIIIA (transcription factor IIIA), DNA binding domain (DBD) of the NF- κ B mediated transcription factor Sp1 [39], formamidopyrimidine DNA glycosylase (Fpg) [66], xeroderma pigmentosum group A protein (XPA), and NF- κ B affecting the expression of downstream genes [39, 66]. Apart from its inhibitory effect on many DNA binding proteins, ebselen can also induce transcription of the antioxidant response element (ARE) [67] and nuclear factor erythroid 2-related factor 2 [68, 69].

Due to its interaction with several classes of biomolecules, it is likely that this molecule will influence a broad range of biological processes. In the next section, we will describe some of these.

Biological pathways affected by ebselen

Recent studies have illustrated that ebselen is a highly pleiotropic molecule that interacts with several molecular targets inside the cell. Ebselen may alter the activity of target proteins by direct binding or regulate the protein function indirectly.

Apoptosis

Apoptosis occurs normally during development and aging, and functions as a homeostatic mechanism to maintain cell populations in tissues [70]. At low doses, a variety of injurious stimuli such as heat, radiation, hypoxia, and cytotoxic anticancer drugs can induce apoptosis [71, 72]. Ebselen is known to prevent apoptosis that might occur due to injury or exposure to toxic agents. For example, it prevents apoptosis induced due to reperfusion of ischemic tissues by reducing the oxidative stress associated with ischemia/reperfusion [73–76]. The alkylating agent nitrogen mustard (HN2) and ionizing radiation can cause apoptosis through the production of oxygen free radicals. Ebselen provides efficient protection against HN2-induced cell death in lymphocytes, and can prevent cellular damage caused by the alkylating agents [77]. Ebselen is equally effective against ionizing radiation. Upon exposure to γ -irradiation, there is a distinct difference between untreated cells and cells pretreated with ebselen with respect to viability, cellular redox status, and oxidative damage [78]. Treatment of cells with ebselen reduces peroxide levels, protecting the thymocytes from radiation-induced apoptosis produced during and after gamma-irradiation [79]. Ebselen attenuate oxidative stress-induced apoptosis by scavenging free radicals in PC12 cell lines [80].

The mechanism through which ebselen prevents apoptotic events remains unclear. One study showed that in

response to γ -irradiation, ebselen effectively suppresses programmed cell death by decreasing the apoptotic events, such as the caspase activation, increased levels of anti-apoptotic molecules (Bcl-2), and decreased levels of proapoptotic molecules (Bax), presumably via the preservation of redox status [78]. Similarly, it also attenuates the H₂O₂-induced cell death through the inhibition of signaling pathways mediated by p38 MAP kinase (p38-MAPK) [81, 82]. The neuroprotective effects of ebselen at low doses are associated with the increased expression of Bcl-2 and inhibition of Bax proteins [83]. Furthermore, ebselen potently inhibits cytochrome *c* release and caspase-3 activation in differentiated PC12 cells in response to nitric oxide (NO)-induced apoptosis. Ebselen also blocks the activation of apoptosis signal-regulating kinase 1 (ASK1) and c-Jun N-terminal protein kinase (JNK) [84]. In addition, it also activate p44/42 MAPK and inhibit the down-regulation of Bcl-2 in SNP-treated PC12 cells [82]. Similar effects have been observed in a mouse model upon ebselen treatment. Furthermore, it has been demonstrated that ebselen attenuates ischemic neuronal apoptosis by inhibiting cytochrome *c* release in ICR mice [28].

Although most studies show that ebselen inhibits apoptosis under different experimental conditions, the mode of action is primarily through the interaction with thiol-containing proteins. Interestingly, ebselen induces apoptosis in HepG2 cells in a dose- and time-dependent manner. Evidence suggests that ebselen exposure causes rapid depletion of intracellular thiols leading to apoptotic cell death [37].

Genomic Integrity

DNA is constantly damaged by exogenous and endogenous factors. One of the main causes of DNA damage is the production of ROS, which are highly reactive molecules that are constantly generated due to metabolic activities or exposure to environmental factors [85, 86]. ROS can cause single- or double-stranded breaks. Ebselen offers strong protection against ROS-mediated cytotoxicity and DNA damage [22, 87, 88]. Ebselen exhibits protective actions against H₂O₂-induced cytotoxicity and DNA damage in HepG2 cells [87]. In vitro, ebselen efficiently protects against peroxynitrite-induced DNA single-strand breaks through free radical scavenging [89, 90]. Low concentrations of ebselen (5–10 μ M) do not induce DNA damage in V79 cells and budding yeast, while the same concentrations diminish the extent of DNA damage induced by H₂O₂ [36]. Ebselen exhibits DNA protective effects against several toxic compounds. For example, it attenuates the cyclophosphamide-induced DNA damage in mice [91], and it also significantly reduces formation of the hepatic aflatoxin

B1 (AFB1)–DNA adducts and 8-hydroxydeoxyguanosine following administration of AFB1, a potent hepatocarcinogen, in a rat model [92].

However, at higher doses, ebselen acts as genotoxic agent and induces DNA damage [36]. Significant growth arrest have been reported in yeast cells exposed to ebselen [93]. Further, it has been demonstrated that ebselen causes the derepression of DNA repair genes through the activation of the Mec1-Rad53-Dun1 kinase pathway [38], this kinase pathway regulates DNA damage response in yeast cells [94]. We have noted that ZF (zinc finger) motifs are highly reactive towards oxidizing selenium compounds including ebselen. Potentially, ebselen-induced disruption of the ZF motif adversely affects gene expression, DNA repair and, therefore, genomic stability [66].

Immune system

Ebselen exhibits pleiotropic modes of action as it targets different biological pathways. Ebselen is known as a modest cytokine inducer or immunostimulant [95]. Ebselen exposure has been reported to induce several interleukins including interleukin-1 (IL-1), IL-6, IL-10, and IL-18. Furthermore, ebselen stimulates production of the interferon (IFN)-gamma, tumor necrosis factor (TNF)- α , intercellular adhesion molecule (ICAM)-1, affecting immune cell adhesion and migration and contributing to its anti-inflammatory properties [96–99]. Conversely, ebselen has been shown to inhibit the TNF- α production and mRNA expression in rat primary Kupffer cells. In addition, ebselen neutralizes the liver injury induced by the superantigen staphylococcal enterotoxin B [96], suggesting that ebselen has an immunomodulatory function. Ebselen also inhibited inhibits the lipopolysaccharide (LPS)-induced phosphorylation of JNKs (c-Jun amino-terminal kinases), leading to decreased transcription of TNF- α [100].

Ebselen administration reduces the serum levels of TNF- α , IL-18/IGIF (Interferon-gamma-inducing factor), and interferon (IFN)- γ after LPS administration in mice [101]. It has also been reported that ebselen attenuates liver injury induced by TNF- α and enhances the release of IL-10 [79]. Similarly, treatment of rats with ebselen lowers plasma IL-6 levels in glial cells, leading to decreased levels of heme oxygenase-1, an inducible heat shock protein [97]. Ebselen also inhibits ICAM-1 expression in leukocytes [102] and decreases TNF- α levels, leading to suppression of bronchiolar inflammation [103]. Additionally, ebselen is found to inhibit the adhesion of polymorphonuclear leukocytes (PMNL) to IL-1-activated endothelium and to inhibit the transendothelial PMNL migration [104]. Ebselen is equally effective as a suitable anti-inflammatory agent in ocular tissues [105]. One interesting study show that ebselen treatment enhances the viability of bone

marrow-derived cells (BM-DCs) during transplantation by decreasing oxidative stress during this procedure [106]. Lung function in the ebselen-treated rats is also significantly improved at 24 h after transplantation [26].


Collectively, the results of the above studies suggest that ebselen modulates the host immune system and exhibits anti-inflammatory effects in response to various stimuli.

Cellular transport

Different genetic screening assays have shown that ebselen inhibits activity of the divalent metal transporter 1 (DMT1) [107–110]. DMT1, a proton-coupled metal-ion transport protein expressed in neurons is known to actively transport several different divalent cations such as Fe²⁺, Zn²⁺, Mn²⁺, Co²⁺, Cd²⁺, Cu²⁺, Ni²⁺, and Pb²⁺ [111, 112]. Interestingly, ebselen has been found to be specific for inhibiting uptake of iron but not manganese [109]. Surprisingly, DTT supplementation fails to reverse inhibition of DMT1-mediated iron uptake, suggesting that ebselen does not block DMT1 transporter function by direct modification of protein thiols. It has been proposed that ebselen might influence the activity of specific factors involved in the intracellular targeting of iron [109]. Recently, it has been demonstrated that DMT1 inhibition by ebselen leads to reduction in ferrous iron-induced activation of CDK5 (cyclin-dependent kinase 5) and GSK3 β (glycogen synthase kinase 3 beta) in human neuroblastoma SH-SY5Y cells [113]. However, ebselen not only interferes with iron metabolism in cells, but also affects the intracellular calcium homeostasis. One study shows that ebselen hampers an agonist-triggered increase in intracellular calcium by inhibiting inositol 1,4,5-trisphosphate (IP₃)-induced calcium release [114, 115]. Ebselen has also been shown to increase free cytosolic Ca²⁺ concentrations in rat hippocampal astrocytes [116]. Additionally, ebselen has been found to degrade mitochondrial function by stimulating Ca²⁺ release from mitochondria [33]. As calcium is involved in several signaling pathways, these results suggest that alteration of calcium homeostasis by ebselen might affect the downstream pathways. For example, ebselen inhibits the kinase activity of partially purified Ca²⁺- and phospholipid-dependent protein kinase C [117]. Taken together, these studies indicate that ebselen targets iron and calcium homeostasis in the cells by modulating the transporter and storage of these ions.

Thus far, we have described the molecular targets and biological pathways targeted by ebselen. A considerable body of research has revealed that ebselen has a surprisingly wide range of beneficial properties, including anti-inflammatory, antioxidant, anti-mutagenic, and many more as shown in Fig. 3. These activities have been demonstrated both in cultured cells and in animal models. In the next section, we will explain the therapeutic uses of ebselen.

Fig. 3 Schematic representation of biological activities displayed by ebselen. Based on the studies reported, ebselen is a bioactive molecule, which exhibits myriads of properties as described in the figure



Ebselen in therapeutics

ROS such as superoxide anion, hydroxyl radicals, and hydrogen peroxide are responsible for lipid peroxidation, protein oxidation, and DNA damage. These effects are deleterious to cells because they accelerate cell senescence and aging, and lead to various age-related diseases, diabetes and its associated disorders, neurologic disorders including stroke, Parkinson's disease, Alzheimer's disease, and diseases associated with cartilage degradation [118–120]. Oxidative stress resulting from ROS production is also one of the main causes of apoptosis [121, 122]. Antioxidant compounds and enzymes that have the potential to scavenge free radicals are of great pharmacological importance. Ebselen appears to be a promising drug because of the multiplicity of its mechanisms for inhibiting free radical-induced injury, coupled with its lack of side effects, good blood–brain permeability, and rapid absorption following oral administration [123]. A summary of pharmacological applications of ebselen are shown in Table 2.

Neuroprotective effects of ebselen

Various studies have demonstrated that ebselen attenuates neuronal cell death induced by ischemia/reperfusion [74, 75]. It has been reported that ebselen affects the morphological structure of neurons, and reduces the degree of oxidative stress [124] through its GPx(glutathione peroxidase)-like and anti-lipoperoxidative activities [125]. Direct interaction of ebselen with the glutamatergic system has been shown in rats [13], suggesting that ebselen reduces glutamate-induced oxidative stress in different regions of the rat brain [126]. The mechanism of action for this effect has not been established, but preliminary studies suggest

that ebselen protects neurons from ischemic damage via control of the expression of γ -aminobutyric acid (GABA) shunt enzymes to supply the tricarboxylic acid cycle (TCA) cycle [24]. Ebselen also significantly inhibits acetylcholinesterase (AChE) activity [127], demonstrating its involvement in the cholinergic system. In rodent models, ebselen has been shown to provide significant protection against ischemic damage in both gray and white matter [8] and the ventroposterior nucleus (VPN) of the brain [128]. Additionally, the administration of ebselen in gerbils significantly reduces neuronal death induced by ischemia/reperfusion in the hippocampal CA1 region [74].

Lithium is the most effective mood-stabilizing drug for the treatment of bipolar disorder [129]. Unfortunately, lithium usage is associated with a high degree of toxicity and has many undesirable side effects at therapeutic doses [130]. Lithium is active in bipolar disorder primarily by inhibiting inositol monophosphatase (IMPase) [131, 132]. Recently, it has been reported that ebselen inhibits IMPase and acts as a lithium mimetic in mouse models of bipolar disorder [133]. It was also found that inhibition of IMPase by ebselen is covalent and irreversible. This offers several advantages, since irreversible inhibition cannot be overcome by accumulation of substrate, prolonging the activity of ebselen and increasing its selectivity for IMPase [133]. Therefore, ebselen represents an alternative to lithium that might serve as a safer treatment for bipolar disorder.

Ebselen as a potential treatment for diabetes-related disorders

The ability of ebselen to mimic glutathione peroxidase (GPx) activity makes it a potential drug for the treatment of diabetes-

Table 2 List showing potential use of ebselen in Pharmacology

| S. No. | Complications | Ebselen mode of action | Reference |
|--------|--|---|------------|
| 1 | Eye problems due to alcohol | Decreases neurotoxicity induced by ethanol | [31] |
| 2 | Brain disorders associated with demyelinating events | Inhibit AChE activity | [122] |
| 4 | Arthritic joints | Blocks cartilage proteoglycan breakdown | [30] |
| 5 | Acute lung injuries(ozone exposure) | Modulates the oxidant-related inflammatory process | [29] |
| 6 | MeHg-induced cytotoxicity | Suppresses oxidative stress | [141, 142] |
| 7 | Nitrogen mustards toxicity | Reduces membrane damage | [148, 149] |
| 8 | Cadmium-induced testicular damage | Reduces the oxidative stress | [143] |
| 9 | Cisplatin-induced nephrotoxicity | Reduces the oxidative stress | [144] |
| 10 | Inflammation-associated carcinogenesis | regulates expression of pro-inflammatory mediators | [7, 74] |
| 11 | Mn-induced toxicity | | [146] |
| 12 | Diabetes-related atherosclerosis and nephropathy | Effectively repletes the lack of GPx1 | [11] |
| 13 | Vascular complications in patients with Diabetes-mellitus (DM) | Scavenges free radicals | [134] |
| 14 | Opportunistic pathogens such as <i>Cryptococcus neoformans</i> and <i>Candida albicans</i> infection | Block the activity of Pma1p | [149, 155] |
| 15 | Anti-malarial activity | Selective inhibition of plasmodia | [154] |
| 16 | Anti-asthma activity | Suppression of oxidant formation and iNOS induction | [32] |
| 18 | Anti-thrombotic agent | Reduces oxidative stress | [12] |
| 19 | Chronic iron overload | Reduces oxidative stress | [140] |
| 20 | Lung oedema | Reduces BALTNF-a and ET-1 levels | [98] |
| 21 | Oxidant stress-associated thrombotic events | Reduces blood cell aggregate formation and vessel occlusion | [12] |

related disorders associated with reduced GPx levels. Ebselen effectively compensates for the lack of GPx and acts as an effective therapeutic molecule for the treatment of diabetes-related atherosclerosis and nephropathy [11]. In diabetic rats, ebselen considerably reduces nitrotyrosine accumulation, preventing tissues from oxidative damage [134]. Furthermore, in insulin-dependent diabetes mellitus (IDDM), high levels of nitric oxide synthase (iNOS) activity result in the accumulation of nitric oxide (NO), which is deleterious to insulin-producing cells. Ebselen antagonizes the effect of NO through scavenging [135]. In a diabetic mouse model, ebselen exhibits beneficial effects on β -cell mass and function [136]. Ebselen also protects from vascular complications in patients with diabetes-mellitus (DM) [137].

Protective role of ebselen against heart disorders

Heart disease (cardiopathy) and cardiovascular diseases are a group of numerous pathological disorders such as heart failure, cardiac arrhythmias; coronary heart disease, hypertension and so forth, in which signaling processes of ROS, RNS and oxidative stress are important features [138, 139]. The novel antioxidant therapies, specifically, use of Gpx1-mimetics holds promise as a targeted antioxidant approach for the treatment of oxidative stress-induced heart diseases. It has been shown that ebselen treatment increases glutathione peroxidase activity in the rat heart [140]. Ebselen is also

known to prevent apoptosis induced due to reperfusion of the ischemic heart by reducing the oxidative stress associated with ischemia/reperfusion through GSH preservation [141, 142]. Ebselen also reduces apoptotic cell death during an open-heart surgery by preventing free radical formation [73]. Another report suggests that the ebselen inhibits the blood cell aggregate formation and vessel occlusion in vivo [12]. It also reduces the neutrophil infiltration into the ischemic myocardium [143]. In stroke-prone spontaneously hypertensive rats (SHRsp), ebselen is shown to protect the endothelium and vascular structure during the chronic process of hypertension [144]. In diabetes mellitus, heart associated complications are common. It has been found that in diabetes patients hyperglycemia triggers increased ROS levels can be attenuated by ebselen, leading to protection of endothelial dysfunction [145]. It has been observed that ebselen reduces nitration and restores voltage-gated potassium channel function in small coronary arteries of diabetic rats [137].

Ebselen is very effective drug against chronic iron overload that is a major cause of cardiac failure throughout the world. Ebselen gives the cardioprotective effects from iron overload in the heart of in B6D2F1 mice by inducing increase in GPx activity level [146]. Ebselen also protects from daunorubicin-induced cardiomyopathy. It has been found that subcutaneous administration of ebselen to daunorubicin-treated rats shows significant improvement in serum cardiac indices including creatine kinase isoenzyme

and lactate dehydrogenase as well as serum glutathione (GSH) peroxidase [147]. Ebselen also protects from adriamycin (Adr)-induced cardiotoxicity by attenuating oxidative stress in heart and liver tissue of rats [148]. Altogether, these studies show that ebselen is very effective drug against the cardiopathy.

Ebselen acts as an effective detoxifying agent

Xenobiotic compounds exhibits toxicity through various mechanisms. Xenobiotics that induce the production of ROS are readily attenuated by ebselen due to its anti-oxidative properties. For example, ebselen is highly effective in reducing metal-induced cytotoxicity, such as toxicity caused by cadmium, iron, methylmercury (MeHg), or cisplatin [146, 149–153]. Treatment with ebselen effectively suppresses oxidative stress and protects cells against MeHg-induced cytotoxicity [154]. Ebselen also inhibits MeHg-induced phosphorylation of extracellular-signal-regulated kinase (ERK) in rat primary astrocyte cultures, leading to a block in activation of caspase-3 and protection from apoptosis [153]. Furthermore, ebselen attenuates cadmium-induced testicular damage in mice by reducing oxidative stress parameters [149]. Ebselen is equally effective against toxicity of cisplatin exposure [150, 155]. It has been proposed that the GPx-like activity of ebselen might prevent chronic iron overload in heart tissue [146]. Treatment of B6D2F1 mice with iron plus ebselen decreases both cytotoxic aldehyde and iron concentrations in heart tissue, compared to iron-only treated mice. Similarly, ebselen has been shown to exhibit promising efficacy in attenuating the neurotoxic effects of manganese (Mn) in an in vivo rat model [156].

Ebselen is effective against several toxic compounds and protects against acute lung injuries caused by ozone exposure through modulation of oxidant-related inflammatory processes [29]. Ebselen has also been shown to protect against the effects of diphenyl ditelluride poisoning [157], nitrogen mustards [158] such as mechlorethamine (HN2), and 2-chloroethyl ethyl sulfide (CEES) [159]. It has been demonstrated that pre-treatment of lymphocytes with ebselen reduces DNA damage caused by alloxan [160] or the antibiotic streptozotocin [161]. Additionally, pre-treatment of cells with the ebselen minimizes the extent of DNA damage induced by amsacrine [162] or oxysterol [163]. All together, these studies suggest that ebselen might be a promising detoxifying agent.

Ebselen possesses anti-microbial properties

Ebselen displays antimicrobial activity against several microorganisms. Ebselen has shown anti-malarial activity in vitro against *Plasmodium berghei* and *P. falciparum*

[164]. The uptake and incorporation of methionine and adenosine in *P. berghei* was inhibited by ebselen, leading to a reduction in infectivity. Furthermore, ebselen is quite effective against chloroquine-resistant *P. falciparum* strains, inhibiting the development of asexual stages of the parasite and blocking the invasion of erythrocytes by merozoites [164]. Recently, through high-throughput screening (HTS) ebselen has been identified as a potent inhibitor of *P. falciparum* hexokinase (PfHK). The results of biochemical assays suggest that the basis for enzyme inactivation by ebselen is not due to covalent association with the inhibitor, as addition of reducing agent is not sufficient to overcome the inhibition [165]. Ebselen is also shown as a potent inhibitor of the plasma membrane H⁺-ATPase Pma1p. Pma1p is required by a variety of pathogens for their survival. Ebselen blocks the activity of Pma1p of pathogenic *Cryptococcus neoformans* and *Candida albicans* as well as of nonpathogenic yeasts such as *S. cerevisiae* [166, 167]. Recently, ebselen has been shown to inhibit *Pseudomonas aeruginosa* by covalently reacting with cysteine residues in diguanylate cyclases [168]. Interestingly, ebselen also shows potent inhibition of *Mycobacterium tuberculosis*. The antigen 85 (Ag85) complex has emerged as an mycobacterial drug target due to its central role in membrane formation. Mass spectrometry reveals that ebselen binds covalently to a cysteine residue in the Ag85C active site to inhibit of Ag85 complex formation [169]. These studies indicate that ebselen is a potent inhibitor of several pathogenic microorganisms, and suggest that this compound might be useful for the treatment of infectious diseases.

Ebselen as a chemotherapeutic agent

Oxidative stress leads to DNA damage, resulting in the accumulation of genetic instability. Selenium compounds including ebselen show potential chemopreventive activity to protect against carcinogenesis [7, 170]. Ebselen has been shown to reduce the expression of pro-inflammatory cytokines in glioma cells, and to overcome the resistance of these cells to TNF- α -induced apoptosis. The cotreatment of glioblastoma cells with ebselen and TNF- α stimulated glioma cells to undergo apoptosis, suggesting a potential role for ebselen in cancer prevention. Furthermore, ebselen abrogates the invasive potential of glioblastoma cells [171]. Some cancers have been shown to resist TNF- α -induced apoptosis through the stimulation of NF- κ B transcriptional activity. Ebselen inhibits TNF- α -mediated NF- κ B activity, and sensitizes tumor cells to undergo apoptosis [172]. Additionally, ebselen ameliorates the progression of microvasculopathy and partially restores angiogenesis [173].

Recently, ebselen has been used in several clinical trials. In one of the clinical trials, ebselen has been administered

in the form of granules suspended in water for oral administration in patients. A placebo-controlled clinical trial in 286 SAH patients has revealed that ebselen might be a prophylactic neuroprotective agent [174]. Another clinical trial with 300 acute ischemia patients has also revealed that ebselen treatment has achieved a significantly better therapeutic outcome if patients are administered with ebselen within 24 h of stroke onset [75]. All together, the outcome of these clinical trials is promising and expected that in future ebselen might be used for the treatment of various diseases.

Future directions

The recent growth in the understanding of free radicals biology is leading to a medical revolution that assures a new age of health and disease management [175]. Free radicals and antioxidants are frequently used terms in current discussions of disease mechanisms [176]. ROS are the most common intermediates generated due to metabolic reactions inside the cell or due to interaction with environmental agents. ROS can damage proteins, lipids, DNA, and RNA, and are associated with a number of cell pathologies. Several synthetic antioxidants are recently being reported to be dangerous to human health. Thus, the hunt for effective, non-toxic compounds with antioxidative activity has been intensifying in recent years. As noted above, ebselen is an organoselenium compound that shows considerable antioxidant activity. This compound has the ability to scavenge free radicals, and may be a valuable supplement for the treatment or prevention of many diseases that arise due to dysfunctional cellular redox mechanisms.

The versatile chemical structure of ebselen enables it to interact with a large number of molecules inside the cell, leading to a variety of biological actions. Advances in microarray and mass spectrometry technologies have yielded large output datasets of potential target proteins and genes for therapeutic interventions. Extensive, genome-wide studies on ebselen have not been reported, however, and should be of significant value for the discovery of novel targets for this compound, and target identification from genome-wide information remains a challenge. Additional analytical techniques such as bioinformatics and/or molecular biology tools will be necessary for target validation. This will significantly assist in our understanding of the biological pathways affected by ebselen. Recently, chimeric compounds based on ebselen have been synthesized. For example, selenpezil, which was developed by the fusion of donepezil and ebselen, was seen to exhibit promising activity against Alzheimer's disease [177]. Additionally, a fusion molecule of heptamethine cyanine conjugated to ebselen was used to measure the redox status of cells by

fluorescence [178]. Similar approaches that combine ebselen with existing drugs may allow for the discovery of new compounds with enhanced medicinal properties. In pharmacology, it is essential to identify molecular mechanisms of drug action in order to understand adverse side effects. To this end, appropriate efforts should be made so that the pharmacological properties of ebselen can be harnessed, subsequently, diseases associated with oxidative stress may be treated with this interesting compound.

Conclusions

Ebselen is a biologically active compound that affects several cellular processes. Studies with model organisms such as yeast and mice, and in vitro experiments with liposomes, microsomes, and cell lines have shown that the protective effects of ebselen against oxidative stress are due largely to its activity as a GPx mimic. Ebselen has both protective as well as harmful effects on biological systems under different conditions. A number of novel ebselen-based pharmaceutical agents are under intense investigation. Continued research on ebselen and its mechanisms of action are needed to develop potent derivatives of the compound for future pharmacological and therapeutic uses. Innovative approaches employing collaborative research and modern technology in combination with the established health principles will be advantageous in near future for improving health.

Acknowledgments We apologize to the authors whose work is not cited due to space limitation. Council of Scientific and Industrial Research (CSIR), Govt. of India is acknowledged for fellowship support to GKA. This work was financially supported by the fund from Council of Scientific and Industrial Research (CSIR), Govt. of India to RST. Members of the Laboratory of Chromatin Biology are acknowledged for helpful discussions throughout the preparation of this manuscript.

References

1. Sakurai T, Kanayama M, Shibata T, Itoh K, Kobayashi A, Yamamoto M, Uchida K (2006) Ebselen, a seleno-organic antioxidant, as an electrophile. *Chem Res Toxicol* 19(9):1196–1204. doi:[10.1021/tx0601105](https://doi.org/10.1021/tx0601105)
2. Ullrich V, Weber P, Meisch F, von Appen F (1996) Ebselen-binding equilibria between plasma and target proteins. *Biochem Pharmacol* 52(1):15–19
3. Zhao R, Masayasu H, Holmgren A (2002) Ebselen: a substrate for human thioredoxin reductase strongly stimulating its hydroperoxide reductase activity and a superfast thioredoxin oxidant. *Proc Natl Acad Sci USA* 99(13):8579–8584. doi:[10.1073/pnas.122061399](https://doi.org/10.1073/pnas.122061399)
4. Antony S, Bayse CA (2011) Modeling the mechanism of the glutathione peroxidase mimic ebselen. *Inorg Chem* 50(23):12075–12084. doi:[10.1021/ic201603v](https://doi.org/10.1021/ic201603v)

5. Sies H, Sharov VS, Klotz LO, Briviba K (1997) Glutathione peroxidase protects against peroxynitrite-mediated oxidations. A new function for selenoproteins as peroxynitrite reductase. *J Biol Chem* 272(44):27812–27817
6. Zhao R, Holmgren A (2004) Ebselen is a dehydroascorbate reductase mimic, facilitating the recycling of ascorbate via mammalian thioredoxin systems. *Antioxid Redox Signal* 6(1):99–104. doi:[10.1089/152308604771978390](https://doi.org/10.1089/152308604771978390)
7. Nakamura Y, Feng Q, Kumagai T, Torikai K, Ohigashi H, Osawa T, Noguchi N, Niki E, Uchida K (2002) Ebselen, a glutathione peroxidase mimetic seleno-organic compound, as a multifunctional antioxidant. Implication for inflammation-associated carcinogenesis. *J Biol Chem* 277(4):2687–2694. doi:[10.1074/jbc.M109641200](https://doi.org/10.1074/jbc.M109641200)
8. Imai H, Masayasu H, Dewar D, Graham DI, Macrae IM (2001) Ebselen protects both gray model of focal and white matter in a rodent cerebral ischemia. *Stroke* 32(9):2149–2156. doi:[10.1161/hs0901.095725](https://doi.org/10.1161/hs0901.095725)
9. Noguchi N, Yoshida Y, Kaneda H, Yamamoto Y, Niki E (1992) Action of ebselen as an antioxidant against lipid peroxidation. *Biochem Pharmacol* 44(1):39–44
10. Schewe T (1995) Molecular actions of ebselen—an antiinflammatory antioxidant. *Gen Pharmacol* 26(6):1153–1169
11. Chew P, Yuen DY, Stefanovic N, Pete J, Coughlan MT, Jandeleit-Dahm KA, Thomas MC, Rosenfeldt F, Cooper ME, de Haan JB (2010) Antiatherosclerotic and renoprotective effects of ebselen in the diabetic apolipoprotein E/GPx1-double knockout mouse. *Diabetes* 59(12):3198–3207. doi:[10.2337/db10-0195](https://doi.org/10.2337/db10-0195)
12. Lindenblatt N, Schareck W, Belusa L, Nickels RM, Menger MD, Vollmar B (2003) Anti-oxidant ebselen delays microvascular thrombus formation in the rat cremaster muscle by inhibiting platelet P-selectin expression. *Thromb Haemost* 90(5):882–892. doi:[10.1267/THRO03050882](https://doi.org/10.1267/THRO03050882)
13. Nogueira CW, Rotta LN, Zeni G, Souza DO, Rocha JB (2002) Exposure to ebselen changes glutamate uptake and release by rat brain synaptosomes. *Neurochem Res* 27(4):283–288
14. Kasraee B, Nikolic DS, Salomon D, Carraux P, Fontao L, Piguet V, Omrani GR, Sorg O, Saurat JH (2012) Ebselen is a new skin depigmenting agent that inhibits melanin biosynthesis and melanosomal transfer. *Exp Dermatol* 21(1):19–24. doi:[10.1111/j.1600-0625.2011.01394.x](https://doi.org/10.1111/j.1600-0625.2011.01394.x)
15. Morin D, Zini R, Ligeret H, Neckameyer W, Labidalle S, Tillement JP (2003) Dual effect of ebselen on mitochondrial permeability transition. *Biochem Pharmacol* 65(10):1643–1651
16. Batna A, Fuchs C, Spittler G (1997) Lipid peroxidation in presence of ebselen. *Chem Phys Lipids* 87(2):149–158
17. Chan PH (1996) Role of oxidants in ischemic brain damage. *Stroke* 27(6):1124–1129
18. Muller A, Cadenas E, Graf P, Sies H (1984) A novel biologically active seleno-organic compound-I. Glutathione peroxidase-like activity in vitro and antioxidant capacity of PZ 51 (Ebselen). *Biochem Pharmacol* 33(20):3235–3239
19. Sies H (1993) Ebselen, a selenoorganic compound as glutathione peroxidase mimic. *Free Radical Biol Med* 14(3):313–323
20. Sies H, de Groot H (1992) Role of reactive oxygen species in cell toxicity. *Toxicol Lett* 64–65:547–551
21. Masumoto H, Sies H (1996) The reaction of ebselen with peroxynitrite. *Chem Res Toxicol* 9(1):262–267. doi:[10.1021/tx950115u](https://doi.org/10.1021/tx950115u)
22. Briviba K, Roussyn I, Sharov VS, Sies H (1996) Attenuation of oxidation and nitration reactions of peroxynitrite by selenomethionine, selenocystine and ebselen. *Biochem J* 319(Pt 1):13–15
23. Masumoto H, Kissner R, Koppenol WH, Sies H (1996) Kinetic study of the reaction of ebselen with peroxynitrite. *FEBS Lett* 398(2–3):179–182
24. Seo JY, Lee CH, Cho JH, Choi JH, Yoo KY, Kim DW, Park OK, Li H, Choi SY, Hwang IK, Won MH (2009) Neuroprotection of ebselen against ischemia/reperfusion injury involves GABA shunt enzymes. *J Neurol Sci* 285(1–2):88–94. doi:[10.1016/j.jns.2009.05.029](https://doi.org/10.1016/j.jns.2009.05.029)
25. Gabryel B, Malecki A (2006) Ebselen attenuates oxidative stress in ischemic astrocytes depleted of glutathione. Comparison with glutathione precursors. *Pharmacol Rep* 58(3):381–392
26. Hamacher J, Stammberger U, Weber E, Lucas R, Wendel A (2009) Ebselen improves ischemia-reperfusion injury after rat lung transplantation. *Lung* 187(2):98–103. doi:[10.1007/s00408-009-9134-x](https://doi.org/10.1007/s00408-009-9134-x)
27. Pawlas N, Malecki A (2007) Effects of ebselen on glutathione level in neurons exposed to arachidonic acid and 4-hydroxynonenal during simulated ischemia in vitro. *Pharmacol Rep* 59(6):708–714
28. Namura S, Nagata I, Takami S, Masayasu H, Kikuchi H (2001) Ebselen reduces cytochrome c release from mitochondria and subsequent DNA fragmentation after transient focal cerebral ischemia in mice. *Stroke* 32(8):1906–1911
29. Ishii Y, Hashimoto K, Hirano K, Morishima Y, Mochizuki M, Masuyama K, Nomura A, Sakamoto T, Uchida Y, Sagai M, Sekizawa K (2000) Ebselen decreases ozone-induced pulmonary inflammation in rats. *Lung* 178(4):225–234
30. Pratta MA, Ackerman NR, Arner EC (1998) Effect of ebselen on IL-1-induced alterations in cartilage metabolism. *Inflamm Res* 47(3):115–121
31. Johnsen-Soriano S, Genoves JM, Romero B, Garcia-Delpech S, Muriach M, Sancho-Tello M, Bosch-Morell F, Romero FJ (2007) Chronic ethanol feeding induces oxidative stress in the rat retina: treatment with the antioxidant ebselen. *Arch Soc Esp Oftalmol* 82(12):757–762
32. Zhang M, Nomura A, Uchida Y, Iijima H, Sakamoto T, Ishii Y, Morishima Y, Mochizuki M, Masuyama K, Hirano K, Sekizawa K (2002) Ebselen suppresses late airway responses and airway inflammation in guinea pigs. *Free Radical Biol Med* 32(5):454–464
33. Gogvadze V, Klein SD, Shigenaga M, Ames BN, Richter C (2000) Effect of ebselen on Ca^{2+} transport in mitochondria. *Redox Rep* 5(6):359–363. doi:[10.1179/135100000101535924](https://doi.org/10.1179/135100000101535924)
34. Yang CF, Shen HM, Ong CN (2000) Intracellular thiol depletion causes mitochondrial permeability transition in ebselen-induced apoptosis. *Arch Biochem Biophys* 380(2):319–330. doi:[10.1006/abbi.2000.1939](https://doi.org/10.1006/abbi.2000.1939)
35. Guerin PJ, Gauthier ER (2003) Induction of cellular necrosis by the glutathione peroxidase mimetic ebselen. *J Cell Biochem* 89(1):203–211. doi:[10.1002/jcb.10500](https://doi.org/10.1002/jcb.10500)
36. Miorelli ST, Rosa RM, Moura DJ, Rocha JC, Lobo LA, Henriques JA, Saffi J (2008) Antioxidant and anti-mutagenic effects of ebselen in yeast and in cultured mammalian V79 cells. *Mutagenesis* 23(2):93–99. doi:[10.1093/mutage/gem048](https://doi.org/10.1093/mutage/gem048)
37. Yang CF, Shen HM, Ong CN (2000) Ebselen induces apoptosis in HepG(2) cells through rapid depletion of intracellular thiols. *Arch Biochem Biophys* 374(2):142–152. doi:[10.1006/abbi.1999.1574](https://doi.org/10.1006/abbi.1999.1574)
38. Azad GK, Balkrishna SJ, Sathish N, Kumar S, Tomar RS (2012) Multifunctional Ebselen drug functions through the activation of DNA damage response and alterations in nuclear proteins. *Biochem Pharmacol* 83(2):296–303. doi:[10.1016/j.bcp.2011.10.011](https://doi.org/10.1016/j.bcp.2011.10.011)
39. Larabee JL, Hocker JR, Hanas JS (2009) Mechanisms of inhibition of zinc-finger transcription factors by selenium compounds ebselen and selenite. *J Inorg Biochem* 103(3):419–426. doi:[10.1016/j.jinorgbio.2008.12.007](https://doi.org/10.1016/j.jinorgbio.2008.12.007)
40. Jacob C, Maret W, Vallee BL (1998) Ebselen, a selenium-containing redox drug, releases zinc from metallothionein.

- Biochem Biophys Res Commun 248(3):569–573. doi:[10.1006/bbrc.1998.9026](https://doi.org/10.1006/bbrc.1998.9026)
41. Wagner G, Schuch G, Akerboom TP, Sies H (1994) Transport of ebselen in plasma and its transfer to binding sites in the hepatocyte. *Biochem Pharmacol* 48(6):1137–1144
 42. Christison J, Sies H, Stocker R (1994) Human blood cells support the reduction of low-density-lipoprotein-associated cholesterol ester hydroperoxides by albumin-bound ebselen. *Biochem J* 304(Pt 2):341–345
 43. Aitken JB, Lay PA, Duong TT, Aran R, Witting PK, Harris HH, Lai B, Vogt S, Giles GI (2012) Synchrotron radiation induced X-ray emission studies of the antioxidant mechanism of the organoselenium drug ebselen. *J Biol Inorg Chem* 17(4):589–598. doi:[10.1007/s00775-012-0879-y](https://doi.org/10.1007/s00775-012-0879-y)
 44. Boireau A, Marechal PM, Meunier M, Dubedat P, Moussaoui S (2000) The anti-oxidant ebselen antagonizes the release of the apoptogenic factor cytochrome c induced by Fe^{2+} /citrate in rat liver mitochondria. *Neurosci Lett* 289(2):95–98
 45. Tiano L, Fedeli D, Santoni G, Davies I, Wakabayashi T, Falconi G (2003) Ebselen prevents mitochondrial ageing due to oxidative stress: in vitro study of fish erythrocytes. *Mitochondrion* 2(6):428–436. doi:[10.1016/S1567-7249\(03\)00032-1](https://doi.org/10.1016/S1567-7249(03)00032-1)
 46. Moussaoui S, Obinu MC, Daniel N, Reibaud M, Blanchard V, Imperato A (2000) The antioxidant ebselen prevents neurotoxicity and clinical symptoms in a primate model of Parkinson's disease. *Exp Neurol* 166(2):235–245. doi:[10.1006/exnr.2000.7516](https://doi.org/10.1006/exnr.2000.7516)
 47. Nagi MN, Laguna JC, Cook L, Cinti DL (1989) Disruption of rat hepatic microsomal electron transport chains by the selenium-containing anti-inflammatory agent Ebselen. *Arch Biochem Biophys* 269(1):264–271
 48. Narayanaswami V, Sies H (1990) Oxidative damage to mitochondria and protection by ebselen and other antioxidants. *Biochem Pharmacol* 40(7):1623–1629
 49. Santofimia-Castano P, Salido GM, Gonzalez A (2013) Ebselen alters mitochondrial physiology and reduces viability of rat hippocampal astrocytes. *DNA Cell Biol* 32(4):147–155. doi:[10.1089/dna.2012.1939](https://doi.org/10.1089/dna.2012.1939)
 50. Xu KH, Zhang Y, Tang B, Laskin J, Roach PJ, Chen H (2010) Study of highly selective and efficient thiol derivatization using selenium reagents by mass spectrometry. *Anal Chem* 82(16):6926–6932. doi:[10.1021/AC1011602](https://doi.org/10.1021/AC1011602)
 51. Schewe C, Schewe T, Wendel A (1994) Strong inhibition of mammalian lipoxygenases by the antiinflammatory seleno-organic compound ebselen in the absence of glutathione. *Biochem Pharmacol* 48(1):65–74
 52. Walther M, Holzthutter HG, Kuban RJ, Wiesner R, Rathmann J, Kuhn H (1999) The inhibition of mammalian 15-lipoxygenases by the anti-inflammatory drug ebselen: dual-type mechanism involving covalent linkage and alteration of the iron ligand sphere. *Mol Pharmacol* 56(1):196–203
 53. Hattori R, Yui Y, Shinoda E, Inoue R, Aoyama T, Masayasu H, Kawai C, Sasayama S (1996) Effect of ebselen on bovine and rat nitric oxide synthase activity is modified by thiols. *Jpn J Pharmacol* 72(2):191–193
 54. Smith SM, Min J, Ganesh T, Diebold B, Kawahara T, Zhu Y, McCoy J, Sun A, Snyder JP, Fu H, Du Y, Lewis I, Lambeth JD (2012) Ebselen and congeners inhibit NADPH oxidase 2-dependent superoxide generation by interrupting the binding of regulatory subunits. *Chem Biol* 19(6):752–763. doi:[10.1016/j.chembiol.2012.04.015](https://doi.org/10.1016/j.chembiol.2012.04.015)
 55. Tabuchi Y, Ogasawara T, Furuhashi K (1994) Mechanism of the inhibition of hog gastric H⁺, K⁺-ATPase by the seleno-organic compound ebselen. *Arzneimittelforschung* 44(1):51–54
 56. Mishra B, Priyadarsini KI, Mohan H, Mughesh G (2006) Horseradish peroxidase inhibition and antioxidant activity of ebselen and related organoselenium compounds. *Bioorg Med Chem Lett* 16(20):5334–5338. doi:[10.1016/j.bmcl.2006.07.085](https://doi.org/10.1016/j.bmcl.2006.07.085)
 57. Azad GK, Singh V, Mandal P, Singh P, Golla U, Baranwal S, Chauhan S, Tomar RS (2014) Ebselen induces reactive oxygen species (ROS)-mediated cytotoxicity in *Saccharomyces cerevisiae* with inhibition of glutamate dehydrogenase being a target. *FEBS Open Bio* 4:77–89. doi:[10.1016/j.fob.2014.01.002](https://doi.org/10.1016/j.fob.2014.01.002)
 58. Lugokenski TH, Muller LG, Taube PS, Rocha JB, Pereira ME (2011) Inhibitory effect of ebselen on lactate dehydrogenase activity from mammals: a comparative study with diphenyl diselenide and diphenyl ditelluride. *Drug Chem Toxicol* 34(1):66–76. doi:[10.3109/01480541003782294](https://doi.org/10.3109/01480541003782294)
 59. Borges VC, Rocha JB, Nogueira CW (2005) Effect of diphenyl diselenide, diphenyl ditelluride and ebselen on cerebral Na⁺, K⁺-ATPase activity in rats. *Toxicology* 215(3):191–197. doi:[10.1016/j.tox.2005.07.002](https://doi.org/10.1016/j.tox.2005.07.002)
 60. Terentis AC, Freewan M, Sempertegui Plaza TS, Raftery MJ, Stocker R, Thomas SR (2010) The selenazal drug ebselen potently inhibits indoleamine 2,3-dioxygenase by targeting enzyme cysteine residues. *Biochemistry* 49(3):591–600. doi:[10.1021/bi901546e](https://doi.org/10.1021/bi901546e)
 61. Xia R, Ganther HE, Egge A, Abramson JJ (2004) Selenium compounds modulate the calcium release channel/ryanodine receptor of rabbit skeletal muscle by oxidizing functional thiols. *Biochem Pharmacol* 67(11):2071–2079. doi:[10.1016/j.bcp.2004.02.018](https://doi.org/10.1016/j.bcp.2004.02.018)
 62. Klug A (2010) The discovery of zinc fingers and their applications in gene regulation and genome manipulation. *Annu Rev Biochem* 79:213–231. doi:[10.1146/annurev-biochem-010909-095056](https://doi.org/10.1146/annurev-biochem-010909-095056)
 63. Laity JH, Lee BM, Wright PE (2001) Zinc finger proteins: new insights into structural and functional diversity. *Curr Opin Struct Biol* 11(1):39–46. doi:[10.1016/S0959-440x\(00\)00167-6](https://doi.org/10.1016/S0959-440x(00)00167-6)
 64. Antony S, Bayse CA (2013) Density functional theory study of the attack of ebselen on a zinc-finger model. *Inorg Chem*. doi:[10.1021/ic401429z](https://doi.org/10.1021/ic401429z)
 65. Sekirnik R, Rose NR, Thalhammer A, Seden PT, Mecinovic J, Schofield CJ (2009) Inhibition of the histone lysine demethylase JMJD2A by ejection of structural Zn(II). *Chem Commun (Camb)* 42:6376–6378. doi:[10.1039/b916357c](https://doi.org/10.1039/b916357c)
 66. Blessing H, Kraus S, Heindl P, Bal W, Hartwig A (2004) Interaction of selenium compounds with zinc finger proteins involved in DNA repair. *Eur J Biochem/FEBS* 271(15):3190–3199. doi:[10.1111/j.1432-1033.2004.04251.x](https://doi.org/10.1111/j.1432-1033.2004.04251.x)
 67. Wood-Allum CA, Barber SC, Kirby J, Heath P, Holden H, Mead R, Higginbottom A, Allen S, Beaujeux T, Alexson SE, Ince PG, Shaw PJ (2006) Impairment of mitochondrial anti-oxidant defence in SOD1-related motor neuron injury and amelioration by ebselen. *Brain* 129(Pt 7):1693–1709. doi:[10.1093/brain/awl118](https://doi.org/10.1093/brain/awl118)
 68. Kim SJ, Park C, Han AL, Youn MJ, Lee JH, Kim Y, Kim ES, Kim HJ, Kim JK, Lee HK, Chung SY, So H, Park R (2009) Ebselen attenuates cisplatin-induced ROS generation through Nrf2 activation in auditory cells. *Hear Res* 251(1–2):70–82. doi:[10.1016/j.heares.2009.03.003](https://doi.org/10.1016/j.heares.2009.03.003)
 69. Tamasi V, Jeffries JM, Arteel GE, Falkner KC (2004) Ebselen augments its peroxidase activity by inducing nrf-2-dependent transcription. *Arch Biochem Biophys* 431(2):161–168. doi:[10.1016/j.abb.2004.07.030](https://doi.org/10.1016/j.abb.2004.07.030)
 70. Meier P, Finch A, Evan G (2000) Apoptosis in development. *Nature* 407(6805):796–801. doi:[10.1038/35037734](https://doi.org/10.1038/35037734)
 71. Milleron RS, Bratton SB (2006) Heat shock induces apoptosis independently of any known initiator caspase-activating complex. *J Biol Chem* 281(25):16991–17000. doi:[10.1074/jbc.M512754200](https://doi.org/10.1074/jbc.M512754200)
 72. Verheij M, Bartelink H (2000) Radiation-induced apoptosis. *Cell Tissue Res* 301(1):133–142

73. Maulik N, Yoshida T (2000) Oxidative stress developed during open heart surgery induces apoptosis: reduction of apoptotic cell death by ebselen, a glutathione peroxidase mimic. *J Cardiovasc Pharmacol* 36(5):601–608
74. Koizumi H, Fujisawa H, Suehiro E, Shirao S, Suzuki M (2011) Neuroprotective effects of ebselen following forebrain ischemia: involvement of glutamate and nitric oxide. *Neurol Med Chir* 51(5):337–343
75. Yamaguchi T, Sano K, Takakura K, Saito I, Shinohara Y, Asano T, Yasuhara H, Grp ES (1998) Ebselen in acute ischemic stroke—a placebo-controlled, double-blind clinical trial. *Stroke* 29(1):12–17
76. Parnham M, Sies H (2000) Ebselen: prospective therapy for cerebral ischaemia. *Expert Opin Investig Drugs* 9(3):607–619. doi:[10.1517/13543784.9.3.607](https://doi.org/10.1517/13543784.9.3.607)
77. Holl V, Coelho D, Silbernagel L, Keyser JF, Waltzinger C, Dufour P, Bischoff PL (2000) Prevention of nitrogen mustard-induced apoptosis in normal and transformed lymphocytes by ebselen. *Biochem Pharmacol* 60(11):1565–1577
78. Tak JK, Park JW (2009) The use of ebselen for radioprotection in cultured cells and mice. *Free Radical Biol Med* 46(8):1177–1185. doi:[10.1016/j.freeradbiomed.2009.01.023](https://doi.org/10.1016/j.freeradbiomed.2009.01.023)
79. Ramakrishnan N, Kalinich JF, McClain DE (1996) Ebselen inhibition of apoptosis by reduction of peroxides. *Biochem Pharmacol* 51(11):1443–1451
80. Yoshizumi M, Kogame T, Suzuki Y, Fujita Y, Kyaw M, Kirima K, Ishizawa K, Tsuchiya K, Kagami S, Tamaki T (2002) Ebselen attenuates oxidative stress-induced apoptosis via the inhibition of the c-Jun N-terminal kinase and activator protein-1 signalling pathway in PC12 cells. *Br J Pharmacol* 136(7):1023–1032. doi:[10.1038/sj.bjp.0704808](https://doi.org/10.1038/sj.bjp.0704808)
81. Ali N, Yoshizumi M, Tsuchiya K, Kyaw M, Fujita Y, Izawa Y, Abe S, Kanematsu Y, Kagami S, Tamaki T (2004) Ebselen inhibits p38 mitogen-activated protein kinase-mediated endothelial cell death by hydrogen peroxide. *Eur J Pharmacol* 485(1–3):127–135
82. Sarker KP, Biswas KK, Rosales JL, Yamaji K, Hashiguchi T, Lee KY, Maruyama I (2003) Ebselen inhibits NO-induced apoptosis of differentiated PC12 cells via inhibition of ASK1-p38 MAPK-p53 and JNK signaling and activation of p44/42 MAPK and Bcl-2. *J Neurochem* 87(6):1345–1353
83. Xu JH, Hu HT, Liu Y, Qian YH, Liu ZH, Tan QR, Zhang ZJ (2006) Neuroprotective effects of ebselen are associated with the regulation of Bcl-2 and Bax proteins in cultured mouse cortical neurons. *Neurosci Lett* 399(3):210–214. doi:[10.1016/j.neulet.2006.02.024](https://doi.org/10.1016/j.neulet.2006.02.024)
84. Yoshizumi M, Fujita Y, Izawa Y, Suzuki Y, Kyaw M, Ali N, Tsuchiya K, Kagami S, Yano S, Sone S, Tamaki T (2004) Ebselen inhibits tumor necrosis factor- α -induced c-Jun N-terminal kinase activation and adhesion molecule expression in endothelial cells. *Exp Cell Res* 292(1):1–10
85. Cooke MS, Evans MD, Dizdaroglu M, Lunec J (2003) Oxidative DNA damage: mechanisms, mutation, and disease. *FASEB J* 17(10):1195–1214. doi:[10.1096/fj.02-0752rev](https://doi.org/10.1096/fj.02-0752rev)
86. Wiseman H, Halliwell B (1996) Damage to DNA by reactive oxygen and nitrogen species: role in inflammatory disease and progression to cancer. *Biochem J* 313(Pt 1):17–29
87. Yang CF, Shen HM, Ong CN (1999) Protective effect of ebselen against hydrogen peroxide-induced cytotoxicity and DNA damage in HepG2 cells. *Biochem Pharmacol* 57(3):273–279
88. Li J, Chen JJ, Zhang F, Zhang C (2000) Ebselen protection against hydrogen peroxide-induced cytotoxicity and DNA damage in HL-60 cells. *Acta Pharmacol Sin* 21(5):455–459
89. Roussyn I, Briviba K, Masumoto H, Sies H (1996) Selenium-containing compounds protect DNA from single-strand breaks caused by peroxynitrite. *Arch Biochem Biophys* 330(1):216–218. doi:[10.1006/abbi.1996.0245](https://doi.org/10.1006/abbi.1996.0245)
90. Li Y, Cao Z (2002) The neuroprotectant ebselen inhibits oxidative DNA damage induced by dopamine in the presence of copper ions. *Neurosci Lett* 330(1):69–73
91. Tripathi DN, Jena GB (2008) Ebselen attenuates cyclophosphamide-induced oxidative stress and DNA damage in mice. *Free Radical Res* 42(11–12):966–977. doi:[10.1080/10715760802566558](https://doi.org/10.1080/10715760802566558)
92. Yang CF, Liu J, Wasser S, Shen HM, Tan CE, Ong CN (2000) Inhibition of ebselen on aflatoxin B(1)-induced hepatocarcinogenesis in Fischer 344 rats. *Carcinogenesis* 21(12):2237–2243
93. Balkrishna SJ, Kumar S, Azad GK, Bhakuni BS, Panini P, Ahalawat N, Tomar RS, Dettly MR (2014) An ebselen like catalyst with enhanced GPx activity via a selenol intermediate. *Org Biomol Chem* 12(8):1215–1219. doi:[10.1039/c4ob00027g](https://doi.org/10.1039/c4ob00027g)
94. Zhao XL, Chabes A, Domkin V, Thelander L, Rothstein R (2001) The ribonucleotide reductase inhibitor Sml1 is a new target of the Mec1/Rad53 kinase cascade during growth and in response to DNA damage. *EMBO J* 20(13):3544–3553. doi:[10.1093/emboj/20.13.3544](https://doi.org/10.1093/emboj/20.13.3544)
95. Inglot AD, Mlochowski J, Zielinska-Jencyzyk J, Piasecki E, Ledwon TK, Kloc K (1996) Seleno-organic compounds as immunostimulants: an approach to the structure-activity relationship. *Arch Immunol Therap Exp* 44(1):67–75
96. Tiegs G, Kusters S, Kunstle G, Hentze H, Kiemer AK, Wendel A (1998) Ebselen protects mice against T cell-dependent, TNF-mediated apoptotic liver injury. *J Pharmacol Exp Therap* 287(3):1098–1104
97. Gladilin S, Bidmon HJ, Divanach A, Arteel GE, Witte OW, Zilles K, Sies H (2000) Ebselen lowers plasma interleukin-6 levels and glial heme oxygenase-1 expression after focal photothrombotic brain ischemia. *Arch Biochem Biophys* 380(2):237–242. doi:[10.1006/abbi.2000.1943](https://doi.org/10.1006/abbi.2000.1943)
98. Matsushita T, Fukuda K, Yamamoto H, Yamazaki K, Tomiyama T, Oh M, Hamanishi C (2004) Effect of ebselen, a scavenger of reactive oxygen species, on chondrocyte metabolism. *Mod Rheumatol* 14(1):25–30. doi:[10.1007/s10165-003-0261-6](https://doi.org/10.1007/s10165-003-0261-6)
99. Jean Harry G, Bruccoleri A, Lefebvre d'Hellencourt C (2003) Differential modulation of hippocampal chemical-induced injury response by ebselen, pentoxifylline, and TNF α -, IL-1 α -, and IL-6-neutralizing antibodies. *J Neurosci Res* 73(4):526–536. doi:[10.1002/jnr.10653](https://doi.org/10.1002/jnr.10653)
100. Shimohashi N, Nakamuta M, Uchimura K, Sugimoto R, Iwamoto H, Enjoji M, Nawata H (2000) Selenoorganic compound, ebselen, inhibits nitric oxide and tumor necrosis factor- α production by the modulation of jun-N-terminal kinase and the NF-kappaB signaling pathway in rat Kupffer cells. *J Cell Biochem* 78(4):595–606
101. Koyanagi T, Nakamuta M, Enjoji M, Iwamoto H, Motomura K, Sakai H, Nawata H (2001) The selenoorganic compound ebselen suppresses liver injury induced by *Propionibacterium acnes* and lipopolysaccharide in rats. *Int J Mol Med* 7(3):321–327
102. Zhao H, Lu HG, Shi YB, Zhao LM, Bai C, Wang X (2006) Role of enteral nutrition supplemented with ebselen and EHEC in pancreatitis-associated multiple organ dysfunction in rats. *Inflamm Res* 55(10):423–429. doi:[10.1007/s00011-006-6008-z](https://doi.org/10.1007/s00011-006-6008-z)
103. Belvisi MG, Haddad EB, Battram C, Birrell M, Foster M, Webber S (2000) Anti-inflammatory properties of ebselen in a model of sephadex-induced lung inflammation. *Eur Respir J* 15(3):579–581
104. Issekutz AC, Lopes N (1992) Effect of Ebselen on polymorphonuclear leukocyte adhesion to and migration through cytokine-activated vascular endothelium. *Int J Immunopharmacol* 14(8):1383–1390

105. Bosch-Morell F, Roma J, Puertas FJ, Marin N, Diaz-Llopis M, Romero FJ (1999) Efficacy of the antioxidant ebselen in experimental uveitis. *Free Radical Biol Med* 27(3–4):388–391
106. Chen J, Li H, Addabbo F, Zhang F, Pelger E, Patschan D, Park HC, Kuo MC, Ni J, Gobe G, Chander PN, Nasjletti A, Goligorsky MS (2009) Adoptive transfer of syngeneic bone marrow-derived cells in mice with obesity-induced diabetes: selenoorganic antioxidant ebselen restores stem cell competence. *Am J Pathol* 174(2):701–711. doi:[10.2353/ajpath.2009.080606](https://doi.org/10.2353/ajpath.2009.080606)
107. Buckett PD, Wessling-Resnick M (2009) Small molecule inhibitors of divalent metal transporter-1. *Am J Physiol Gastrointest Liver Physiol* 296(4):G798–G804. doi:[10.1152/ajpgi.90342.2008](https://doi.org/10.1152/ajpgi.90342.2008)
108. Soe-Lin S, Apte SS, Mikhael MR, Kayembe LK, Nie G, Ponka P (2010) Both Nramp1 and DMT1 are necessary for efficient macrophage iron recycling. *Exp Hematol* 38(8):609–617. doi:[10.1016/j.exphem.2010.04.003](https://doi.org/10.1016/j.exphem.2010.04.003)
109. Wetli HA, Buckett PD, Wessling-Resnick M (2006) Small-molecule screening identifies the selanazol drug ebselen as a potent inhibitor of DMT1-mediated iron uptake. *Chem Biol* 13(9):965–972. doi:[10.1016/j.chembiol.2006.08.005](https://doi.org/10.1016/j.chembiol.2006.08.005)
110. Xie L, Zheng W, Xin N, Xie JW, Wang T, Wang ZY (2012) Ebselen inhibits iron-induced tau phosphorylation by attenuating DMT1 up-regulation and cellular iron uptake. *Neurochem Int*. doi:[10.1016/j.neuint.2012.05.016](https://doi.org/10.1016/j.neuint.2012.05.016)
111. Erikson KM, Aschner M (2006) Increased manganese uptake by primary astrocyte cultures with altered iron status is mediated primarily by divalent metal transporter. *Neurotoxicology* 27(1):125–130. doi:[10.1016/j.neuro.2005.07.003](https://doi.org/10.1016/j.neuro.2005.07.003)
112. Gunshin H, Mackenzie B, Berger UV, Gunshin Y, Romero MF, Boron WF, Nussberger S, Gollan JL, Hediger MA (1997) Cloning and characterization of a mammalian proton-coupled metal-ion transporter. *Nature* 388(6641):482–488. doi:[10.1038/41343](https://doi.org/10.1038/41343)
113. Xie L, Zheng W, Xin N, Xie JW, Wang T, Wang ZY (2012) Ebselen inhibits iron-induced tau phosphorylation by attenuating DMT1 up-regulation and cellular iron uptake. *Neurochem Int* 61(3):334–340. doi:[10.1016/j.neuint.2012.05.016](https://doi.org/10.1016/j.neuint.2012.05.016)
114. Brune B, Diewald B, Ullrich V (1991) Ebselen affects calcium homeostasis in human platelets. *Biochem Pharmacol* 41(12):1805–1811
115. Dimmeler S, Brune B, Ullrich V (1991) Ebselen prevents inositol (1,4,5)-trisphosphate binding to its receptor. *Biochem Pharmacol* 42(5):1151–1153
116. Salazar M, Pariente JA, Salido GM, Gonzalez A (2008) Ebselen increases cytosolic free Ca^{2+} concentration, stimulates glutamate release and increases GFAP content in rat hippocampal astrocytes. *Toxicology* 244(2–3):280–291. doi:[10.1016/j.tox.2007.12.002](https://doi.org/10.1016/j.tox.2007.12.002)
117. Cotgreave IA, Duddy SK, Kass GE, Thompson D, Moldeus P (1989) Studies on the anti-inflammatory activity of ebselen. Ebselen interferes with granulocyte oxidative burst by dual inhibition of NADPH oxidase and protein kinase C? *Biochem Pharmacol* 38(4):649–656
118. Halliwell B (1992) Reactive oxygen species and the central nervous system. *J Neurochem* 59(5):1609–1623
119. Coyle JT, Puttfarcken P (1993) Oxidative stress, glutamate, and neurodegenerative disorders. *Science* 262(5134):689–695
120. Mylonas C, Kouretas D (1999) Lipid peroxidation and tissue damage. *In Vivo* 13(3):295–309
121. Whittemore ER, Loo DT, Cotman CW (1994) Exposure to hydrogen peroxide induces cell death via apoptosis in cultured rat cortical neurons. *NeuroReport* 5(12):1485–1488
122. Greenlund LJ, Deckwerth TL, Johnson EM Jr (1995) Superoxide dismutase delays neuronal apoptosis: a role for reactive oxygen species in programmed neuronal death. *Neuron* 14(2):303–315
123. Takasago T, Peters EE, Graham DI, Masayasu H, Macrae IM (1997) Neuroprotective efficacy of ebselen, an anti-oxidant with anti-inflammatory actions, in a rodent model of permanent middle cerebral artery occlusion. *Br J Pharmacol* 122(6):1251–1256. doi:[10.1038/sj.bjp.0701426](https://doi.org/10.1038/sj.bjp.0701426)
124. Kalayci M, Coskun O, Cagavi F, Kanter M, Armutcu F, Gul S, Acikgoz B (2005) Neuroprotective effects of ebselen on experimental spinal cord injury in rats. *Neurochem Res* 30(3):403–410
125. Porciuncula LO, Rocha JB, Boeck CR, Vendite D, Souza DO (2001) Ebselen prevents excitotoxicity provoked by glutamate in rat cerebellar granule neurons. *Neurosci Lett* 299(3):217–220
126. Dalla Corte CL, Bastos LL, Dobrachinski F, Rocha JB, Soares FA (2012) The combination of organoselenium compounds and guanosine prevents glutamate-induced oxidative stress in different regions of rat brains. *Brain Res* 1430:101–111. doi:[10.1016/j.brainres.2011.10.049](https://doi.org/10.1016/j.brainres.2011.10.049)
127. Mazzanti CM, Spanevello R, Ahmed M, Pereira LB, Goncalves JF, Correa M, Schmatz R, Stefanello N, Leal DB, Mazzanti A, Ramos AT, Martins TB, Danesi CC, Graca DL, Morsch VM, Schetinger MR (2009) Pre-treatment with ebselen and vitamin E modulate acetylcholinesterase activity: interaction with demyelinating agents. *Int J Dev Neurosci* 27(1):73–80. doi:[10.1016/j.ijdevneu.2008.09.005](https://doi.org/10.1016/j.ijdevneu.2008.09.005)
128. He M, Xing S, Yang B, Zhao L, Hua H, Liang Z, Zhou W, Zeng J, Pei Z (2007) Ebselen attenuates oxidative DNA damage and enhances its repair activity in the thalamus after focal cortical infarction in hypertensive rats. *Brain Res* 1181:83–92. doi:[10.1016/j.brainres.2007.08.072](https://doi.org/10.1016/j.brainres.2007.08.072)
129. Geddes JR, Goodwin GM, Rendell J, Azorin JM, Cipriani A, Ostacher MJ, Morriss R, Alder N, Juszczak E (2010) Lithium plus valproate combination therapy versus monotherapy for relapse prevention in bipolar I disorder (BALANCE): a randomised open-label trial. *Lancet* 375(9712):385–395. doi:[10.1016/S0140-6736\(09\)61828-6](https://doi.org/10.1016/S0140-6736(09)61828-6)
130. McKnight RF, Adida M, Budge K, Stockton S, Goodwin GM, Geddes JR (2012) Lithium toxicity profile: a systematic review and meta-analysis. *Lancet* 379(9817):721–728. doi:[10.1016/S0140-6736\(11\)61516-X](https://doi.org/10.1016/S0140-6736(11)61516-X)
131. Berridge MJ, Downes CP, Hanley MR (1989) Neural and developmental actions of lithium—a unifying hypothesis. *Cell* 59(3):411–419. doi:[10.1016/0092-8674\(89\)90026-3](https://doi.org/10.1016/0092-8674(89)90026-3)
132. Belmaker RH, Bersudsky Y, Agam G, Levine J, Kofman O (1996) How does lithium work on manic depression? Clinical and psychological correlates of the inositol theory. *Annu Rev Med* 47:47–56
133. Singh N, Halliday AC, Thomas JM, Kuznetsova OV, Baldwin R, Woon ECY, Aley PK, Antoniadou I, Sharp T, Vasudevan SR, Churchill GC (2013) A safe lithium mimetic for bipolar disorder. *Nat Commun* 4:1332. doi:[10.1038/Ncomms2320](https://doi.org/10.1038/Ncomms2320)
134. Hoshida S, Aoki K, Nishida M, Yamashita N, Igarashi J, Hori M, Kuzuya T, Tada M (1997) Effects of preconditioning with ebselen on glutathione metabolism and stress protein expression. *J Pharmacol Exp Therap* 281(3):1471–1475
135. deMello MAR, Flodstrom M, Eizirik DL (1996) Ebselen and cytokine-induced nitric oxide synthase expression in insulin-producing cells. *Biochem Pharmacol* 52(11):1703–1709. doi:[10.1016/S0006-2952\(96\)00520-5](https://doi.org/10.1016/S0006-2952(96)00520-5)
136. Mahadevan J, Parazzoli S, Oseid E, Hertzog AV, Bernlohr DA, Vallerie SN, Liu CQ, Lopez M, Harmon JS, Robertson RP (2013) Ebselen treatment prevents islet apoptosis, maintains intranuclear Pdx-1 and MafA levels, and preserves beta-cell mass and function in ZDF rats. *Diabetes* 62(10):3582–3588. doi:[10.2337/Db13-0357](https://doi.org/10.2337/Db13-0357)
137. Bubolz AH, Wu Q, Larsen BT, Gutterman DD, Liu Y (2007) Ebselen reduces nitration and restores voltage-gated potassium

- channel function in small coronary arteries of diabetic rats. *Am J Physiol Heart Circ Physiol* 293(4):H2231–H2237. doi:[10.1152/ajpheart.00717.2007](https://doi.org/10.1152/ajpheart.00717.2007)
138. Afanas'ev I (2011) ROS and RNS signaling in heart disorders: could antioxidant treatment be successful? *Oxid Med Cell Longev* 2011:293769. doi:[10.1155/2011/293769](https://doi.org/10.1155/2011/293769)
 139. Sugamura K, Keaney JF Jr (2011) Reactive oxygen species in cardiovascular disease. *Free Radical Biol Med* 51(5):978–992. doi:[10.1016/j.freeradbiomed.2011.05.004](https://doi.org/10.1016/j.freeradbiomed.2011.05.004)
 140. Hermenegildo C, Nies E, Monsalve E, Puertas FJ, Higuera V, Romero FJ (1990) Some aspects of cardiac antioxidant defense—ebselen (Pz51) treatment increases glutathione-peroxidase activity in the rat-heart. *Biochem Soc Trans* 18(6):1193–1194
 141. Baljinnayam E, Hasebe N, Morihira M, Sumitomo K, Matsusaka T, Fujino T, Fukuzawa J, Ushikubi F, Kikuchi K (2006) Oral pretreatment with ebselen enhances heat shock protein 72 expression and reduces myocardial infarct size. *Hypertens Res* 29(11):905–913. doi:[10.1291/hyres.29.905](https://doi.org/10.1291/hyres.29.905)
 142. Maulik N, Yoshida T, Das DK (1998) Oxidative stress developed during the reperfusion of ischemic myocardium induces apoptosis. *Free Radical Biol Med* 24(5):869–875
 143. Hoshida S, Kuzuya T, Nishida M, Yamashita N, Hori M, Kamada T, Tada M (1994) Ebselen protects against ischemia-reperfusion injury in a canine model of myocardial infarction. *Am J Physiol* 267(6 Pt 2):H2342–H2347
 144. Sui H, Wang W, Wang PH, Liu LS (2005) Effect of glutathione peroxidase mimic ebselen (PZ51) on endothelium and vascular structure of stroke-prone spontaneously hypertensive rats. *Blood Press* 14(6):366–372. doi:[10.1080/08037050500210781](https://doi.org/10.1080/08037050500210781)
 145. de Haan JB, Cooper ME (2011) Targeted antioxidant therapies in hyperglycemia-mediated endothelial dysfunction. *Front Biosci (Schol Ed)* 3:709–729
 146. Davis MT, Bartfay WJ (2004) Ebselen decreases oxygen free radical production and iron concentrations in the hearts of chronically iron-overloaded mice. *Biol Res Nurs* 6(1):37–45. doi:[10.1177/1099800403261350](https://doi.org/10.1177/1099800403261350)
 147. Saad SY, Najjar TA, Arafah MM (2006) Cardioprotective effects of subcutaneous ebselen against daunorubicin-induced cardiomyopathy in rats. *Basic Clin Pharmacol Toxicol* 99(6):412–417. doi:[10.1111/j.1742-7843.2006.pto_523.x](https://doi.org/10.1111/j.1742-7843.2006.pto_523.x)
 148. Pritsos CA, Sokoloff M, Gustafson DL (1992) PZ-51 (Ebselen) in vivo protection against adriamycin-induced mouse cardiac and hepatic lipid peroxidation and toxicity. *Biochem Pharmacol* 44(4):839–841
 149. Ardais AP, Santos FW, Nogueira CW (2008) Ebselen attenuates cadmium-induced testicular damage in mice. *J Appl Toxicol* 28(3):322–328. doi:[10.1002/Jat.1282](https://doi.org/10.1002/Jat.1282)
 150. Husain K, Morris C, Whitworth C, Trammell GL, Rybak LP, Somani SM (1998) Protection by ebselen against cisplatin-induced nephrotoxicity: antioxidant system. *Mol Cell Biochem* 178(1–2):127–133. doi:[10.1023/A:1006889427520](https://doi.org/10.1023/A:1006889427520)
 151. Yoshida M, Iizuka K, Terada A, Hara M, Nishijima H, Shimada A, Nakada K, Satoh Y, Akama Y (2000) Prevention of nephrotoxicity of cisplatin by repeated oral administration of ebselen in rats. *Tohoku J Exp Med* 191(4):209–220. doi:[10.1620/Tjem.191.209](https://doi.org/10.1620/Tjem.191.209)
 152. Hardej D, Trombetta LD (2002) The effects of ebselen on cisplatin and diethyldithiocarbamate (DDC) cytotoxicity in rat hippocampal astrocytes. *Toxicol Lett* 131(3):215–226. doi:[10.1016/S0378-4274\(02\)00056-5](https://doi.org/10.1016/S0378-4274(02)00056-5)
 153. Yin Z, Lee E, Ni M, Jiang H, Milatovic D, Rongzhu L, Farina M, Rocha JBT, Aschner M (2011) Methylmercury-induced alterations in astrocyte functions are attenuated by ebselen. *Neurotoxicology* 32(3):291–299. doi:[10.1016/j.neuro.2011.01.004](https://doi.org/10.1016/j.neuro.2011.01.004)
 154. Usuki F, Yamashita A, Fujimura M (2011) Post-transcriptional defects of antioxidant selenoenzymes cause oxidative stress under methylmercury exposure. *J Biol Chem* 286(8):6641–6649. doi:[10.1074/jbc.M110.168872](https://doi.org/10.1074/jbc.M110.168872)
 155. Baldew GS, McVie JG, van der Valk MA, Los G, de Goeij JJ, Vermeulen NP (1990) Selective reduction of cis-diamminedichloroplatinum(II) nephrotoxicity by ebselen. *Cancer Res* 50(21):7031–7036
 156. Marreilha Dos Santos AP, Lucas RL, Andrade V, Mateus ML, Milatovic D, Aschner M, Batoreu MC (2012) Protective effects of ebselen (Ebs) and para-aminosalicylic acid (PAS) against manganese (Mn)-induced neurotoxicity. *Toxicol Appl Pharmacol* 258(3):394–402. doi:[10.1016/j.taap.2011.12.003](https://doi.org/10.1016/j.taap.2011.12.003)
 157. Moretto MB, Funchal C, Zeni G, Rocha JB, Pessoa-Pureur R (2005) Organoselenium compounds prevent hyperphosphorylation of cytoskeletal proteins induced by the neurotoxic agent diphenyl ditelluride in cerebral cortex of young rats. *Toxicology* 210(2–3):213–222. doi:[10.1016/j.tox.2005.02.003](https://doi.org/10.1016/j.tox.2005.02.003)
 158. Hardej D, Billack B (2007) Ebselen protects brain, skin, lung and blood cells from mechlorethamine toxicity. *Toxicol Ind Health* 23(4):209–221
 159. Pino MA, Billack B (2008) Reduction of vesicant toxicity by butylated hydroxyanisole in A-431 skin cells. *Cutan Ocul Toxicol* 27(3):161–172. doi:[10.1080/15569520802092070](https://doi.org/10.1080/15569520802092070)
 160. Blasiak J, Sikora A, Czechowska A, Drzewoski J (2003) Free radical scavengers can modulate the DNA-damaging action of alloxan. *Acta biochimica Polonica* 50(1):205–210
 161. Blasiak J, Sikora A, Wozniak K, Drzewoski J (2004) Genotoxicity of streptozotocin in normal and cancer cells and its modulation by free radical scavengers. *Cell Biol Toxicol* 20(2):83–96
 162. Blasiak J, Gloc E, Drzewoski J, Wozniak K, Zadrozny M, Skorski T, Pertynski T (2003) Free radical scavengers can differentially modulate the genotoxicity of amsacrine in normal and cancer cells. *Mutat Res* 535(1):25–34
 163. Wu Q, Huang K (2006) Protective effect of ebselen on cytotoxicity induced by cholestane-3 beta, 5 alpha, 6 beta-triol in ECV-304 cells. *Biochim Biophys Acta* 1761(3):350–359. doi:[10.1016/j.bbalip.2006.02.017](https://doi.org/10.1016/j.bbalip.2006.02.017)
 164. Huther AM, Zhang Y, Sauer A, Parnham MJ (1989) Antimalarial properties of ebselen. *Parasitol Res* 75(5):353–360
 165. Harris MT, Walker DM, Drew ME, Mitchell WG, Dao K, Schroeder CE, Flaherty DP, Weiner WS, Golden JE, Morris JC (2013) Interrogating a hexokinase-selected small-molecule library for inhibitors of *Plasmodium falciparum* hexokinase. *Antimicrob Agents Chemother* 57(8):3731–3737. doi:[10.1128/AAC.00662-13](https://doi.org/10.1128/AAC.00662-13)
 166. Billack B, Santoro M, Lau-Cam C (2009) Growth inhibitory action of ebselen on fluconazole-resistant *Candida albicans*: role of the plasma membrane H⁺-ATPase. *Microb Drug Resist* 15(2):77–83. doi:[10.1089/mdr.2009.0872](https://doi.org/10.1089/mdr.2009.0872)
 167. Billack B, Pietka-Ottlik M, Santoro M, Nicholson S, Mlochowski J, Lau-Cam C (2010) Evaluation of the antifungal and plasma membrane H⁺-ATPase inhibitory action of ebselen and two ebselen analogs in *S. cerevisiae* cultures. *J Enzyme Inhib Med Chem* 25(3):312–317. doi:[10.3109/14756360903179419](https://doi.org/10.3109/14756360903179419)
 168. Lieberman OJ, Orr MW, Wang Y, Lee VT (2013) High-throughput screening using the differential radial capillary action of ligand assay identifies ebselen as an inhibitor of diguanylate cyclases. *ACS Chem Biol*. doi:[10.1021/cb400485k](https://doi.org/10.1021/cb400485k)
 169. Favrot L, Grzegorzewicz AE, Lajiness DH, Marvin RK, Boucau J, Isailovic D, Jackson M, Ronning DR (2013) Mechanism of inhibition of *Mycobacterium tuberculosis* antigen 85 by ebselen. *Nat Commun* 4:2748. doi:[10.1038/ncomms3748](https://doi.org/10.1038/ncomms3748)
 170. Sinha R, Ei-Bayoumy K (2004) Apoptosis is a critical cellular event in cancer chemoprevention and chemotherapy by

- selenium compounds. *Curr Cancer Drug Tar* 4(1):13–28. doi:[10.2174/1568009043481614](https://doi.org/10.2174/1568009043481614)
171. Tewari R, Sharma V, Koul N, Ghosh A, Joseph C, Hossain SkU, Sen E (2009) Ebselen abrogates TNFalpha induced pro-inflammatory response in glioblastoma. *Mol Oncol* 3(1):77–83. doi:[10.1016/j.molonc.2008.10.004](https://doi.org/10.1016/j.molonc.2008.10.004)
 172. Sharma V, Tewari R, Sk UH, Joseph C, Sen E (2008) Ebselen sensitizes glioblastoma cells to Tumor Necrosis Factor (TNF-alpha)-induced apoptosis through two distinct pathways involving NF-kappaB downregulation and Fas-mediated formation of death inducing signaling complex. *Int J Cancer* 123(9):2204–2212. doi:[10.1002/ijc.23771](https://doi.org/10.1002/ijc.23771)
 173. Gealekman O, Brodsky SV, Zhang F, Chander PN, Friedli C, Nasjletti A, Goligorsky MS (2004) Endothelial dysfunction as a modifier of angiogenic response in Zucker diabetic fat rat: amelioration with Ebselen. *Kidney Int* 66(6):2337–2347. doi:[10.1111/j.1523-1755.2004.66035.x](https://doi.org/10.1111/j.1523-1755.2004.66035.x)
 174. Saito I, Asano T, Sano K, Takakura K, Abe H, Yoshimoto T, Kikuchi H, Ohta T, Ishibashi S (1998) Neuroprotective effect of an antioxidant, ebselen, in patients with delayed neurological deficits after aneurysmal subarachnoid hemorrhage. *Neurosurgery* 42 (2):269–277; discussion 277–268
 175. Aruoma OI (2003) Methodological considerations for characterizing potential antioxidant actions of bioactive components in plant foods. *Mutat Res* 523–524:9–20
 176. Young IS, Woodside JV (2001) Antioxidants in health and disease. *J Clin Pathol* 54(3):176–186. doi:[10.1136/jcp.54.3.176](https://doi.org/10.1136/jcp.54.3.176)
 177. Luo Z, Sheng J, Sun Y, Lu C, Yan J, Liu A, Luo HB, Huang L, Li X (2013) Synthesis and evaluation of multi-target-directed ligands against Alzheimer's disease based on the fusion of donepezil and ebselen. *J Med Chem* 56(22):9089–9099. doi:[10.1021/jm401047q](https://doi.org/10.1021/jm401047q)
 178. Xu KH, Qiang MM, Gao W, Su RX, Li N, Gao Y, Xie YX, Kong FP, Tang B (2013) A near-infrared reversible fluorescent probe for real-time imaging of redox status changes in vivo. *Chem Sci* 4(3):1079–1086. doi:[10.1039/C2sc22076h](https://doi.org/10.1039/C2sc22076h)