

ping, ischemia must be suspected, even without the immediate appearance of clinical symptoms.

Acknowledgment: The Evanston Northwestern Healthcare movement disorders surgical program acknowledges the generous support of the Parkinson's Disease Research Society.

REFERENCES

1. Hoehn MM, Yahr MD. Parkinsonism: onset, progression and mortality. *Neurology* 1967;17:427–442.
2. Laitinen LV, Bergenheim AT, Hariz MI. Leksell's posteroventral pallidotomy in the treatment of Parkinson's disease. *J Neurosurg* 1992;76:53–61.
3. Limousin P, Pollak P, Benazzouz A, et al. Effect of parkinsonian signs and symptoms of bilateral subthalamic nucleus stimulation. *Lancet* 1995;345:91–95.
4. Siegfried J, Lippitz B. Bilateral chronic electrostimulation of ventroposterolateral pallidum: a new therapeutic approach for alleviating all parkinsonian symptoms. *Neurosurgery* 1994;35:1126–1129; discussion, 1129–1130.
5. Limousin P, Krack P, Pollak P, et al. Electrical stimulation of the subthalamic nucleus in advanced Parkinson's disease. *N Engl J Med* 1998;339:1105–1111.
6. Gregory R. Surgery for movement disorders. *J Neurol Neurosurg Psychiatry* 2002;72(Suppl. 1):132–135.
7. Vingerhoets FJ, Villemure JG, Temperli P, et al. Subthalamic DBS replaces levodopa in Parkinson's disease: two-year follow-up. *Neurology* 2002;58:396–401.
8. Umemura A, Jaggi JL, Hurtig HI, et al. Deep brain stimulation for movement disorders: morbidity and mortality in 109 patients. *J Neurosurg* 2003;98:779–784.
9. Schaltenbrand G, Walker AE. *Stereotaxy of the Human Brain: Anatomical, Physiological, and Clinical Applications*. 2nd ed. Stuttgart, Germany: Georg Thieme; 1982.
10. Bosch DA. *Stereotactic Techniques in Clinical Neurosurgery*. Wien, Germany: Springer-Verlag; 1986.
11. Binder DK, Rau GM, Starr PA. Risk factors for hemorrhage during microelectrode-guided deep brain stimulator implantation for movement disorders. *Neurosurgery* 2005;56:722–732.
12. Gorgulho A, De Salles AA, Frighetto L, Behnke E. Incidence of hemorrhage associated with electrophysiological studies performed using macroelectrodes and microelectrodes in functional neurosurgery. *J Neurosurg* 2005;102:888–896.
13. Pollak P, Fraix V, Krack P, et al. Treatment results: Parkinson's disease. *Mov Disord* 2002;17(Suppl. 3):S75–S83.
14. Gancher ST. Scales for the assessment of movement disorders. In: Herndon RM, editor. *Handbook of Neurologic Rating Scales*. New York, NY: Demos Vermande; 1997. p 81–104.
15. Schaltenbrand G, Wahren W. *Atlas for Stereotaxy of the Human Brain; With an Accompanying Guide*. Second, Revised, and Enlarged edition ed. Chicago, IL: Year Book Medical Publishers, Inc; 1977.
16. Cox RW. AFNI: software for analysis and visualization of functional magnetic resonance neuroimages. *Comput Biomed Res* 1996;29:162–173.
17. Sutton JP. Deep brain stimulation (DBS) of the subthalamic nucleus (STN) and thalamic ischemia: a report of two cases. *Mov Disord* 2004;19(Suppl. 9):S324.
18. Christoforidis GA, Spickler EM, Papaioannou G, Junn F. Lacunar infarct during pallidotomy: case report. *Neuroradiology* 2001;43:321–324.
19. Lim JY, De Salles AA, Bronstein J, Masterman DL, Saver JL. Delayed internal capsule infarctions following radiofrequency pallidotomy. Report of three cases. *J Neurosurg* 1997;87:955–960.
20. Arboix A, Morcillo C, Garcia-Eroles L, et al. Different vascular risk factor profiles in ischemic stroke subtypes: a study from the "Sagrat Cor Hospital of Barcelona Stroke Registry". *Acta Neurol Scand* 2000;102:264–270.
21. Rovira A, Grive E, Alvarez-Sabin J. Distribution territories and causative mechanisms of ischemic stroke. *Eur Radiol* 2005;15:416–426.
22. Landau WM. Clinical neuromyology VI. Au clair de lacune: holy, wholly, holey logic. *Neurology* 1989;39:725–730.
23. Dietrich HH, Dacey RG Jr. Molecular keys to the problems of cerebral vasospasm. *Neurosurgery* 2000;46:517–530.

Neuropsychological Deficits in Parkinson's Disease Patients With Visual Hallucinations

Blanca Ramírez-Ruiz, MSc,^{1,2*} Carme Junqué, PhD,^{1,2}
 María-José Martí, MD, PhD,^{2,3}
 Francesc Valldeoriola, MD,^{2,3}
 and Eduardo Tolosa, MD^{2,3}

¹Department of Psychiatry and Clinical Psychobiology, University of Barcelona, Barcelona, Spain; ²Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain; ³Parkinson's Disease and Movement Disorders Unit, Neurology Service, Institut Clínic de Neurociències, Hospital Clínic Universitari, University of Barcelona, Barcelona, Spain

Abstract: Recent neuropathological and neuroimaging studies suggest the involvement of several temporal regions in Parkinson's disease (PD) patients with visual hallucinations (VH). We examined 24 nondemented PD patients with VH, 21 PD patients without VH, and 21 healthy controls using a battery of tests assessing different aspects of temporal lobe function. PD patients with VH showed poorer performance in language, verbal learning, semantic fluency, and visuo-perceptive functions compared to controls and PD patients without VH. Differences in verbal learning and visuo-perceptive functions were independent of general cognitive status, disease severity, and depression. We suggest that a wide range of neuropsychological deficits can contribute to the emergence of VH in PD. © 2006 Movement Disorder Society

Key words: Parkinson's disease; visual hallucinations; language; memory; visuo-perceptive

Visual hallucinations (VH) belong to the most frequent neuropsychiatric symptoms of Parkinson's disease

*Correspondence to: Dr. Carme Junqué, Departament de Psiquiatria i Psicobiologia Clínica, Universitat de Barcelona, IDIBAPS, Casanova 143, 08036 Barcelona, Spain. E-mail: cjunque@ub.edu

Received 16 November 2005; Revised 20 January 2006; Accepted 24 January 2006

Published online 16 May 2006 in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/mds.20965

(PD), affecting 22% to 50% of patients.^{1,2} Although they have been associated with the presence of cognitive impairment or frank dementia,^{1,2} only few studies have addressed specific neuropsychological deficits in nondemented PD patients with VH. Barnes and colleagues^{3,4} found that hallucinating PD patients showed deficits in object perception and visual recognition memory in comparison with both nonhallucinating PD patients and healthy controls. Cognitive deficits related to executive functions have also been reported in hallucinating PD patients.⁵ In terms of the underlying brain mechanisms, neuroimaging and neuropathological studies found an association between temporal and frontal cerebral abnormalities and VH in PD patients.^{6–8} Our study investigates the potential relationship between VH in PD and deficits in different neuropsychological domains, with a particular emphasis on testing of temporal lobe function.

PATIENTS AND METHODS

Subjects

Participants were recruited from the Parkinson Disease and Movement Disorders Unit, Hospital Clinic Universitari, Barcelona, Spain. The sample comprised 24 PD patients who had experienced VH in the previous year, 21 PD patients who never suffered from VH, and 21 healthy elderly controls. The controls comprised spouses of PD patients and community volunteers without any history of psychiatric or neurological disorders who were matched with patients for age and education. All subjects gave informed consent to participate in this study, which was approved by the local Ethics Committee.

Diagnostic Criteria and Clinical Evaluation

The details of the diagnostic criteria and clinical assessment are described elsewhere.⁹ Patients who fulfilled Diagnostic and Statistical Manual of Mental Disorders, Revised Fourth Edition, criteria for dementia or presented clinical depression (Hamilton score ≥ 15) were excluded. All patients passed a visual acuity test consisting of reading from a distance of approximately 40 cm.

The demographic and clinical characteristic of the sample (PD+VH; PD; Controls) are, respectively, as follows: age (74.7 ± 5.4 ; 73.3 ± 6.1 ; 73 ± 6.7), sex distribution ([males/females] 10/14; 9/12; 9/12), years of education (7.3 ± 3.4 ; 7.7 ± 3.4 ; 7.9 ± 4.6), and visual acuity (0.5 ± 0.2 ; 0.6 ± 0.2 ; 0.7 ± 0.2). None of these variables differed significantly among the samples. PD patients with VH obtained a lower Mini-Mental State Examination (MMSE; 26.7 ± 2.1) and a higher Hamilton scores (6.5 ± 4.6) than both non-VH PD patients (MMSE = 29.2 ± 1.4 ; Hamilton = 3.6 ± 2.7) and

healthy controls (MMSE = 29.9 ± 1.6 ; Hamilton = 2.7 ± 2.6) ($P < 0.0005$; $P < 0.05$, respectively). Concerning clinical differences between both PD groups, hallucinating PD patients presented similar motor severity as assessed by the Unified Parkinson's Disease Rating Scale (UPDRS) Motor scale (30.6 ± 14.5) but a more advanced Hoehn & Yahr stage (3.3 ± 1.1) compared to nonhallucinating PD patients (UPDRS = 24.9 ± 13.7 ; H&Y = 2.5 ± 0.7) ($P = 0.21$; $P = 0.015$, respectively). No differences in levodopa daily dose were found between PD groups with and without VH ($P = 0.071$).

The VH in our PD sample consisted of well-formed images of people, faces, or animals. Insight into the hallucinatory nature of the phenomenon was maintained in 63% of the patients. Associated delusions were present in 33% of the patients. These delusions were primarily paranoid in type and involved elementary misbeliefs concerning infidelity or theft. All hallucinating PD patients but 1 were on dopaminergic treatment at the time of evaluation. A modification of the antiparkinsonian treatment preceding the onset of VH was recorded in 33% of the patients. The VH disappeared after dopaminergic drug dosage reduction in 11 cases (45.8%), whereas 5 patients (20.8%) needed antipsychotic medication and 5 (20.8%) required both interventions. In the remaining patients, medication was not modified because the hallucinations were well tolerated.

Neuropsychological Assessment

General intellectual ability was assessed by the Information and Similarities subtests from Wechsler Adult Intelligence Scale-Third Edition (WAIS-III). Language was evaluated by a verbal comprehension test (Token Test) and a picture naming test (Boston Naming Test) and frontal functions by phonological and semantic fluency tests. A modified version of the Rey Auditory-Verbal Learning Test (RAVLT) was used to assess verbal learning, delayed recall, and recognition. To assess visuoperceptive functions, we used the Benton Facial Recognition Test (BFRT) and the standard drawing and multiple-choice version of the Benton Visual Form Discrimination Test (VFDT).¹⁰ Finally, we evaluated visual memory functions using the subtest Memory for Faces from Warrington's Recognition Memory Test. All tests were administered and scored according to conventional procedures.¹¹

Data Analysis

Statistical analysis was carried out using SPSS 11.0. Sex distribution was compared with χ^2 tests. For normally distributed quantitative variables with homogene-

TABLE 1. Neuropsychological test scores

	PD with VH (n = 24)	PD without VH (n = 21)	Controls (n = 21)	<i>P</i> value ^a	<i>P</i> value ^b
General intellectual ability					
Similarities (WAIS-III)	11.7 ± 4.8	12.7 ± 6.0	14.0 ± 5.1	NS	—
Information (WAIS-III)	11.5 ± 6.9	12.4 ± 6.1	12.9 ± 7.3	NS	—
Language					
Token Test	27.8 ± 4.4	30.6 ± 2.8	31.9 ± 3.0	<0.05**	NS
Boston Naming Test	47.3 ± 7.0	51.8 ± 4.3	52.5 ± 5.4	<0.05**	NS
Verbal memory					
RAVLT-learning	24.8 ± 7.2	38.8 ± 7.0	41.9 ± 7.2	<0.0005**	<0.0005
RAVLT-memory loss	8.4 ± 10.8	3.7 ± 4.4	3.0 ± 2.7	0.043*	—
RAVLT-correct recognition	12.5 ± 2.1	12.8 ± 2.1	13.7 ± 1.4	NS	—
RAVLT-false recognition	2.4 ± 2.3	1.3 ± 1.3	0.9 ± 0.8	0.007*	—
Executive functions					
Phonological fluency (P)	7.4 ± 4.5	9.7 ± 3.7	12.7 ± 6.1	0.002*	—
Semantic fluency (animals)	9.4 ± 4.5	13.3 ± 4.7	17.1 ± 5.1	<0.05**	NS
Visuoperceptive functions					
Benton Facial Recognition					
Test	43.7 ± 4.5	49.0 ± 4.3	49.4 ± 2.9	<0.0005**	<0.013
Visual Form Discrimination					
Test	26.4 ± 3.8	28.8 ± 3.1	30.4 ± 2.0	<0.05**	NS
Visual Memory					
Warrington's Recognition					
Test	31.1 ± 5.6	33.5 ± 6.5	40.4 ± 5.6	<0.005*	—

^aAnalysis of variance.

^bAnalysis of covariance between PD groups using Mini-Mental State Evaluation, Hoehn & Yahr, and Hamilton scores as covariates.

*Significant difference between PD patients with VH and controls.

**Significant difference between PD patients with VH and controls and between PD patients with VH and PD patients without VH.

PD, Parkinson's disease; VH, visual hallucinations; WAIS-III, Wechsler Adult Intelligence Scale, Third Edition; RAVLT-learning, sum of words correctly recalled from Trials 1 to 5; RAVLT-memory loss, percentage of memory loss: percentage of words loss after 20 min of interference; RAVLT-correct recognition, number of words correctly recognized from the original list; RAVLT-false recognition, number of words falsely recognized from the original list.

ity of variance, we used an analysis of variance (ANOVA test) and the post hoc Bonferroni test. For non-normally distributed variables, and/or in the case of no equality of variance between the groups, we used the nonparametric Mann-Whitney *U* test. Finally, we performed an analysis of covariance (ANCOVA) using MMSE, Hoehn & Yahr stage, and Hamilton scores to determine whether differences between the two PD groups on cognitive tests persisted after controlling for these variables.

RESULTS

The results of the neuropsychological assessment are shown in Table 1. The groups did not differ in general intellectual ability as evaluated with the Similarities ($P = 0.343$) and Information ($P = 0.777$) subtests from WAIS-III. However, the ANOVA showed group differences in the other domains assessed: language, memory, frontal, and visuoperceptive functions. The post hoc test revealed poorer performance in PD patients with VH than in healthy controls in all neuropsychological measures except RAVLT-recognition. Patients with VH were more impaired than those without VH on language assessed by Token and Boston Naming tests ($P = 0.034$

and $P = 0.032$, respectively), verbal learning ($P < 0.0005$), semantic fluency ($P = 0.022$), and visuoperceptive functions evaluated by Benton Facial Recognition ($P < 0.0005$) and Visual Form Discrimination ($P = 0.042$). The only significant difference between PD patients without VH and healthy controls was observed in visual memory ($P = 0.001$).

Given that the specific neuropsychological deficits may be related to the global cognitive performance (MMSE score) and to the differences in illness stage (Hoehn & Yahr scale) and mood (Hamilton scale), we performed an ANCOVA between the two PD groups using these clinical variables as covariates. After removing the effect of these variables, differences in RAVLT-verbal learning ($P < 0.0005$) and in Benton Facial Recognition Test ($P = 0.013$) remained significant.

DISCUSSION

PD patients with VH showed several cognitive deficits affecting not only visuoperceptive performance, as previously reported, but also language and verbal memory domains. This finding suggests that cerebral dysfunction in PD with VH extends beyond the subcortical-frontal circuits typically described in PD.

On tests of verbal memory, PD patients with VH demonstrate the classic PD pattern, consisting of deficits in free recall but normal recognition.¹² However, in addition, our patients with VH showed deficits in long-term retention (RAVLT-memory loss) and the presence of false intrusions. These findings could reflect an involvement of limbic and temporal cortex, as suggested by neuropathological studies,⁷ in addition to a disruption of nigro-striato-thalamo-cortical circuitry.

The differences on Benton's Facial recognition test in the hallucinating patients are in agreement with the neuropsychological⁴ as well as neuroimaging findings suggesting involvement of visual associative areas in patients with VH.^{6,8} These differences are not likely to be caused by a lower visual acuity, although we cannot exclude a faulty visual input due to primary visual deficits commonly seen in PD, such as reduced color and contrast discrimination.¹³ Differences in BFRT remained significant after controlling for MMSE. In contrast, the differences between hallucinating and nonhallucinating PD patients on the VFDT lost significance after such covariation. The performance on VFDT, in addition to visuo-perceptive skills involved in BFRT, requires additional visuospatial functions. This finding could explain why this test depends to a higher degree on the general cognitive status.

Regarding the face recognition memory, our results are partially in agreement with those reported by Barnes and associates (2003) that found a significant impairment in PD patients with and without VH vs. healthy controls. Nevertheless, we did not observe significant statistical differences on the Warrington Facial Recognition test between hallucinating and nonhallucinating groups, although PD patients with VH scored lower than those without this symptom. Longitudinal studies can clarify the relevance of visual memory impairment on cognitive decline associated with the presence of hallucinations.

In addition to complex visual dysfunction, we observed deficits on language comprehension. The poor performance on the Token Test could be related to an impairment in sentence processing,¹⁴ which has been associated with a reduced activation of anteromedial prefrontal cortex and posterolateral temporal regions.¹⁵ Additionally, the reduced performance on the Token Test could also be related to a visual attention impairment, as alterations of parieto-frontal connections associated primarily with the attentional modulation of visual perception have been implicated in the pathophysiology of VH.⁸

The recruitment of inferior frontal and ventral temporal regions have been observed during confrontation naming and semantic fluency tasks in healthy subjects.^{16,17} The deficits found in these cognitive domains in our hallucinating patients, therefore, could reflect dys-

function in these brain areas. From the clinical point of view, naming impairment has been related to the level of general cognitive dysfunction in PD¹² and prospective studies have identified deficits in verbal fluency as predictors of the development of dementia.¹⁸ This finding is consistent with our finding that the differences between the PD groups in naming and category fluency lost significance once the effects of general cognitive status were removed from the analysis. The reduced efficiency in language and semantic fluency in our PD patients with VH could reflect the starting point of the more general cognitive decline which might later lead to dementia.

In conclusion, hallucinating PD patients were characterized by a wide range of impairments affecting several cognitive domains. The presence of such extensive deficits may escape the compensatory processes and, thus, favor the perception of VH. Given the well-documented relationship between dementia and hallucinations, it remains to be shown in longitudinal studies whether visuo-perceptive and memory impairments found in our hallucinating PD sample are the earliest symptoms of a progressive cognitive decline.

Acknowledgments: This study was supported by the following grants: Red CIEN IDIBAPS- ISCIII RTIC C03/06 (E. Tolosa and C. Junqué), 2001SGR00139 and 2001SGR00387 (Generalitat de Catalunya to C. Junqué and E. Tolosa), and Award "Distinció per a la Promoció de Recerca Universitària Generalitat de Catalunya" to E. Tolosa and C. Junqué. B. Ramírez-Ruiz was funded by a grant AP-2001-0823 from the Ministerio de Educación, Cultura y Deporte.

REFERENCES

1. Fenelon G, Mahieux F, Huon R, Ziegler M. Hallucinations in Parkinson's disease. Prevalence, phenomenology and risk factors. *Brain* 2000;123:733-745.
2. Williams DR, Lees AJ. Visual hallucinations in the diagnosis of idiopathic Parkinson's disease: a retrospective autopsy study. *Lancet Neurol* 2005;4:605-610.
3. Barnes J, David AS. Visual hallucinations in Parkinson's disease: a review and phenomenological survey. *J Neurol Neurosurg Psychiatry* 2001;70:727-733.
4. Barnes J, Boubert L, Harris J, Lee A, David AS. Reality monitoring and visual hallucinations in Parkinson's disease. *Neuropsychologia* 2003;41:565-574.
5. Grossi D, Trojano L, Pellecchia MT, et al. Frontal dysfunction contributes to the genesis of hallucinations in non-demented Parkinsonian patients. *Int J Geriatr Psychiatry* 2005;20:668-673.
6. Okada K, Suyama N, Oguro H, Yamaguchi S, Kobayashi S. Medication-induced hallucination and cerebral blood flow in Parkinson's disease. *J Neurol* 1999;246:365-368.
7. Harding AJ, Broe GA, Halliday GM. Visual hallucinations in Lewy body disease relate to Lewy bodies in the temporal lobe. *Brain* 2002;125:391-403.
8. Stebbins GT, Goetz CG, Carrillo MC, et al. Altered cortical visual processing in PD with hallucinations: an fMRI study. *Neurology* 2004;63:1409-1416.
9. Ramírez-Ruiz B, Martí MJ, Tolosa E, et al. Longitudinal evaluation of cerebral morphological changes in Parkinson's disease with and without dementia. *J Neurol* 2005;252:1345-1352.

10. Benton AL, Sivan AB, Hamsher K, Varney NR, Spreen O. Contributions to Neuropsychological Assessment. 2nd ed. New York: Oxford University Press; 1994.
11. Spreen O, Strauss E, editors. A Compendium of Neuropsychological Tests. 2nd ed. New York: Oxford University Press; 1998.
12. Pahwa R, Paolo A, Tröster A, Koller W. Cognitive impairment in Parkinson's disease. *Eur J Neurol* 1998;5:431–441.
13. Diederich NJ, Goetz CG, Stebbins GT. Repeated visual hallucinations in Parkinson's disease as disturbed external/internal perceptions: focused review and a new integrative model. *Mov Disord* 2005;20:130–140.
14. Grossman M, Lee C, Morris J, Stern MB, Hurtig HI. Assessing resource demands during sentence processing in Parkinson's disease. *Brain Lang* 2002;80:603–616.
15. Grossman M, Cooke A, DeVita C, et al. Grammatical and resource components of sentence processing in Parkinson's disease. An fMRI study. *Neurology* 2003;60:775–781.
16. Henry TR, Buchtel HA, Koepp RA, et al. Absence of normal activation of the left anterior fusiform gyrus during naming in left temporal lobe epilepsy. *Neurology* 1998;50:787–790.
17. Vitali P, Abutalebi J, Tettamanti M, et al. Generating animal and tool names: an fMRI study of effective connectivity. *Brain Lang* 2005;93:32–45.
18. Jacobs DM, Marder K, Cote LJ, et al. Neuropsychological characteristics of preclinical dementia in Parkinson's disease. *Neurology* 1995;45:1691–1696.

OFF-off Rebound Dyskinesia in Subthalamic Nucleus Deep Brain Stimulation of Parkinson's Disease

Matthew A. Brodsky, MD,* Penelope Hogarth, MD, and John G. Nutt, MD

Department of Neurology, Oregon Health & Science University, and Parkinson's Disease Research, Education & Clinical Care Center, Portland VA Medical Center, Portland, Oregon, USA

Video



Abstract: A 61-year-old man with Parkinson's disease (PD), motor fluctuations, and dyskinesias underwent bilateral implantation of deep brain stimulation (DBS) electrodes in

the subthalamic nucleus (STN). One month after surgery, DBS was optimized to bilateral monopolar settings at the most proximal electrode just superior to the STN, which improved motor fluctuations and dyskinesias. At several postoperative evaluations off medications overnight, both stimulators were turned off and within 60 seconds he developed severe dyskinesias. When the stimulators were turned back on, the dyskinesias soon resolved. This article is a first report of a unique pattern of rebound-type dyskinesia that occurred in the off medication state produced by stopping STN DBS. © 2006 Movement Disorder Society

Key words: Parkinson's disease; dyskinesia; deep brain stimulation

Deep brain stimulation (DBS) for Parkinson's disease (PD) has become a widely accepted treatment for patients who suffer from disabling dyskinesia or motor fluctuations and have reached an endpoint of further improvement with medical therapy. DBS improves all of the cardinal signs of PD, including muscle rigidity, bradykinesia, and tremor.¹ Furthermore, an improvement in levodopa-induced dyskinesias is often seen after DBS surgery of the subthalamic nucleus (STN).² It is not certain whether this improvement is due to a decrease in L-dopa dose permitted by the improvement in parkinsonian symptoms² or a direct effect of the stimulation itself.³ Direct stimulation of the STN has also been demonstrated to induce dyskinesia.⁴ Although DBS of the STN has been reported to both increase or decrease dyskinesia while a patient is on medications, we report a case in which dyskinesias were induced by turning off STN stimulation when the subject was pharmacologically off.

CASE REPORT

A 61-year-old right-handed man with PD developed symptoms of left-sided muscle rigidity and bradykinesia 15 years before DBS surgery. He initially was treated with selegiline, with modest benefit. He subsequently was started on L-dopa/carbidopa 12 years before surgery and responded with an improvement in his PD symptoms. Ten years before surgery, a dopamine agonist was added to his drug regimen. Seven years before surgery, he began to develop motor fluctuations and dyskinesias. Over the subsequent 7 years, these motor complications progressively became more disabling. In the year before surgery, he was spending approximately 10% of his waking hours in the on state without disabling dyskinesias. During the month before surgery, his medication regimen was as follows: carbidopa/L-dopa 25/100, 2.5 tablets every 3 hours 8 times per day (around the clock); entacapone 200 mg with each dose of carbidopa/L-dopa; pramipexole 1.5 mg three times per day (t.i.d.); and

This article includes Supplementary Video, available online at <http://www.interscience.wiley.com/jpages/0885-3185/suppmat>

*Correspondence to: Dr. Matthew A. Brodsky, Parkinson Center of Oregon, Oregon Health & Science University, 3181 SW Sam Jackson Park Road, Mailcode OP-32, Portland, OR 97239.
E-mail: brodskym@ohsu.edu

Received 4 February 2005; Revised 28 December 2005; Accepted 24 January 2006

Published online 23 May 2006 in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/mds.20964