

See discussions, stats, and author profiles for this publication at:
<https://www.researchgate.net/publication/19118408>

Expression of abl and other oncogenes is independent of metastatic potential in Abelson virus-transformed malignant murine large lymphoma

ARTICLE *in* CLINICAL AND EXPERIMENTAL METASTASIS · MARCH 1985

Impact Factor: 3.49 · DOI: 10.1007/BF01758957 · Source: PubMed

CITATIONS

15

READS

8

4 AUTHORS, INCLUDING:



Varda Rotter

Weizmann Institute of Science

281 PUBLICATIONS 12,871 CITATIONS

SEE PROFILE



Garth Nicolson

The Institute for Molecular Medi...

664 PUBLICATIONS 32,262 CITATIONS

SEE PROFILE

Expression of *abl* and other oncogenes is independent of metastatic potential in Abelson virus-transformed malignant murine large cell lymphoma

VARDA ROTTER†, DAVID WOLF†, MARK BLICK‡
and GARTH L. NICOLSON§||

† Department of Cell Biology, The Weizmann Institute of Science,
Rehovot 76100, Israel

‡ Departments of §Clinical Immunology and Biological Therapy and §Tumor
Biology, The University of Texas M. D. Anderson Hospital and Tumor
Institute, Houston, TX 77030, U.S.A.

(Received 16 February 1985; accepted 2 April 1985)

The role of oncogene expression in tumor metastasis was examined using the Abelson leukemia virus-transformed murine large cell lymphoma RAW117. Cell sublines of low and high metastatic potential expressed equally *abl* oncogene-coded mRNA and its phosphoprotein product p160, and the capacity of p160 to become autophosphorylated with γ -[^{32}P]ATP was the same among low and high metastatic cells. The expression of other oncogene-coded mRNAs (*fos*, *myc*, *myb*), if present, was also similar in low and high metastatic RAW117 cells. Although oncogene expression is thought to be important in initiating, and in some cases maintaining, the transformed phenotype, its expression in RAW117 lymphoma cells appears to be unrelated to metastatic phenotype.

Introduction

The spread of malignant tumors from primary to near and distant secondary sites occurs via a complex series of sequential steps [19, 34, 35, 44, 45]. At each step of the metastatic process certain tumor cell characteristics are required [34, 38-40, 44], and maintenance of these characteristics necessitates the expression of certain genes that programme for or control metastasis-associated gene products [36, 37, 43].

The role of viral oncogenes or their cellular prototypes in the control of metastatic behavior is largely unknown, although oncogenes appear to be important in the maintenance of the transformed phenotype in some cells [5, 10, 14]. Insight into how cellular processes involved in transformation might be controlled by certain oncogenes has come from studies in which a cellular oncogene was activated by insertion of a viral gene promoter at a DNA site upstream from the cellular oncogene [24]. Other mechanisms of activation of cellular oncogenes, such as point mutation [16, 53], rearrangement [7, 18, 25] and amplification [2, 13] could individually or collectively affect the malignant potential of tumor cells.

We have examined the expression of the *abl* and other oncogenes in murine large cell lymphoma cells transformed by Abelson leukemia virus (AbLV) [46]. Using parental AbLV-transformed RAW117 lymphoma cells (RAW117-P), highly meta-

|| To whom correspondence should be addressed at Department of Tumor Biology,
Box 108, The University of Texas M. D. Anderson Hospital and Tumor Institute,
Houston, Texas 77030, U.S.A.

static sublines have been established by sequential selection, such as the subline RAW117-H10 selected ten times *in vivo* for liver colonization [6]. RAW117-H10 subline forms 200 to 250 times more gross liver tumor nodules than the RAW117-P line within two weeks after intravenous or subcutaneous injection of tumor cells [48].

Materials and methods

Cells

RAW117 cells were grown and maintained in suspension cultures in Dulbecco-modified Eagle's medium (DME, Gibco, Grant Island, NY, U.S.A.) containing 10 per cent fetal bovine serum (FBS, Reheis Division, Armour Pharmaceutical, Kankakee, IL, U.S.A.). Cell cultures were used within 10 passages from frozen stocks of low passage cells to eliminate possible phenotypic drift in metastatic properties [41]. Cultures were tested for the presence of *Mycoplasma* using Hoechst 33258 staining [8] and were found to be negative.

Animals and metastasis assays

RAW117 sublines were assayed for organ colonization by intravenous injection of $5\text{--}10 \times 10^3$ viable tumor cells in 0.1 ml Hank's balanced salt solution into groups of 10 female Balb/c mice [6]. Animals were fed on a diet of chow without silicon dioxide, and were given spring water (chlorine < 1 p.p.m.) *ad libitum*.

Oncogene-encoded mRNA

Total RNA was prepared according to Auffray and Rougeon [3] or Glisin *et al.* [22]. Polyadenylated mRNA was selected by oligo(dT)-cellulose chromatography [4]. Aliquots of 5 μ g polyadenylated (pA^+) or nonadenylated (pA^-) mRNA from RAW117 cells were heated for 10 min at 60°C in 50 per cent formamide, 6 per cent formaldehyde, 20 mM MOPS, 5 mM sodium acetate, 1 mM EDTA, pH 7.0. The samples were electrophoresed in a 1 per cent agarose gel containing 6 per cent formaldehyde. The RNA was transferred onto a nitrocellulose sheet [54] and hybridized to nick-translated oncogene plasmid inserts [51]. Oncogenes probed included *abl* [29], *fos* [11], *myb* [20], *myc* [12] and *B-lym* [17]. Hybridization was for 16 h at 43°C in 50 per cent formamide, 5X Denhart's solution [15], 5X SCC and 20 mM sodium phosphate, 100 μ g salmon sperm DNA, 10 per cent dextran sulfate, pH 7.0 [56]. Hybridized filters were washed at 60°C with $0.1 \times$ SCC and 0.1 per cent sodium dodecyl sulfate (SDS), and autoradiograms were prepared. Molecular weight markers were [^{32}P]-labeled 18S and 28S ribosomal RNA.

abl-oncogene phosphoprotein p160 and Moloney virus products

Tumor cells (10^7) were washed several times by centrifugation and resuspended in PBS. The cells were suspended in 1.5 ml DME (without methionine) containing 250 μCi [^{35}S]methionine (Amersham, U.K.) and 10 per cent dialysed, heat-inactivated FBS, and were incubated for 1 h at 37°C. The cells were then washed by centrifugation and resuspension into PBS, and extracted with 5 ml lysis buffer (10 mM sodium phosphate, 100 mM sodium chloride, 1 per cent Triton X-100, 0.5 per cent SDS, pH 7.5). Cell lysates were treated with non-immune serum and absorbed with *Staphylococcus aureus* [26]. Equivalent amounts of radioactive protein were immunoprecipitated with various antibody preparations, absorbed with *S. aureus*, and separated by polyacrylamide slab gel electrophoresis in SDS [30].

Antibody preparations Ab-T1 and Ab-T2 were made against AbLV-transformed L12 tumor cells and obtained as described previously [59].

Phosphorylation experiments were conducted using immunoprecipitated complexes from unlabeled cell lysates. The immunoprecipitates were collected with *S. aureus*, washed by centrifugation and resuspended in 10 mM magnesium chloride containing 1 μ Ci γ -[32 P]ATP, and incubated for 20 min at 0°C. The complexes were then washed in the same buffer without isotope and subjected to SDS-polyacrylamide slab gel electrophoresis as described [9, 60].

Immunoprecipitation of Moloney leukemia virus (MoLV) gene products was accomplished with anti-MoLV. Cells were labeled as described above with [32 S]methionine and immunoprecipitated with goat anti-MoLV sera (Resources Branch, U.S. National Cancer Institute). Analysis of immunoprecipitated proteins was by SDS-polyacrylamide slab gel electrophoresis.

Results

AbLV-transformed cells express the *abl* oncogene-coded proteins, which are 90–160 kd in size, depending on transformed cell type [50, 52, 58]. The *abl* oncogene product p160 of RAW117 cells can be immunoprecipitated from RAW117 cell lysates by antibodies directed against the shared portion of p160 molecule that is coded by the normal *c-abl* cellular oncogene [59]. Antiserum Ab-T1 contains antibodies directed against p160, and antiserum Ab-T2 contains antibodies directed against p160 determinants plus additional MoLV precursors for the *env* (p80) and *gag* (p65) proteins [52]. Equivalent amounts of the *abl*-encoded p160 is precipitated by Ab-T1 and Ab-T2 from cell lysates of low (RAW117-P) or high (RAW117-H10) metastatic potential cells (figure 1). As a positive control for these experiments, the *abl*-encoded protein p120, synthesized by a different AbLV-transformed cell line, 2M3/M [54], is also immunoprecipitated by these reagents (figure 1).




Figure 1. Synthesis and phosphorylation of *abl*-coded p160 in murine large cell lymphoma cells of low (RAW117-P) and high (RAW117-H10) metastatic potential and in AbLV-transformed fibroblasts (2M3/M) by immunoprecipitation with Ab-T1 (lanes 1–6) and Ab-T2 (lanes 7–12). Lanes 1, 4, 7, 10, 2M3/M; lanes 2, 5, 8, 11, RAW117-P; lanes 3, 6, 9, 12, RAW117-H10; lanes 1–3, 7–9, γ -[32 P]ATP; lanes 4–6, 10–12, [35 S]methionine.




Figure 2. Immunoprecipitation of [^{35}S]methionine-labeled proteins in murine lymphoma cells of low (RAW117-P) and high (RAW117-H10) metastatic potential by goat anti-MoLV. Serial dilutions of goat anti-MoLV were used to precipitate: RAW117-P (lanes 1-5); RAW117-H10 (lanes 6-10).

The amounts of p160 synthesized in low and high metastatic potential RAW117 cells can be compared by pulse-labeling the cells with [^{35}S]methionine and immunoprecipitating p160 from the radiolabeled cell lysates. Similar amounts of [^{35}S]methionine label in p160 are immunoprecipitated with Ab-T1 or Ab-T2 antibodies from RAW117 cell lysates (figure 1; only RAW117-P and RAW117-H10 are shown). Moreover, in pulse chase experiments the p160 from RAW117 cells of low or high metastatic potential exhibit the same half-lives (data not shown).

One of the unique properties of many oncogene-coded proteins is their capacity to be phosphorylated *in vitro* at tyrosine residues [9,60]. We therefore tested whether p160 would be differentially phosphorylated in the presence of γ -[^{32}P]ATP in low and high metastatic potential RAW117 cells. When γ -[^{32}P]ATP is added to p160 immunoprecipitated by Ab-T1 or Ab-T2 antibodies from cell lysates of RAW117-P or RAW117-H10, and p120 immunoprecipitated from cell lysates of control 2M3/M cells, equivalent autophosphorylation of the *abl* oncogene product occurs (figure 1).

To further compare p160 from low and high metastatic RAW117 cells, we labeled cells with [^{35}S]methionine and immunoprecipitated cell lysates with goat anti-MoLV antibodies that bind to the *gag*-encoded region of the p160 molecule [58]. In agreement with the previous experiment using Ab-T1 and Ab-T2 antibodies, excess anti-MoLV antibodies immunoprecipitate similar amounts of p160 from low and high metastatic RAW117 cells (figure 2). Although lower concentrations of the anti-MoLV appear to precipitate more p160 from the high metastasis RAW117-H10 than the low metastasis RAW117-P cells, this discrepancy is probably due to the higher levels of *gag* precursor product and p30 in RAW117-P cells [48]. These latter two components are known to share antigenic determinants with p160 and are recognized by the goat anti-MoLV reagent. Therefore, saturation of these components plus p160 in the RAW117-P cell lysates requires higher antibody concentrations (figure 2).




Figure 3. Analysis of *abl*-specific mRNA in murine large cell lymphoma cells of low (RAW117-P) and high (RAW117-H10) metastatic potential and AbLV-transformed fibroblasts. Lane 1, pA⁺ RNA of 2M3/M; lane 2, pA⁺ RNA of 2M3/M; lane 3, pA⁺ RNA of RAW117-H10; lane 4, pA⁻ RNA of RAW117-H10; lane 5, pA⁺ RNA of RAW117-P; lane 6, pA⁻ RNA of RAW117-P. 18S and 28S ribosomal RNA serve as molecular weight markers.

In addition to the Abelson leukemia virus sequences, RAW117 cells also contain MoLV helper virus. Immunoprecipitation of cell lysates with the goat anti-MoLV antibody will, therefore, yield MoLV-related products and their precursors. Such an experiment reveals that RAW117-P cells synthesize greater amounts of the precursor envelope product (p80) than RAW117-H10 cells (figure 2). This agrees with the observation that RAW117-P cells possess higher levels of gp70 than RAW117-H10 cells [48]. In addition, RAW117-P cells contain more *gag* precursor, detected as a $M_r \sim 65\,000$ product, than RAW117-H10 cells. A mature p30 *gag* product could only be detected in RAW117-P cells (figure 2).

We also compared the expression of specific *abl*-encoded mRNA in RAW117 cells. Polyadenylated and nonadenylated mRNA can be separated on formaldehyde-treated agarose gels, transferred onto nitrocellulose paper, and hybridized with radiolabeled pAbl sub 9 plasmid containing the entire v-*abl* sequence. Both low and high metastatic RAW117 cells express a major species of mRNA of 6.2 kd that hybridizes specifically with the pAbl sub 9 probe, indicating the existence of *abl*-encoded mRNA. Equal amounts of *abl* encoded mRNA are present in low and high metastatic RAW117 cells, suggesting that RAW117 cells of low and high metastatic potential transcribe equally *abl*-specific mRNA (figure 3). Analyses of oncogene expression using probes for the oncogenes *fos* (no expression), *myc* (low expression),

and *myb* (low expression) indicate that these oncogenes are also expressed at similar levels, if expressed, in low and high metastatic potential RAW117 cells (data not shown).

Discussion

We have found that the capacity of AbLV-transformed RAW117 cells to metastasize is unrelated to their expression of a functional *abl* oncogene protein. Although the initial step in cellular transformation in this system may involve switching on the *abl* oncogene, and its expression may be related to the maintenance of transformation, *abl* expression is not required additionally for further steps in malignant progression, such as metastasis. This notion is in agreement with the data of Whitlock *et al.* [57], who showed that the progression of early passages of AbLV-transformed cell clones to more malignant growth behavior does not involve detectable changes in the concentration, phosphorylation or kinase activity of the *abl* oncogene protein. We did not detect differences in the electrophoretic, immunologic or enzymatic properties of *abl*-coded p160 in low and high metastatic potential RAW117 cells; therefore, differences in the amounts of p160 molecules or their activities in cell sublines of widely disparate metastatic phenotypes could not be detected in this system. However, our analyses cannot exclude minor changes in p160 sequence or activity.

Tumors induced by AbLV and cells transformed *in vitro* by AbLV can lose expression of the *abl* oncogene while maintaining their transformed phenotypes [23]. However, here we found that several RAW117 cell sublines and clones all express the *abl* oncogene at equivalent levels. In addition, AbLV-transformed cells can show expression of other oncogenes [28]. Therefore, we examined the expression of several other oncogenes in AbLV-transformed RAW117 cells. If expressed, these oncogenes were also expressed equivalently in low and high metastatic RAW117 cells, suggesting that differential oncogene expression is not involved in regulating RAW117 metastatic phenotype. Similar conclusions were recently reached independently by Kris *et al.* [27] using other murine tumor cell sublines of differing metastatic potentials. They found that B16 melanoma and UV-2237 fibrosarcoma cells express similar amounts of *Ki-ras*, the major oncogene in these cells, and *p21^{ras}* in both low and high metastatic potential cells.

In RAW117 cells the *abl*-coded product p160 does not vary in expression, but the MoLV coded products, such as gp70, p30, p15 etc. are progressively diminished in their expression with increasing metastatic potential. In highly metastatic RAW117-H10 cells these MoLV-coded proteins are barely detectable [47, 48] and their expression correlates inversely ($r=0.93$) with metastasis [41]. Therefore, the *abl* gene must be controlled differently from the MoLV genes, even though the *abl* gene is inserted into a helper MoLV genome to form the AbLV. In RAW117 cells the loss of cell surface antigenic components [33, 42, 48], decreased sensitivity to macrophage-mediated cytolysis and cytostasis [32, 49] and increased amounts of cell surface receptors for organ recognition [31] are related to high metastatic potential. The products that confer these properties do not appear to be controlled by the transformation-related *abl* oncogene in RAW117 lymphoma cells.

The question of whether oncogene expression at any level is required for the appearance of metastatic cells within a tumor cell population could not be answered by our experiments or those of Kris *et al.* [27]. However, Thorgeirsson *et al.* [55] have used transfection of 3T3 cells with the *ras* oncogene to produce *ras*-expressing

metastatic sarcomas. In contrast, oncogene expression in metastases of some human cancers, such as colorectal carcinomas, is low or non-existent compared to primary tumors from the same patients [1, 21]. It has been proposed that oncogenes may set in motion the generation of cellular phenotypic diversity and heterogeneity by generating changes in expression of regulatory genes or their promoters similar to the generation of diversity found in the multigene families that code for immunoglobulin molecules [36, 37]. Once tumor cells become malignant and undergo diversification, further changes in oncogene expression may not be critical for achieving the metastatic phenotype.

Acknowledgments

We thank H. Friedman for technical assistance. These studies were supported by grants from the Israel Cancer Research Foundation (to V. Rotter) and the U.S. National Cancer Institute (RO1-CA29571 to G. L. Nicolson). V. Rotter is recipient of the Norman and Helen Ashes Career Development Award.

References

- [1] ALBINO, A. P., LESTRANGE, R., OLIFF, A. I., FURTH, M. E., and OLD, L. J., 1984, Transforming *ras* genes from human melanoma: A manifestation of tumor heterogeneity? *Nature*, **308**, 69–72.
- [2] ALITALO, K., SCHWAB, M., LIN, C. C., VARMUS, H. E., and BISHOP, J. M., 1983, Homogeneously staining chromosomal regions contain amplified copies of an abundantly expressed cellular oncogene (*c-myc*) in malignant neuroendocrine cells from a human colon carcinoma. *Proceedings of the National Academy of Sciences, U.S.A.*, **80**, 1707–1711.
- [3] AUFRAY, D., and ROUGEON, F., 1980, Purification of mouse immunoglobulin heavy chain messenger RNAs from total myeloma tumor RNA. *European Journal of Biochemistry*, **107**, 303–314.
- [4] AVIV, H., and LEDER, P., 1972, Purification of biologically active globin messenger RNA by chromatography on oligothymidylic acid cellulose. *Proceedings of the National Academy of Sciences, U.S.A.*, **69**, 1408–1412.
- [5] BLAIR, D. G., OSKARSSON, M., WOOD, T. G., MCCLEMENTS, W. L., FISCHINGER, P. J., and VAN DE WOUDE, G., 1981, Activation of the transforming potential of a normal cell sequence: A molecular model for oncogenesis. *Science*, **212**, 941–943.
- [6] BRUNSON, K. W., and NICOLSON, G. L., 1978, Selection and biologic properties of malignant variants of a murine lymphosarcoma. *Journal of the National Cancer Institute*, **61**, 1499–1503.
- [7] CHAGANTI, R. S. K., 1983, The significance of chromosome change to neoplastic development. *Chromosome Mutation and Neoplasia*, edited by J. German (New York: Liss), pp. 359–396.
- [8] CHEN, T. R., 1977, *In situ* detection of mycoplasma contamination in cell cultures by fluorescent Hoechst 33258 stain. *Experimental Cell Research*, **104**, 255–262.
- [9] COLLET, M. S., and ERIKSON, R. L., 1978, Protein kinase activity associated with the avian sarcoma virus *Src* gene product. *Proceedings of the National Academy of Sciences, U.S.A.*, **75**, 2012–2024.
- [10] COOPER, G. M., 1982, Cellular transforming genes. *Science*, **218**, 801–806.
- [11] CURRAN, T., PETTERS, G., VAN BEVOREN, C., TEICH, N. M., and VERMA, I. M., 1982, FBJ murine osteosarcoma virus: Identification and molecular cloning of biologically active proviral DNA. *Journal of Virology*, **44**, 674–682.
- [12] DALLA-FAVERA, R., MARTINOTTI, S., GALLO, R. C., ERIKSON, J., and CROCE, C. M., 1983, Translocation and rearrangements of the *c-myc* oncogenes locus in human undifferentiated B-cell lymphomas. *Science*, **219**, 963–967.
- [13] DALLA-FAVERA, R., WONG-STAAI, F., and GALLO, R. C., 1982, Oncogene amplification in promyelocytic leukemia cell line HL-60 and primary leukaemic cells of the same patient. *Nature*, **298**, 61–63.

- [14] DEFEO, D., GONDA, M. A., YOUNG, H. A., CHANG, E. H., LOWY, D. R., SCOLNICK, E. L. M., and ELLIS, R. W., 1981, Analysis of divergent rat genomic clones homologous to the transforming gene of Harvey murine sarcoma virus. *Proceedings of the National Academy of Sciences, U.S.A.*, **78**, 3328–3332.
- [15] DENHARDT, D., 1966, A membrane-filter technique for the detection of complementary DNA. *Biophysical and Biochemical Research Communication*, **23**, 641–646.
- [16] DER, C. J., KRONIRIS, T. G., and COOPER, G. M., 1982, Transforming genes of human bladder and lung carcinoma cell lines are homologous to the *ras* genes of Harvey and Kisten sarcoma viruses. *Proceedings of the National Academy of Sciences, U.S.A.*, **79**, 3637–3640.
- [17] DIAMOND, A., DEVINE, J. M., and COOPER, G. M., 1983, Nucleotide sequence of a human *B-lym* transforming gene activated in Burkitt's lymphoma. *Science*, **225**, 516–519.
- [18] ERIKSON, J., AR-RUSHDI, A., DRWINGA, H. L., NOWELL, P. C., and CROCE, C. M., 1983, Transcriptional activation of the translocated *c-myc* oncogenes in Burkitt lymphoma. *Proceedings of the National Academy of Sciences, U.S.A.*, **80**, 820–824.
- [19] FIDLER, I. J., and HART, I. R., 1982, Biological diversity in metastatic neoplasms origins and implications. *Science*, **217**, 998–1003.
- [20] FRANCHINI, G., WONG-STAAI, F., BALUDA, M. A., LENGEL, C., and TRONICK, S. R., 1983, Structural organization and expression of human DNA sequences related to the transforming gene of avian myeloblastosis virus. *Proceedings of the National Academy of Sciences, U.S.A.*, **80**, 7385–7389.
- [21] GALLICK, G. E., KURZROCK, R., KLOETZER, W. S., ARLINGHAUS, R. B., and GUTTERMAN, J. U., 1985, Expression of the p21 *ras* gene products in fresh primary and metastatic human colorectal tumors. *Proceedings of the National Academy of Sciences, U.S.A.*, **82**, 1795–1799.
- [22] GLISIN, W., CRKVENJAKOV, R., and BYUS, C., 1974, Ribonucleic acid isolated by cesium chloride centrifugation. *Biochemistry*, **13**, 2633–2637.
- [23] GRUNWALD, D. J., DALE, B., DUDLEY, J., LAMPH, W., SUGDEN, B., OZANNE, B., and KISSER, R., 1982, Loss of viral gene expression and retention of tumorigenicity by Abelson lymphoma cells. *Journal of Virology*, **43**, 92–103.
- [24] HAYWARD, W. S., NEEL, B. G., and ASTRIN, S. M., 1981, Activation of a cellular *onc* gene by promoter insertion in ALV-induced lymphoid leukemia. *Nature*, **290**, 475–480.
- [25] HEISTERKAMP, N., STEPHENSON, J. R., GROFFEN, J., HANSEN, P. F., DEKLEIN, A., BARTRAM, C. T., and GROSVELD, G., 1983, Localization of the *c-abl* oncogene adjacent to a translocation break point in chronic myelocytic leukemia. *Nature*, **306**, 239–242.
- [26] KESSLER, S. W., 1975, Rapid isolation of antigens from cells with a staphylococcal protein A antibody adsorbent: Parameters of the interaction of antibody-antigen complex with protein A. *Journal of Immunology*, **115**, 1617–1624.
- [27] KRIS, R. M., AVIVI, A., BAR-ELI, M., ALON, Y., CARNI, P., SCHLESSINGER, J., and RAZ, A., 1985, The expression of *KI-RAS* oncogene in tumor cell variants exhibiting different metastatic capabilities. *International Journal of Cancer*, **35**, 227–230.
- [28] LANE, M. A., NEARY, D., and COOPER, G. M., 1982, Activation of a cellular transforming gene in tumours induced by Abelson murine leukemia virus. *Nature*, **300**, 659–660.
- [29] LAIT, S. A., GOFF, S. P., TABIN, C. J., PASKIND, M., WANG, J. Y. J., and BALTIMORE, D., 1983, Cloning and analysis of reverse transcript p160 genomes of Abelson murine leukemia virus. *Journal of Virology*, **45**, 1195–1199.
- [30] LAEMMLI, U. K., 1970, Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, **227**, 680–685.
- [31] MCGUIRE, E. J., MASCALI, J. J., GRADY, S. R., and NICOLSON, G. L., 1984, Involvement of cell-cell adhesion molecules in liver colonization by metastatic murine lymphoma/lymphosarcoma variants. *Clinical and Experimental Metastasis*, **2**, 213–222.
- [32] MINER, K. M., and NICOLSON, G. L., 1983, Differences in the sensitivities of murine metastatic lymphoma/lymphosarcoma cells to macrophage-mediated cytolysis and/or cytostasis. *Cancer Research*, **43**, 2063–2071.
- [33] MINER, K. M., WALTER, H., and NICOLSON, G. L., 1981, Subfractionation of malignant variants of metastatic murine lymphosarcoma cells by countercurrent distribution in two-polymer aqueous phases. *Biochemistry*, **20**, 6244–6250.
- [34] NICOLSON, G. L., 1982, Cancer metastasis: Organ colonization and the cell surface properties of malignant cells. *Biochimica et Biophysica Acta*, **695**, 113–176.

- [35] NICOLSON, G. L., 1984, Cell surface molecules and tumor metastasis. Regulation of metastatic diversity. *Experimental Cell Research*, **150**, 3–22.
- [36] NICOLSON, G. L., 1984, Generation of phenotypic diversity and progression in metastatic tumors. *Cancer Metastasis Review*, **3**, 25–42.
- [37] NICOLSON, G. L., 1984, Tumor progression, oncogenes and the evolution of metastatic phenotypic diversity. *Clinical and Experimental Metastasis*, **2**, 85–105.
- [38] NICOLSON, G. L., and POSTE, G., 1982, Tumor cell diversity and host responses in cancer metastasis. I. Properties of metastatic cells. *Current Problems in Cancer*, **7(6)**, 1–83.
- [39] NICOLSON, G. L., and POSTE, G., 1983, Tumor cell diversity and host responses in cancer metastasis. II. Host immune responses and therapy of metastases. *Current Problems in Cancer*, **7(7)**, 1–43.
- [40] NICOLSON, G. L., and POSTE, G., 1983, Tumor implantation and invasion at metastatic sites. *International Review of Experimental Pathology*, **25**, 77–181.
- [41] NICOLSON, G. L., MASCALI, J. J., and MCGUIRE, E. J., 1982, Metastatic RAW117 lymphosarcoma as a model for malignant–normal cell interactions. Possible roles for cell surface antigens in determining the quantity and location of secondary tumors. *Oncodevelopmental Biology and Medicine*, **4**, 149–159.
- [42] NICOLSON, G. L., READING, C. L., and BRUNSON, K. W., 1980, Blood-borne tumor metastasis: Some properties of selected tumor cell variants of differing malignancies. *Tumor Progression*, edited by R. G. Crispen (Amsterdam: Elsevier/North Holland), pp. 31–48.
- [43] NOWELL, P. C., 1983, Tumor progression and clonal evolution: the role of genetic instability. In: J. German (ed.), *Chromosome Mutation and Neoplasia*, edited by J. German (New York: Liss), pp. 413–432.
- [44] POSTE, G., 1982, Experimental systems for analysis of the malignant phenotype. *Cancer Metastasis Reviews*, **1**, 141–199.
- [45] POSTE, G., and FIDLER, I. J., 1980, The pathogenesis of cancer metastasis. *Nature*, **283**, 139–146.
- [46] RASCHKE, W. C., RALPH, P., WATSON, J., SKLAR, M., and COON, H., 1975, Oncogenic transformation of murine lymphoid cells by *in vitro* infection with Abelson leukemia virus. *Journal of the National Cancer Institute*, **54**, 1249–1253.
- [47] READING, C. L., BELLONI, P. N., and NICOLSON, G. L., 1980, Selection and *in vivo* properties of lectin-attachment variants of malignant murine lymphosarcoma cell lines. *Journal of the National Cancer Institute*, **64**, 1241–1249.
- [48] READING, C. L., BRUNSON, K. W., TORRAINI, M., and NICOLSON, G. L., 1980, Malignancies of metastatic murine lymphosarcoma cell lines and clones correlate with decreased cell surface display of RNA tumor virus envelope glycoprotein gp70. *Proceedings of the National Academy of Sciences, U.S.A.*, **77**, 5943–5947.
- [49] READING, C. L., KRAEMER, P. M., MINER, K. M., and NICOLSON, G. L., 1983, *In vivo* and *in vitro* properties of malignant variants of RAW117 metastatic murine lymphoma/lymphosarcoma. *Clinical and Experimental Metastasis*, **1**, 135–151.
- [50] REYNOLDS, F. H., SACKS, T. I., DEOBAGHAR, D. H., and STEPHENSON, J. R., 1978, Cells nonproductively transformed by Abelson–Murine Leukemia Virus express a high molecular weight polypeptide containing structural and nonstructural components. *Proceedings of the National Academy of Sciences, U.S.A.*, **75**, 3974–3978.
- [51] RIGBY, P. W., DIECKMANN, M., RHODES, C., and BERG, P., 1977, Labeling deoxyribonucleic acid to high specific activity *in vitro* by nick translation with DNA polymerase I. *Journal of Molecular Biology*, **113**, 237–251.
- [52] ROSENBERG, N., and WITTE, O. N., 1980, Abelson murine leukemia virus mutants with alteration in the virus specific p120 molecule. *Journal of Virology*, **33**, 340–348.
- [53] SANTOS, E., TRONICK, S. R., AARONSON, S. A., PULCIANI, S., and BARBACID, M., 1982, T24 human bladder carcinoma oncogene is an activated form of the normal human homologue of BALB- and Harvey-MSV transforming genes. *Nature*, **298**, 343–347.
- [54] THOMAS, P. S., 1980, Hybridization of denatured RNA and small DNA fragments transferred to nitrocellulose. *Proceedings of the National Academy of Sciences, U.S.A.*, **77**, 5201–5205.
- [55] THORGEIRSSON, U. P., TURPEENNIEMI-HUJANEN, T., WILLIAMS, J. E., WESTEIN, E. H., HEILMAN, C. A., TALMADGE, J. E., and LIOTTA, L. A., 1985, NIH/3T3 cells transfected

- with human tumor DNA containing activated *ras* oncogenes express the metastatic phenotype in nude mice. *Molecular and Cellular Biology*, **5**, 259–262.
- [56] WANG, J. Y. J., and BALTIMORE, D., 1983, Cellular RNA homologous to the Abelson murine leukemia virus transforming gene: Expression and relationship to the viral sequence. *Molecular and Cellular Biology*, **3**, 773–779.
- [57] WHITLOCK, C. A., ZIEGLER, S. F., and WITTE, O. N., 1983, Progression of the transformed phenotype in clonal lines of Abelson virus infected lymphocytes. *Molecular Cellular Biology*, **3**, 596–604.
- [58] WITTE, O. N., ROSENBERG, N., PASKIND, M., SHIELDS, A., and BALTIMORE, D., 1978, Identification of an Abelson murine leukemia virus-encoded protein present in transformed fibroblast and lymphoid cells. *Proceedings of the National Academy of Sciences, U.S.A.*, **75**, 2488–2492.
- [59] WITTE, O. N., ROSENBERG, N., and BALTIMORE, D., 1979, Preparation of syngeneic tumor regressor serum reactive with the unique determinants of the Abelson murine leukemia virus-encoded p120 protein at the cell surface. *Journal of Virology*, **31**, 776–784.
- [60] WITTE, O. N., DASGUPTA, A., and BALTIMORE, D., 1980, Abelson murine leukemia virus protein is phosphorylated *in vitro* to form phosphotyrosine. *Nature*, **283**, 826–831.