

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/5577788>

Recent overview of the Mg branch of the tetrapyrrole biosynthesis leading to chlorophylls

ARTICLE *in* PHOTOSYNTHESIS RESEARCH · JUNE 2008

Impact Factor: 3.5 · DOI: 10.1007/s11120-008-9291-4 · Source: PubMed

CITATIONS

60

READS

18

1 AUTHOR:



[Tatsuru Masuda](#)

The University of Tokyo

103 PUBLICATIONS **3,847** CITATIONS

SEE PROFILE

Recent overview of the Mg branch of the tetrapyrrole biosynthesis leading to chlorophylls

Tatsuru Masuda

Received: 11 October 2007 / Accepted: 29 January 2008 / Published online: 14 February 2008
© Springer Science+Business Media B.V. 2008

Abstract In plants, chlorophylls (chlorophyll *a* and chlorophyll *b*) are the most abundant tetrapyrrole molecules and are essential for photosynthesis. The first committed step of chlorophyll biosynthesis is the insertion of Mg²⁺ into protoporphyrin IX, and thus subsequent steps of the biosynthesis are called the Mg branch. As the Mg branch in higher plants is complex, it was not until the last decade—after many years of intensive research—that most of the genes encoding the enzymes for the pathway were identified. Biochemical and molecular genetic analyses have certainly modified the classic metabolic map of tetrapyrrole biosynthesis, and only recently have the molecular mechanisms of regulatory pathways governing chlorophyll metabolism been elucidated. As a result, novel functions of tetrapyrroles and biosynthetic enzymes have been proposed. In this review, I summarize the recent findings on enzymes involved in the Mg branch, mainly in higher plants.

Keywords Chlorophyll · Chloroplast · Mg branch · Tetrapyrrole · Photosynthesis

Abbreviations

AAA	ATPase associated with various cellular activities
ALA	5-aminolevulinic acid
Bchl	Bacteriochlorophyll
CAO	Chlorophyllide <i>a</i> oxygenase
Chl	Chlorophyll
DVR	3,8-Divinyl Pchlide <i>a</i> 8-vinyl reductase

DPOR	Light-independent protochlorophyllide oxidoreductase
GFP	Green fluorescent protein
<i>gun</i>	Genome uncoupled
LHC	Light-harvesting chlorophyll <i>a/b</i> -protein complex
LHCII	Light-harvesting chlorophyll <i>a/b</i> -protein complex of photosystem II
LHPP	Light-harvesting protochlorophyllide <i>a/b</i> -binding protein complex
Pchlde	Protochlorophyllide
PLB	Prolamellar body
POR	Light-dependent NADPH:protochlorophyllide oxidoreductase
Ptc	Protochlorophyllide translocon complex
SAM	<i>S</i> -adenosyl-L-methionine
ZnPP	Zinc-protopheophorbide

Introduction

Chlorophylls (Chls) and bacteriochlorophylls (Bchls) are essential molecules for photosynthesis. They are responsible for harvesting and transferring solar energy in antenna systems, and for charge separation and electron transport in reaction centers. Chls and Bchls are Mg²⁺-containing tetrapyrrole compounds. In plants, algae and many bacteria, all of the tetrapyrroles, including Chls and heme, originate from a common biosynthetic pathway. They are synthesized from glutamyl-tRNA^{glu} via 5-aminolevulinic acid (ALA), a linear five-carbon molecule. The formation of ALA is the first committed step of tetrapyrrole biosynthesis, and is therefore assumed to be a regulatory point for the synthesis of total amounts of

T. Masuda (✉)
Graduate School of Arts and Sciences, The University of Tokyo,
Komaba 3-8-1, Meguro-ku, Tokyo 153-8902, Japan
e-mail: ctmasuda@mail.ecc.u-tokyo.ac.jp

tetrapyrroles (Beale 1999). Actually, the ALA synthesizing activity is subjected to feedback regulation by heme and repressed by a regulatory protein, FLU, in the dark (Meskauskiene et al. 2001; Goslings et al. 2004). Two molecules of ALA are then condensed to form a pyrrole molecule. Subsequently, four pyrrole molecules are polymerized to form a linear pyrrole that subsequently undergoes cyclization to yield a macrocyclic tetrapyrrole. The branch point of the Chl and heme biosynthetic pathway is protoporphyrin IX, a closed macrocycle lacking a chelated ion. The first committed step of Chl/Bchl biosynthesis is the insertion of Mg^{2+} into protoporphyrin IX, and thus subsequent steps in this pathway are called the Mg branch (Fig. 1).

The enzymes responsible for the Mg branch have been identified and characterized primarily via molecular genetic analyses of the photosynthetic gene cluster in *Rhodobacter (Rba.) capsulatus* (Bollivar et al. 1994b) and *Rba. sphaeroides* (Coomber et al. 1990). These gene clusters comprise approximately 45 kbp and contain most of the genes encoding structural and regulatory proteins for assembly of the photosynthetic apparatus as well as enzymes involved in Bchl biosynthesis from protoporphyrin IX. Directed mutagenesis and subsequent heterologous expression in *E. coli* have identified many genes encoding these Bchl biosynthetic enzymes. In addition, molecular genetic studies in plants have identified many genes encoding Chl biosynthetic enzymes, many of which are orthologous to the bacterial genes. Owing to the complexity of this pathway in higher plants, only in the last decade have the genes encoding the enzymes of the Mg branch been identified—and then only after many years of intensive research. Biochemical and molecular genetic analyses have modified the classical metabolic map of tetrapyrrole biosynthesis, but only recently have the molecular mechanisms of regulatory pathways governing Chl metabolism been elucidated. These insights have enabled researchers to propose novel functions for tetrapyrroles and their associated biosynthetic enzymes. Thus, it is an opportune time to review and redraw the metabolic pathway for Chl biosynthesis. The most recently updated biosynthetic pathway is shown in Fig. 1.

In this review, I summarize the recent findings on enzymes involved in the Mg branch, focusing mainly in higher plants. According to the order of function of each enzyme in the metabolic pathway, I review the gene identification, enzyme biochemistry, expression and localization, and regulation. Readers may also refer to recent comprehensive reviews on this field (Beale 1999; Vavilin and Vermaas 2002; Rebiez et al. 2003; Eckhardt et al. 2004; Moulin and Smith 2005; Bollivar 2006; Tanaka and Tanaka 2006, 2007).


Mg-chelatase

Mg-chelatase catalyzes the insertion of Mg^{2+} into protoporphyrin IX. This is the first committed step in Chl biosynthesis with the enzymes prior to this step also being shared with the heme biosynthetic pathway. Although Mg-chelatase and ferrochelatase catalyze virtually the same reaction—for heme, the insertion of Fe^{2+} into protoporphyrin IX to form protoheme—the structures of these enzymes are completely different. Ferrochelatase is a single-subunit enzyme encoded by a single gene and does not require a cofactor for catalysis. By contrast, Mg-chelatase requires ATP for catalysis and is a heterologous complex consisting of 40-, 70-, and 140-kDa subunits, designated I, D and H subunits, respectively. These respective subunits are known as BchI, BchD, and BchH in Bchl-producing organisms and ChlI, ChlD, and ChlH in Chl-producing organisms (Walker and Willows 1997). Mg-chelatase, a key enzyme at the branch point between the heme and Chl biosynthetic pathways, has been extensively studied during the last decade. Structural and functional analyses of individual subunits have provided overall mechanisms for the reaction and enzyme regulation (Walker and Willows 1997; Reid and Hunter 2002; Willows 2003).

Identification and characterization of genes encoding subunits of Mg-chelatase

Genes encoding subunits of Mg-chelatase were first proposed by analysis of directional mutagenesis of the photosynthetic gene cluster of *Rhodobacter*, which accumulated protoporphyrin IX (Coomber et al. 1990; Bollivar et al. 1994b). Reconstitution of Mg-chelatase activity with recombinant proteins demonstrated that the chelatase is composed by three subunits both in Bchl-producing prokaryotes (Gibson et al. 1995; Petersen et al. 1998; Willows and Beale 1998) and Chl-producing cyanobacterium *Synechocystis* PCC6803 (Jensen et al. 1996a). In 1996, natural photosynthesis using Zn-containing Bchl *a* was discovered in the aerobic acidophilic purple photosynthetic bacterium, *Acidiphilium rubrum* (Wakao et al. 1996). This bacterium predominantly produces Zn-containing Bchl *a*, resulting from Zn^{2+} insertion (instead of Mg^{2+}) as the central metal of the tetrapyrrole ring; this Bchl *a* is associated with fully active reaction centers and light-harvesting complexes analogous to those of other purple photosynthetic bacteria (Wakao et al. 1996). Analysis of the photosynthetic gene cluster of *A. rubrum* revealed that this bacterium contains genes encoding three subunits of the functional Mg-chelatase, and thus, substitution with Zn^{2+} may occur after Mg-insertion probably for acclimation to an acidic environment (Masuda et al. 1999). These prokaryotic

genes encoding subunits of Mg-chelatase have been used to analyze the early evolution of photosynthesis. Phylogenetic analysis of these genes suggested that purple bacteria are the earliest photosynthetic lineage to have emerged (Xiong et al. 2000).



amounts of Chl using a second gene, *ChlI2* (Rissler et al. 2002). Owing to a Chl-deficient yellow phenotype and ALA-dependent accumulation of protoporphyrin IX, barley mutants at three genetic loci, termed *xantha-f*, *xantha-g*, and *xantha-h*, were characterized as Mg-chelatase mutants corresponding to the gene locus encoding the H, D, and I subunits, respectively (Jensen et al. 1996b; Petersen et al. 1999b). Three semidominant alleles of the *xantha-h* locus, originally isolated as pale green *chlorina* mutants, each have single missense mutations (Hansson et al. 1999). *ChlI* was identified from soybean as a stromal protein of chloroplasts (Nakayama et al. 1995). In maize, the gene *Oil Yellow1* (*Oy1*) was identified as subunit I (Sawers et al. 2006). In the *Oy1* mutant, amino acid substitutions resulted in a semidominant phenotype probably via competitive inhibition of wild type I subunit. In rice, map-based cloning identified *Chlorina-1* and *Chlorina-9* as genes encoding subunits ChlD and ChlI, respectively (Zhang et al. 2006). These genes occur in the nuclear genome. *ChlI* genes have also been identified from plastid genomes of green (Orsat et al. 1992), red, and cryptomonad algae (Douglas and Reith 1993).

Mutants of *ChlH* have been described in *Antirrhinum majus* (Hudson et al. 1993). *ChlH* was identified from soybean (Nakayama et al. 1998). *ChlD* was identified from tobacco by homology, and recombinant ChlD along with ChlH and ChlI subunits reconstituted active Mg-chelatase (Papenbrock et al. 1997). In *Chlamydomonas reinhardtii*, *ChlH* was identified and defective mutants were characterized (Chekounova et al. 2001). *Arabidopsis* mutants, *cch*, and *gun5* (genome uncoupled 5), were identified as containing single missense mutations in the H subunit, *ChlH* (Mochizuki et al. 2001). The mutant *gun5* was found to impart aberrant regulation of chloroplast-to-nucleus signal transduction, suggesting that Mg-porphyrins are involved in intracellular signaling (see below).

On the basis of functional and genetic studies and results from sequencing projects, chelatases have been categorized into three main classes: (I) ATP-dependent heterotrimeric chelatases, (II) ATP-independent chelatases, and (III) multifunctional homodimeric chelatases (Brindley et al. 2003; Al-Karadaghi et al. 2006). Co-chelatase inserts Co^{2+} into hydrogenobyrinic acid *a,c*-diamide, a tetrapyrrole precursor of vitamin B₁₂. Both Co-chelatase and Mg-chelatase are type I chelatases, whereas ferrochelatase is a type II chelatase. Co-chelatase consists of a 140-kDa subunit (CobN) that binds tetrapyrrole, and a 450-kDa tetramer of the subunits CobS and CobT (38 and 80 kDa, respectively) that binds ATP (Debussche et al. 1992). As such, Co-chelatase is somewhat similar to Mg-chelatase; there is a significant sequence similarity between CobN and subunit H, although no similarity between other subunits have been identified apart from their respective

molecular masses and a consensus ATP-binding site in CobS and subunit I. More recently, a metal ion-dependent adhesion site (MIDAS motif), which is involved in binding of subunits (see below), was found in both CobT and subunit D, suggesting important similarity between these two subunits as well as between the Mg- and Co-chelatases (Fodje et al. 2001).

Surprisingly, recent work proposed that ChlH is a receptor for abscisic acid, which regulates stomata opening and seed development (Shen et al. 2006). It was shown that *A. thaliana* ChlH specifically binds abscisic acid and mediates its signaling as a positive regulator of seed germination, post-germination growth, and stomatal movement.


Characterization of Mg-chelatase activity

A model of the catalytic cycle of Mg-chelatase is shown in Fig. 2. Earlier work on chloroplast extracts suggested that Mg chelation proceeded by a two-step reaction: an enzyme-activation step, followed by a Mg^{2+} insertion step (Walker and Weinstein 1994). Subsequent work using recombinant subunits encoded by genes from photosynthetic bacteria showed that the activation step requires ATP, Mg^{2+} and subunits I and D for the formation of a ternary I–D–Mg–ATP complex (Willows et al. 1996; Gibson et al. 1999). This step does not require the hydrolysis of ATP, as ADP or nonhydrolyzable ATP analogs such as 5'-[γ -thio]triphosphate (ATP γ S) can substitute for ATP (Jensen et al. 1999a; Lake et al. 2004). The Mg-insertion step also requires ATP, although at a lower concentration than needed for activation, and this cannot be replaced by ATP γ S. Thus, ATP hydrolysis is required for the Mg insertion step (Jensen et al. 1999a). The I–D–Mg–ATP complex uses Mg^{2+} and the porphyrin-bound H subunit as the substrate for the Mg insertion step, with ATP hydrolysis driving the reaction.

The I subunits are classified as members of the AAA⁺-proteins (ATPase associated with various cellular activities), by sequence (Neuwall et al. 1999), by conservation of a characteristic fold (Lecroff et al. 2000), and by structure analysis (Fodje et al. 2001). The I subunit is an ATPase (Gibson et al. 1999; Jensen et al. 1999a). The ATPase activity of subunit I is repressed when this subunit forms a complex with subunit D (Gibson et al. 1999; Jensen et al. 1999a), although the opposite observation also has been made (Hansson and Kannangara 1997). The N-terminal halves of subunits D and I share high sequence similarity, whereas the C-terminal half of subunit D includes a metal ion coordination motif (MIDAS) characteristic of the integrin I domain (Fodje et al. 2001). The similarity between the D and I subunit N termini suggests that the D subunit is also an

Fig. 2 Model of the catalytic cycle of Mg-chelatase.

Activation step: In a Mg^{2+} and ATP-dependent process, six I subunits are assembled into a hexameric ring structure, while six D subunits form a hexameric ring in an ATP-independent manner. The two-tiered hexameric ring forms the Mg-ATP-I-D complex. It is suggested that the ATPase activity of subunit I is inhibited in this complex due to binding of the integrin I domain of subunit D to the integrin I binding domain of subunit I. Meanwhile, subunit H binds to protoporphyrin IX and most likely also the Mg^{2+} substrate—as this subunit is considered the catalytic subunit—to form the Mg-H-protoporphyrin IX complex. Chelation step: ATP hydrolysis is triggered upon switching the binding of the integrin I domain of subunit D to the integrin I binding domain of subunit H. After the formation of Mg-protoporphyrin IX, the complex disassembles. Here, subunit H is shown docking to the D subunits, although it remains unknown if subunit H docks to the I or D side of the complex



AAA⁺-protein, although no ATPase activity has been detected from this subunit (Hansson and Kannangara 1997; Jensen et al. 1999a; Petersen et al. 1999a). An acidic, proline-rich region linking the N- and C-terminal domains of the D subunit may contribute to the association of subunit D with subunit I (Fodje et al. 2001). In tobacco, a special motif (EK-X18-R) has been identified on both sides of the linker domain of subunit D. A region of approximately 110 amino acid residues spanning these special motifs and the linker domain enables interaction of the D subunit with the I subunit and maintenance of the Mg-chelatase activity (Gräfe et al. 1999). Meanwhile, the integrin I domain that is located on the C terminus of subunit D is proposed to bind to the integrin I domain-binding sequence motifs of subunits I and H (Fodje et al. 2001). The binding of subunit D to subunit I may block the ATPase activity of subunit I. It has been proposed that a conformational transition of subunit D is induced upon I-D complex formation and subsequent binding to subunit H may trigger porphyrin metallation, and also relieve the block of the ATP-binding site of subunit I by subunit D, thereby

allowing ATP hydrolysis (Fodje et al. 2001). AAA⁺-proteins generally form oligomeric ring structures, and subunit I forms high-molecular-mass aggregates in the presence of ATP and $MgCl_2$ (Jensen et al. 1998; Gibson et al. 1999). Negative staining electron microscopy showed that I subunits from *Rba. capsulatus* form a hexameric ring (Willows et al. 2004) and that I subunits from *Synechocystis* PCC6803 form a heptameric ring (Reid et al. 2003).

In *Arabidopsis*, most of the homozygous *CHL1* mutants of *Arabidopsis thaliana*, such as *ch42-1* (Fisherova 1975), *cs* (*ch42-2*) (Koncz et al. 1990), *ch42-3* (Rissler et al. 2002), are recessive and have a pale-green phenotype. The *cs* mutant has a T-DNA insertion in the 3'-terminal end of *CHL1*, resulting in *CHL1* malfunction because the C-terminal end is extended (Koncz et al. 1990). These mutants could still make small amounts of Chl probably using a second gene *Chl2* (Rissler et al. 2002). In wild-type plants, *CHL1* and *CHL2* mRNAs accumulate to similar levels, but *CHL2* protein is undetectable in wild type and the *ch42-3* mutant. It has been proposed that unusual post-translational protein instability

limits the function of CHLI2 in the Mg-chelatase complex (Rissler et al. 2002). Meanwhile, a semidominant mutant of *CHLI1* (*aci5*) has been identified as a point mutant with a single amino acid substitution (D240N) (Soldatova et al. 2005). A heterozygous *cs/aci5* hybrid plant is completely *albino*, although the plant expresses the mutant genes encoded by both *cs* and *aci5* alleles (Apchelimov et al. 2007). Thus, *cs* allele-induced C-terminal extension of CHLI may perturb the assembly of the CHLI ring structure, whereas displacement by *aci5*-encoded CHLI causes inactivation of the Mg-chelatase complex because an essential residue in the catalytic site of CHLI1 has been substituted (Apchelimov et al. 2007). Since *CHLI2* cannot compensate for the *albino* phenotype in this hybrid plant, it was proposed that CHLI2 is not functional in the Mg-chelatase complex. It was suggested that CHLI2 lost the ability to form effective ring-like structures because of mutations that changed amino acids at the C-terminal end (Apchelimov et al. 2007). Very recently, the physiological contribution of CHLI2 was examined using a *chli2* T-DNA knockout line (Kobayashi et al. submitted). This article shows that the *chli2* mutation is semidominant on a homozygous *cs* background, revealing that although *CHLI2* plays a limited role in chlorophyll biosynthesis, this subunit certainly contributes to the assembly of the Mg-chelatase complex.

Semidominant and recessive mutants deficient in subunit I have been identified in barley *xantha-h* (Hansson et al. 1999; Hansson et al. 2002) and maize (Sawers et al. 2006). The semidominant mutations were all point mutations resulting in changes in single amino acid residues, whereas the recessive mutations were deficient in transcription. In vitro analysis of semidominant mutants of subunit I in expression systems for *Rba. capsulatus* BchI and *Synechocystis* ChII showed reduced or no ATPase activity. The defective protein could not contribute to the Mg-chelatase activity and rather inhibited when mixed with wild-type I subunits, explaining the dominant effect of the mutated protein (Hansson et al. 2002; Sawers et al. 2006). Thus, dominant mutations in the I subunits function in a cooperative manner within the hexameric complex (Hansson et al. 2002). Subunit D could not be detected in any barley *xantha-h* mutants, suggesting that a functional I subunit is required to maintain the D subunit (Lake et al. 2004). Subunit D also forms ATP-independent oligomeric structures and is classified as an AAA⁺ protein (Axelsson et al. 2006). All barley *xantha-g* mutants defective in D subunits that have been identified were recessive. An in-vitro study of mutant proteins of subunit D that can be cooperatively incorporated into the oligomeric structure of *Rba. capsulatus* BchD protein resulted in lower Mg-chelatase activity (Axelsson et al. 2006). This cooperative behavior suggests that the D oligomer takes an active part

in the conformational dynamics among the subunits of the enzyme (Axelsson et al. 2006).

The H subunit is regarded as the catalytic subunit since Mg chelation takes place on this subunit (Reid and Hunter 2002). Porphyrin binding analysis has demonstrated that subunit H is the porphyrin binding subunit (Willows et al. 1996; Jensen et al. 1998; Willows and Beale 1998). The binding of deuteroporphyrin IX, a more water-soluble analog of the biological substrate, to subunit H caused quenching of Trp fluorescence, allowing the binding constant to be determined (4.0 μ M for *Synechocystis* ChIH and 0.75 μ M for *Rba. sphaeroides* BchH) (Karger et al. 2001). Compared to free deuteroporphyrin IX, the red shifts and peak broadening of the spectra of the H subunit-bound deuteroporphyrin IX implied a slightly deformed porphyrin configuration (Karger et al. 2001) that may permit ready chelation (Reid and Hunter 2002). ATPase activities for the various H subunits have been reported for *Rba. sphaeroides* (Hansson and Kannangara 1997), *Chlorobium vibrioforme* (Petersen et al. 1999a) and *Synechocystis* PCC6803 (Jensen et al. 1999a), but a recent study suggests that the H subunit ATPase activity is an artifact of a contaminating *E. coli* protein (Sirijovski et al. 2006).

In *Synechocystis*, the Mg-chelatase-catalyzed reaction requires hydrolysis of ~ 15 Mg-ATP, and therefore the chelation reaction is energetically unfavorable (Reid and Hunter 2004). The ATPase activity of isolated I subunit is substantially lower than that of the holo-chelatase. Mg²⁺ activates the Mg-chelatase by increasing both the velocity of its reaction and specificity for the substrates, Mg-ATP and porphyrin, possibly through a conformational change (Reid and Hunter 2004).

Regulation of Mg-chelatase

Mg-chelatase is subjected to several forms of regulation. The major ones are transcriptional regulation of the H subunit and post-translational modifications of the subunit that affect the overall activity.

For the expression of each subunit of Mg-chelatase, upregulation of *ChII* and *ChIH* expression during greening of etiolated tissues has been reported in barley (Jensen et al. 1996b) and soybean (Nakayama et al. 1998). *ChII* is constitutively expressed in mature green tissues of barley (Jensen et al. 1996b) and *A. thaliana* (Gibson et al. 1996), and the levels of ChII mRNA and protein also remain constant in soybean (Nakayama et al. 1998). *ChIH* expression follows a circadian rhythm in barley (Jensen et al. 1996b) and in soybean with concomitant dramatic changes of ChIH protein, which peaks during the light phase (Nakayama et al. 1998). Transcriptome analysis that covers the majority of

tetrapyrrole biosynthetic genes of *Arabidopsis* shows that *ChlH* is one of the key regulatory genes, which is coordinately regulated by light and circadian rhythm with other key regulatory genes, such as *HEMA1* (encoding glutamyl-tRNA reductase), *CRD1* (*CHL27*) (encoding a subunit of Mg-protoporphyrin IX monomethyl ester cyclase), and *CAO* (encoding chlorophyllide *a* oxygenase) (Matsumoto et al. 2004). From the gene expression profiles, *CHL11*, *CHL12* and *CHLD* were involved in the same group, which are gradually upregulated by light during greening and obeyed diurnal rhythm but not circadian oscillation in matured leaves (Matsumoto et al. 2004). Co-expression of these genes are confirmed by recently developed databases of *Arabidopsis* microarray data, such as the AtGenExpress Visualization Tool (AVT) (<http://jsp.weigelworld.org/expviz/expviz.jsp>) and the ATTED-II (*A. thaliana* trans-factor and cis-element prediction database; <http://www.atted.bio.titech.ac.jp>) (Obayashi et al. 2007). Therefore, in *Arabidopsis*, it is assumed that the transcriptional control *CHLH* is a key regulatory point of Mg-chelatase activity, and a primary determinant of the level of active Mg-chelatase holo-complex, as well as orchestrating activities with other regulatory enzymes of tetrapyrrole biosynthesis. Meanwhile, the expression of *CHL11*, *CHL12*, and *CHLD* is regulated in a similar manner: they are induced by light during chloroplast development and are then under the control of a diurnal rhythm to maintain constant protein levels in chloroplasts. Recently, Shen et al. (2006) showed that *Arabidopsis* ChlH is ubiquitously expressed in both green and non-green tissues to perceive the abscisic acid signal at the whole-plant level, an expression profile that is quite different from previously published results. Future work must clarify whether the CHLH protein exists alone in non-photosynthetic plastids and functions as a receptor of abscisic acid, separate from its role in tetrapyrrole biosynthesis. In tobacco, *ChlH* and *ChlI* follow a similar circadian profile of expression, but *ChlD* has an inverse expression profile with maximal mRNA levels in the dark phase (Papenbrock et al. 1999). The difference of gene expression profiles are attributable to distinct plant species or developmental conditions.

Mg-chelatase activity is regulated by GUN4, a recently identified porphyrin-binding protein (Larkin et al. 2003). *GUN4* was identified as the defective gene in the *gun4* mutant, which exhibits aberrant regulation of chloroplast-to-nucleus signal transduction like *gun5*. *GUN4* is regarded to be involved in intracellular signaling (see below), and it binds both protoporphyrin IX and Mg-protoporphyrin IX and stimulates the activity of Mg-chelatase in *Synechocystis* (Verdecia et al. 2005) and *Thermosynechococcus elongatus* (Davison et al. 2005). Disruption of *GUN4*

reduces the cellular levels of heme, suggesting that *GUN4* is also involved in heme biosynthesis and thus may control the flow of substrate into the heme or Chl branch (Wilde et al. 2004).

In addition to regulation via rhythmic gene expression, Mg-chelatase activity is also regulated by diurnal cycling, which is attributed to fluctuations in the stromal concentration of free Mg^{2+} and the [ATP]:[ADP][Pi] ratio. Mg^{2+} triggers a conformational change in Mg-chelatase, resulting in a more active form that promotes cooperative binding of the other substrates, ATP and porphyrin (Reid and Hunter 2004). The free Mg^{2+} concentration in stroma increases from 0.5 to 2.0 mM upon shift from darkness to light in spinach (Ishijima et al. 2003). In the presence of *GUN4*, the Mg^{2+} concentration required for full activation of Mg-chelatase is lowered from 6 to 2 mM, suggesting that *GUN4* controls Mg-chelatase activity at a physiologically relevant Mg^{2+} concentration (Davison et al. 2005). Stromal Mg^{2+} concentration may also affect subcellular localization of ChlH. ChlH associates with the envelope membrane when chloroplasts are disrupted in the presence of 5 mM Mg^{2+} , whereas it is detected in the soluble stromal fraction in the presence of 1 mM Mg^{2+} (Gibson et al. 1996; Nakayama et al. 1998). This suggests that translocation of ChlH from stroma to the envelope membrane is required for formation of active Mg-chelatase complex as well as to bind the substrate, protoporphyrin IX, such that the reaction may proceed.

In chloroplasts, the regulation of the activity of a number of enzymes involved in the photosynthetic reactions is coupled to photosynthetic electron transport via the thio-redoxin system in which a light-induced change in enzyme activity is linked to the redox state of a disulfide bond located within each enzyme (Buchanan and Balmer 2005). Using a thioredoxin-affinity chromatography technique, ChlI was identified as a potential thioredoxin-target protein (Balmer et al. 2003). In fact, Mg-chelatase activity is sensitive to thiol-modifying reagents (Fuesler et al. 1984a, b; Walker and Weinstein 1991a, b), suggesting that thiol groups in the molecule are essential for proper function and optimal activity of this enzyme. The thiol-modifying reagent *N*-ethylmaleimide specifically binds to ChlI and inhibits its ATPase activity (Jensen et al. 2000). The ATPase activity of *A. thaliana* ChlI1 is fully inactivated by oxidation but is easily recovered by thioredoxin-assisted reduction, suggesting that ChlI1 is a target of regulation by thioredoxin (Ikegami et al. 2007). Moreover, the in vivo redox state of ChlI is light dependent (Ikegami et al. 2007).

Thus, taken together, the current data indicate that Mg-chelatase activity is tightly regulated by a diurnal cycle and photosynthetic electron transport in order to coordinate Chl biosynthesis with the photosynthetic activities in chloroplasts.

S-Adenosyl-L-methionine:Mg-protoporphyrin IX methyltransferase

S-Adenosyl-L-methionine:Mg-protoporphyrin IX methyltransferase (EC 2.1.1.11) catalyzes the transfer of the methyl group from S-adenosyl-L-methionine (SAM) to the carboxyl group of the 13-propionate side chain of Mg-protoporphyrin IX. Initially, the gene *BchH* was mistakenly assigned to code for SAM:Mg-protoporphyrin IX methyltransferase in *Rhodobacter* because Mg chelation is obligatorily coupled to the methyltransferase reaction in purple bacteria (Gorchein et al. 1993). Later, heterologous expression of the *BchM* genes of *Rba. capsulatus* and *Rba. sphaeroides* in *E. coli* demonstrated that the gene products had the SAM:Mg-protoporphyrin IX methyltransferase activity in vitro (Bollivar and Bauer 1992; Bollivar et al. 1994a). The enzyme belongs to the broad family of SAM-dependent methyltransferases (Kagan and Clarke 1994). The *Rba. capsulatus* SAM:Mg-protoporphyrin IX methyltransferase activity is stimulated by the BchH subunit of Mg-chelatase (Hinchigeri et al. 1997), suggesting that Mg-protoporphyrin IX may be channeled directly to this enzyme from Mg-chelatase through the BchH subunit. Assembly of functional Mg-chelatase and SAM:Mg-protoporphyrin IX methyltransferase in *E. coli* by heterologous expression of *Rba. sphaeroides* *BchM*, together with *BchI*, *BchD*, and *BchH*, resulted in the introduction of a new branch point in the tetrapyrrole biosynthetic pathway of *E. coli*, causing the accumulation of Mg-protoporphyrin IX monomethyl ester (Jensen et al. 1999b).

In Chl-producing organisms, the gene *chlM* was cloned from *Synechocystis* PCC6803 by functional complementation of the *bchM* mutant of *Rba. capsulatus* (Smith et al. 1996). The reaction mechanism of SAM:Mg-protoporphyrin IX methyltransferase was examined using a recombinant protein of *Synechocystis* ChlM. Transient kinetic analysis showed that a rapid porphyrin binding step to ChlM is preceded by a slower enzyme conformational change (isomerization step). For the subsequent partial reaction, a lag phase attributable to the formation of stable intermediate has been identified (Shepherd and Hunter 2004).

The SAM:Mg-protoporphyrin IX methyltransferase gene has also been cloned from *A. thaliana* (Block et al. 2002) and tobacco (Alawady et al. 2005). In *Arabidopsis*, the enzyme was found to localize to both the thylakoid and envelope membranes within chloroplasts (Block et al. 2002). In antisense transgenic tobacco, reduction of SAM:Mg-protoporphyrin IX methyltransferase activity was directly linked to a reduction of Mg-chelatase activity and ALA synthesis; ferrochelatase activity, however, was elevated. Moreover, overexpression of the enzyme in sense

plants showed the opposite phenotypic profile. Due to changes in ALA synthesis and Mg-chelatase activity correlate with transcript levels of *HemA* (encoding glutamyl-tRNA reductase), *Gsa* (encoding glutamate-1-semialdehyde aminotransferase), and *ChlH*, it has been proposed that changes in methyltransferase activity are communicated to the cytoplasm to coordinate transcriptional activities of these regulatory enzymes (Alawady and Grimm 2005).

Involvement of Mg-porphyrins in intracellular signaling

The photosynthetic apparatus is composed of proteins that are encoded in the nuclear and chloroplast genomes. The mechanism that evolved to coordinate nuclear and organellar gene expression includes communication between the nucleus and chloroplasts. It is believed that chloroplasts send signals to the nucleus in various ways—so called retrograde signaling. Mg-protoporphyrin IX was identified as one of the plastid-derived retrograde signals involved in intracellular communication between chloroplasts and the nucleus (Rodermel 2001; Rodermel and Park 2003; Strand 2004; Beck 2005; Nott et al. 2006). In *Chlamydomonas reinhardtii*, administration of Mg-protoporphyrin IX and Mg-protoporphyrin IX monomethyl ester induces the expression of the light-responsive nuclear gene *HSP70*, which encodes a heat shock protein (Kropat et al. 1997, 2000). Analysis of *cis* elements in the promoter region of *HSP70A* identified two enhancer regions that are involved in Mg-protoporphyrin IX and light inducibility (von Gromoff et al. 2006).

In *A. thaliana*, mutants termed *gun* (genome uncoupled) were identified in which intracellular signaling was disrupted (Susek et al. 1993). These mutants express nuclear-encoded photosynthesis genes even when chloroplast function is disrupted by treatment with norflurazon, an inhibitor of carotenoid biosynthesis. Among five *gun* mutants (*gun1–5*), four of them (*gun2–5*) have mutations in tetrapyrrole biosynthetic enzymes. The *gun2* and *gun3* mutants are alleles of *hy1* and *hy2*, which encode heme oxygenase and phytylchromobilin synthase, respectively. The gene *GUN5* encodes the H subunit of Mg-chelatase (Mochizuki et al. 2001), and *GUN4* encodes a chloroplast porphyrin-binding protein that activates Mg-chelatase (Larkin et al. 2003). Wild type plants accumulate high amounts of Mg-protoporphyrin IX when grown on norflurazon, whereas this accumulation occurs only partially or is absent in *gun2* and *gun5* mutants (Strand et al. 2003). Analysis of nuclear gene expression in an *A. thaliana* knockout mutant of SAM:Mg-protoporphyrin IX methyltransferase also suggested that Mg-protoporphyrin IX is a negative effector of nuclear photosynthetic gene expression (Pontier et al. 2007). Based on this observation, Mg-

protoporphyrin IX has been proposed to accumulate under stress conditions and act as a negative regulator of photosynthetic gene regulation.

In barley, mutants defective in all three subunits of Mg-chelatase displayed the *gun* phenotype, whereas a mutant defective in Mg-protoporphyrin IX monomethyl ester cyclase (see below) did not, although this mutant accumulated Mg-protoporphyrin IX monomethyl ester (Gadjieva et al. 2005). For this reason, it was suggested that Chl intermediates might only function as signal molecules early in chloroplast development (Gadjieva et al. 2005). Taking advantage of the fact that tetrapyrroles have intrinsic fluorescence, the accumulation of Mg-protoporphyrin IX was visualized in vivo using confocal laser scanning microscopy (Ankele et al. 2007). Under stress conditions, Mg-protoporphyrin IX accumulated both in the chloroplast and in the cytosol upon feeding with ALA, suggesting that this intermediate is exported from chloroplasts to the cytosol. Since the accumulation of Mg-protoporphyrin IX was only observed in ALA-treated samples that might be involved in artificial effects, the physiological significance of this phenomenon should be carefully examined. The stress-induced accumulation of Mg-protoporphyrin IX coordinates nuclear- and plastid-encoded photosynthesis genes by repressing the expression of nuclear-encoded sigma factors, which are essential components of plastid-encoded RNA polymerase (Ankele et al. 2007).

However, contradictory observations have been made; *cs* and *ch-42* mutants with a defect in *ChlI* did not display the *gun* phenotype, although the Mg-chelatase activity was severely impaired (Mochizuki et al. 2001). Reduction of the endogenous level of Mg-protoporphyrin IX by over-expressing SAM:Mg-protoporphyrin IX methyltransferase did not alter the expression of a nuclear-encoded photosynthesis gene (Alawady and Grimm 2005).

Very recently, the protein GUN1, the mutation of which caused *gun* phenotype but was independent of tetrapyrrole biosynthesis, was identified as a chloroplast-localized pentatricopeptide-repeat protein (Koussevitzky et al. 2007). Surprisingly, GUN1 and ABI4, an Apetala 2 (AP2)-type transcription factor, are common to retrograde signaling pathways that mediate GUN5-dependent modulation of Mg-protoporphyrin IX levels, as well as the reduction/oxidation (redox) state of the photosynthetic electron transfer chain. It was thus proposed that multiple indicators of aberrant plastid function in *Arabidopsis* are integrated upstream of GUN1 within plastids, which leads to ABI4-mediated repression of nuclear-encoded genes (Koussevitzky et al. 2007).

Moreover, it has been recently proposed that the *gun* phenotype is not directly (or not proportionally) correlated with the levels of Mg-protoporphyrin IX in mutants of tetrapyrrole biosynthesis. Several groups independently

show the contradict results of the accumulation of Mg-protoporphyrin IX under stress conditions. Previously, Strand et al. (2003) showed that norflurazon-treated *Arabidopsis* seedlings accumulate more than 6 nmol g⁻¹ FW of Mg-protoporphyrin IX, which is more than 15-fold higher than untreated wild type seedlings. This level is unusually high without taking into account of up-regulation or activation of upstream enzymes of tetrapyrrole biosynthesis including ALA biosynthesis. However, gene expression analyses of upstream tetrapyrrole biosynthetic genes showed that the treatment of norflurazon to *Arabidopsis* seedlings substantially repressed the transcripts levels of almost all genes (Masuda et al. unpublished data; M.J. Terry, personal communication). Furthermore, Mochizuki et al. (manuscript in preparation) found that Mg-protoporphyrin IX and its methyl ester do not accumulate in norflurazon-treated mutants defective in genes involved in Mg-chelatase, SAM:Mg-protoporphyrin IX methyltransferase, or Mg-protoporphyrin IX monomethyl ester cyclase (see below), and the endogenous levels of these intermediates in wild type and these mutants (including *gun*) are not correlated with the *LHC* expression. Independently, using a sensitive LC/MS method, Moulin and Smith (2008) also showed that no accumulation of Mg-protoporphyrin IX under conditions in which the *LHC* genes are repressed. These observations suggest that the levels of Mg protoporphyrins do not simply account for the regulation of *LHC* transcript level. Thus, future work must clarify how defect of tetrapyrrole biosynthesis in *gun* mutants correlates to the control of expression of nuclear-encoded photosynthetic genes.

Mg-protoporphyrin IX monomethyl ester cyclase

Mg-protoporphyrin IX monomethyl ester cyclase catalyzes the incorporation of atomic oxygen into Mg-protoporphyrin IX monomethyl ester to form 3,8-divinyl protochlorophyllide (Pchl_{id}). This oxidative cyclization reaction creates the fifth ring of Chl. Two types of Mg-protoporphyrin IX monomethyl ester oxidative cyclase have been identified in photosynthetic organisms. One is the anaerobic cyclase, which incorporates atomic oxygen from water; the activity of this enzyme depends on a functional *bchE* gene. Directed mutagenesis of *Rba. capsulatus* revealed that Mg-protoporphyrin IX monomethyl ester accumulated in a *bchE* mutant, suggesting that the *bchE* gene product is required for Mg-protoporphyrin IX monomethyl ester oxidative cyclase activity (Bollivar et al. 1994b). Similarities of the deduced sequence of *bchE* to that of the methylcobalamin-requiring P-methylase from *Streptomyces hygroscopicus* (Kuzuyama et al. 1995) indicated that the *bchE* gene product uses a cobalamin cofactor, and indeed two vitamin

B₁₂-requiring *Rba. capsulatus* mutants (*bluE* and *bluB*) accumulated Mg-protoporphyrin IX monomethyl ester (Gough et al. 2000). Supplementation with adenosylcobalamin or methylcobalamin restored the cyclase activity in permeabilized *bluE* and *bluB* cells. Another type of the cyclase is the an aerobic cyclase, which incorporates atomic oxygen from O₂. The activity of this enzyme depends on the gene *acsF*, homologs of which are found only in photosynthetic organisms. The *acsF* gene was first identified from *Rubrivivax gelatinosus* as a protein containing a putative binuclear iron cluster (Pinta et al. 2002).

The strict aerobic photosynthetic organisms, including higher plants, seem to contain the aerobic Mg-protoporphyrin IX monomethylester cyclase, whereas the strict anaerobes examined so far contain only the anaerobic cyclase. In facultative aerobic photosynthetic bacteria, aerobic and anaerobic cyclases coexist in the same organism (Porra et al. 1998; Pinta et al. 2002), apparently providing the competitive advantage of producing Chls under both aerobic and anaerobic conditions (Ouchane et al. 2004). Although *C. reinhardtii* can grow and produce photosystems under oxygen-limited conditions, this alga lacks a *bchE* gene. In *C. reinhardtii*, two homologs of *acsF*, *CRD1* and *CTH1*, were identified. The *CRD1* was shown to be induced under low oxygen tension and copper deficiency, and a *crd1* mutant is chlorotic under such conditions. In contrast, *CTH1* was expressed primarily during copper sufficiency and hence could not cover the loss of *CRD1* function in *crd1* mutants (Moseley et al. 2000; Eriksson et al. 2004). Therefore, it is probable that Chl biosynthesis is supplied by *CRD1* under low oxygen tension in *Chlamydomonas*. It is possible that lack of *bchE* in *Chlamydomonas* is related to oxygen-evolving photosynthesis in this organism. In purple bacteria, *puc* and *puf* operons, encoding proteins of the photosynthetic apparatus, are regulated by oxygen level. Ouchane et al. (2004) suggested that the existence of two types of Mg-protoporphyrin IX monomethyl ester cyclase encoded by *bchE* and *acsF* in purple bacteria may be advantageous to rapidly build up the photosystem under aerobic or anaerobic conditions, enabling an immediate start of photosynthetic growth following exposure to light.

In higher plants, an *acsF* homolog, *CHL27*, has been found in the *Arabidopsis* genome. Antisense *Arabidopsis* plants with reduced amounts of *CHL27* accumulate Mg-protoporphyrin IX monomethyl ester upon ALA feeding, and the plants have chlorotic leaves with reduced levels of all Chl-binding proteins (Tottey et al. 2003). An in vitro assay for the aerobic cyclase reaction required membrane-bound and soluble components from the chloroplasts (Walker et al. 1988). It was proposed that *CHL27* is membrane-bound catalytic subunit of the aerobic cyclase (Tottey et al. 2003). In barley, the gene *Xantha-I*, a

homolog of *acsF* and *CHL27*, also was identified as a membrane-bound subunit. In addition to the *Xantha-I* gene product, it was proposed that the aerobic cyclase requires at least one soluble and two membrane-bound components (Rzeznicka et al. 2005). Biochemical characterization showed that the aerobic cyclase requires the cofactors NADPH, Fe²⁺, and O₂ (Walker et al. 1989).

3,8-divinyl Pchlide *a* 8-vinyl reductase (DVR)

According to the number of vinyl side chains, chlorophylls of oxygenic photosynthetic organisms are classified into two groups: 3,8-divinyl chlorophyll (divinyl chlorophyll) and 3-vinyl chlorophyll (monovinyl chlorophyll). DVR catalyzes reduction of the 8-vinyl group on the tetrapyrrole to an ethyl group (Parham and Rebeiz 1995) using NADPH as the reductant. Reduction of the 8-vinyl group can probably occur at various steps of Chl biosynthesis before and after the reduction of the D pyrrole ring. Therefore, it is likely that the monovinyl and divinyl Chl biosynthesis reactions may occur in parallel rather than by separate linear pathways (Parham and Rebeiz 1995).

Almost all of the oxygenic photosynthetic organisms contain monovinyl chlorophylls, regardless of the variation in their indigenous environments (Porra 1997). The exceptions are species of *Prochlorococcus marinus*, marine picophytoplanktons that contain divinyl chlorophylls as their photosynthetic pigments (Chisholm et al. 1992; Coleman and Chisholm 2007). The advantage for *Prochlorococcus* sp. to have divinyl Chls is that the absorbance maximum of the pigments in the Soret region red shifts by ~10 nm compared with monovinyl Chls, enabling more efficient absorption of green light that is enriched in deep water layers.

Two independent groups recently isolated the *DVR* gene by characterizing *A. thaliana* mutants that exhibit a predominant replacement of monovinyl Chl with divinyl Chl and another mutant that is pale green and has a reduced level of Chl *b* (Nagata et al. 2005; Nakanishi et al. 2005). Interestingly, a *DVR* homolog was identified in marine *Synechococcus* but not in *Prochlorococcus* (Nagata et al. 2005). It has been proposed that the common ancestor of *Prochlorococcus* lost *DVR* and acquired divinyl Chls, which allow photosynthetic growth in deep-sea waters by absorbing green light, which predominates in this niche.

Arabidopsis *DVR* can reduce 3,8-divinyl Pchlide as well as 3,8-divinyl chlorophyllide, but the efficiency of the former substrate is substantially lower than that of the latter in vitro and in vivo (Nagata et al. unpublished results). Thus, it is possible that the *DVR* reaction occurs after the NADPH:Pchlide oxidoreductase (POR) reaction, and thus

should be designated as 3,8-divinyl chlorophyllide *a* 8-vinyl reductase.

DVR has been identified in higher plants and green algae, but no homologs have been found in the genome of the unicellular red alga *Cyanidioscyzoon merolae*, a eukaryotic photosynthetic organism, although this organism synthesizes monovinyl Chl *a*. Among cyanobacteria, five *Synechococcus* species were found to contain a DVR homolog, but no other species, including eleven *Synechococcus* species, *Synechocystis* sp PCC6803, *Nostoc* sp. PCC7120, and *Gloeobacter violaceus* PCC7421, contain a DVR homolog although all of these organisms synthesize monovinyl Chl *a*. The gene *bchJ* has been also reported to be essential in the reduction of the 8-vinyl group in *Rba. capsulatus* (Bollivar et al. 1994b), but neither *bchJ* nor the DVR homolog has been found in the majority of the cyanobacterial genomes. Interestingly, recent analysis of the green sulfur bacterium *Chlorobium tepidum* led to the proposal that BchJ may play an important role in substrate channeling and/or regulation of Chl biosynthesis but showed that it is not a vinyl reductase (Chew and Bryant 2007). These observations imply the existence of a third type of DVR that is not related to *A. thaliana* DVR or to *Rba. capsulatus* BchJ (Nagata et al. 2005). Very recently a novel cyanobacterial DVR was identified by bioinformatic survey of whole-genome comparisons (personal communication from Dr. Ayumi Tanaka).

Pchlde oxidoreductase

Pchlde oxidoreductase catalyzes the reduction of the C₁₇–C₁₈ double bond of the D pyrrole ring of the tetrapyrrole macrocycle. Two types of Pchlde oxidoreductase have been identified in photosynthetic organisms. One is light-dependent NADPH:Pchlde oxidoreductase (POR; EC 1.3.1.33 or EC 1.6.99.1), which has been the subject of a number of recent reviews (Aronsson et al. 2003b; Schoefs and Franck 2003; Masuda and Takamiya 2004; Heyes and Hunter 2005). Thus, here I discuss only the most recent findings on POR. POR is a single-subunit enzyme that requires light for catalysis and is present in all Chl-synthesizing organisms but is not found in Bchl-synthesizing organisms. Another type is light-independent Pchlde oxidoreductase (DPOR) (Armstrong 1998; Fujita and Bauer 2003). DPOR is a multisubunit enzyme consisting of three subunits related to nitrogenase. DPOR is not found in angiosperms but is present in most other Chl- and Bchl-synthesizing organisms; this allows these organisms to synthesize Chl or Bchl in darkness. When angiosperms are germinated in darkness, they accumulate small amounts of Pchlde in a ternary complex with NADPH and POR, and they are unable to synthesize Chl

until this bound Pchlde is converted to chlorophyllide upon exposure to light.

One aspect of POR on which recent research has focused is the import mechanism for the POR precursor (pPOR) and its assembly into the prolamellar body (PLB) in plastids. Two types of POR, PORA and PORB, have been identified in *A. thaliana* (Armstrong et al. 1995) and barley (Holtorf et al. 1995). PORA and PORB are highly similar except in the N-terminal region that contains structural features such as the chloroplast transit peptide, but their gene expression profiles are substantially different. Both *POR* mRNAs are formed in etiolated seedlings, but only *PORB* mRNA continues to accumulate in light-grown plants, whereas *PORA* mRNA rapidly disappears after illumination. These results suggest that PORB operates throughout the greening process and in light-adapted mature plants, whereas PORA is active only in etiolated seedlings at the beginning of illumination. Thereafter, a third POR, PORC, the expression of which is induced by light, was identified in *A. thaliana* (Oosawa et al. 2000). The in vitro POR activity of PORC is recently confirmed (Pattanayak and Tripathy 2002).

Reinbothe et al. (1995a, b) originally suggested that the import of the precursor of barley pPORA into chloroplasts was totally dependent on envelope-bound Pchlde, the enzyme's substrate. It was suggested that the pPORA transit peptide directly interacts with Pchlde in the plastid envelope for translocation (Reinbothe et al. 1996, 1997). Moreover, Reinbothe et al. (1999) performed in vitro reconstitution experiments with the two barley POR enzymes and synthetic zinc analogs of Pchlde *b* and Pchlde *a*, zinc-protopheophorbide *b* (ZnPPb) and zinc-protopheophorbide *a* (ZnPPa). These experiments led to the hypothesis of a novel light-harvesting Pchlde *a/b*-binding protein complex, named LHPP, and a new proposal for distinct functions of PORA and PORB in vivo. The LHPP complex is thought to consist of a 5:1 ratio of the dark-stable ternary complexes of PORA and PORB, which specifically bind to Pchlde *b* and *a*, respectively, and are embedded in the lipid bilayers of PLB of etioplasts. Only the PORB-bound Pchlde *a* in the LHPP complex appears to be reduced immediately upon illumination, whereas the PORA-bound Pchlde *b* is proposed to function initially as a light-harvesting pigment. Energy transfer from Pchlde *b* to Pchlde *a* is speculated to provide a mechanism for photoprotection during the early stage of seedling greening. Subsequently, several papers from the same group have been published (Reinbothe et al. 2000, 2004a, b, 2005; Pollmann et al. 2007; Schemenewitz et al. 2007). The basic concept of their proposal is that pPORA is imported into plastids with a specific translocon, designated Pchlde translocon complex (Ptc), in an envelope-bound Pchlde *b*-dependent manner to form a PORA-Pchlde *b* complex

with LHPP, whereas PORB is imported with the general translocon. However, several papers dispute this concept. Thus, the following sections first review a series of recent studies—mainly from Reinbothe's group—that support the LHPP concept, followed by a discussion of criticisms of this concept.

Reinbothe et al. (2003c) reported that Pchl $ide\ b$ is abundant in barley etioplasts but metabolically unstable because it is rapidly converted to Pchl $ide\ a$ by the activity of 7-formyl reductase, an enzyme involved in the Chl b -to-Chl a reaction. They proposed that 7-formyl reductase finely tunes the levels of Pchl $ide\ b$ and Pchl $ide\ a$, and thereby may regulate the steady-state level of LHPP (Reinbothe et al. 2003c). Reconstitution of LHPP complexes in vitro from barley PORA and PORB, chemically synthesized Pchl $ide\ a$, Pchl $ide\ b$, and lipids showed pigment binding characteristics of PORA and PORB specific to Pchl $ide\ b$ and Pchl $ide\ a$, respectively (Reinbothe et al. 2003a). The reconstituted ~480-kDa complex displayed the same characteristics as photoactive Pchl ide . Upon illumination, only the PORB-bound Pchl $ide\ a$ was photoactive and it was converted to Chl $ide\ a$, whereas Pchl $ide\ b$ bound to PORA remained photoinactive. Fractionation of isolated barley etioplasts identified POR-pigment complexes that are similar in size, stoichiometry, and photochemical properties to the reconstituted LHPP (Reinbothe et al. 2003a). The authors further analyzed deletion mutants of barley POR and found that the central region of the polypeptide, the so-called “extra-loop” that distinguishes POR from the structurally related short-chain alcohol dehydrogenase, is dispensable for pigment binding but required for the assembly of LHPP (Reinbothe et al. 2003b).

Pchl ide -dependent import of the transit peptide of barley pPORA has been shown in chloroplasts of tobacco, *A. thaliana*, and five other monocotyledonous and dicotyledonous plant species (Reinbothe et al. 2000). A competition assay with the precursors of the small subunit of ribulose-1,5-bisphosphate carboxylase/oxygenase (pSSU) and ferredoxin as well as an antibody-blocking experiment with anti-TOC86 and anti-TOC75 indicated that the import site is specific for pPORA (Reinbothe et al. 2000). By chemical crosslinking using a reagent that interacts with the transit peptide of pPORA, the outer plastid envelope protein Oep16 was identified as the translocase for pPORA (Reinbothe et al. 2004b). Furthermore, analysis of components of the Ptc complex identified a putative Pchl $ide\ a$ oxygenase (Ptc52), a tyrosine aminotransferase, GTP-binding proteins Toc33/34, and Oep16 (Reinbothe et al. 2004a). As Pchl $ide\ b$ was detected as part of the Ptc complex, it was proposed that the putative Pchl $ide\ a$ oxygenase, which is homologous to Lls1, Toc55, and chlorophyll $ide\ a$ oxygenase (CAO), provides Pchl $ide\ b$

as import substrate for pPORA to form the major component, PORA-Pchl $ide\ b$, of the LHPP complex (Reinbothe et al. 2004a). Oep16 forms a larger complex with TOC33 but not TOC34 in *Arabidopsis* chloroplasts (Reinbothe et al. 2005). Plastids of the *ppi1* mutant of *Arabidopsis* lacking Toc33 were unable to import pPORA but imported pSSU, pPORB, and a precursor of ferredoxin (Reinbothe et al. 2005). A loss-of-function mutation in *Arabidopsis* Oep16 led to defects in import and assembly of PORA, which caused excess accumulation of Pchl ide in the dark. Then, the pigment operated as a photosensitizer and provoked cell death during greening (Pollmann et al. 2007). In the Oep16 mutant, the PLB was completely lacking as compared with wild-type etioplasts. Schemenewitz et al. (2007) reported that pPORA can use two different plastid import pathways that differ in their requirements for cytosolic 14:3:3 proteins and Hsp70. pPORA synthesized in vitro in wheat germ lysate segregated into two different fractions by size exclusion chromatography, one of which bound with 14:3:3 proteins and Hsp70, and pPORA in this fraction was imported into Pchl ide -free chloroplasts by the standard translocon, whereas other free and Hsp70-bound pPORA were imported in Pchl ide -dependent manner. They also showed that Ptc52, encoding the putative Pchl $ide\ b$ oxygenase, but not CAO, is indispensable for Pchl $ide\ b$ -dependent transport (Schemenewitz et al. 2007).

Their hypothesis is supported by several reports by other research groups. Smith et al. (2004) reported that the main receptor of transit peptides that are imported into chloroplasts, Toc159, interacts with precursors to photosynthetic proteins; however, this protein had little activity in binding assays with nonphotosynthetic precursors, including pPORA. Using stable *Arabidopsis* transformants expressing pPOR-green fluorescent protein (GFP) fusion constructs, Kim and Apel (2004) showed that substrate-dependent import of pPORA into plastids of cotyledons occurred in the *flu* mutant, which over-accumulates Pchl ide in the dark, but not in the *xantha2* mutant, which is devoid of Pchl ide . In true leaves, PORA uptake does not require the presence of Pchl ide . However, Kim et al. (2005) showed that in *ppi3*, *ppi1* and *ppi2* mutants, which are deficient in atToc34, atToc33 and atToc159, respectively, none of these translocon constituents is required for substrate-dependent and organ-specific import of PORA, whereas atToc33 is necessary for the import of PORB.

On the contrary, other studies have reported results that conflict with the Pchl ide -dependent import and the LHPP model (Armstrong et al. 2000). The import into chloroplasts in a homologous system from *Arabidopsis* (Jarvis et al. 1998; Jarvis and Soll 2001), pea (Dahlin et al. 1995; Aronsson et al. 2000, 2001, 2003a) and wheat (Teakle and Griffiths 1993) was not dependent on Pchl ide . The absence or presence of Pchl ide did not significantly affect the

capacity to import pPORA and pPORB in barley (Dahlin et al. 1995; Aronsson et al. 2000). A competition assay with pSSU and crosslinking analysis indicated that pPORA uses the general import pathway into chloroplasts (Aronsson et al. 2000). Re-examination of pigment extraction from etiolated barley failed to detect Pchl $ide\ b$ irrespective of the extraction protocol, and disputed the existence of LHPP complexes in barley etioplasts (Kollosov and Rebeiz 2003). Phillipar et al. (2007) recently analyzed homozygous loss-of-function mutant lines of Oep16 and showed that etioplasts from *oep16* mutants contained PORA protein, as assessed by mass spectrometry. Strikingly, even though they analyzed the same T-DNA insertional SALK line of Oep16 that was analyzed by Pollmann et al. (2007), they showed that mutants had a normally structured PLB that contained the protochlorophyllide holochrome.

There is ongoing disagreement/discussion regarding the presence of Pchl $ide\ b$ in vivo, Pchl $ide\ b$ -dependent import of pPORA, and the resultant formation of PORA-Pchl $ide\ b$ in the LHPP complex. For the presence of Pchl $ide\ b$ in vivo, although Pchl $ide\ b$ was reported in several green plants (Shedbalkar et al. 1991), the abundant accumulation of Pchl $ide\ b$ in etioplasts observed by Reinbothe et al. (2003c) was not reproduced by the same extraction protocol that inhibits putative 7-formyl reductase (Kollosov and Rebeiz 2003). Furthermore, molecular nature of this 7-formyl reductase, which may have unusually high activity to convert Pchl $ide\ b$ to Pchl $ide\ a$ is still unidentified. For production of Pchl $ide\ b$, Ptc52 is proposed as the putative Pchl $ide\ a$ oxygenase (Reinbothe et al. 2004a), however in vitro activity of this protein has not been demonstrated. In addition, the Pchl $ide\ b$ added to etioplast membrane was photoreduced to Chl $ide\ b$ by POR (Scheumann et al. 1999). Scheumann et al. (1999) also detected a substantial reduction of exogenously added ZnPP b to zinc-71-hydroxy-protopheoporphide a in darkness, which is catalyzed by Chl b reductase (see below).

For discrepancy of the Pchl ide -dependent import of pPORA, instability of the protein import receptor and the involvement of a high concentration of urea in the reaction mixtures have been considered (Aronsson et al. 2000; Phillipar et al. 2007). A major protein import receptor, TOC160, is extremely sensitive to proteolysis upon cell lysis, and degrades with a half-life of a few minutes if strict precautions are not taken (Botler et al. 1998). The degradation of TOC160 lowers the import efficiency of pSSU, although the degraded 86-kDa form still supports the import reaction. High concentrations of urea used in Reinbothe's experiments seem to influence chloroplast integrity and import competence. Aronsson et al. (2000, 2003b) proposed that differential functional integrity of the TOC complex might influence the requirement for import of pPOR into plastid in vitro. Very recently, Schemenewitz

et al. (2007) proposed that cytosolic targeting factors, 14:3:3 proteins, present in wheat germ lysates, which were used to in vitro translation in other research groups, may target pPORA to an import site other than the Ptc complex. They proposed the level of the 14:3:3 proteins may physiologically and developmentally regulate the import of PORA. Further analysis is necessary to identify the effects of urea and the involvement of such cytosolic targeting factor on the substrate-dependent import of pPORA. Moreover, why the same T-DNA knockout line of Oep16, a putative component of Ptc complex, showed very distinct phenotypes should be clarified (Phillipar et al. 2007; Pollmann et al. 2007).

For the formation of the LHPP, it is still difficult to generalize as the central structural determinant of the PLBs in etioplasts of angiosperms. The given 5:1 stoichiometry of PORA to PORB is not consistent with the levels of PORA and PORB mRNA in etiolated seedlings of *Arabidopsis* (Armstrong et al. 1995). Furthermore, the total quantities of either PORA or PORB mRNA in *Arabidopsis* can be specifically manipulated by their constitutive overexpression or knockout by DNA tagging (Sperling et al. 1997, 1998; Franck et al. 2000; Frick et al. 2003; Masuda et al. 2003a). Such studies provide no evidence that alterations in the PORA to PORB ratio, independent of the total quantity of POR, dramatically influence the extent of PLB formation and the photoactive Pchl ide . More strikingly, overexpression of cyanobacterial *POR* from *Synechocystis* PCC6803 or *Gloeobacter violaceus* in *Arabidopsis* PORA knockdown mutant showed significant complementation of Chl deficiency, with concomitant formation of PLB in etioplasts (Shinji Masuda et al. unpublished data). These cyanobacteria, producing only Chl a but not Chl b , contain single *POR* gene but do not form PLB in the dark, since they also contain DPOR. Thus, it is likely that the PLB forming capacity is instinct nature of POR proteins, irrespective of species of photosynthetic organisms. Moreover, cucumber (Fusada et al. 2000) and pea (Spano et al. 1992; Sundqvist and Dahlin 1997) provide an example of angiosperms that apparently contain only one *POR* gene. Therefore, the in vivo data from *Arabidopsis*, cucumber and pea are not consistent with specific functions for PORA and PORB within an LHPP complex of the type postulated especially for barley. One possibility is that the LHPP complex is specific to monocotyledonous plants, since Reinbothe et al. (2003a) found a similar solubilization of PORA during the isolation of etioplast inner membranes from oat and wheat etioplasts. However, the predominance of non-photoactive Pchl ide , which is one essential prediction of the LHPP model, is not fulfilled in these plants, since most of the Pchl ide was phototransformed by flash illumination of wheat etioplast (Ryberg and Sundqvist 1988), etiolated wheat seedlings

(Franck et al. 1999), and oat seedlings (Domanski and Rüdiger 2001). Thus, although the proposed LHPP model is intriguing, it is generally untenable in view of many opposing arguments.

DPOR

Many gymnosperms, ferns, mosses, cyanobacteria, and photosynthetic bacteria have the ability to synthesize Chl in the dark, indicating the existence of a light-independent mechanism for Pchlde reduction. Analysis of mutants of *Rhodobacter* species identified three genetic loci, known as *BchL*, *BchB*, and *BchN*, that are required for light-independent Pchlde reduction in a photosynthetic gene cluster in photosynthetic bacteria (Zsebo and Hearst 1984; Coomber et al. 1990; Yang and Bauer 1990; Burke et al. 1993; Bollivar et al. 1994b). Fujita and Bauer (2000) subsequently reconstituted the activity of light-independent Pchlde reduction in vitro with recombinant BchL, BchN, and BchB from *Rba. capsulatus*.

DPOR is an oxygen-sensitive nitrogenase-like enzyme consisting of the L protein (BchL/ChlL-dimer) as the reductant component and the NB proteins (BchN/ChlN–BchB/ChlB-heterotetramer) as the catalytic component (Fujita and Bauer 2003). In the cyanobacteria, *Synechocystis* sp. PCC 6803 (Wu and Vermaas 1995) and *Leptolyngbya boryana* (formerly *Plectonema boryanum*) (Fujita et al. 1998), analysis of *ChlL*-disrupted mutants revealed that DPOR contributes to Chl synthesis in cells growing in light, suggesting that DPOR is tolerant of oxygen to some extent or has a mechanism of protection from the oxygen evolved in oxygenic phototrophs. The extent of the contribution by DPOR decreases with increasing light intensity, suggesting inefficient catalysis when oxygen evolution is maximal (Fujita et al. 1998). Yamazaki et al. (2006) showed that anaerobic conditions are required for DPOR to compensate for the loss of POR in a mutant of *L. boryana*. In the POR-lacking mutant, DPOR activity was stimulated by an increase in levels of the DPOR subunits, especially ChlL and ChlN, suggesting that this increase stimulates DPOR activity that is protected efficiently from oxygen by anaerobic environments, resulting in complementation of the loss of POR. The metalcenter of BchL was identified as an oxygen-sensitive [4Fe–4S] cluster similar to the nitrogenase Fe protein (Nomata et al. 2006). Considering the oxygen sensitivity of DPOR, the evolution of light-dependent POR is considered to have been critical for the survival of ancestral cyanobacteria that were growing in oxidative environments exposed to high light (Yamazaki et al. 2006). Furthermore, dual Pchlde reductases are assumed to operate differentially in oxygenic photosynthetic cells

grown under natural environments where oxygen levels change dynamically.

The Chl cycle

It is now considered that the chlorophyllide *a* produced by Pchlde reductase is subsequently converted to Chl *a* by a Chl synthase-catalyzed phytol chain attachment. A portion of chlorophyllide *a* is converted to chlorophyllide *b* by a Reiske-type monooxygenase, CAO (Oster et al. 2000) and subsequently is converted to Chl *b*—one of the major light-harvesting pigments essential for optimal light harvesting. The phytol chain of Chl *a* can be removed by chlorophyllase, and the resultant chlorophyllide *a* can be used as a substrate of CAO. Chl *b* that is produced by the action of CAO can be reversibly converted to Chl *a* through the intermediate 7-hydroxymethyl Chl *a* (Ito et al. 1993, 1994) via the action of Chl *b* reductase and 7-hydroxymethyl Chl *a* reductase (Scheumann et al. 1998; Kusaba et al. 2007). Thus, Chl *a* and Chl *b* are interconvertible, with the entire interconversion system being designated as the Chl cycle (Ito et al. 1996; Rüdiger 2002). The Chl cycle is considered to be important for altering the size of light-harvesting antennae to allow plant acclimation to various light intensities (Tanaka et al. 2001; Tanaka and Tanaka 2005). In addition, the conversion of Chl *b* to Chl *a* is crucial because this is the first step in Chl degradation (Kusaba et al. 2007).

Chl synthase

Chl synthetase, encoded by *ChlG*, catalyzes the esterification of chlorophyllide *a* with phytol. For the substrates of Chl synthetase, early evidence indicated that during greening alcohol moiety is provided by geranylgeranyl diphosphate but not phytol (Shioi and Sasa 1983; Maloney et al. 1989). It has been suggested that chlorophyll *a* phytol is formed in two steps: the esterification of chlorophyllide *a* and geranylgeraniol and followed by three successive hydrogenations of the alcohol moiety. The corresponding enzyme in photosynthetic bacteria catalyzes the esterification of bacteriochlorophyllide *a* to Bchl *a*. Chl synthase activity was originally detected in etioplast membranes of *Avena sativa* L. (Rüdiger et al. 1980). The activity and substrate specificity of Chl synthase was demonstrated with the recombinant enzymes from *Rba. capsulatus*, *Synechocystis* sp. PCC6803 (Oster et al. 1997), *A. thaliana* (Oster and Rüdiger 1997) and *Avena sativa* (Schmid et al. 2001). *ChlG* is constitutively expressed as the same transcript level is found in dark-grown and light-grown seedlings (Schmid et al. 2001). Analysis of esterification kinetics of

chlorophyllide suggested that POR and Chl synthase form a complex having the ratio 7:1 (Domanskii et al. 2003). Experiments with chemically modified substrates suggested that tetrapyrroles bind with the same orientation on both enzymes; this excludes simultaneous binding of substrate to both enzymes (Rüdiger et al. 2005).

Chlorophyllide *a* oxygenase (CAO)

CAO catalyzes the conversion of chlorophyllide *a* to chlorophyllide *b*. The gene *CAO* was first identified from a Chl *b*-less insertional mutant of *C. reinhardtii* (Tanaka et al. 1998). The *CAO* gene encodes an oxygenase with a Rieske-type [2Fe–2S] cluster and a mononuclear iron-binding site. *CAO* genes have since been cloned from *A. thaliana* (Espineda et al. 1999; Tomitani et al. 1999), *Oryza sativa*, *Marchantia polymorpha*, *Dunaliella salina*, and Chl *b*-producing prokaryotes *Prochlorothrix hollandica* (Nagata et al. 2004) and *Prochloron didemni* (Tomitani et al. 1999). *CAO* has not been found in the genomes of *Prochlorococcus* species, although these organisms also produce Chl *b*. Using bioinformatics and molecular genetic techniques, Satoh and Tanaka (2006) identified the Chl *b* synthesis gene from *Prochlorococcus marinus* MIT9313, and this gene is distantly related to *CAO* genes. *CAO* is a single nuclear gene in *Chlamydomonas* (Tanaka et al. 1998) and *Arabidopsis* (Espineda et al. 1999), and mutants lacking *CAO*, such as *chlorina1*, fail to accumulate Chl *b* (Espineda et al. 1999; Oster et al. 2000). The recombinant *Arabidopsis* CAO protein requires oxygen and reduced ferredoxin for its activity, and it catalyzes a two-step oxygenation reaction of chlorophyllide *a* to chlorophyllide *b* via 7-hydroxymethyl chlorophyllide (Oster et al. 2000; Reinbothe et al. 2006). The enzyme can also use Zn-chlorophyllide *a* as a substrate but not pheophorbide *a* or Chl *a*. Heterologous expression of *Arabidopsis* CAO in *Synechocystis* PCC6803 caused the accumulation of Chl *b* (Satoh et al. 2001).

In the antenna systems of higher plants and green algae, Chl *b* binds to peripheral light-harvesting complexes (LHC). The antenna size can vary in response to growth at different light intensities, with concomitant changes in the Chl *a/b* ratio. Overexpression of *CAO* in transgenic *Arabidopsis* (Tanaka et al. 2001) and tobacco (Pattanayak et al. 2005) decreased the Chl *a/b* ratio, implying an enlargement of antenna size. When low-light-grown *Dunaliella salina* was transferred to high-light conditions, the *CAO* transcript level decreased in concert with an increased Chl *a/b* ratio (Masuda et al. 2002, 2003b). Such light intensity-dependent changes of *CAO* mRNA levels have been observed in tobacco (Pattanayak et al. 2005) and

Arabidopsis (Harper et al. 2004; Tanaka and Tanaka 2005). These results suggested that Chl *b* synthesis is regulated in part at the *CAO* mRNA level.

The level of the CAO protein is also regulated by protein turnover. CAO is almost undetectable in wild type plants, even though *CAO* mRNA can readily be detected. CAO is comprised of three domains, designated A, B, and C (Nagata et al. 2004). The catalytic domain C can catalyze the reaction independently of domains A and B. Domain A senses the presence of Chl *b* and controls the overall level of the CAO protein because this domain is recognized by protease. When domain C alone was expressed in *Arabidopsis*, the levels of CAO dramatically increased, resulting in extremely low Chl *a/b* ratios (Yamasato et al. 2005). It was proposed that such a feedback mechanism at the protein level effectively controls the level of Chl *b* in response to cellular demand for this protein (Tanaka and Tanaka 2006). An *Arabidopsis* line overexpressing a CAO-GFP fusion was used to identify mutants that had defects in the control of CAO accumulation (Nakagawara et al. 2007). This analysis revealed that mutation of the gene *ClpC1*, encoding the chloroplast protease ClpC1, yielded stronger CAO-GFP fluorescence with decreased Chl *a/b* ratio, suggesting that this protease is involved in regulating Chl *b* synthesis through the destabilization of CAO in response to the accumulation of Chl *b*.

CAO was found on the chloroplast envelope and thylakoid membranes in chloroplasts (Eggink et al. 2004; Reinbothe et al. 2006). In dark-grown *Chlamydomonas* cells, CAO was detected only on the envelope inner membrane (Eggink et al. 2004). CAO was identified as a part of a tranlocon involved in the regulated import and stabilization of LHC proteins Lhcb1 (LHCII) and Lhcb4 (CP29) (Reinbothe et al. 2006).

Chl *b* reductase

Chl *b* is converted to Chl *a* by Chl *b* reductase and 7-hydroxymethyl Chl *a* reductase. The rice *non-yellow coloring 1* (*nyc1*) mutant, which displays the “stay-green” phenotype, retained Chl *b* and LHCII in the light as well as in the dark during leaf senescence. Map-based cloning showed that *NYC1* encodes a chloroplast-localized short-chain dehydrogenase (Kusaba et al. 2007). The *NYC1*-like protein (*NOL*) showed Chl *b* reductase activity, suggesting that *NYC1* and *NOL* encode Chl *b* reductases having divergent functions and physiological roles (Kusaba et al. 2007). It was suggested that 7-hydroxymethyl Chl *a* reductase is a ferredoxin-dependent enzyme (Scheumann et al. 1998), but the gene encoding this enzyme has not been identified.

Chlorophyllase

Chlorophyllase catalyzes the hydrolysis of Chl *a* to yield chlorophyllide *a* and phytol (Takamiya et al. 2000). Chlorophyllase is not only the first enzyme in the Chl degradation pathway but also the enzyme involved in the Chl cycle because CAO uses chlorophyllide *a* as substrate (Oster et al. 2000). The gene encoding chlorophyllase was originally identified in *Chenopodium album* and *Arabidopsis* (Tsuchiya et al. 1999), and in ethylene-treated citrus fruit (Jacob-Wilk et al. 1999). Cloning of this gene from *Chenopodium album* revealed that this enzyme contained a lipase motif (Tsuchiya et al. 1999). Site-directed mutagenesis of recombinant chlorophyllase of *C. album* revealed that the catalytic mechanism is analogous to that of serine hydrolase, which has the catalytic triad composed of Ser, His, and Asp in its active site (Tsuchiya et al. 2003).

One of the two isoforms of *Arabidopsis* chlorophyllase (AtCLH1) was originally isolated as the COR11 (coronatine-induced) protein from the mutant *coil*. AtCLH1 is rapidly induced by methyl jasmonate, which promotes senescence and Chl degradation in plants (Tsuchiya et al. 1999). The in vivo function of AtCLH1 was shown in wild type and *coil* plants overexpressing AtCLH1, and these plants had increased levels of Chlide without a substantial change in the total amount of extractable Chl, resulting in high Chlide to Chl ratios in leaves (Benedetti and Arruda 2002). Furthermore, Kariola et al. (2005) reported that AtCLH1 is involved in plant damage control and can modulate the balance among defense pathways. In an AtCLH1 RNAi-silenced line, failure to degrade free Chl upon damage caused hydrogen peroxide to accumulate, with concomitant induction of defense genes, under both pathogen inoculation and high-light conditions. Downregulation of AtCLH1 resulted in resistance to the bacterial necrotroph, *Erwinia carotovora*, but increased susceptibility to the necrotrophic fungus, *Alternaria brassicicola*, resistance to which requires jasmonate signaling (Kariola et al. 2005).

Meanwhile, the expression of citrus chlorophyllase in tobacco protoplast or squash leaves suggested post-translational regulation of this enzyme through cleavage of its N-terminal peptide, which targets the enzyme for degradation (Harpaz-Saad et al. 2007). Heterologous expression of chlorophyllase lacking N-terminal 21 residues resulted in a chlorotic phenotype, whereas no visible phenotype was observed in plants expressing full-length chlorophyllase. As processing of the N-terminal peptide appeared to occur in tobacco, it was proposed that chlorophyllase functions as a rate-limiting enzyme in Chl catabolism controlled via post-translational regulation (Harpaz-Saad et al. 2007).

Concluding remarks

During the last decade, except for 7-hydroxymethyl Chl *a* reductase and unidentified subunits of Mg-protoporphyrin IX monomethyl ester cyclase, all enzymes involved in the Mg branch of Chl biosynthetic pathway have been identified. Analyses of enzymes in vitro and in vivo have modified the classical biosynthetic pathway as redrawn in Fig. 1. Based on the identified enzymes and pathway, novel molecular mechanisms for regulatory pathways governing Chl metabolism have been proposed such as those involving newly identified regulatory proteins like FLU and GUN4, a new protein import and assembly mechanism for POR, and the degradation mechanism of CAO that is triggered by pigments. Further refinement of these mechanisms will provide insight as to how Chl biosynthesis is controlled to meet the exact cellular demands for this essential but potentially toxic molecule in plants. Moreover, recent data indicate that Chl intermediates and degradation products have various functions in diverse physiological processes such as retrograde signaling and programmed cell death (Hörtensteiner 2004; Pruzinska et al. 2005). The complete elucidation of the network comprising these processes linked to Chl biosynthesis will enhance our basic understanding of the photosynthetic system as well as other cellular processes.

Acknowledgments I am grateful to Dr. Ryouichi Tanaka for critical reading of the manuscript. I acknowledge the financial support from a Grant-in-Aid for Scientific Research (18570034) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

- Alawady AE, Grimm B (2005) Tobacco Mg protoporphyrin IX methyltransferase is involved in inverse activation of Mg porphyrin and protoheme synthesis. *Plant J* 41:282–290
- Alawady A, Reski R, Yaronskaya E, Grimm B (2005) Cloning and expression of the tobacco *CHLM* sequence encoding Mg protoporphyrin IX methyltransferase and its interaction with Mg-chelatase. *Plant Mol Biol* 57:679–691
- Al-Karadaghi S, Franco R, Hansson M, Shelnutt JA, Isaya G, Ferreira GC (2006) Chelataes: distort to select? *Trends Biochem Sci* 31:135–142
- Ankele E, Kindgren P, Pesquet E, Strand A (2007) In vivo visualization of Mg-protoporphyrin IX, a coordinator of photosynthetic gene expression in the nucleus and the chloroplast. *Plant Cell* 19:1964–1979
- Apchelimov AA, Soldatova OP, Ezhova TA, Grimm B, Shestakov SV (2007) The analysis of the *ChlI 1* and *ChlI 2* genes using acifluorfen-resistant mutant of *Arabidopsis thaliana*. *Planta* 225:935–943
- Armstrong GA (1998) Greening in the dark: light-independent chlorophyll biosynthesis from anoxygenic photosynthetic bacteria to gymnosperms. *J Photochem Photobiol B* 43:87–100
- Armstrong GA, Runge S, Frick G, Sperling U, Apel K (1995) Identification of NADPH:protochlorophyllide oxidoreductases A

- and B: A branch pathway for light-dependent chlorophyll biosynthesis in *Arabidopsis thaliana*. *Plant Physiol* 108: 1505–1517
- Armstrong GA, Apel K, Rüdiger W (2000) Does a light-harvesting protochlorophyllide *a/b*-binding protein complex exist? *Trend Plant Sci* 5:40–44
- Aronsson H, Sohrt K, Soll J (2000) NADPH:Protochlorophyllide oxidoreductase uses the general import route into chloroplasts. *Biol Chem* 381:1263–1267
- Aronsson H, Sundqvist C, Timko MP, Dahlin C (2001) Characterization of the assembly pathway of the pea NADPH: protochlorophyllide (Pchl) oxidoreductase (POR), with emphasis on the role of its substrate, Pchl. *Physiol Plant* 111:239–244
- Aronsson H, Sundqvist C, Dahlin C (2003a) POR—import and membrane association of a key element in chloroplast development. *Physiol Plant* 118:1–9
- Aronsson H, Sundqvist C, Dahlin C (2003b) POR hits the road: import and assembly of a plastid protein. *Plant Mol Biol* 51:1–7
- Axelsson E, Lundqvist J, Sawicki A, Nilsson S, Schröder I, Al-Karadaghi S, Willows RD, Hansson M (2006) Recessiveness and dominance in barley mutants deficient in Mg-chelatase subunit D, an AAA protein involved in chlorophyll biosynthesis. *Plant Cell* 18:3606–3616
- Balmer Y, Koller A, del Val G, Manieri W, Schurmann P, Buchanan BB (2003) Proteomics gives insight into the regulatory function of chloroplast thioredoxins. *Proc Natl Acad Sci USA* 100:370–375
- Beale SI (1999) Enzymes of chlorophyll biosynthesis. *Photosynth Res* 60:43–73
- Beck CF (2005) Signaling pathways from the chloroplast to the nucleus. *Planta* 222:743–756
- Benedetti CE, Arruda P (2002) Altering the expression of the chlorophyllase gene *ATHCOR1* in transgenic *Arabidopsis* caused changes in the chlorophyll-to-chlorophyllide ratio. *Plant Physiol* 128:1255–1263
- Block MA, Tewari AK, Albrieux C, Marechal E, Joyard J (2002) The plant *S*-adenosyl-L-methionine:Mg-protoporphyrin IX methyltransferase is located in both envelope and thylakoid chloroplast membranes. *Eur J Biochem* 269:240–248
- Bollivar DW (2006) Recent advances in chlorophyll biosynthesis. *Photosynth Res* 90:173–194
- Bollivar DW, Bauer CE (1992) Nucleotide sequence of *S*-adenosyl-L-methionine: magnesium protoporphyrin methyltransferase from *Rhodobacter capsulatus*. *Plant Physiol* 98:408–410
- Bollivar DW, Jiang ZY, Bauer CE, Beale SI (1994a) Heterologous expression of the *bchM* gene product from *Rhodobacter capsulatus* and demonstration that it encodes *S*-adenosyl-L-methionine:Mg-protoporphyrin IX methyltransferase. *J Bacteriol* 176:5290–5296
- Bollivar DW, Suzuki JY, Beatty JT, Dobrowolski JM, Bauer CE (1994b) Directed mutational analysis of bacteriochlorophyll *a* biosynthesis in *Rhodobacter capsulatus*. *J Mol Biol* 237:622–640
- Botler B, May T, Soll J (1998) A protein import receptor in pea chloroplasts, TOC86, is only a proteolytic fragment of a larger polypeptide. *FEBS Lett* 441:59–62
- Brindley AA, Raux E, Leech HK, Schubert HL, Warren MJ (2003) A story of chelatase evolution: identification and characterization of a small 13–15-kDa “ancestral” cobaltochelatase (CbiXS) in the archaea. *J Biol Chem* 278:22388–22395
- Buchanan BB, Balmer Y (2005) Redox regulation: a broadening horizon. *Annu Rev Plant Biol* 56:187–220
- Burke DH, Alberti M, Hearst JE (1993) *bchFNBH* bacteriochlorophyll synthesis genes of *Rhodobacter capsulatus* and identification of the third subunit of light-independent protochlorophyllide reductase in bacteria and plants. *J Bacteriol* 175:2414–2422
- Chekounova E, Voronetskaya V, Papenbrock J, Grimm B, Beck CF (2001) Characterization of *Chlamydomonas* mutants defective in the H subunit of Mg-chelatase. *Mol Genet Genomics* 266: 363–373
- Chew AG, Bryant DA (2007) Characterization of a plant-like protochlorophyllide a divinyl reductase in green sulfur bacteria. *J Biol Chem* 282:2967–2975
- Chisholm SW, Frankel SL, Goericke R, Olson RJ, Palenik B, Waterbury JB, West-Johnsrud L, Zettler ER (1992) *Prochlorococcus marinus* nov. gen. nov. sp.: A marine prokaryote containing divinylchlorophyll *a* and *b*. *Arch Microbiol* 157: 297–300
- Coleman ML, Chisholm SW (2007) Code and context: *Prochlorococcus* as a model for cross-scale biology. *Trends Microbiol* 15:398–407
- Coomer SA, Chaudhri M, Connor A, Britton G, Hunter CN (1990) Localized transposon *Tn5* mutagenesis of the photosynthetic gene cluster of *Rhodobacter sphaeroides*. *Mol Microbiol* 4:977–989
- Dahlin C, Sundqvist C, Timko MP (1995) The in vitro assembly of the NADPH-protochlorophyllide oxidoreductase in pea chloroplasts. *Plant Mol Biol* 29:317–330
- Davison PA, Schubert HL, Reid JD, Iorg CD, Heroux A, Hill CP, Hunter CN (2005) Structural and biochemical characterization of Gun4 suggests a mechanism for its role in chlorophyll biosynthesis. *Biochemistry* 44:7603–7612
- Debussche L, Couder M, Thibaut D, Cameron B, Crouzet J, Blanche F (1992) Assay, purification, and characterization of cobaltochelatase, a unique complex enzyme catalyzing cobalt insertion in hydrogenobyrinic acid *a,c*-diamide during coenzyme B12 biosynthesis in *Pseudomonas denitrificans*. *J Bacteriol* 174: 7445–7451
- Domanski VP, Rüdiger W (2001) On the nature of the two pathways in chlorophyll formation from protochlorophyllide. *Photosynth Res* 68:131–139
- Domanski V, Rassadina V, Gus-Mayer S, Wanner G, Schoch S, Rüdiger W (2003) Characterization of two phases of chlorophyll formation during greening of etiolated barley leaves. *Planta* 216:475–483
- Douglas SE, Reith M (1993) A *bchI* homolog, encoding a subunit of Mg-chelatase, is located on the plastid genomes of red and cryptomonad algae. *J Mar Biotechnol* 1:135–141
- Eckhardt U, Grimm B, Hörtensteiner S (2004) Recent advances in chlorophyll biosynthesis and breakdown in higher plants. *Plant Mol Biol* 56:1–14
- Eggink LL, LoBrutto R, Brune DC, Brusslan J, Yamasato A, Tanaka A, Hooper JK (2004) Synthesis of chlorophyll *b*: localization of chlorophyllide *a* oxygenase and discovery of a stable radical in the catalytic subunit. *BMC Plant Biol* 4:5
- Eriksson M, Moseley JL, Tottey S, Del Campo JA, Quinn J, Kim Y, Merchant S (2004) Genetic dissection of nutritional copper signaling in *Chlamydomonas* distinguishes regulatory and target genes. *Genetics* 168:795–807
- Espineda CE, Linford AS, Devine D, Brusslan JA (1999) The *AtCAO* gene, encoding chlorophyll *a* oxygenase, is required for chlorophyll *b* synthesis in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 96:10507–10511
- Fisherova H (1975) Linkage relationships of recessive chlorophyll mutants in *Arabidopsis thaliana*. *Biol Plant* 17:182–188
- Fodje MN, Hansson A, Hansson M, Olsen JG, Gough S, Willows RD, Al-Karadaghi S (2001) Interplay between an AAA module and an integrin I domain may regulate the function of magnesium chelatase. *J Mol Biol* 311:111–122
- Franck F, Bereza B, Böddi B (1999) Protochlorophyllide-NADP⁺ and protochlorophyllide⁻ NADPH complexes and their regeneration after flash illumination in leaves and etioplast membranes of dark-grown wheat. *Photosynth Res* 59:53–61

- Franck F, Sperling U, Frick G, Pochert B, van Cleve B, Apel K, Armstrong GA (2000) Regulation of etioplast pigment–protein complexes, inner membrane architecture, and protochlorophyllide a chemical heterogeneity by light-dependent NADPH:protochlorophyllide oxidoreductases A and B. *Plant Physiol* 124:1678–1696
- Frick G, Su Q, Apel K, Armstrong GA (2003) An *Arabidopsis* *porB porC* double mutant lacking light-dependent NADPH:protochlorophyllide oxidoreductases B and C is highly chlorophyll-deficient and developmentally arrested. *Plant J* 35:141–153
- Fuesler TP, Castelfranco PA, Wong Y-S (1984a) Formation of Mg-containing chlorophyll precursors from protoporphyrin IX, δ -aminolevulinic acid, and glutamate in isolated, photosynthetically competent, developing chloroplasts. *Plant Physiol* 74:928–933
- Fuesler TP, Wong Y-S, Castelfranco PA (1984b) Localization of Mg-chelatase and Mg-protoporphyrin IX monomethyl ester (oxidative) cyclase activities within isolated, developing cucumber chloroplasts. *Plant Physiol* 75:662–664
- Fujita Y, Bauer CE (2000) Reconstitution of light-independent protochlorophyllide reductase from purified bchl and BchN-BchB subunits. In vitro confirmation of nitrogenase-like features of a bacteriochlorophyll biosynthesis enzyme. *J Biol Chem* 275:23583–23588
- Fujita Y, Bauer CE (2003) The light-independent protochlorophyllide reductase: a nitrogenase-like enzyme catalyzing a key reaction for greening in the dark. In: Kadish KM, Smith KM, Guillard R (eds) *The porphyrin handbook*, vol 13. Elsevier Science, Amsterdam, pp 71–108
- Fujita Y, Takagi H, Hase T (1998) Cloning of the gene encoding a protochlorophyllide reductase: the physiological significance of the co-existence of light-dependent and -independent protochlorophyllide reduction systems in the cyanobacterium *Plectonema boryanum*. *Plant Cell Physiol* 39:177–185
- Fusada N, Masuda T, Kuroda H, Shiraishi T, Shimada H, Ohta H, Takamiya K (2000) NADPH-protochlorophyllide oxidoreductase in cucumber is encoded by a single gene and its expression is transcriptionally enhanced by illumination. *Photosynth Res* 64:147–154
- Gadjieva R, Axelsson E, Olsson U, Hansson M (2005) Analysis of *gun* phenotype in barley magnesium chelatase and Mg-protoporphyrin IX monomethyl ester cyclase mutants. *Plant Physiol Biochem* 43:901–908
- Gibson LC, Willows RD, Kannangara CG, von Wettstein D, Hunter CN (1995) Magnesium-protoporphyrin chelatase of *Rhodobacter sphaeroides*: reconstitution of activity by combining the products of the *bchH*, *-I*, and *-D* genes expressed in *Escherichia coli*. *Proc Natl Acad Sci USA* 92:1941–1944
- Gibson LCD, Marrison JL, Leech RM, Jensen PE, Bassham DC, Gibson M, Hunter CN (1996) A putative Mg-chelatase subunit from *Arabidopsis thaliana* cv C24. *Plant Physiol* 111:61–71
- Gibson LC, Jensen PE, Hunter CN (1999) Magnesium chelatase from *Rhodobacter sphaeroides*: initial characterization of the enzyme using purified subunits and evidence for a BchI-BchD complex. *Biochem J* 337:243–251
- Gorchein A, Gibson LCD, Hunter CN (1993) Gene expression and control of enzymes for synthesis of magnesium protoporphyrin monomethyl ester in *Rhodobacter sphaeroides*. *Biochem Soc Trans* 21:201S
- Goslings D, Meskauskiene R, Kim C, Lee KP, Nater M, Apel K (2004) Concurrent interactions of heme and FLU with Glu tRNA reductase (HEMA1), the target of metabolic feedback inhibition of tetrapyrrole biosynthesis, in dark- and light-grown *Arabidopsis* plants. *Plant J* 40:957–967
- Gough SP, Petersen BO, Duus JO (2000) Anaerobic chlorophyll isocyclic ring formation in *Rhodobacter capsulatus* requires a cobalamin cofactor. *Proc Natl Acad Sci USA* 97:6908–6913
- Gräfe S, Saluz H-P, Grimm B, Hänel F (1999) Mg-chelatase of tobacco: the role of the subunit CHL D in the chelation step of protoporphyrin IX. *Proc Natl Acad Sci USA* 96:1941–1946
- Hansson M, Kannangara CG (1997) ATPases and phosphate exchange activities in magnesium chelatase subunits of *Rhodobacter sphaeroides*. *Proc Natl Acad Sci USA* 94:13351–13356
- Hansson A, Kannangara CG, von Wettstein D, Hansson M (1999) Molecular basis for semidominance of missense mutations in the XANTHA-H (42-kDa) subunit of magnesium chelatase. *Proc Natl Acad Sci USA* 96:1744–1749
- Hansson A, Willows RD, Roberts TH, Hansson M (2002) Three semidominant barley mutants with single amino acid substitutions in the smallest magnesium chelatase subunit form defective AAA⁺ hexamers. *Proc Natl Acad Sci USA* 99:13944–13949
- Harpaz-Saad S, Azoulay T, Arazi T, Ben-Yaakov E, Mett A, Shibolet Y, Hörtensteiner S, Gidoni D, Gal-On A, Goldschmidt EE, Eyal Y (2007) Chlorophyllase is a rate-limiting enzyme in chlorophyll catabolism and is posttranslationally regulated. *Plant Cell* 19:1007–1022
- Harper AL, von Gesjen SE, Linford AS, Peterson MP, Faircloth RS, Thissen MM, Brusslan JA (2004) Chlorophyllide *a* oxygenase mRNA and protein levels correlate with the chlorophyll *a/b* ratio in *Arabidopsis thaliana*. *Photosynth Res* 79:149–159
- Heyes DJ, Hunter CN (2005) Making light work of enzyme catalysis: protochlorophyllide oxidoreductase. *Trends Biochem Sci* 30:642–649
- Hinchigeri SB, Hundle B, Richards WR (1997) Demonstration that the BchH protein of *Rhodobacter capsulatus* activates S-adenosyl-L-methionine:magnesium protoporphyrin IX methyltransferase. *FEBS Lett* 407:337–342
- Holtorf H, Reinbothe S, Reinbothe C, Bereza B, Apel K (1995) Two routes of chlorophyllide synthesis that are differentially regulated by light in barley (*Hordeum vulgare* L.). *Proc Natl Acad Sci USA* 92:3254–3258
- Hörtensteiner S (2004) The loss of green color during chlorophyll degradation—a prerequisite to prevent cell death? *Planta* 219:191–194
- Hudson A, Carpenter R, Doyle S, Coen ES (1993) *Olive*: a key gene required for chlorophyll biosynthesis in *Antirrhinum majus*. *EMBO J* 12:3711–3719
- Ikegami A, Motohashi K, Yoshimura N, Takahashi S, Romano PGN, Hisabori T, Takamiya K, Masuda T (2007) The CHL1 subunit of *Arabidopsis thaliana* magnesium chelatase is a target protein of the chloroplast thioredoxin. *J Biol Chem* 282:19282–19291
- Ishijima S, Uchibori A, Takagi H, Maki R, Ohnishi M (2003) Light-induced increase in free Mg²⁺ concentration in spinach chloroplasts: measurement of free Mg²⁺ by using a fluorescent probe and necessity of stromal alkalinization. *Arch Biochem Biophys* 412:126–132
- Ito H, Tanaka Y, Tsuji H, Tanaka A (1993) Conversion of chlorophyll *b* to chlorophyll *a* by isolated cucumber etioplasts. *Arch Biochem Biophys* 306:148–151
- Ito H, Takaichi S, Tsuji H, Tanaka A (1994) Properties of synthesis of chlorophyll *a* from chlorophyll *b* in cucumber etioplasts. *J Biol Chem* 269:22034–22038
- Ito H, Ohtsuka T, Tanaka A (1996) Conversion of chlorophyll *b* to chlorophyll *a* via 7-hydroxymethyl chlorophyll. *J Biol Chem* 271:1475–1479
- Jacob-Wilk D, Holland D, Goldschmidt EE, Riou J, Eyal Y (1999) Chlorophyll breakdown by chlorophyllase: isolation and functional expression of the Chlase 1 gene from ethylene-treated Citrus fruit and its regulation during development. *Plant J* 20:653–661
- Jarvis P, Soll J (2001) Toc, Tic, and chloroplast protein import. *Biochim Biophys Acta* 1541:64–79
- Jarvis P, Chen L-J, Li H-M, Peto CA, Fankhauser C, Chory J (1998) An *Arabidopsis* mutant defective in the plastid general protein import apparatus. *Science* 282:100–103

- Jensen PE, Gibson LCD, Henningsen KW, Hunter CN (1996a) Expression of the *chlI*, *chlD*, and *chlH* genes from the cyanobacterium *Synechocystis* PCC6803 in *Escherichia coli* and demonstration that the three cognate proteins are required for magnesium-protoporphyrin chelatase activity. *J Biol Chem* 271:16662–16667
- Jensen PE, Willows RD, Petersen BL, Vothknecht UC, Stummann BM, Kannangara CG, von Wettstein D, Henningsen KW (1996b) Structural genes for Mg-chelatase subunits in barley: *Xantha-f*, *-g* and *-h*. *Mol Gen Genet* 250:383–394
- Jensen PE, Gibson LCD, Hunter CN (1998) Determinants of catalytic activity with the use of purified I, D and H subunits of the magnesium protoporphyrin IX chelatase from *Synechocystis* PCC6803. *Biochem J* 334:335–344
- Jensen PE, Gibson LC, Hunter CN (1999a) ATPase activity associated with the magnesium-protoporphyrin IX chelatase enzyme of *Synechocystis* PCC6803: evidence for ATP hydrolysis during Mg^{2+} insertion, and the $MgATP$ -dependent interaction of the ChII and ChID subunits. *Biochem J* 339:127–134
- Jensen PE, Gibson LCD, Shephard F, Smith V, Hunter CN (1999b) Introduction of a new branchpoint in tetrapyrrole biosynthesis in *Escherichia coli* by co-expression of genes encoding the chlorophyll-specific enzymes magnesium chelatase and magnesium protoporphyrin methyltransferase. *FEBS Lett* 455:349–354
- Jensen PE, Reid JD, Hunter CN (2000) Modification of cysteine residues in the ChII and ChIH subunits of magnesium chelatase results in enzyme inactivation. *Biochem J* 352:435–441
- Kagan RM, Clarke S (1994) Widespread occurrence of three sequence motifs in diverse *S*-adenosyl methionine-dependent methyltransferases suggests a common structure for these enzymes. *Arch Biochem Biophys* 310:417–427
- Karger GA, Reid JD, Hunter CN (2001) Characterization of the binding of deuteroporphyrin IX to the magnesium chelatase H subunit and spectroscopic properties of the complex. *Biochemistry* 40:9291–9299
- Kariola T, Brader G, Li J, Palva ET (2005) Chlorophyllase 1, a damage control enzyme, affects the balance between defense pathways in plants. *Plant Cell* 17:282–294
- Kim C, Apel K (2004) Substrate-dependent and organ-specific chloroplast protein import in planta. *Plant Cell* 16:88–98
- Kim C, Ham H, Apel K (2005) Multiplicity of different cell- and organ-specific import routes for the NADPH-protochlorophyllide oxidoreductases A and B in plastids of *Arabidopsis* seedlings. *Plant J* 42:329–340
- Kollosov VL, Rebeiz CA (2003) Chloroplast biogenesis 88: protochlorophyllide *b* occurs in green but not in etiolated plants. *J Biol Chem* 278:49675–49678
- Koncz C, Mayerhofer R, Koncz-Kalman Z, Nawrath C, Reiss B, Redei GP, Schell J (1990) Isolation of a gene encoding a novel chloroplast protein by T-DNA tagging in *Arabidopsis thaliana*. *EMBO J* 9:1337–1346
- Koussevitzky S, Nott A, Mockler TC, Hong F, Sachetto-Martins G, Surpin M, Lim J, Mittler R, Chory J (2007) Signals from chloroplasts converge to regulate nuclear gene expression. *Science* 316:715–719
- Kropat J, Oster U, Rüdiger W, Beck CF (1997) Chlorophyll precursors are signals of chloroplast origin involved in light induction of nuclear heat-shock genes. *Proc Natl Acad Sci USA* 94:14168–14172
- Kropat J, Oster U, Rüdiger W, Beck CF (2000) Chloroplast signalling in the light induction of nuclear *HSP70* genes requires the accumulation of chlorophyll precursors and their accessibility to cytoplasm/nucleus. *Plant J* 24:523–531
- Kusaba M, Ito H, Morita R, Iida S, Sato Y, Fujimoto M, Kawasaki S, Tanaka R, Hirochika H, Nishimura M, Tanaka A (2007) Rice NON-YELLOW COLORING1 is involved in light-harvesting complex II and grana degradation during leaf senescence. *Plant Cell* 19:1362–1375
- Kuzuyama T, Seki T, Dairi T, Hidaka T, Seto H (1995) Nucleotide sequence of fortimicin KL1 methyltransferase gene isolated from *Micromonospora olivasterospora*, and comparison of its deduced amino acid sequence with those of methyltransferases involved in the biosynthesis of bialaphos and fosfomycin. *J Antibiot* 48:1191–1193
- Lake V, Olsson U, Willows RD, Hansson M (2004) ATPase activity of magnesium chelatase subunit I is required to maintain subunit D in vivo. *Eur J Biochem* 271:2182–2188
- Larkin RM, Alonso JM, Ecker JR, Chory J (2003) GUN4, a regulator of chlorophyll synthesis and intracellular signaling. *Science* 299:902–906
- Lecerof D, Fodje M, Hansson A, Hansson M, Al-Karadaghi S (2000) Structural and mechanistic basis of porphyrin metallation by ferrochelatase. *J Mol Biol* 297:221–232
- Maloney MA, Hoober JK, Marks DB (1989) Kinetics of chlorophyll accumulation and formation of chlorophyll–protein complexes during greening of *Chlamydomonas reinhardtii* y-1 at 38°C. *Plant Physiol* 91:1100–1106
- Masuda T, Takamiya K (2004) Novel insights into the enzymology, regulation and physiological functions of light-dependent protochlorophyllide oxidoreductase in angiosperms. *Photosynth Res* 81:1–29
- Masuda T, Inoue K, Masuda M, Nagayama M, Tamaki A, Ohta H, Shimada H, Takamiya K (1999) Magnesium insertion by magnesium chelatase in the biosynthesis of zinc bacteriochlorophyll *a* in an aerobic acidophilic bacterium *Acidiphilium rubrum*. *J Biol Chem* 274:33594–33600
- Masuda T, Polle JE, Melis A (2002) Biosynthesis and distribution of chlorophyll among the photosystems during recovery of the green alga *Dunaliella salina* from irradiance stress. *Plant Physiol* 128:603–614
- Masuda T, Fusada N, Oosawa N, Takamatsu K, Yamamoto YY, Ohto M, Nakamura K, Goto K, Shibata D, Shirano Y, Hayashi H, Kato T, Tabata S, Shimada H, Ohta H, Takamiya K (2003a) Functional analysis of isoforms of NADPH:protochlorophyllide oxidoreductase (POR), PORB and PORC, in *Arabidopsis thaliana*. *Plant Cell Physiol* 44:963–974
- Masuda T, Tanaka A, Melis A (2003b) Chlorophyll antenna size adjustments by irradiance in *Dunaliella salina* involve coordinate regulation of chlorophyll *a* oxygenase (CAO) and *Lhcb* gene expression. *Plant Mol Biol* 51:757–771
- Matsumoto F, Obayashi T, Sasaki-Sekimoto Y, Ohta H, Takamiya K, Masuda T (2004) Gene expression profiling of the tetrapyrrole metabolic pathway in *Arabidopsis* with a mini-array system. *Plant Physiol* 135:2379–2391
- Meskauskiene R, Nater M, Goslings D, Kessler F, op den Camp R, Apel K (2001) FLU: a negative regulator of chlorophyll biosynthesis in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 98:12826–12831
- Mochizuki N, Brusslan JA, Larkin R, Nagatani A, Chory J (2001) *Arabidopsis* genomes uncoupled 5 (GUN5) mutant reveals the involvement of Mg-chelatase H subunit in plastid-to-nucleus signal transduction. *Proc Natl Acad Sci USA* 98:2053–2058
- Moseley J, Quinn J, Eriksson M, Merchant S (2000) The *CrdI* gene encodes a putative di-iron enzyme required for photosystem I accumulation in copper deficiency and hypoxia in *Chlamydomonas reinhardtii*. *EMBO J* 19:2139–2151
- Moulin M, Smith AG (2005) Regulation of tetrapyrrole biosynthesis in higher plants. *Biochem Soc Trans* 33:737–742
- Moulin M, Smith AG (2008) A robust method for determination of chlorophyll intermediates by tandem mass spectrometry. In: Allen J, Gantt E, Golbeck J, Osmond B (eds) *Energy from the sun*. Springer (in press)

- Nagata N, Satoh S, Tanaka R, Tanaka A (2004) Domain structures of chlorophyllide *a* oxygenase of green plants and *Prochlorothrix hollandica* in relation to catalytic functions. *Planta* 218: 1019–1025
- Nagata N, Tanaka R, Satoh S, Tanaka A (2005) Identification of a vinyl reductase gene for chlorophyll synthesis in *Arabidopsis thaliana* and implications for the evolution of *Prochlorococcus* species. *Plant Cell* 17:233–240
- Nakagawara E, Sakuraba Y, Yamasato A, Tanaka R, Tanaka A (2007) Clp protease controls chlorophyll *b* synthesis by regulating the level of chlorophyllide *a* oxygenase. *Plant J* 49:800–809
- Nakanishi H, Nozue H, Suzuki K, Kaneko Y, Taguchi G, Hayashida N (2005) Characterization of the *Arabidopsis thaliana* mutant *pcb2* which accumulates divinyl chlorophylls. *Plant Cell Physiol* 46:467–473
- Nakayama M, Masuda T, Sato N, Yamagata H, Bowler C, Ohta H, Shioi Y, Takamiya K (1995) Cloning, subcellular localization and expression of CHL1, a subunit of magnesium-chelatase in soybean. *Biochem Biophys Res Commun* 215:422–428
- Nakayama M, Masuda T, Bando T, Yamagata H, Ohta H, Takamiya K (1998) Cloning and expression of the soybean *chlH* gene encoding a subunit of Mg-chelatase and localization of the Mg²⁺ concentration-dependent ChlH protein within the chloroplast. *Plant Cell Physiol* 39:275–284
- Neuwald AF, Aravind L, Spouge JL, Koonin EV (1999) AAA⁺: a class of chaperone-like ATPases associated with the assembly, operation, and disassembly of protein complexes. *Genome Res* 9:27–43
- Nomata J, Kitashima M, Inoue K, Fujita Y (2006) Nitrogenase Fe protein-like Fe-S cluster is conserved in L-protein (BchL) of dark-operative protochlorophyllide reductase from *Rhodobacter capsulatus*. *FEBS Lett* 580:6151–6154
- Nott A, Jung HS, Koussevitzky S, Chory J (2006) Plastid-to-nucleus retrograde signaling. *Annu Rev Plant Biol* 57:739–759
- Obayashi T, Kinoshita K, Nakai K, Shibaoka M, Hayashi S, Saeki M, Shibata D, Saito K, Ohta H (2007) ATTED-II: a database of co-expressed genes and cis elements for identifying co-regulated gene groups in *Arabidopsis*. *Nucleic Acids Res* 35:D863–D869
- Oosawa N, Masuda T, Awai K, Fusada N, Shimada H, Ohta H, Takamiya K (2000) Identification and light-induced expression of a novel gene of NADPH-protochlorophyllide oxidoreductase isoform in *Arabidopsis thaliana*. *FEBS Lett* 474:133–136
- Orsat B, Monfort A, Chatellard P, Stutz E (1992) Mapping and sequencing of an actively transcribed *Euglena gracilis* chloroplast gene (*ccsA*) homologous to the *Arabidopsis thaliana* nuclear genome *cs* (*ch-42*). *FEBS Lett* 303:181–184
- Oster U, Rüdiger W (1997) The *G4* gene of *Arabidopsis thaliana* encodes a chlorophyll synthase of etiolated plants. *Bot Acta* 110:420–423
- Oster U, Bauer CE, Rüdiger W (1997) Characterization of chlorophyll *a* and bacteriochlorophyll *a* synthases by heterologous expression in *Escherichia coli*. *J Biol Chem* 272:9671–9676
- Oster U, Tanaka R, Tanaka A, Rüdiger W (2000) Cloning and functional expression of the gene encoding the key enzyme for chlorophyll *b* biosynthesis (CAO) from *Arabidopsis thaliana*. *Plant J* 21:305–310
- Ouchane S, Steunou AS, Picaud M, Astier C (2004) Aerobic and anaerobic Mg-protoporphyrin monomethyl ester cyclases in purple bacteria: a strategy adopted to bypass the repressive oxygen control system. *J Biol Chem* 279:6385–6394
- Papenbrock J, Gräfe S, Kruse E, Hänel F, Grimm B (1997) Mg-chelatase of tobacco: identification of a *Chl D* cDNA sequence encoding a third subunit, analysis of the interaction of the three subunits with the yeast two-hybrid system, and reconstitution of the enzyme activity by co-expression of recombinant CHL D, CHL H and CHL I. *Plant J* 12:981–990
- Papenbrock J, Mock H-P, Kruse E, Grimm B (1999) Expression studies in tetrapyrrole biosynthesis: inverse maxima of magnesium chelatase and ferrochelatase activity during cyclic photoperiods. *Planta* 208:264–273
- Parham R, Rebeiz CA (1995) Chloroplast biogenesis 72: a [4-vinyl]chlorophyllide *a* reductase assay using divinyl chlorophyllide *a* as an exogenous substrate. *Anal Biochem* 231:164–169
- Pattanayak GK, Tripathy BC (2002) Catalytic function of a novel protein protochlorophyllide oxidoreductase C of *Arabidopsis thaliana*. *Biochem Biophys Res Commun* 291:921–924
- Pattanayak GK, Biswal AK, Reddy VS, Tripathy BC (2005) Light-dependent regulation of chlorophyll *b* biosynthesis in chlorophyllide *a* oxygenase overexpressing tobacco plants. *Biochem Biophys Res Commun* 326:466–471
- Petersen BL, Jensen PE, Gibson LC, Stummann BM, Hunter CN, Henningsen KW (1998) Reconstitution of an active magnesium chelatase enzyme complex from the bchl, -D, and -H gene products of the green sulfur bacterium *Chlorobium vibrioforme* expressed in *Escherichia coli*. *J Bacteriol* 180:699–704
- Petersen BL, Kannangara CG, Henningsen KW (1999a) Distribution of ATPase and ATP-to-ADP phosphate exchange activities in magnesium chelatase subunits of *Chlorobium vibrioforme* and *Synechocystis* PCC6803. *Arch Microbiol* 171:146–150
- Petersen BL, Moller MG, Jensen PE, Henningsen KW (1999b) Identification of the *Xan-g* gene and expression of the Mg-chelatase encoding genes *Xan-f*, -g and -h in mutant and wild type barley (*Hordeum vulgare* L.). *Hereditas* 131:165–170
- Philippart K, Geis T, Ilkavets I, Oster U, Schwenkert S, Meurer J, Soll J (2007) Chloroplast biogenesis: the use of mutants to study the etioplast-chloroplast transition. *Proc Natl Acad Sci USA* 104:678–683
- Pinta V, Picaud M, Reiss-Husson F, Astier C (2002) *Rubrivivax gelatinosus* acsF (previously orf358) codes for a conserved, putative binuclear-iron-cluster-containing protein involved in aerobic oxidative cyclization of Mg-protoporphyrin IX monomethylester. *J Bacteriol* 184:746–753
- Pollmann S, Springer A, Buhr F, Lahroussi A, Samol I, Bonneville JM, Tichtinsky G, von Wettstein D, Reinbothe C, Reinbothe S (2007) A plant *Porphyrin* related to defects in plastid import of protochlorophyllide oxidoreductase A. *Proc Natl Acad Sci USA* 104:2019–2023
- Pontier D, Albrieux C, Joyard J, Lagrange T, Block MA (2007) Knock-out of the magnesium protoporphyrin IX methyltransferase gene in *Arabidopsis*. Effects on chloroplast development and on chloroplast-to-nucleus signaling. *J Biol Chem* 282:2297–2304
- Porra RJ (1997) Recent progress in porphyrin and chlorophyll biosynthesis. *Photochem Photobiol* 65:492–516
- Porra RJ, Urzinger M, Winkler J, Bubenzer C, Scheer H (1998) Biosynthesis of the 3-acetyl and 13(1)-oxo groups of bacteriochlorophyll *a* in the facultative aerobic bacterium, *Rhodovulum sulfidophilum*-the presence of both oxygenase and hydratase pathways for isocyclic ring formation. *Eur J Biochem* 257: 185–191
- Pruzinska A, Tanner G, Aubry S, Anders I, Moser S, Muller T, Ongania KH, Krautler B, Youn JY, Liljegen SJ, Hörtensteiner S (2005) Chlorophyll breakdown in senescent *Arabidopsis* leaves. Characterization of chlorophyll catabolites and of chlorophyll catabolic enzymes involved in the degreening reaction. *Plant Physiol* 139:52–63
- Rebeiz CA, Kolosov VL, Briskin D, Gawienowski M (2003) Chloroplast biogenesis: chlorophyll biosynthetic heterogeneity, multiple biosynthetic routes and biological spin-offs. In: Nalwa HS (ed) handbook of photochemistry and photobiology, vol 4. American Scientific Publishers, Los Angeles, pp 183–248
- Reid JD, Hunter CN (2002) Current understanding of the function of magnesium chelatase. *Biochem Soc Trans* 30:643–645

- Reid JD, Hunter CN (2004) Magnesium-dependent ATPase activity and cooperativity of magnesium chelatase from *Synechocystis* sp. PCC6803. *J Biol Chem* 279:26893–26899
- Reid JD, Siebert CA, Bullough PA, Hunter CN (2003) The ATPase activity of the ChlI subunit of magnesium chelatase and formation of a heptameric AAA⁺ ring. *Biochemistry* 42:6912–6920
- Reinbothe S, Reinbothe C, Runge S, Apel K (1995a) Enzymatic product formation impairs both the chloroplast receptor-binding function as well as translocation competence of the NADPH: protochlorophyllide oxidoreductase, a nuclear-encoded plastid precursor protein. *J Cell Biol* 129:299–308
- Reinbothe S, Runge S, Reinbothe C, van Cleve B, Apel K (1995b) Substrate-dependent transport of the NADPH:protochlorophyllide oxidoreductase into isolated plastids. *Plant Cell* 7:161–172
- Reinbothe S, Reinbothe C, Neumann D, Apel K (1996) A plastid enzyme arrested in the step of precursor translocation in vivo. *Proc Natl Acad Sci USA* 93:12026–12030
- Reinbothe C, Lebedev N, Apel K, Reinbothe S (1997) Regulation of chloroplast protein import through a protochlorophyllide-responsive transit peptide. *Proc Natl Acad Sci USA* 94:8890–8894
- Reinbothe C, Lebedev N, Reinbothe S (1999) A protochlorophyllide light-harvesting complex involved in de-etiolation of higher plants. *Nature* 397:80–84
- Reinbothe S, Mache R, Reinbothe C (2000) A second, substrate-dependent site of protein import into chloroplasts. *Proc Natl Acad Sci USA* 97:9795–9800
- Reinbothe C, Buhr F, Pollmann S, Reinbothe S (2003a) In vitro reconstitution of light-harvesting POR-protochlorophyllide complex with protochlorophyllides *a* and *b*. *J Biol Chem* 278:807–815
- Reinbothe C, Lepinat A, Deckers M, Beck E, Reinbothe S (2003b) The extra loop distinguishing POR from the structurally related short-chain alcohol dehydrogenases is dispensable for pigment binding but needed for the assembly of light-harvesting POR-protochlorophyllide complex. *J Biol Chem* 278:816–822
- Reinbothe S, Pollmann S, Reinbothe C (2003c) *In situ* conversion of protochlorophyllide *b* to protochlorophyllide *a* in barley. Evidence for a novel role of 7-formyl reductase in the prolamellar body of etioplasts. *J Biol Chem* 278:800–806
- Reinbothe S, Quigley F, Gray J, Schemenewitz A, Reinbothe C (2004a) Identification of plastid envelope proteins required for import of protochlorophyllide oxidoreductase A into the chloroplast of barley. *Proc Natl Acad Sci USA* 101:2197–2202
- Reinbothe S, Quigley F, Springer A, Schemenewitz A, Reinbothe C (2004b) The outer plastid envelope protein Oep16: role as precursor translocase in import of protochlorophyllide oxidoreductase A. *Proc Natl Acad Sci USA* 101:2203–2208
- Reinbothe S, Pollmann S, Springer A, James RJ, Tichtinsky G, Reinbothe C (2005) A role of Toc33 in the protochlorophyllide-dependent plastid import pathway of NADPH:protochlorophyllide oxidoreductase (POR). *Plant J* 42:1–12
- Reinbothe C, Bartsch S, Eggink LL, Hooper JK, Brusslan J, Andrade-Paz R, Monnet J, Reinbothe S (2006) A role for chlorophyllide *a* oxygenase in the regulated import and stabilization of light-harvesting chlorophyll *a/b* proteins. *Proc Natl Acad Sci USA* 103:4777–4782
- Rissler HM, Collakova E, DellaPenna D, Whelan J, Pogson BJ (2002) Chlorophyll biosynthesis. Expression of a second chl I gene of magnesium chelatase in *Arabidopsis* supports only limited chlorophyll synthesis. *Plant Physiol* 128:770–779
- Rodermel S (2001) Pathways of plastid-to-nucleus signaling. *Trends Plant Sci* 6:471–478
- Rodermel S, Park S (2003) Pathways of intracellular communication: tetrapyrroles and plastid-to-nucleus signaling. *Bioessays* 25: 631–636
- Rüdiger W (2002) Biosynthesis of chlorophyll *b* and the chlorophyll cycle. *Photosynth Res* 74:187–193
- Rüdiger W, Benz J, Guthoff C (1980) Detection and partial characterization of activity of chlorophyll synthetase in etioplast membranes. *Eur J Biochem* 109:193–200
- Rüdiger W, Böhm S, Helfrich M, Schulz S, Schoch S (2005) Enzymes of the last steps of chlorophyll biosynthesis: modification of the substrate structure helps to understand the topology of the active centers. *Biochemistry* 44:10864–10872
- Ryberg M, Sundqvist C (1988) The regular ultrastructure of isolated prolamellar bodies depends on the presence of membrane-bound NADPH-protochlorophyllide oxidoreductase. *Physiol Plant* 73:216–226
- Rzeznicka K, Walker CJ, Westergren T, Kannangara CG, von Wettstein D, Merchant S, Gough SP, Hansson M (2005) *Xanthal* encodes a membrane subunit of the aerobic Mg-protoporphyrin IX monomethyl ester cyclase involved in chlorophyll biosynthesis. *Proc Natl Acad Sci USA* 102:5886–5891
- Satoh S, Tanaka A (2006) Identification of chlorophyllide *a* oxygenase in the *Prochlorococcus* genome by a comparative genomic approach. *Plant Cell Physiol* 47:1622–1629
- Satoh S, Ikeuchi M, Mimuro M, Tanaka A (2001) Chlorophyll *b* expressed in *Cyanobacteria* functions as a light-harvesting antenna in photosystem I through flexibility of the proteins. *J Biol Chem* 276:4293–4297
- Sawers RJ, Viney J, Farmer PR, Bussey RR, Olsefski G, Anufrikova K, Hunter CN, Brutnell TP (2006) The maize *Oil yellow1* (*Oy1*) gene encodes the I subunit of magnesium chelatase. *Plant Mol Biol* 60:95–106
- Schemenewitz A, Pollmann S, Reinbothe C, Reinbothe S (2007) A substrate-independent, 14:3:3 protein-mediated plastid import pathway of NADPH:protochlorophyllide oxidoreductase A. *Proc Natl Acad Sci USA* 104:8538–8543
- Scheumann V, Schoch S, Rüdiger W (1998) Chlorophyll *a* formation in the chlorophyll *b* reductase reaction requires reduced ferredoxin. *J Biol Chem* 273:35102–35108
- Scheumann V, Klement H, Helfrich M, Oster U, Schoch S, Rüdiger W (1999) Protochlorophyllide *b* does not occur in barley etioplasts. *FEBS Lett* 445:445–448
- Schmid HC, Oster U, Kogel J, Lenz S, Rüdiger W (2001) Cloning and characterisation of chlorophyll synthase from *Avena sativa*. *Biol Chem* 382:903–911
- Schoefs B, Franck F (2003) Protochlorophyllide reduction: mechanisms and evolutions. *Photochem Photobiol* 78:543–557
- Shen YY, Wang XF, Wu FQ, Du SY, Cao Z, Shang Y, Wang XL, Peng CC, Yu XC, Zhu SY, Fan RC, Xu YH, Zhang DP (2006) The Mg-chelatase H subunit is an abscisic acid receptor. *Nature* 443:823–826
- Shedbalkar VP, Ioannides IM, Rebeiz CA (1991) Chloroplast biogenesis. Detection of monovinyl protochlorophyll(ide) *b* in plants. *J Biol Chem* 266:17151–17157
- Shepherd M, Hunter CN (2004) Transient kinetics of the reaction catalysed by magnesium protoporphyrin IX methyltransferase. *Biochem J* 382:1009–1013
- Shioi Y, Sasa T (1983) Esterification of chlorophyllide *b* in higher plants. *Biochim Biophys Acta* 756:127–131
- Sirijovski N, Olsson U, Lundqvist J, Al-Karadaghi S, Willows RD, Hansson M (2006) ATPase activity associated with the magnesium chelatase H-subunit of the chlorophyll biosynthetic pathway is an artifact. *Biochem J* 400:477–484
- Smith CA, Suzuki JY, Bauer CE (1996) Cloning and characterization of the chlorophyll biosynthesis gene *chlM* from *Synechocystis* PCC 6803 by complementation of a bacteriochlorophyll biosynthesis mutant of *Rhodospirillum rubrum*. *Plant Mol Biol* 30:1307–1314
- Smith MD, Rounds CM, Wang F, Chen K, Afithile M, Schnell DJ (2004) atToc159 is a selective transit peptide receptor for the import of nucleus-encoded chloroplast proteins. *J Cell Biol* 165:323–334

- Soldatova O, Apchelimov A, Radukina N, Ezhova T, Shestakov S, Ziemann V, Hedtke B, Grimm B (2005) An Arabidopsis mutant that is resistant to the protoporphyrinogen oxidase inhibitor acifluorfen shows regulatory changes in tetrapyrrole biosynthesis. *Mol Genet Genomics* 273:311–318
- Spano AJ, He Z, Michel H, Hunt DF, Timko MP (1992) Molecular cloning, nuclear gene structure, and developmental expression of NADPH: protochlorophyllide oxidoreductase in pea (*Pisum sativum* L.). *Plant Mol Biol* 18:967–972
- Sperling U, van Cleve B, Frick G, Apel K, Armstrong GA (1997) Overexpression of light-dependent PORA or PORB in plants depleted of endogenous POR by far-red light enhances seedling survival in white light and protects against photooxidative damage. *Plant J* 12:649–658
- Sperling U, Franck F, van Cleve B, Frick G, Apel K, Armstrong GA (1998) Etioplast differentiation in *Arabidopsis*: both PORA and PORB restore the prolamellar body and photoactive protochlorophyllide-F655 to the cop1 photomorphogenic mutant. *Plant Cell* 10:283–296
- Strand A (2004) Plastid-to-nucleus signalling. *Curr Opin Plant Biol* 7:621–625
- Strand A, Asami T, Alonso J, Ecker JR, Chory J (2003) Chloroplast to nucleus communication triggered by accumulation of Mg-protoporphyrin IX. *Nature* 421:79–83
- Sundqvist C, Dahlin C (1997) With chlorophyll from prolamellar bodies to light-harvesting complexes. *Physiol Plant* 100:748–759
- Susek RE, Ausubel FM, Chory J (1993) Signal transduction mutants of Arabidopsis uncouple nuclear *CAB* and *RBCS* gene expression from chloroplast development. *Cell* 74:787–799
- Takamiya KI, Tsuchiya T, Ohta H (2000) Degradation pathway(s) of chlorophyll: what has gene cloning revealed? *Trends Plant Sci* 5:426–431
- Tanaka R, Tanaka A (2005) Effects of chlorophyllide *a* oxygenase overexpression on light acclimation in *Arabidopsis thaliana*. *Photosynth Res* 85:327–340
- Tanaka A, Tanaka R (2006) Chlorophyll metabolism. *Curr Opin Plant Biol* 9:248–255
- Tanaka R, Tanaka A (2007) Tetrapyrrole biosynthesis in higher plants. *Annu Rev Plant Biol* 58:321–346
- Tanaka A, Ito H, Tanaka R, Tanaka NK, Yoshida K, Okada K (1998) Chlorophyll *a* oxygenase (CAO) is involved in chlorophyll *b* formation from chlorophyll *a*. *Proc Natl Acad Sci USA* 95:12719–12723
- Tanaka R, Koshino Y, Sawa S, Ishiguro S, Okada K, Tanaka A (2001) Overexpression of chlorophyllide *a* oxygenase (CAO) enlarges the antenna size of photosystem II in *Arabidopsis thaliana*. *Plant J* 26:365–373
- Teakle GR, Griffiths WT (1993) Cloning, characterization and import studies on protochlorophyllide reductase from wheat (*Triticum aestivum*). *Biochem J* 296:225–230
- Tomitani A, Okada K, Miyashita H, Matthijs HCP, Ohno T, Tanaka A (1999) Chlorophyll *b* and phycobilins in the common ancestor of cyanobacteria and chloroplasts. *Nature* 400:159–162
- Totter S, Block MA, Allen M, Westergren T, Albrieux C, Scheller HV, Merchant S, Jensen PE (2003) Arabidopsis CHL27, located in both envelope and thylakoid membranes, is required for the synthesis of protochlorophyllide. *Proc Natl Acad Sci USA* 100:16119–16124
- Tsuchiya T, Ohta H, Okawa K, Iwamatsu A, Shimada H, Masuda T, Takamiya K (1999) Cloning of chlorophyllase, the key enzyme in chlorophyll degradation: finding of a lipase motif and the induction by methyl jasmonate. *Proc Natl Acad Sci USA* 96:15362–15367
- Tsuchiya T, Suzuki T, Yamada T, Shimada H, Masuda T, Ohta H, Takamiya K (2003) Chlorophyllase as a serine hydrolase: identification of a putative catalytic triad. *Plant Cell Physiol* 44:96–101
- Vavilin DV, Vermaas WF (2002) Regulation of the tetrapyrrole biosynthetic pathway leading to heme and chlorophyll in plants and cyanobacteria. *Physiol Plant* 115:9–24
- Verdecia MA, Larkin RM, Ferrer JL, Riek R, Chory J, Noel JP (2005) Structure of the Mg-chelatase cofactor GUN4 reveals a novel hand-shaped fold for porphyrin binding. *PLoS Biol* 3:e151
- von Gromoff ED, Schroda M, Oster U, Beck CF (2006) Identification of a plastid response element that acts as an enhancer within the *Chlamydomonas HSP70A* promoter. *Nucleic Acids Res* 34:4767–4779
- Wakao N, Yokoi N, Isoyama N, Hiraishi A, Shimada K, Kobayashi M, Kise H, Iwaki M, Itoh S, Takaichi S, Sakurai Y (1996) Discovery of natural photosynthesis using Zn-containing bacteriochlorophyll in an aerobic bacterium *Acidiphilium rubrum*. *Plant Cell Physiol* 37:889–893
- Walker CJ, Weinstein JD (1991a) Further characterization of the magnesium chelatase in isolated developing cucumber chloroplasts. *Plant Physiol* 95:1189–1196
- Walker CJ, Weinstein JD (1991b) In vitro assay of the chlorophyll biosynthetic enzyme Mg-chelatase: resolution of the activity into soluble and membrane-bound fractions. *Proc Natl Acad Sci USA* 88:5789–5793
- Walker CJ, Weinstein JD (1994) The magnesium-insertion step of chlorophyll biosynthesis is a two-stage reaction. *Biochem J* 299:277–284
- Walker CJ, Willows RD (1997) Mechanism and regulation of Mg-chelatase. *Biochem J* 327:321–333
- Walker CJ, Mansfield KE, Rezzano IN, Hanamoto CM, Smith KM, Castelfranco PA (1988) The magnesium-protoporphyrin IX (oxidative) cyclase system. Studies on the mechanism and specificity of the reaction sequence. *Biochem J* 255:685–692
- Walker CJ, Mansfield KE, Smith KM, Castelfranco PA (1989) Incorporation of atmospheric oxygen into the carbonyl functionality of the protochlorophyllide isocyclic ring. *Biochem J* 257:599–602
- Wilde A, Mikolajczyk S, Alawady A, Lokstein H, Grimm B (2004) The *gun4* gene is essential for cyanobacterial porphyrin metabolism. *FEBS Lett* 571:119–123
- Willows RD (2003) Biosynthesis of chlorophylls from protoporphyrin IX. *Nat Prod Rep* 20:327–341
- Willows RD, Beale SI (1998) Heterologous expression of the *Rhodobacter capsulatus Bchl*, *-D*, and *-H* genes that encode magnesium chelatase subunits and characterization of the reconstituted enzyme. *J Biol Chem* 273:34206–34213
- Willows RD, Gibson LCD, Kanangara CG, Hunter CN, von Wettstein D (1996) Three separate proteins constitute the magnesium chelatase of *Rhodobacter sphaeroides*. *Eur J Biochem* 235:438–443
- Willows RD, Hansson A, Birch D, Al-Karadaghi S, Hansson M (2004) EM single particle analysis of the ATP-dependent Bchl complex of magnesium chelatase: an AAA⁺ hexamer. *J Struct Biol* 146:227–233
- Wu Q, Vermaas WF (1995) Light-dependent chlorophyll *a* biosynthesis upon *chlL* deletion in wild-type and photosystem I-less strains of the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Mol Biol* 29:933–945
- Xiong J, Fischer WM, Inoue K, Nakahara M, Bauer CE (2000) Molecular evidence for the early evolution of photosynthesis. *Science* 289:1724–1730
- Yamasato A, Nagata N, Tanaka R, Tanaka A (2005) The N-terminal domain of chlorophyllide *a* oxygenase confers protein instability in response to chlorophyll *B* accumulation in Arabidopsis. *Plant Cell* 17:1585–1597

- Yamazaki S, Nomata J, Fujita Y (2006) Differential operation of dual protochlorophyllide reductases for chlorophyll biosynthesis in response to environmental oxygen levels in the cyanobacterium *Leptolyngbya boryana*. *Plant Physiol* 142:911–922
- Yang ZM, Bauer CE (1990) *Rhodobacter capsulatus* genes involved in early steps of the bacteriochlorophyll biosynthetic pathway. *J Bacteriol* 172:5001–5010
- Zhang H, Li J, Yoo JH, Yoo SC, Cho SH, Koh HJ, Seo HS, Paek NC (2006) Rice *Chlorina-1* and *Chlorina-9* encode ChlD and ChII subunits of Mg-chelatase, a key enzyme for chlorophyll synthesis and chloroplast development. *Plant Mol Biol* 62:325–337
- Zsebo KM, Hearst JE (1984) Genetic-physical mapping of a photosynthetic gene cluster from *R. capsulata*. *Cell* 38:937–947