

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/50228857>

Genomics of the NF- κ B signaling pathway: Hypothesized role in ovarian cancer

ARTICLE in CANCER CAUSES AND CONTROL · FEBRUARY 2011

Impact Factor: 2.74 · DOI: 10.1007/s10552-011-9745-4 · Source: PubMed

CITATIONS

13

READS

39

6 AUTHORS, INCLUDING:



Kairi Kalli

119 PUBLICATIONS 3,002 CITATIONS

SEE PROFILE



Ellen L Goode

Mayo Foundation for Medical Education and ...

262 PUBLICATIONS 8,219 CITATIONS

SEE PROFILE

Published in final edited form as:

Cancer Causes Control. 2011 May ; 22(5): 785–801. doi:10.1007/s10552-011-9745-4.

Genomics of the NF- κ B signaling pathway: hypothesized role in ovarian cancer

Kristin L. White,

Department of Health Sciences Research, Mayo Clinic College of Medicine, 200 First Street SW, Rochester, MN 55905, USA

David N. Rider,

Department of Health Sciences Research, Mayo Clinic College of Medicine, 200 First Street SW, Rochester, MN 55905, USA

Kimberly R. Kalli,

Department of Oncology, Mayo Clinic College of Medicine, 200 First Street SW, Rochester, MN 55905, USA

Keith L. Knutson,

Department of Immunology, Mayo Clinic College of Medicine, 200 First Street SW, Rochester, MN 55905, USA

Gail P. Jarvik, and

University of Washington, Seattle, WA, USA

Ellen L. Goode

Department of Health Sciences Research, Mayo Clinic College of Medicine, 200 First Street SW, Rochester, MN 55905, USA

Ellen L. Goode: egoode@mayo.edu

Abstract

Objective—We sought to review evidence linking nuclear factor-kappa B (NF- κ B) to ovarian cancer and to identify genetic variants involved in NF- κ B signaling.

Methods—PubMed was reviewed to inform on ovarian cancer biology and NF- κ B signaling and to identify key genes. Public linkage disequilibrium (LD) data were analyzed to identify informative inherited variants (tagSNPs) using ldSelect.

Results—We identified 319 key NF- κ B genes including five NF- κ B subunits, 167 activating genes, and 55 inhibiting genes. We found that the 1000 Genomes Project was the most informative LD source for most genes (92.8%), and we identified 13,027 LD bins ($r^2 \geq 0.9$, minor allele frequency ≥ 0.05) and 1,018 putative-functional variants worthy of investigation. We also report that reliance on a commonly used genome-wide SNP array and genotype imputation with HapMap Phase II data provides data on only 74% of the common inherited NF- κ B SNPs of interest.

Conclusions—Compelling evidence suggests that NF- κ B plays a critical role in ovarian cancer, yet inherited variation in these genes has not been thoroughly assessed in relation to disease risk or

outcome. We present a collection of variants in key genes and suggest creation of a custom genotyping array as an optimal approach.

Keywords

Association studies; Genetic variation; NF-kappaB; Etiology; tagSNPs; Single nucleotide polymorphisms

Introduction

Ovarian cancer is the fifth leading cause for cancer death among women in the United States [1]. Due to a lack of effective screening strategies and the non-specific nature of early signs and symptoms associated with this disease, most patients are diagnosed with advanced disease. For the approximate 25% of women who are diagnosed with disease confined to the ovary or ovaries, 5-year survival rates are high (75–90%); however, for the 75% of women diagnosed with stage III and IV disease, the likelihood of long-term disease-free survival is low (15–20%) [2–4]. Identification of factors associated with risk and clinical outcome is essential to understand the pathogenesis of ovarian cancer, enabling preventative and therapeutic advances to be made in this rare but deadly disease.

Recent studies indicate that there are two distinct types of ovarian cancer, Type I and Type II [5], and that a proportion of cases of both types likely originate in pelvic tissues outside the ovary. Ovarian cancer is originally thought to develop de novo from the ovarian surface epithelium that is derived from mesodermal epithelium and contains no müllerian-type tissue. However, serous, endometrioid, and clear cell ovarian carcinomas are all derived from müllerian-type tissue indicating that ovarian tumors may not develop de novo in the ovary, but instead develop in müllerian-derived tissues outside ovary (such as the cervix, endometrium, and fallopian tubes) and involve the ovary secondarily [6–8]. Type I ovarian cancers (clear cell, mucinous carcinomas, and low-grade serous and endometrioid) are usually confined to the ovaries at diagnosis, develop from borderline tumor precursors, and are relatively genetically stable with typical somatic mutations, including *KRAS*, *BRAF*, *ERBB2*, and *PIK3CA* but not *TP53* [9]. Type II ovarian cancers (high-grade serous, high-grade endometrioid, malignant mixed mesodermal tumors, and undifferentiated carcinomas) are more aggressive, are believed to develop from intraepithelial carcinomas in the fallopian tube, and are genetically unstable with characteristic somatic *TP53* mutations but rarely mutations characteristic of Type I tumors [7].

Previous data also suggest that ovarian cancer pathogenesis may be dependent on the inflammatory response within the ovarian surface epithelium. This hypothesis is supported by an etiological hypothesis of “incessant ovulation,” which postulates that increased lifetime ovulations are associated with increased risk of ovarian cancer [10]. Ovulation causes repeated trauma to the ovarian surface epithelium that requires repair mediated by the inflammatory response through the recruitment of leukocytes, cytokines, and chemokines, release of nitrous oxide, DNA repair, and tissue restructuring [10–14]. This restructuring can result in epithelial cells trapped internally in inclusion cysts, where the local microenvironment appears to induce them to gain quasi-neoplastic characteristics such as increased p53 accumulation and a ninefold higher proliferation index than surface epithelial cells [15].

Nuclear factor-kappaB (NF-κB) refers to a family of transcription factors (also called the Rel family) that regulate the expression of over 400 genes. These regulated genes are critical to numerous cellular processes and include a multitude of cytokines and chemokines, pro-inflammatory genes, adhesion molecules, acute phase response proteins, complement

factors, stress response genes, oxidative enzymes, apoptosis genes, growth factors, and other transcription factors [16–18]. Due to this regulatory role, NF- κ B has been implicated in several diseases including cardiovascular disease and various cancers [19, 20]. Evidence indicates that NF- κ B is constitutively activated in many cancer cell lines, influencing cancer progression and response to chemotherapy [21, 22]. Therefore, we set out to review the hypothesis that NF- κ B may play a role in ovarian cancer specifically and to identify the optimal approach for a large-scale study of inherited variation in this pathway and ovarian cancer risk.

A role for NF- κ B in ovarian cancer

The NF- κ B family of transcription factors has received much attention for its role in cancer since the discovery that c-rel is the human homologue of the avian retroviral oncogene v-rel [23]. Subsequent work has demonstrated roles for NF- κ B-mediated signaling in promoting survival from pro-apoptotic stimuli, angiogenesis, proliferation, and inflammation—all characteristics of malignancy [24]. All of these NF- κ B-influenced functions appear to play a role in ovarian cancer, underlying the importance of further understanding NF- κ B-mediated events in order to better identify women at risk of developing ovarian cancer and to facilitate development of more effective therapeutic strategies.

NF- κ B has been shown to be important directly in tumor cells. For example, the mitogen lysophosphatidic acid that is found in ascites fluid may induce proliferation by increasing levels of cyclin D1, an important cell cycle control protein [25]. LPA-induced increases in activity of a cyclin D1 promoter-driven reporter gene required the NF- κ B site. Ovarian tumors can acquire enhanced NF- κ B functions that allow them to circumvent apoptotic pathways and afford protection against environmental insults such as anti-tumor immune effectors or chemotherapy. Inhibition of NF- κ B activity in vitro or in human cell line xenografts increases the sensitivity of the cells to apoptosis induced by both cisplatin [26] and paclitaxel [27], drugs representing the platinum/taxane combination most commonly used as first-line treatment for ovarian cancer. Ovarian cancers are resistant to tumor necrosis factor alpha (TNF α)-induced apoptosis because tumor cells can upregulate pathways leading to the activation of NF- κ B and induction of apoptosis inhibitory proteins [28]. In addition, using a dominant-negative I κ B α construct to block NF- κ B activity in the SKOV3 ovarian cancer cell line resulted in reduced expression of VEGF and IL-8, leading to reduced tumorigenicity, invasiveness, and angiogenesis in athymic mice [29]. Similar reductions were seen in cell line invasiveness, IL-8 production, and anchorage-independent growth after treatment with an IKK β inhibitor in experiments performed after determining that elevated IKK β levels and a 9-gene “IKK β signature” were associated with worse outcome in ovarian cancer patients [30].

In addition to studies that assess all ovarian cancer cells as one population, cells phenotypically and functionally characterized as potential ovarian cancer stem cells have been shown to have constitutively activated NF- κ B [31]. The group characterizing these cells further determined that expression of transcripts from the microRNA cluster miRNA199A2/214 are lower in the CD44-positive stem cell-like population than in CD44-negative, “mature” ovarian cancer cells, and that decreases in expression of this specific miRNA cluster are associated with decreased expression of Twist1, a transcription factor that controls IKK β levels [32]. Because Twist1 has an antagonistic function in NF- κ B-mediated cytokine expression, it appears that the result of reduced expression of this miRNA cluster and decreased Twist 1 levels is increased NF- κ B activity. Due to the strong potential for chemoresistant stem cells to contribute to ovarian cancer recurrence after initially effective chemotherapy, it is clear that understanding the potential role of NF- κ B in driving stem cell activity is a critical area of study.

In addition to direct actions in tumor cells, NF- κ B-mediated events also contribute to the microenvironment and peritoneal inflammation associated with ovarian cancer. For example, activated thrombin, found in the inflammatory environment of the peritoneum in ovarian cancer patients, appears to induce monocytes to develop a tumor-associated macrophage phenotype. Monocytes treated with activated thrombin have elevated levels of activated NF- κ B; supernatants from these cells increase ovarian cancer cell invasion in vitro. This increased invasiveness can be blocked with a neutralizing antibody targeting IL-8, an NF- κ B-driven cytokine [33]. The central role of NF- κ B in inflammatory responses has developed increased interest in the context of ovarian cancer over the last decade because of the clear association of anti-tumor immune responses with improved outcome [34]. Ovarian tumors are infiltrated with a variety of immune effectors, including anti-tumor T-cells [35–38]. The type of immune infiltrate influences the disease course, with elevated levels of suppressive T regulatory cells associated with poor outcome [38] and higher CD8/Treg ratios associated with improved survival [39]. The type of anti-tumor immune response that develops is controlled by chemokines and cytokines produced by infiltrating tumor macrophages, dendritic cells, and the tumors themselves to control the mobilization of the various T-cell subsets. For example, one mechanism by which ovarian cancer modulates immunity is through the production of interleukin-10 (IL-10). Synthesized directly by the tumor and infiltrating immune effectors, IL-10 blocks NF- κ B activation in T-cells and dendritic cells and prevents activation of effective ovarian cancer anti-tumor immunity [40–42].

Because of the vast body of evidence supporting the importance of NF- κ B-regulated events to ovarian cancer cells and the anti-tumor immune response, we hypothesize that much of the unexplained risk of this inflammation-related cancer is due to genetic variations in the NF- κ B pathway [43].

Nuclear factor-kappaB subunits and primary signaling pathways

NF- κ B is present in cells in an uninduced state, requiring activation to initiate nuclear translocation and the ability to influence gene expression. It consists of five members, p50, p52, p65, Rel-B, and c-Rel [44], that can form multiple homo- and hetero-dimers (p65/p50, p65/p65, p50/c-Rel, p50/p50, c-Rel/c-Rel, and p52/p52), the most common of which are p65/p50 and p52/Rel-B [45, 46]. The NF- κ B dimers have specific non-redundant functions in various tissue and cell types [22, 46]. p50, p65, and p52 are found in most cell types and at the highest concentration in the thymus, whereas Rel-B and c-Rel are primarily found in lymphoid tissues [46].

The primary NF- κ B signaling pathway is referred to as the canonical pathway (Fig. 1). Stimulation by TNF α and other ligands initiates a signaling cascade that assembles the I κ B kinase (IKK) complex and promotes the degradation of the inhibitors of κ B (I κ B), resulting in the activation and nuclear translocation of NF- κ B dimers [16]. The canonical pathway is particularly important for the ability of the myeloid (innate) cells of the immune system, including monocytes, macrophages, and dendritic cells, to communicate with the adaptive immune system [47]. NF- κ B is also important for the innate immune cells to sense “danger” through a family of molecules known as the toll-like receptors (TLRs) [48].

Much less is understood about the non-canonical NF- κ B pathway. This signaling cascade involves ligation by CD40, B-cell activating factor (BAFF), and lymphotoxin beta (LT β), subsequent IKK α activation, and poly-ubiquitination and partial degradation of p100, which yields activated p52 and Rel-B dimers (Fig. 1) [45, 46, 49, 50]. In contrast to the canonical pathway, the non-canonical NF- κ B receptor-mediated pathway is important for lymphoid organ development and the adaptive immune response [47]. Targeted disruptions in the individual NF- κ B subunits can lead to abnormal splenic architecture. Due to the spleen’s

important role in both human and mouse immunity, it is likely that these splenic abnormalities play a role in some forms of immunodeficiency. NF- κ B is also known to play key roles in the architecture of other immune organs such as the lymph nodes, which are important for the eradication of both infection and malignancy. One of the best-characterized roles of the non-canonical pathway is in the development of an adaptive immune response. For example, targeted disruption of IKK α expression blocks maturation of B-cells in the germinal centers of lymph nodes, indicating the important role of IKK α in the acquisition of humoral immunity [51].

A role for polymorphisms

Inherited variation in key genes may account for some portion of ovarian cancer susceptibility. Familial aggregation of ovarian and breast cancer has long been established, and account for an estimated 5–10% of epithelial ovarian cancer cases [52, 53]. Highly penetrant mutations in *BRCA1*, *BRCA2*, and DNA mismatch repair genes explain less than 40% of familial epithelial ovarian cancer risk [54]; however, approximately two-thirds of families with three or more cases of familial breast or ovarian cancer had *BRCA1* or *BRCA2* mutations [55, 56]. The remaining inherited risk is most likely due to combinations of common, less-penetrant alleles [57–61]. The most powerful method for identification of moderate risk alleles is a genetic association study examining frequencies of variants such as single-nucleotide polymorphisms (SNPs) in candidate genes of key biological pathways, which has been successful in identifying new risk alleles for a variety of cancers [62–64], or via genome-wide association studies (GWAS). Thus far, variants in the 2q31, 3q25, 8q24, 9p22, 17q21, and 19p13 chromosomal regions [65–67] and in genes involved in cell cycle control [68], steroid hormone metabolism [69], DNA repair [70], and one-carbon metabolism [71] have been associated with ovarian cancer susceptibility. Additional evidence is accumulating that inherited factors, including those in inflammation, relate to clinical outcome in ovarian cancer [72–75].

The identification of polymorphisms worthy of study in given genes can draw on putative-functional or location data (reviewed by Bhatti et al. [76]) or can rely on informative tagSNPs selected on the basis of linkage disequilibrium (LD, the correlation of alleles along a chromosome). The structure of LD for common variants has been mapped by the International HapMap Project using genome-wide genotyping [77], and the LD structure of both common and rarer variants is improved by the 1000 Genomes Project [78] using genome-wide sequencing. These projects allow investigators to harness the power of LD for common disease prediction [79–83]. Due to the strong LD within a region, only a few SNPs need to be assayed to successfully predict all of the common genetic variation once SNP correlations are estimated for a core of individuals. If LD structure within a particular gene is known, careful selection of a reduced set of SNPs that identify or “tag” a particular haplotype may provide as much information as genotyping all available SNPs [84–86]. These tagSNPs then theoretically represent all genetic variation exceeding a specified minor allele frequency (MAF). Thus, by genotyping the LD-selected subset of common variants, it is possible to perform a comprehensive analysis of all common variants in a candidate gene.

The purpose of the current investigation is to identify and classify the highest priority genes within the NF- κ B signaling pathway, to evaluate publicly available LD data in the encoding genes, and to select a panel of SNPs representing genetic variation within these genes. The resulting set of annotated genes and variants is now available to the scientific community.

Methods

Classification of genes

Using the PubMed1 search engine in December 2009, we identified genes involved in NF- κ B signaling through a literature search including the phrases “NF-kappaB,” “pathway,” and “gene.” Genes were categorized into NF- κ B subpathways (canonical, non-canonical, or both canonical and non-canonical) and biological function within the subpathways. Biological function definitions were as follows: subunits (NF- κ B proteins), activation (role in signaling cascade leading to NF- κ B subunit activation), inhibition (inhibitory role), degradation (involved in proteasomal degradation), and nuclear function (nuclear proteins involved in transcription).

Evaluation of genotype data sources

We assessed three types of large-scale LD characterization projects: (1) genome-wide sequencing efforts by the 1000 Genomes Project^{2,3}; (2) genome-wide SNP genotyping projects, including those of the International HapMap Project⁴ and Perlegen Sciences⁵; (3) gene-centric resequencing projects, such as several NHLBI Program for Genomic Applications (PGAs),^{6,7,8} SeattleSNPs,⁹ the National Institute of Environmental Health Sciences Environmental Genome Project SNPs Program (NIEHS SNPs),¹⁰ Pharmacogenetics and Risk of Cardiovascular Disease (PARC),¹¹ and Molecular Diversity and Epidemiology of Common Disease (MDECODE).¹² Based on European or European-American data availability for NF- κ B genes, we identified, from among these sources, those sources which were appropriate for LD-based SNP selection and applied the tagSNP selection approach described below.

LD binning and choice of genotype data source

Within each genotyped data source and for each gene, we binned SNPs with $MAF \geq 0.05$ that were pairwise correlated at $r^2 \geq 0.9$ using the algorithm implemented in ldSelect [87]. For genome-wide projects (the 1000 Genomes Project, the International HapMap Project, and Perlegen Sciences), gene regions were defined as 5 kb upstream and downstream using genomic coordinates based on RefSeq release 29 (NCBI build 36). For genes with data available in multiple LD sources, we then identified the “best source” of data as the source with the highest number of LD bins after tagSNP selection. It is conservative to include a greater number of tagSNPs rather than a smaller number of tagSNPs in the event that one source overestimated LD. For example, even with an equal number of underlying SNPs, if Source A indicated a larger number of LD bins than Source B, we proceed with Source A’s tagSNPs as analysis of data from Source A samples suggested lower LD within the gene and thus the need for more LD bins than Source B. More often, however, the source with the greatest number of LD bins was also the source with the greatest number of underlying SNPs, thus clearly capturing the greatest amount of genetic variation.

¹<http://www.ncbi.nlm.nih.gov/pubmed/>.

²<http://www.1000genomes.org/>; accessed April 2010.

³ftp://ftp-trace.ncbi.nih.gov/1000genomes/ftp/pilot_data/release/2010_03/pilot1/CEU.SRP000031.2010_03.genotypes.vcf.gz.

⁴<http://www.hapmap.org/>; accessed November 2008.

⁵<http://genome.perlegen.com/>; accessed November 2008.

⁶<http://pga.swmed.edu/>; accessed November 2008.

⁷<http://cardiogenomics.med.harvard.edu/home>; accessed November 2008.

⁸<http://innateimmunity.net/>; accessed November 2008.

⁹<http://pga.mbt.washington.edu/>; accessed November 2008.

¹⁰<http://egp.gs.washington.edu/>; accessed November 2008.

¹¹<http://droog.gs.washington.edu/parc/>; accessed November 2008.

¹²<http://droog.gs.washington.edu/mdecode/>; accessed November 2008.

SNP selection

After LD binning using the best data source for each gene, we identified particular tagSNPs to assay within each bin. Using Illumina iSelect HD Custom Genotyping annotation (San Diego, CA), we prioritized tagSNPs first based on expected assay performance. One tagSNP was selected in bins with less than ten SNPs, two tagSNPs were selected in bins with 10–29 SNPs, and three tagSNPs were selected in bins with 30 or more SNPs. Within each gene, additional putative-functional SNPs with $MAF \geq 0.05$ (that were not tagSNPs) were also identified as non-synonymous, 5' UTR, 3' UTR, 1 kb 5' upstream, residing in a splice site (dbSNP version 129), or residing in an miRNA-binding site.^{13,14}

Comparison to genome-wide panels

In order to assess the utility of creating a custom NF- κ B SNP panel versus use of a commercially available platform, we compared our prioritized SNPs to: (1) SNPs available on the Illumina Infinium HD Human610-Quad BeadChip genome-wide SNP panel (San Diego, CA) that has been used in prior ovarian cancer GWAS [65] and (2) SNPs in the International HapMap Project Phase II which prior ovarian cancer GWAS have used as the basis for imputation of up to 2.5 million SNPs [65]. We estimated “coverage” of SNPs within NF- κ B genes versus those available in two ways: (1) as the proportion of NF- κ B LD bins with at least one tagSNP included on these panels and (2) as the proportion of NF- κ B SNPs either tagged or directly assessed in bins included on these panels.

Results

Identification of the key components of NF- κ B signaling

Our review identified 319 key genes in the NF- κ B signaling pathway; these are classified in Table 1 by subpathway (canonical, non-canonical, or both) and biological function (subunits, activation, inhibition, degradation, and nuclear function). We excluded the 26S proteasome (degradatory role) and RNA polymerase II (nuclear role) from consideration due to a lack of specificity to NF- κ B signaling and an excessive number of genes encoding the many subunits required for the functional protein complexes. Supplemental Table 1 provides detailed information for each gene including protein descriptions, location, size, and references attesting to each gene's relevance in NF- κ B signaling. The major components of the NF- κ B signaling pathway are summarized below.

Activation of NF- κ B subunits—Components of NF- κ B signaling cascades include ligands, receptors, and intracellular target molecules that lead to IKK complex activation, I κ B degradation, and activation of NF- κ B dimers. The primary canonical NF- κ B signaling pathway is initiated by binding of TNF α , a proinflammatory cytokine produced mainly by macrophages, but also by mast cells, lymphoid cells, endothelial cells, and fibroblasts (Fig. 2) [88, 89]. Interleukin-1 (IL-1), B-cell activating factor belonging to the TNF family BAFF [90], CD40 ligand [18], lymphotoxin beta (LT β) [91], tumor necrosis factor receptor superfamily, member 11a, NF- κ B activator (called RANK) [92], and T-cell receptor (TCR) [18] and lipopolysaccharide (LPS) ligands [88] binding also initiate NF- κ B signaling (Figs. 2, 3, 4; Table 1, Supplemental Table 1). After ligation, specific receptors recruit and activate intracellular target molecules, eventually leading to activation of the IKK complex [88, 93, 94]. In the canonical pathway, the IKK complex contains three kinases, IKK α , IKK β , and NF- κ B essential modulator (NEMO or IKK γ), with various roles in the activation of NF- κ B. IKK β is responsible for phosphorylating I κ B, marking it for proteolysis [95]. NEMO is a

¹³<http://www.microrna.org/microrna/home.do>.

¹⁴<http://www.targetscan.org/>.

non-enzymatic regulatory subunit that modulates and aids in IKK α and IKK β activation. In the non-canonical pathway, IKK α initiates processing of inhibitory p100 to functional p52 [46]. Activation of the IKK complex leads to processing of p100 and p105 and degradation of I κ B, freeing NF- κ B to enter the nucleus [46, 95]. For more details, see Fig. 2 for TNF α [88, 96–103] and TCR ligand signaling pathways [18, 104–114], Fig. 3 for IL-1 [88, 96–98, 115–125] and LPS signaling pathways [18, 88, 126–129], and Fig. 4 for the non-canonical signaling pathway [50, 130, 131].

Inhibition—Several molecules play inhibitory roles by seizing key molecules, preventing apoptosis, or interfering with proteasomal degradation (Table 1, Supplemental Table 1) [93]. The primary NF- κ B inhibitors are the seven members of the I κ B family, I κ B α , I κ B β , I κ B ϵ , I κ B γ , p100, p105, and BCL-3 [44]. The IKK complex phosphorylates I κ Bs at two serine residues, marking the inhibitors for poly-ubiquitination by the SCF $^{\beta}$ -TrCP complex and subsequent degradation by the 26S proteasome. Without ubiquitination and proteolysis of I κ B, NF- κ B complexes remain inactive in the cytoplasm [44, 46]. Preliminary evidence indicates that each I κ B family member has a unique and specific role in the NF- κ B pathway [132–135]. I κ B α , I κ B β , I κ B ϵ , and I κ B γ mask NF- κ B's nuclear localization signal (NLS), which sequesters NF- κ B in the cytoplasm, preventing nuclear translocation and transcription [132]. I κ B α and I κ B β have a similar binding affinity for NF- κ B; however, I κ B α also contains a nuclear export signal (NES) that enables I κ B α to shuttle NF- κ B complexes out of the nucleus. I κ B β does not contain a NES, but is more effective than I κ B α at masking NF- κ B's NLS and preventing p65 and p50 translocation [46, 132–135]. p100 and p105 are inhibitory precursors of p52 and p50, respectively, that are partially degraded, yielding functional p52 and p50. Although evidence is unclear, BCL-3 does not appear to have an inhibitory role, but instead may facilitate activation of p50 and p52 in the nucleus [44].

Degradation—Proteolysis of inhibitory molecules is critical in regulating activation of the NF- κ B signaling cascade [93]. The SCF $^{\beta}$ -TrCP complex marks the majority of inhibitors in the NF- κ B pathway for degradation through poly-ubiquitination. The main components of this complex are beta-transducin repeat containing (β -TrCP) [136–138], S-phase kinase-associated protein 1 (Skp1), cullin 1 (Cul1), ring-box 1 (RBX1 or Roc1), and ubiquitin-conjugating enzyme E2D 1 (UBE2D1 or HbcH5; Table 1, Supplemental Table 1) [139]. This complex recognizes phosphorylated inhibitors, and, through a series of events, it ubiquitinates these inhibitors, thus marking them for partial or complete degradation by the 26S proteasome [93]. Figure 5 provides more detail on the SCF $^{\beta}$ -TrCP complex partial degradation of p105/p100 and poly-ubiquitination of I κ B [18, 93, 139–143].

Nuclear function—After nuclear translocation, DNA-bound, transcription-promoting NF- κ B complexes associate with the RNA polymerase II transcription machinery to initiate gene expression. RNA polymerase II can only initiate transcription after recruitment and assembly of the replication complex components. The general transcription factors, transcription factor for RNA polymerase II (TFII)-A, -B, -E, -F, and -H, contain either a TATA box region, a binding site for nuclear proteins, or an Inr element with a downstream promoter region that functions in a similar manner as the TATA box [93, 144, 145]. Other critical components for transcription of NF- κ B-regulated genes include TATA-binding protein (TBP), 14 TBP-associated factors (TAFs) [93, 145], K(lysine) acetyltransferase 2B (called PCAF) [146], cAMP-responsive element-binding protein 1 (CREB) [147], CREB-binding protein (CBP), and E1A-binding protein p300 (p300) [146, 148]. Figure 6 provides more detail on the components of the RNA polymerase II transcription complex [146–148].

SNP selection

Evaluation of LD data sources—Among the LD sources listed in “Methods”, we found that data from the following were optimal at the time of analysis (Table 2): the genome-wide sequencing 1000 Genomes Project Low-coverage Pilot on 60 unrelated individuals of European descent,^{15,16} the two genome-wide genotyping projects the International Hapmap Project Phase II Release 2417 and Perlegen Sciences,¹⁸ and the three gene resequencing programs SeattleSNPs,¹⁹ NIEHS SNPs,²⁰ and Innate Immunity SNPs.²¹ For 319 NF-κB identified candidate genes, 1000 Genomes Project data were available for all 302 autosomal genes (and none of the 17 chromosome X genes), International HapMap Project data were available for all but one gene (*GTF2H2*), and Perlegen Sciences data were available for 297 candidate genes; SeattleSNPs had resequenced 42 genes, Innate Immunity SNPs had resequenced 21 genes, and NIEHS SNPs had resequenced nine genes. Using data from each source, tagSNP selection was performed for all genes to bin SNPs with $MAF \geq 0.05$ and $r^2 \geq 0.9$. The 1000 Genomes Project was found to be the most informative source for 296 genes (92.8%), the International HapMap Project Phase II for 16 genes (5.0%, including 15 on chromosome X), Innate Immunity SNPs for three genes (0.9%, including one on chromosome X), SeattleSNPs for three genes (0.9%), and NIEHS SNPs for one gene (0.3%, including one on chromosome X); Perlegen Sciences data was not the most informative source for any gene. Supplemental Table 2 provides the most informative source identified for each gene.

TagSNPs and putative-functional SNPs—Summarizing across all genes and using the optimal LD source, 13,027 LD bins covering 36,181 common SNPs (mean 81 bins [range 0–905 bins] per gene, mean 226 SNPs [range 0–3,069 SNPs] per gene) were identified. Putative-functional SNPs, some of which were also tagSNPs, were identified; putative-functional SNPs were identified in 260 genes including 173 non-synonymous SNPs, 81 5' UTR SNPs, 389 3' UTR SNPs, 304 SNPs 1 kb 5' upstream, and 165 SNPs residing in an miRNA-binding site. No splice site SNPs were identified. Supplemental Table 2 provides detailed information about the number of bins within each gene as well as the number of SNPs tagged by tagSNPs. In total, a collection of 14,231 SNPs of interest to genotype in 319 NF-κB genes has been assembled representing common underlying genetic variation among populations of European descent. These SNPs do not attempt to cover rare variants ($MAF < 5\%$).

Coverage on genome-wide arrays—We evaluated coverage of our 13,027 identified NF-κB tagSNPs on the Illumina Infinium HD Human610-Quad BeadChip and in HapMap Phase II data (often used for imputation). Details per gene are provided in Supplemental Table 2, and Fig. 7 summarizes coverage across genes. Generally, use of the Illumina 610-Quad without additional imputation would directly assess 22% of the identified NF-κB bins and provide data on 50% of the total SNPs with $MAF > 0.05$ in each gene. Thus, the NF-κB tagSNPs included on the Illumina 610-Quad tend to tag larger LD bins, reflecting a preference in Illumina's SNP selection for the most informative tagSNPs and placement of lower priority on singleton bins. The approach of dropping smaller bins is reasonable due to cost constraints, because smaller bins convey information on a minimal set of underlying SNPs. However, this coverage is not optimal for many applications. With additional

¹⁵<http://www.1000genomes.org/>; accessed April 2010.

¹⁶ftp-trace.ncbi.nih.gov/1000genomes/ftp/pilot_data/release/2010_03/pilot1/CEU.SRP000031.2010_03.genotypes.vcf.gz.

¹⁷<http://www.hapmap.org>; accessed November 2008.

¹⁸<http://genome.perlegen.com/>; accessed November 2008.

¹⁹<http://pga.mbt.washington.edu/>; accessed November 2008.

²⁰<http://egp.gs.washington.edu>; accessed November 2008.

²¹<http://innateimmunity.net/>; accessed November 2008.

imputation using HapMap Phase II data, coverage increases; on average, 43% of NF- κ B bins would be assessed, providing data on 74% of the included SNPs. Again, larger bins have been identified by HapMap Phase II; however, the current availability of 1000 Genomes Project data indicates that less than optimal NF- κ B coverage is provided by the use of HapMap Phase II imputation. We note that this is not due to the discovery of novel rare variants in 1000 Genomes Project data, as NF- κ B LD binning was limited to SNPs with $MAF > 0.05$. Coverage and statistical power is further reduced with use of HapMap Phase II imputation due to the use of imputation rather than directly genotyping SNPs of interest. To address these issues, GWAS panels are currently under commercial development, which are based on 1000 Genomes Project data, and these will fill a critical void in commercial arrays. In the interim, current ovarian cancer GWAS are not expected to have optimal inclusion of NF- κ B tagSNPs. Custom SNP arrays based on the NF- κ B variants identified here will provide maximal information about these key genes of interest to ovarian cancer and other phenotypes.

Discussion

NF- κ B is a key regulator of expression of hundreds of genes involved in crucial cellular processes related to normal cellular function and carcinogenesis. Its signaling pathway involves a complex cascade of events initiated by immune and other signaling proteins, regulated by several inhibitory molecules, and that eventually leads to NF- κ B-mediated regulation of gene expression. There is an abundance of evidence linking a prolonged, aberrant inflammatory response with ovarian cancer. As the most lethal gynecologic malignancy, the quest for a better understanding of ovarian cancer genomics is critical. Due to NF- κ B's vital role in responding to inflammatory signals, initiating expression of pro-inflammatory genes, and proven links between ovarian carcinogenesis and inflammation, a study examining variations in genes involved in the NF- κ B pathway and ovarian cancer risk is warranted.

Here, we summarized the relevance of the NF- κ B signaling pathway in ovarian cancer and identified 319 genes encoding critical components categorized by subpathway and biological function. Additionally, we identified 14,231 tagSNPs and putative-functional SNPs within each of these genes. Of LD data sources available in April 2010, we found that the data from the 1000 Genomes Project was more informative for the vast majority of genes than that from the International HapMap Project and other projects. We also found that coverage of variation in these genes on the commonly used [65] commercial panel Illumina Infinium HD Human610-Quad BeadChip was suboptimal, even considering imputation of missing genotypes using the International HapMap Project Phase II data. Thus, in order to comprehensively assess inherited variation in the NF- κ B signaling pathway, a customized SNP panel, such as an Illumina iSelect HD Custom Genotyping, is preferable. As new versions of data from the 1000 Genomes Project are unveiled, it will be critical to re-evaluate genomic coverage of these candidate genes. The collection of SNPs we have currently identified is available to the scientific community studying association between variation in NF- κ B signaling and any phenotype, including ovarian and other cancers [149–152] as well as autoimmune [153], vascular [154], and muscular disorders [155].

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was made possible by NCI funding R01-CA122443, Mayo Clinic SPORE in Ovarian Cancer (P50 CA136393), and support from the Mayo Foundation.

References

1. Jemal A, Siegel R, Xu J, Ward E. Cancer statistics, 2010. *CA Cancer J Clin.* 2010; 60(5):277–300. [PubMed: 20610543]
2. Hoskins WJ, Bundy BN, Thigpen JT, Omura GA. The influence of cytoreductive surgery on recurrence-free interval and survival in small-volume stage III epithelial ovarian cancer: a Gynecologic Oncology Group study. *Gynecol Oncol.* 1992; 47(2):159–166. [PubMed: 1468693]
3. McGuire V, Jessor CA, Whitemore AS. Survival among U.S. women with invasive epithelial ovarian cancer. *Gynecol Oncol.* 2002; 84(3):399–403. [PubMed: 11855877]
4. Barnholtz-Sloan JS, Schwartz AG, Qureshi F, Jacques S, Malone J, Munkarah AR. Ovarian cancer: changes in patterns at diagnosis and relative survival over the last three decades. *Am J Obstet Gynecol.* 2003; 189(4):1120–1127. [PubMed: 14586365]
5. Shih Ie M, Kurman RJ. Ovarian tumorigenesis: a proposed model based on morphological and molecular genetic analysis. *Am J Pathol.* 2004; 164(5):1511–1518. [PubMed: 15111296]
6. Dubeau L. The cell of origin of ovarian epithelial tumours. *Lancet Oncol.* 2008; 9(12):1191–1197. [PubMed: 19038766]
7. Kurman RJ, Shih Ie M. The origin and pathogenesis of epithelial ovarian cancer: a proposed unifying theory. *Am J Surg Pathol.* 2010; 34(3):433–443. [PubMed: 20154587]
8. Lauchlan SC. The secondary Mullerian system. *Obstet Gynecol Surv.* 1972; 27(3):133–146. [PubMed: 4614139]
9. Cho KR, Shih Ie M. Ovarian cancer. *Annu Rev Pathol.* 2009; 4:287–313. [PubMed: 18842102]
10. Fathalla MF. Incessant ovulation—a factor in ovarian neoplasia? *Lancet.* 1971; 2(7716):163. [PubMed: 4104488]
11. Bonello N, McKie K, Jasper M, Andrew L, Ross N, Braybon E, et al. Inhibition of nitric oxide: effects on interleukin-1 beta-enhanced ovulation rate, steroid hormones, and ovarian leukocyte distribution at ovulation in the rat. *Biol Reprod.* 1996; 54(2):436–445. [PubMed: 8788197]
12. Zackrisson U, Mikuni M, Wallin A, Delbro D, Hedin L, Brannstrom M. Cell-specific localization of nitric oxide synthases (NOS) in the rat ovary during follicular development, ovulation and luteal formation. *Hum Reprod.* 1996; 11(12):2667–2673. [PubMed: 9021370]
13. Schildkraut JM, Bastos E, Berchuck A. Relationship between lifetime ovulatory cycles and overexpression of mutant p53 in epithelial ovarian cancer. *J Natl Cancer Inst.* 1997; 89(13):932–938. [PubMed: 9214672]
14. Fleming JS, Beaugie CR, Haviv I, Chenevix-Trench G, Tan OL. Incessant ovulation, inflammation and epithelial ovarian carcinogenesis: revisiting old hypotheses. *Mol Cell Endocrinol.* 2006; 247(1–2):4–21. [PubMed: 16297528]
15. Pothuri B, Leitao MM, Levine DA, Viale AS, Olshen AB, Arroyo C, et al. Genetic analysis of the early natural history of epithelial ovarian carcinoma. *PLoS ONE.* 2010; 5(4):e10358. [PubMed: 20436685]
16. Pahl HL. Activators and target genes of Rel/NF-kappaB transcription factors. *Oncogene.* 1999; 18(49):6853–6866. [PubMed: 10602461]
17. Kutuk O, Basaga H. Inflammation meets oxidation: NF-kappaB as a mediator of initial lesion development in atherosclerosis. *Trends Mol Med.* 2003; 9(12):549–557. [PubMed: 14659470]
18. Hayden MS, Ghosh S. Signaling to NF-kappaB. *Genes Dev.* 2004; 18(18):2195–2224. [PubMed: 15371334]
19. Courtois G, Gilmore TD. Mutations in the NF-kappaB signaling pathway: implications for human disease. *Oncogene.* 2006; 25(51):6831–6843. [PubMed: 17072331]
20. Cambien F. Coronary heart disease and polymorphisms in genes affecting lipid metabolism and inflammation. *Curr Atheroscler Rep.* 2005; 7(3):188–195. [PubMed: 15811252]

21. Shishodia S, Aggarwal BB. Nuclear factor-kappaB activation: a question of life or death. *J Biochem Mol Biol.* 2002; 35(1):28–40. [PubMed: 16248967]
22. Sethi G, Sung B, Aggarwal BB. Nuclear factor-kappaB activation: from bench to bedside. *Exp Biol Med (Maywood).* 2008; 233(1):21–31. [PubMed: 18156302]
23. Gilmore TD, et al. Multiple mutations contribute to the oncogenicity of the retroviral oncoprotein v-Rel. *Oncogene.* 1999; 18(49):6925–6937. [PubMed: 10602467]
24. Karin M. Nuclear factor-kappaB in cancer development and progression. *Nature.* 2006; 441(7092): 431–436. [PubMed: 16724054]
25. Hu YL, Albanese C, Pestell RG, Jaffe RB. Dual mechanisms for lysophosphatidic acid stimulation of human ovarian carcinoma cells. *J Natl Cancer Inst.* 2003; 95(10):733–740. [PubMed: 12759391]
26. Mabuchi S, Ohmichi M, Nishio Y, Hayasaka T, Kimura A, Ohta T, et al. Inhibition of NFkappaB increases the efficacy of cisplatin in in vitro and in vivo ovarian cancer models. *J Biol Chem.* 2004; 279(22):23477–23485. [PubMed: 15026414]
27. Mabuchi S, Ohmichi M, Nishio Y, Hayasaka T, Kimura A, Ohta T, et al. Inhibition of inhibitor of nuclear factor-kappaB phosphorylation increases the efficacy of paclitaxel in in vitro and in vivo ovarian cancer models. *Clin Cancer Res.* 2004; 10(22):7645–7654. [PubMed: 15569997]
28. Xiao CW, Yan X, Li Y, Reddy SA, Tsang BK. Resistance of human ovarian cancer cells to tumor necrosis factor alpha is a consequence of nuclear factor kappaB-mediated induction of Fas-associated death domain-like interleukin-1beta-converting enzyme-like inhibitory protein. *Endocrinology.* 2003; 144(2):623–630. [PubMed: 12538625]
29. Huang S, Robinson JB, DeGuzman A, Bucana CD, Fidler IJ. Blockade of Nuclear Factor-kB Signaling Inhibits Angiogenesis and Tumorigenicity of Human Ovarian Cancer Cells by Suppressing Expression of Vascular Endothelial Growth Factor and Interleukin 81. *Cancer Res.* 2000; 60:5334–5339. [PubMed: 11034066]
30. Hernandez L, Hsu SC, Davidson B, Birrer MJ, Kohn EC, Annunziata CM. Activation of NF-kappaB signaling by inhibitor of NF-kappaB kinase beta increases aggressiveness of ovarian cancer. *Cancer Res.* 2010; 70(10):4005–4014. [PubMed: 20424119]
31. Alvero AB, Chen R, Fu H, Montagna M, Schwartz PE, Rutherford T, et al. Molecular phenotyping of human ovarian cancer stem cells unravels the mechanisms for repair and chemoresistance. *Cell Cycle.* 2009; 8(1):158–166. [PubMed: 19158483]
32. Yin G, Chen R, Alvero AB, Fu HH, Holmberg J, Glackin C, et al. TWISTing stemness, inflammation and proliferation of epithelial ovarian cancer cells through MIR199A2/214. *Oncogene.* 2010; 29(24):3545–3553. [PubMed: 20400975]
33. Zhang T, Ma Z, Wang R, Wang Y, Wang S, Cheng Z, et al. Thrombin facilitates invasion of ovarian cancer along peritoneum by inducing monocyte differentiation toward tumor-associated macrophage-like cells. *Cancer Immunol Immunother.* 2010; 59(7):1097–1108. [PubMed: 20352429]
34. Zhang L, Conejo-Garcia JR, Katsaros D, Gimotty PA, Massobrio M, Regnani G, et al. Intratumoral T cells, recurrence, and survival in epithelial ovarian cancer. *N Engl J Med.* 2003; 348(3):203–213. [PubMed: 12529460]
35. Santin AD, Hermonat PL, Ravaggi A, Bellone S, Roman JJ, Smith CV, et al. Phenotypic and functional analysis of tumor-infiltrating lymphocytes compared with tumor-associated lymphocytes from ascitic fluid and peripheral blood lymphocytes in patients with advanced ovarian cancer. *Gynecol Obstet Invest.* 2001; 51(4):254–261. [PubMed: 11408737]
36. Curiel TJ, Wei S, Dong H, Alvarez X, Cheng P, Mottram P, et al. Blockade of B7-H1 improves myeloid dendritic cell-mediated antitumor immunity. *Nat Med.* 2003; 9(5):562–567. [PubMed: 12704383]
37. Negus RP, Stamp GW, Hadley J, Balkwill FR. Quantitative assessment of the leukocyte infiltrate in ovarian cancer and its relationship to the expression of C-C chemokines. *Am J Pathol.* 1997; 150(5):1723–1734. [PubMed: 9137096]
38. Curiel TJ, Coukos G, Zou L, Alvarez X, Cheng P, Mottram P, et al. Specific recruitment of regulatory T cells in ovarian carcinoma fosters immune privilege and predicts reduced survival. *Nat Med.* 2004; 10(9):942–949. [PubMed: 15322536]

39. Sato E, Olson SH, Ahn J, Bundy B, Nishikawa H, Qian F, et al. Intraepithelial CD8 + tumor-infiltrating lymphocytes and a high CD8 +/regulatory T cell ratio are associated with favorable prognosis in ovarian cancer. *Proc Natl Acad Sci USA*. 2005; 102(51):18538–18543. [PubMed: 16344461]
40. Pestka S, Krause CD, Sarkar D, Walter MR, Shi Y, Fisher PB. Interleukin-10 and related cytokines and receptors. *Annu Rev Immunol*. 2004; 22(1):929–979. [PubMed: 15032600]
41. Romano MF, Lamberti A, Petrella A, Bisogni R, Tassone PF, Formisano S, et al. IL-10 inhibits nuclear factor-kappa B/Rel nuclear activity in CD3- stimulated human peripheral T lymphocytes. *J Immunol*. 1996; 156(6):2119–2123. [PubMed: 8690900]
42. Matsuzaki J, Gnjatic S, Mhawech-Fauceglia P, Beck A, Miller A, Tsuji T, et al. Tumor-infiltrating NY-ESO-1-specific CD8 + T cells are negatively regulated by LAG-3 and PD-1 in human ovarian cancer. *Proc Natl Acad Sci USA*. 2010; 107(17):7875–7880. [PubMed: 20385810]
43. Li Q, Withoff S, Verma IM. Inflammation-associated cancer: NF-kappaB is the lynchpin. *Trends Immunol*. 2005; 26(6):318–325. [PubMed: 15922948]
44. Lee CH, Jeon YT, Kim SH, Song YS. NF-kappaB as a potential molecular target for cancer therapy. *Biofactors*. 2007; 29(1):19–35. [PubMed: 17611291]
45. Ahn KS, Sethi G, Aggarwal BB. Nuclear factor-kappa B: from clone to clinic. *Curr Mol Med*. 2007; 7(7):619–637. [PubMed: 18045141]
46. Liou HC, Hsia CY. Distinctions between c-Rel and other NF-kappaB proteins in immunity and disease. *Bioessays*. 2003; 25(8):767–780. [PubMed: 12879447]
47. Bonizzi G, Karin M. The two NF-kappaB activation pathways and their role in innate and adaptive immunity. *Trends Immunol*. 2004; 25(6):280–288. [PubMed: 15145317]
48. Hopkins PA, Sriskandan S. Mammalian Toll-like receptors: to immunity and beyond. *Clin Exp Immunol*. 2005; 140(3):395–407. [PubMed: 15932500]
49. Senftleben U, Cao Y, Xiao G, Greten FR, Krahn G, Bonizzi G, et al. Activation by IKKalpha of a second, evolutionary conserved, NF-kappa B signaling pathway. *Science*. 2001; 293(5534):1495–1499. [PubMed: 11520989]
50. Inoue J, Gohda J, Akiyama T, Semba K. NF-kappaB activation in development and progression of cancer. *Cancer Sci*. 2007; 98(3):268–274. [PubMed: 17270016]
51. Weih DS, Yilmaz ZB, Weih F. Essential role of RelB in germinal center and marginal zone formation and proper expression of homing chemokines. *J Immunol*. 2001; 167(4):1909–1919. [PubMed: 11489970]
52. Piver MS, Mettlin CJ, Tsukada Y, Nasca P, Greenwald P, McPhee ME. Familial ovarian cancer registry. *Obstet Gynecol*. 1984; 64(2):195–199. [PubMed: 6738953]
53. Tung KH, Goodman MT, Wu AH, McDuffie K, Wilkens LR, Nomura AM, et al. Aggregation of ovarian cancer with breast, ovarian, colorectal, and prostate cancer in first-degree relatives. *Am J Epidemiol*. 2004; 159(8):750–758. [PubMed: 15051584]
54. Antoniou AC, Easton DF. Risk prediction models for familial breast cancer. *Future Oncol*. 2006; 2(2):257–274. [PubMed: 16563094]
55. Szabo CI, King MC. Population genetics of BRCA1 and BRCA2. *Am J Hum Genet*. 1997; 60(5):1013–1020. [PubMed: 9150148]
56. Lindor NM, McMaster ML, Lindor CJ, Greene MH. Concise handbook of familial cancer susceptibility syndromes—second edition. *J Natl Cancer Inst Monogr*. 2008; (38):1–93. [PubMed: 18559331]
57. Fasching PA, Gayther S, Pearce L, Schildkraut JM, Goode E, Thiel F. Role of genetic polymorphisms and ovarian cancer susceptibility. *Molecular Oncology*. 2009; 3(2):171–181. [PubMed: 19383379]
58. Risch N, Merikangas K. The future of genetic studies of complex human diseases. *Science*. 1996; 273(5281):1516–1517. [PubMed: 8801636]
59. Pharoah PD, Antoniou A, Bobrow M, Zimmern RL, Easton DF, Ponder BA. Polygenic susceptibility to breast cancer and implications for prevention. *Nat Genet*. 2002; 31(1):33–36. [PubMed: 11984562]
60. Ramus SJ, Vierkant RA, Johnatty SE, Pike MC, Van Den Berg DJ, Wu AH, et al. Consortium analysis of 7 candidate SNPs for ovarian cancer. *Int J Cancer*. 2008; 123(2):380–388.

61. Pearce CL, Near AM, Van Den Berg DJ, Ramus SJ, Gentry-Maharaj A, Menon U, et al. Validating genetic risk associations for ovarian cancer through the international Ovarian Cancer Association Consortium. *Br J Cancer*. 2009; 100(2):412–420. [PubMed: 19127255]
62. Dong LM, Potter JD, White E, Ulrich CM, Cardon LR, Peters U. Genetic susceptibility to cancer: the role of polymorphisms in candidate genes. *Jama*. 2008; 299(20):2423–2436. [PubMed: 18505952]
63. Easton DF, Pooley KA, Dunning AM, Pharoah PD, Thompson D, Ballinger DG, et al. Genome-wide association study identifies novel breast cancer susceptibility loci. *Nature*. 2007; 447(7148):1087–1093. [PubMed: 17529967]
64. Tomlinson I, Webb E, Carvajal-Carmona L, Broderick P, Kemp Z, Spain S, et al. A genome-wide association scan of tag SNPs identifies a susceptibility variant for colorectal cancer at 8q24.21. *Nat Genet*. 2007; 39(8):984–988. [PubMed: 17618284]
65. Song H, Ramus SJ, Tyrer J, Bolton KL, Gentry-Maharaj A, Wozniak E, et al. A genome-wide association study identifies a new ovarian cancer susceptibility locus on 9p22.2. *Nat Genet*. 2009; 41(9):996–1000. [PubMed: 19648919]
66. Goode EL, Chenevix-Trench G, Song H, Ramus SJ, Notaridou M, Lawrenson K, et al. A genome-wide association study identifies susceptibility loci for ovarian cancer at 2q31 and 8q24. *Nat Genet*. 2010; 42(10):874–879. [PubMed: 20852632]
67. Bolton KL, Tyrer J, Song H, Ramus SJ, Notaridou M, Jones C, et al. Common variants at 19p13 are associated with susceptibility to ovarian cancer. *Nat Genet*. 2010; 42(10):880–884. [PubMed: 20852633]
68. Gayther SA, Song H, Ramus SJ, Kjaer SK, Whittemore AS, Quaye L, et al. Tagging single nucleotide polymorphisms in cell cycle control genes and susceptibility to invasive epithelial ovarian cancer. *Cancer Res*. 2007; 67(7):3027–3035. [PubMed: 17409409]
69. Pearce CL, Wu AH, Gayther SA, Bale AE, Beck PA, Beesley J, et al. Progesterone receptor variation and risk of ovarian cancer is limited to the invasive endometrioid subtype: results from the ovarian cancer association consortium pooled analysis. *Br J Cancer*. 2008; 98(2):282–288. [PubMed: 18219286]
70. Schildkraut JM, Goode EL, Clyde MA, Iversen ES, Moorman PG, Berchuck A, et al. Single nucleotide polymorphisms in the TP53 region and susceptibility to invasive epithelial ovarian cancer. *Cancer Res*. 2009; 69(6):2349–2357. [PubMed: 19276375]
71. Kelemen LE, Sellers TA, Schildkraut JM, Cunningham JM, Vierkant RA, Pankratz VS, et al. Genetic variation in the one-carbon transfer pathway and ovarian cancer risk. *Cancer Res*. 2008; 68(7):2498–2506. [PubMed: 18381459]
72. Goode EL, Maurer MJ, Sellers TA, Phelan CM, Kalli KR, Fridley BL, et al. Inherited determinants of ovarian cancer survival. *Clin Cancer Res*. 2010; 16(3):995–1007. [PubMed: 20103664]
73. Spitz MR, Wu X, Mills G. Integrative epidemiology: from risk assessment to outcome prediction. *J Clin Oncol*. 2005; 23(2):267–275. [PubMed: 15637390]
74. Quaye L, Gayther SA, Ramus SJ, Di Cioccio RA, McGuire V, Hogdall E, et al. The effects of common genetic variants in oncogenes on ovarian cancer survival. *Clin Cancer Res*. 2008; 14(18):5833–5839. [PubMed: 18794094]
75. Schultheis AM, Lurje G, Rhodes KE, Zhang W, Yang D, Garcia AA, et al. Polymorphisms and clinical outcome in recurrent ovarian cancer treated with cyclophosphamide and bevacizumab. *Clin Cancer Res*. 2008; 14(22):7554–7563. [PubMed: 19010874]
76. Bhatti P, Church DM, Rutter JL, Struwing JP, Sigurdson AJ. Candidate single nucleotide polymorphism selection using publicly available tools: a guide for epidemiologists. *Am J Epidemiol*. 2006; 164(8):794–804. [PubMed: 16923772]
77. International HapMap Consortium. A haplotype map of the human genome. *Nature*. 2005; 437(7063):1299–1320. [PubMed: 16255080]
78. The 1000 Genomes Project Consortium. A map of human genome variation from population-scale sequencing. *Nature*. 2010; 467(7319):1061–1073. [PubMed: 20981092]
79. Gabriel SB, Schaffner SF, Nguyen H, Moore JM, Roy J, Blumenstiel B, et al. The structure of haplotype blocks in the human genome. *Science*. 2002; 296(5576):2225–2229. [PubMed: 12029063]

80. Daly MJ, Rioux JD, Schaffner SF, Hudson TJ, Lander ES. High-resolution haplotype structure in the human genome. *Nat Genet.* 2001; 29(2):229–232. [PubMed: 11586305]
81. Helmuth L. Genome research: map of the human genome 3.0. *Science.* 2001; 293(5530):583–585. [PubMed: 11474073]
82. Patil N, Berno AJ, Hinds DA, Barrett WA, Doshi JM, Hacker CR, et al. Blocks of limited haplotype diversity revealed by high-resolution scanning of human chromosome 21. *Science.* 2001; 294(5547):1719–1723. [PubMed: 11721056]
83. Cao Y, Bonizzi G, Seagroves TN, Greten FR, Johnson R, Schmidt EV, et al. IKK α provides an essential link between RANK signaling and cyclin D1 expression during mammary gland development. *Cell.* 2001; 107(6):763–775. [PubMed: 11747812]
84. Bader JS. The relative power of SNPs and haplotype as genetic markers for association tests. *Pharmacogenomics.* 2001; 2(1):11–24. [PubMed: 11258193]
85. Morris RW, Kaplan NL. On the advantage of haplotype analysis in the presence of multiple disease susceptibility alleles. *Genet Epidemiol.* 2002; 23(3):221–233. [PubMed: 12384975]
86. Johnson GC, Esposito L, Barratt BJ, Smith AN, Heward J, Di Genova G, et al. Haplotype tagging for the identification of common disease genes. *Nat Genet.* 2001; 29(2):233–237. [PubMed: 11586306]
87. Carlson CS, Eberle MA, Rieder MJ, Yi Q, Kruglyak L, Nickerson DA. Selecting a maximally informative set of single-nucleotide polymorphisms for association analyses using linkage disequilibrium. *Am J Hum Genet.* 2004; 74(1):106–120. [PubMed: 14681826]
88. Verstrepen L, Bekaert T, Chau TL, Tavernier J, Chariot A, Beyaert R. TLR-4, IL-1R and TNF-R signaling to NF κ B: variations on a common theme. *Cell Mol Life Sci.* 2008; 65(19):2964–2978. [PubMed: 18535784]
89. Li H, Lin X. Positive and negative signaling components involved in TNF α -induced NF- κ B activation. *Cytokine.* 2008; 41(1):1–8. [PubMed: 18068998]
90. Hauer J, Puschner S, Ramakrishnan P, Simon U, Bongers M, Federle C, et al. TNF receptor (TNFR)-associated factor (TRAF) 3 serves as an inhibitor of TRAF2/5-mediated activation of the noncanonical NF- κ B pathway by TRAF-binding TNFRs. *Proc Natl Acad Sci USA.* 2005; 102(8):2874–2879. [PubMed: 15708970]
91. DeJardin E, Droin NM, Delhase M, Haas E, Cao Y, Makris C, et al. The lymphotoxin-beta receptor induces different patterns of gene expression via two NF- κ B pathways. *Immunity.* 2002; 17(4):525–535. [PubMed: 12387745]
92. Bharti AC, Takada Y, Shishodia S, Aggarwal BB. Evidence that receptor activator of nuclear factor (NF)- κ B ligand can suppress cell proliferation and induce apoptosis through activation of a NF- κ B-independent and TRAF6-dependent mechanism. *J Biol Chem.* 2004; 279(7):6065–6076. [PubMed: 14645259]
93. Cooper, GM.; Hausman, RE. *The cell: a molecular approach.* 4th edn.. Sunderland: Sinauer Associates, Inc.; 2007.
94. Xiao G, Rabson AB, Young W, Qing G, Qu Z. Alternative pathways of NF- κ B activation: a double-edged sword in health and disease. *Cytokine Growth Factor Rev.* 2006; 17(4):281–293. [PubMed: 16793322]
95. Li ZW, Chu W, Hu Y, Delhase M, Deerinc T, Ellisman M, et al. The IKK β subunit of IkappaB kinase (IKK) is essential for nuclear factor kappaB activation and prevention of apoptosis. *J Exp Med.* 1999; 189(11):1839–1845. [PubMed: 10359587]
96. Ea CK, Deng L, Xia ZP, Pineda G, Chen ZJ. Activation of IKK by TNF α requires site-specific ubiquitination of RIP1 and polyubiquitin binding by NEMO. *Mol Cell.* 2006; 22(2):245–257. [PubMed: 16603398]
97. Morlon A, Munnich A, Smahi A. TAB 2, TRAF6 and TAK1 are involved in NF- κ B activation induced by the TNF-receptor, Edar and its adaptator Edaradd. *Hum Mol Genet.* 2005; 14(23):3751–3757. [PubMed: 16251197]
98. Yamaguchi K, Nagai S, Ninomiya-Tsuji J, Nishita M, Tamai K, Irie K, et al. XIAP, a cellular member of the inhibitor of apoptosis protein family, links the receptors to TAB 1-TAK1 in the BMP signaling pathway. *EMBO J.* 1999; 18(1):179–187. [PubMed: 9878061]

99. Hsu H, Shu HB, Pan MG, Goeddel DV. TRADD-TRAF2 and TRADD-FADD interactions define two distinct TNF receptor 1 signal transduction pathways. *Cell*. 1996; 84(2):299–308. [PubMed: 8565075]
100. Beere HM. Death versus survival: functional interaction between the apoptotic and stress-inducible heat shock protein pathways. *J Clin Invest*. 2005; 115(10):2633–2639. [PubMed: 16200196]
101. Tada K, Okazaki T, Sakon S, Koburai T, Kurosawa K, Yamaoka S, et al. Critical roles of TRAF2 and TRAF5 in tumor necrosis factor-induced NF-kappa B activation and protection from cell death. *J Biol Chem*. 2001; 276(39):36530–36534. [PubMed: 11479302]
102. Yang J, Lin Y, Guo Z, Cheng J, Huang J, Deng L, et al. The essential role of MEKK3 in TNF-induced NF-kappaB activation. *Nat Immunol*. 2001; 2(7):620–624. [PubMed: 11429546]
103. Blonska M, You Y, Geleziunas R, Lin X. Restoration of NF-kappaB activation by tumor necrosis factor alpha receptor complex-targeted MEKK3 in receptor-interacting protein-deficient cells. *Mol Cell Biol*. 2004; 24(24):10757–10765. [PubMed: 15572679]
104. Ouellet M, Roy J, Barbeau B, Geleziunas R, Tremblay MJ. NF-kappaB induction by bisperoxovanadium compounds requires CD45, p36(LAT), PKC, and IKK activity and exhibits kinetics of activation comparable to those of TCR/CD28 coengagement. *Biochemistry*. 2003; 42(27):8260–8271. [PubMed: 12846575]
105. Livolsi A, Busuttill V, Imbert V, Abraham RT, Peyron JF (2001) Tyrosine phosphorylation-dependent activation of NF-kappa B. Requirement for p56 LCK and ZAP-70 protein tyrosine kinases. *Eur J Biochem*. 2001; 268(5):1508–1515. [PubMed: 11231305]
106. Dienz O, Moller A, Strecker A, Stephan N, Krammer PH, Droge W, et al. Src homology 2 domain-containing leukocyte phosphoprotein of 76 kDa and phospholipase C gamma 1 are required for NF-kappa B activation and lipid raft recruitment of protein kinase C theta induced by T cell costimulation. *J Immunol*. 2003; 170(1):365–372. [PubMed: 12496421]
107. Wang D, Matsumoto R, You Y, Che T, Lin XY, Gaffen SL, et al. CD3/CD28 costimulation-induced NF-kappaB activation is mediated by recruitment of protein kinase C-theta, Bcl10, and IkappaB kinase beta to the immunological synapse through CARMA1. *Mol Cell Biol*. 2004; 24(1):164–171. [PubMed: 14673152]
108. Bertin J, Wang L, Guo Y, Jacobson MD, Poyet JL, Srinivasula SM, et al. CARD11 and CARD14 are novel caspase recruitment domain (CARD)/membrane-associated guanylate kinase (MAGUK) family members that interact with BCL10 and activate NF-kappa B. *J Biol Chem*. 2001; 276(15):11877–11882. [PubMed: 11278692]
109. Wang L, Guo Y, Huang WJ, Ke X, Poyet JL, Manji GA, et al. Card10 is a novel caspase recruitment domain/membrane-associated guanylate kinase family member that interacts with BCL10 and activates NF-kappa B. *J Biol Chem*. 2001; 276(24):21405–21409. [PubMed: 11259443]
110. Bertin J, Guo Y, Wang L, Srinivasula SM, Jacobson MD, Poyet JL, et al. CARD9 is a novel caspase recruitment domain-containing protein that interacts with BCL10/CLAP and activates NF-kappa B. *J Biol Chem*. 2000; 275(52):41082–41086. [PubMed: 11053425]
111. Hara H, Ishihara C, Takeuchi A, Xue L, Morris SW, Penninger JM, et al. Cell type-specific regulation of ITAM-mediated NF-kappaB activation by the adaptors, CARMA1 and CARD9. *J Immunol*. 2008; 181(2):918–930. [PubMed: 18606643]
112. Baens M, Fevery S, Sagaert X, Noels H, Hagens S, Broeckx V, et al. Selective expansion of marginal zone B cells in Emicro-API2-MALT1 mice is linked to enhanced IkappaB kinase gamma polyubiquitination. *Cancer Res*. 2006; 66(10):5270–5277. [PubMed: 16707452]
113. Che T, You Y, Wang D, Tanner MJ, Dixit VM, Lin X. MALT1/paracaspase is a signaling component downstream of CARMA1 and mediates T cell receptor-induced NF-kappaB activation. *J Biol Chem*. 2004; 279(16):15870–15876. [PubMed: 14754896]
114. Ruland J, Mak TW. Transducing signals from antigen receptors to nuclear factor kappaB. *Immunol Rev*. 2003; 193:93–100. [PubMed: 12752674]
115. Huang J, Gao X, Li S, Cao Z. Recruitment of IRAK to the interleukin 1 receptor complex requires interleukin 1 receptor accessory protein. *Proc Natl Acad Sci USA*. 1997; 94(24):12829–12832. [PubMed: 9371760]

116. Greenfeder SA, Nunes P, Kwee L, Labow M, Chizzonite RA, Ju G. Molecular cloning and characterization of a second subunit of the interleukin 1 receptor complex. *J Biol Chem.* 1995; 270(23):13757–13765. [PubMed: 7775431]
117. Wesche H, Henzel WJ, Shillinglaw W, Li S, Cao Z. MyD88: an adapter that recruits IRAK to the IL-1 receptor complex. *Immunity.* 1997; 7(6):837–847. [PubMed: 9430229]
118. Cao Z, Henzel WJ, Gao X. IRAK: a kinase associated with the interleukin-1 receptor. *Science.* 1996; 271(5252):1128–1131. [PubMed: 8599092]
119. Burns K, Clatworthy J, Martin L, Martinon F, Plumpton C, Maschera B, et al. Tollip, a new component of the IL-1RI pathway, links IRAK to the IL-1 receptor. *Nat Cell Biol.* 2000; 2(6): 346–351. [PubMed: 10854325]
120. Suzuki N, Suzuki S, Yeh WC. IRAK-4 as the central TIR signaling mediator in innate immunity. *Trends Immunol.* 2002; 23(10):503–506. [PubMed: 12297423]
121. Kollewe C, Mackensen AC, Neumann D, Knop J, Cao P, Li S, et al. Sequential autophosphorylation steps in the interleukin-1 receptor-associated kinase-1 regulate its availability as an adapter in interleukin-1 signaling. *J Biol Chem.* 2004; 279(7):5227–5236. [PubMed: 14625308]
122. Jiang Z, Ninomiya-Tsuji J, Qian Y, Matsumoto K, Li X. Interleukin-1 (IL-1) receptor-associated kinase-dependent IL-1-induced signaling complexes phosphorylate TAK1 and TAB 2 at the plasma membrane and activate TAK1 in the cytosol. *Mol Cell Biol.* 2002; 22(20):7158–7167. [PubMed: 12242293]
123. Jiang Z, Johnson HJ, Nie H, Qin J, Bird TA, Li X. Pellino 1 is required for interleukin-1 (IL-1)-mediated signaling through its interaction with the IL-1 receptor-associated kinase 4 (IRAK4)-IRAK-tumor necrosis factor receptor-associated factor 6 (TRAF6) complex. *J Biol Chem.* 2003; 278(13):10952–10956. [PubMed: 12496252]
124. Jensen LE, Whitehead AS. Pellino3, a novel member of the Pellino protein family, promotes activation of c-Jun and Elk-1 and may act as a scaffolding protein. *J Immunol.* 2003; 171(3): 1500–1506. [PubMed: 12874243]
125. Jensen LE, Whitehead AS. Pellino2 activates the mitogen activated protein kinase pathway. *FEBS Lett.* 2003; 545(2–3):199–202. [PubMed: 12804775]
126. Janeway CA Jr, Medzhitov R. Innate immune recognition. *Annu Rev Immunol.* 2002; 20:197–216. [PubMed: 11861602]
127. Oshiumi H, Matsumoto M, Funami K, Akazawa T, Seya T. TICAM-1, an adaptor molecule that participates in Toll-like receptor 3-mediated interferon-beta induction. *Nat Immunol.* 2003; 4(2): 161–167. [PubMed: 12539043]
128. Chen R, Alvero AB, Silasi DA, Steffensen KD, Mor G. Cancers take their Toll—the function and regulation of Toll-like receptors in cancer cells. *Oncogene.* 2008; 27(2):225–233. [PubMed: 18176604]
129. Barton GM, Medzhitov R. Toll-like receptor signaling pathways. *Science.* 2003; 300(5625): 1524–1525. [PubMed: 12791976]
130. Song HY, Regnier CH, Kirschning CJ, Goeddel DV, Rothe M. Tumor necrosis factor (TNF)-mediated kinase cascades: bifurcation of nuclear factor-kappaB and c-jun N-terminal kinase (JNK/SAPK) pathways at TNF receptor-associated factor 2. *Proc Natl Acad Sci USA.* 1997; 94(18):9792–9796. [PubMed: 9275204]
131. Bradley JR, Pober JS. Tumor necrosis factor receptor-associated factors (TRAFs). *Oncogene.* 2001; 20(44):6482–6491. [PubMed: 11607847]
132. Tergaonkar V, Correa RG, Ikawa M, Verma IM. Distinct roles of IkappaB proteins in regulating constitutive NF-kappaB activity. *Nat Cell Biol.* 2005; 7(9):921–923. [PubMed: 16136188]
133. Phelps CB, Sengchanthalangsy LL, Huxford T, Ghosh G. Mechanism of I kappa B alpha binding to NF-kappa B dimers. *J Biol Chem.* 2000; 275(38):29840–29846. [PubMed: 10882738]
134. Tam WF, Lee LH, Davis L, Sen R. Cytoplasmic sequestration of rel proteins by IkappaBalpha requires CRM1-dependent nuclear export. *Mol Cell Biol.* 2000; 20(6):2269–2284. [PubMed: 10688673]
135. Tam WF, Sen R. IkappaB family members function by different mechanisms. *J Biol Chem.* 2001; 276(11):7701–7704. [PubMed: 11152669]

136. Bouwmeester T, Bauch A, Ruffner H, Angrand PO, Bergamini G, Coughton K, et al. A physical and functional map of the human TNF-alpha/NF-kappa B signal transduction pathway. *Nat Cell Biol.* 2004; 6(2):97–105. [PubMed: 14743216]
137. Thoms HC, Dunlop MG, Stark LA. CDK4 inhibitors and apoptosis: a novel mechanism requiring nucleolar targeting of RelA. *Cell Cycle.* 2007; 6(11):1293–1297. [PubMed: 17525529]
138. Fan S, Meng Q, Laterra JJ, Rosen EM. Ras effector pathways modulate scatter factor-stimulated NF-kappaB signaling and protection against DNA damage. *Oncogene.* 2007; 26(33):4774–4796. [PubMed: 17297451]
139. Ben-Neriah Y. Regulatory functions of ubiquitination in the immune system. *Nat Immunol.* 2002; 3(1):20–26. [PubMed: 11753406]
140. Karin M, Ben-Neriah Y. Phosphorylation meets ubiquitination: the control of NF-[kappa]B activity. *Annu Rev Immunol.* 2000; 18:621–663. [PubMed: 10837071]
141. Spencer E, Jiang J, Chen ZJ. Signal-induced ubiquitination of IkappaBalpha by the F-box protein Slimb/beta-TrCP. *Genes Dev.* 1999; 13(3):284–294. [PubMed: 9990853]
142. Winston JT, Strack P, Beer-Romero P, Chu CY, Elledge SJ, Harper JW. The SCFbeta-TRCP-ubiquitin ligase complex associates specifically with phosphorylated destruction motifs in IkappaBalpha and beta-catenin and stimulates IkappaBalpha ubiquitination in vitro. *Genes Dev.* 1999; 13(3):270–283. [PubMed: 9990852]
143. Hoppe T, Matuschewski K, Rape M, Schlenker S, Ulrich HD, Jentsch S. Activation of a membrane-bound transcription factor by regulated ubiquitin/proteasome-dependent processing. *Cell.* 2000; 102(5):577–586. [PubMed: 11007476]
144. Buratowski S, Hahn S, Guarente L, Sharp PA. Five intermediate complexes in transcription initiation by RNA polymerase II. *Cell.* 1989; 56(4):549–561. [PubMed: 2917366]
145. Leurent C, Sanders SL, Demeny MA, Garbett KA, Ruhlmann C, Weil PA, et al. Mapping key functional sites within yeast TFIID. *EMBO J.* 2004; 23(4):719–727. [PubMed: 14765106]
146. Vermeulen L, De Wilde G, Van Damme P, Vanden Berghe W, Haegeman G. Transcriptional activation of the NF-kappaB p65 subunit by mitogen- and stress-activated protein kinase-1 (MSK1). *EMBO J.* 2003; 22(6):1313–1324. [PubMed: 12628924]
147. Viatour P, Merville MP, Bours V, Chariot A. Phosphorylation of NF-kappaB and IkappaB proteins: implications in cancer and inflammation. *Trends Biochem Sci.* 2005; 30(1):43–52. [PubMed: 15653325]
148. Uranishi H, Tetsuka T, Yamashita M, Asamitsu K, Shimizu M, Itoh M, et al. Involvement of the pro-oncoprotein TLS (translocated in liposarcoma) in nuclear factor-kappa B p65-mediated transcription as a coactivator. *J Biol Chem.* 2001; 276(16):13395–13401. [PubMed: 11278855]
149. Reikvam H, Olsnes AM, Gjertsen BT, Ersvar E, Bruserud O. Nuclear factor-kappaB signaling: a contributor in leukemogenesis and a target for pharmacological intervention in human acute myelogenous leukemia. *Crit Rev Oncog.* 2009; 15(1):1–36.
150. Lauscher JC, Gröne J, Dullat S, Hotz B, Ritz JP, Steinhoff U, et al. Association between activation of atypical NF-κB1 p105 signaling pathway and nuclear β-catenin accumulation in colorectal carcinoma. *Mol Carcinog.* 2010; 49(2):121–129. [PubMed: 20027638]
151. Zoubeydi A, Ettinger S, Beraldi E, Hadaschik B, Zardan A, Klomp LWJ, et al. Clusterin Facilitates COMMD1 and I-Î°B Degradation to Enhance NF-Î°B Activity in Prostate Cancer Cells. *Molecular Cancer Research.* 2010; 8(1):119–130. [PubMed: 20068069]
152. Nehra R, Riggins RB, Shajahan AN, Zwart A, Crawford AC, Clarke R. BCL2 and CASP8 regulation by NF-{kappa}B differentially affect mitochondrial function and cell fate in antiestrogen-sensitive and -resistant breast cancer cells. *Faseb J.* 2010; 24(6):2040–2055. [PubMed: 20154269]
153. Nair RP, Duffin KC, Helms C, Ding J, Stuart PE, Goldgar D, et al. Genome-wide scan reveals association of psoriasis with IL-23 and NF-[kappa]B pathways. *Nat Genet.* 2009; 41(2):199–204. [PubMed: 19169254]
154. Escudero C, Sobrevia L. A hypothesis for preeclampsia: adenosine and inducible nitric oxide synthase in human placental microvascular endothelium. *Placenta.* 2008; 29(6):469–483. [PubMed: 18359514]

155. Bakkar N, Guttridge DC. NF-kappaB signaling: a tale of two pathways in skeletal myogenesis. *Physiol Rev.* 2010; 90(2):495–511. [PubMed: 20393192]




Fig. 1.

Key components of NF-κB signaling pathways. Color-coding of molecules: activators = yellow, NF-κB subunits = blue, inhibitors = red, degradators = green, nuclear function = cream. Canonical pathway: Binding of extracellular signaling molecules to receptors leads to recruitment of activation proteins, phosphorylation of IKKβ, activation of the IKK complex (IKKα, IKKβ, and NEMO), and processing of p105 and p100 into functional p50 and p52 by the ubiquitin ligase SCF^{β-TrCP} complex. Activated IKK complex phosphorylates IκB, marking it for ubiquitination by the SCF^{β-TrCP} complex and degradation. The main canonical NF-κB dimer, p65/p50, is activated and translocates to the nucleus to initiate gene transcription with the RNA polymerase II preinitiation complex (PIC). Non-canonical pathway: Ligase and receptor coupling leads to NIK and IKKα activation, which initiates processing of p100 into functional p52 by the SCF^{β-TrCP} complex. The p52/Rel-B dimer translocates to the nucleus to initiate gene expression through the RNA polymerase II PIC





Fig. 2.

TNF α and TCR activation pathways. **A** TNF α binding to its receptors TNF-R1 and TNF-R2 triggers a coupling between TNF-R1/2 and TRADD via their death domains (DD). TRADD recruits TRAF2, RIPK1, RIPK2, and RIPK3, and TRAF5. Following TNF-R1 endocytosis, TRADD dissociates from the receptor complex and associates with FADD and caspase-8, initiating pro-apoptotic signaling pathways. After TRAF2 auto-poly-ubiquitination, RIPK1 is poly-ubiquitinated, which recruits TAB 1, TAB 2, TAB 3, TAK1, and TRAF6 to the complex. TAB 2 forms a bridge between TRAF6 and TAK1, allowing TAK1 to activate the IKK complex by phosphorylating IKK β . **B** Activation of the TCR-CD3 complex induces phosphorylation of ITAMs on TCR ζ and CD3. ZAP70 and LCK are recruited, LCK phosphorylates ZAP70, PLC γ 1, and the scaffold protein SLP-76, and recruits PKC θ to the complex. PKC θ recruits CARMA1, CARD9, CARD10, CARD14, BCL-10, and MALT1, leading to activation of the IKK complex

**Fig. 3.**

IL-1 and TLR activation pathways. **A** Ligand binding to IL-1R1 leads to coupling of IL-1RAPcP1, IL-1RAPcP2, and MYD88 with receptors. MYD88 recruits IRAK-1 and the adaptor proteins Tollip, TRAF6, and IRAK-4 to the receptor complex. After IRAK-1 phosphorylation by IRAK-4, IRAK-1 and TRAF6 dissociate from the receptor complex and associate with TAB 1, TAB 2, TAB 3, and TAK1. Pellino proteins phosphorylate IRAK-1, initiating IRAK-1 and Pellino poly-ubiquitination and degradation, freeing the TRAF6/TAB 1/TAB 2/TAB 3/TAK1 complex. TAB 2 phosphorylates TAK1, which phosphorylates and activates IKK β . **B** Toll-like receptors (TLRs) are critical components of the innate immune response as they are conserved pattern recognition receptors (PRRs) that detect conserved pathogen-associated microbial patterns (PAMPs) present on many antigens. There are over ten TLRs that have been identified, all of which recognize a variety of different PAMPs and lead to NF- κ B activation. TLRs and IL-1R1 have a shared Toll-IL-1R (TIR) domain that mediates interactions with downstream molecules. Due to this shared TIR domain, TLRs initiate a similar signaling cascade as IL-1 involving MYD88, IRAK-1, IRAK-4, TRAF6, TAB 2, and TAK1, but also involve other cascades that include TIRAP, TRIF, TANK, and TBK1




Fig. 4.

Non-canonical activation pathways. Receptor binding of LT β , BAFF, CD40, and RANK initiate the non-canonical signaling pathway. TRAF proteins and other intracellular target molecules are recruited, eventually leading to the activation of NIK through an unknown mechanism. NIK contains binding domains that interact with TRAF2, TRAF3, TRAF5, and TRAF6, but the specific roles of TRAF and other proteins in the non-canonical pathway is still controversial. After NIK phosphorylation, IKK α phosphorylates p100, marking it for poly-ubiquitination and partial degradation by the SCF $^{\beta}$ -TrCP complex and 26S proteasome. Partial degradation of p100 yields functional p52, which most commonly forms heterodimers with Rel-B and translocates to the nucleus to initiate gene transcription






Fig. 5.

Ubiquitination and proteasomal degradation of IκBs. The E3 protein-ubiquitin ligase SCF^{β-TrCP} complex consists of β-TrCP, an F-box protein with a WD repeat region that serves as the docking subunit for the SCF^{β-TrCP} complex, the adaptor proteins Skp1 and Cull1, and the RING protein Roc1 that recruits the E2 ubiquitin-conjugating enzyme UbcH5. The precursor proteins p100 and p105 are partially degraded to yield functional p52 and p50, respectively. p105 contains a glucine-rich region (*GRR*), which separates the functional from inhibitory portions of the protein and is essential for its processing. After being marked for ubiquitination by the SCF^{β-TrCP} complex, the proteasome degrades p105 and p100 until it hits the *GRR* region signaling the completion of processing and the functional subunit is released, or the proteasome is unable to properly unfold the *GRR* region, forcing it to complete processing prematurely and release p50/p65. Additionally, the SCF^{β-TrCP} complex processes IκB after IKKβ phosphorylates IκBα, IκBβ, and IκBε. β-TrCP recognizes the degradation signal and UbcH5 and the SCF^{β-TrCP} complex polyubiquitinate lysine residues on IκBα, which marks it for degradation by the 26S proteasome

**Fig. 6.**

RNA polymerase II transcription complex. Formation of the transcription pre-initiation complex (*PIC*) begins with the general transcription factor TFII-D, which is composed of the TATA-binding protein TBP and 14 TBP-associated factors (*TAFs*). The transcription factor, TBP binds to the TATA box in the promoter region and is necessary for completion of the *PIC* complex. TAF1 and TAF5 appear to be the “organizers” of the complex with TAF1 serving as the scaffold for TAF2, TAF4, TAF5, TAF6, TAF9, TAF11, and TAF12 and branching across two of three lobes in TFII-D. TAF5 binds to TAF1, TAF7, TAF11, TAF12, TAF13, and TBP and contains connections to all three of the TFII-D lobes. Creation of the TFII-D complex recruits TFII-A to bind to both the DNA and TFII-D, stabilizing their interaction. TFII-B binds to TBP and the DNA, serving as a scaffold for RNA polymerase II. TFII-F enters the complex with RNA polymerase II, both of which bind to the TFII-B/TBP complex. After RNA polymerase II is bound, TFII-E and TFII-H enter, completing the *PIC* and initiating transcription

**Fig. 7.**

Coverage of inherited variation in 319 NF-κB signaling genes. Coverage calculations are based on tagSNPs in NF-κB genes ($r^2 \geq 0.9$ and $MAF \geq 0.05$) using the most informative source available, including 1000 Genomes Project Low-coverage Pilot data for 92.8% of genes (see “Methods”). LD Bins reflects the proportion of LD bins with at least one tagSNP on the Infinium HD Human610-Quad BeadChip genome-wide SNP panel (Illumina 610-Quad) or imputable based on HapMap Phase II data. SNPs reflect the proportion of SNPs interrogated, either directly or indirectly, via genotyping the included tagSNPs on the Illumina 610-Quad or imputing based on HapMap Phase II data (representing SNPs directly assessed or in the covered LD bins). Results for the Illumina 610-Quad and HapMap Phase II are presented due to use in current ovarian cancer genome-wide association studies (e.g., Song et al. [65])

Table 1

NF-κB genes classified by function and subpathway

Function	Subpathway			
	Canonical	Non-Canonical	Both	Unknown
Subunits	<i>NFKB1, REL, RELA</i>	<i>NFKB2, RELB</i>	–	–
Activation	<i>AHR, AKT1, AKT2, ATM, AZI2, BCL10, BIRC3, BRCA1, CARD10, CARD11, CARD14, CARD4, CARD9, CASP1, CASP2, CASP8, CC2D1A, CD247, CD28, CD3D, CD3E, CD3G, CDC37, CDK4, CFLAR, CRADD, CRP, CSNK2A2, CSNK2B, ECSIT, EGF, EGFR, EIF2AK2, ERBB2, FADD, FRAP1, HRAS, HSP90AA1, HSP90AA2, HSP90AB1, IKBKB, IKBKE, IKBKG, IL12A, IL15, IL17A, IL18, IL1A, IL1B, IL1R1, IL1R2, IL2, IL6, IL8, IL1RAPL1, IL1RAPL2, ILK, IRAK1, IRAK2, IRAK4, KRAS, LAT, LCK, LCP2, LBP, MALT1, MAP2K1, MAP2K2, MAP2K3, MAP2K4, MAP2K5, MAP2K6, MAP2K7, MAP3K1, MAP3K11, MAP3K2, MAP3K3, MAP3K7, MAP3K7IP1, MAP3K7IP2, MAP3K7IP3, MAP3K8, MAP4K1, MAPK14, MAPK8, MTDH, MYD88, NCAM1, NOD2, NRK, PDPK1, PELI1, PELI2, PELI3, PIK3CA, PIK3CB, PIK3CG, PIK3R1, PLCG1, PRKACA, PRKACB, PRKCA, PRKCQ, PRKDC, PTPRC, RAF1, RHOA, RIPK1, RIPK2, RIPK3, SPAG9, STAT1, SYK, TBK1, TBKBP1, TICAM1, TICAM2, TIFA, TIRAP, TNF, TNFRSF10A, TNFRSF10B, TNFRSF10D, TNFRSF1A, TNFSF10, TOLLIP, TP53, TRA@, TRADD, TRAF2, TRAF7, TRB@, TRD@, TRG@, TRIP6, VISA, ZAP70</i>	<i>CD40, CD40LG, DYNLL1, LTA, LTβ, LTBR, MAP3K14, TNFRSF11A, TNFRSF13B, TNFRSF13C, TNFSF11</i>	<i>CHUK, FKBP5, IKBKAP, TANK, TLR1, TLR10, TLR2, TLR3, TLR4, TLR5, TLR6, TLR7, TLR8, TLR9, TNFRSF1B, TRAF5, TRAF6, TRAPPC9, XIAP</i>	–
Inhibition	<i>ARRB1, ARRB2, ATR, BAG4, BIRC2, CARD6, CARD8, CDKN2A, CHEK1, CLU, CYLD, EGR1, ESR1, ESR2, G3BP2, HEXIM1, HMOX1, IL10, IL1RN, IL3, IRAK3, ITGA3, MCC, MDM2, NALP12, NFKBIA, NFKBIB, NFKBIE, NFKBIZ, NKIRAS1, NKIRAS2, NR1D1, OTUD7B, PGR, PKN1, PPARA, PPARG, PPP1R13L, PPP6C, PTPN13, PYCARD, RNF216, SOD1, TAL1, TNFAIP3, TNIP1, TNIP2, TNIP3, TRAF1, TRIB3</i>	<i>IFNG, TRAF3, TRAF3IP2</i>	<i>TTRAP</i>	<i>NKRF</i>
Degradation	<i>CSNK2A1, GSK3A, GSK3B, HNRNPAB, RPS6KA1, VCP</i>	–	<i>BTRC, CAPN1, CUL1, FBXW11, FBXW7, RBX1, SKP1, SUMO4, UBE2D1, UBE2D2, UBE2 N, UBE2V1, USP11</i>	–
Nuclear function	<i>BCL3, DHX9, F2R, KPNA3, NCOA3, NKAP, PRKCZ, RNF25, RPS6KA4, RPS6KA5, STAT3</i>	<i>KPNA2, KPNA6</i>	<i>ATF1, CREB1, CREBBP, EP300, FUS, GTF2A1, GTF2A2, GTF2B, GTF2E1, GTF2E2, GTF2F1, GTF2F2, GTF2H1, GTF2H2, GTF2H3, GTF2H4, GTF2H5, HDAC1, HDAC3, HDAC6, JUN, KAT2B, MAPK1, MAPK3, MCM5, MCM7, POLR2A, POLR2B, POLR2C, POLR2D, POLR2E, POLR2F, POLR2G, POLR2H, POLR2I, POLR2 J, POLR2 K, POLR2L, RPS6KA3, SP1,</i>	–

Function	Subpathway			
	Canonical	Non-Canonical	Both	Unknown
			TAF1, TAF10, TAF11, TAF12, TAF13, TAF15, TAF2, TAF3, TAF4, TAF4B, TAF5, TAF6, TAF7, TAF8, TAF9, TBP	
Unknown	–	–	–	NFKBIL1, NFKBIL2, NFRKB, TCEAL7

Columns represent a NF-κB subpathway categorization (canonical, non-canonical, both, or unknown); rows represent the biological function of the gene (subunit, activation, inhibition, degradation, nuclear function, or unknown)

Table 2

Characteristics of linkage disequilibrium data sources

Data source	Samples	<i>n</i> Total SNPs/genes	N NF-κB genes	Ref.
Genome-wide sequencing				
1000 Genomes Project Low-coverage Pilot	60 unrelated CEU	7.7 million SNPs	302	1
Genome-wide genotyping				
International HapMap Project Phase II	60 unrelated CEU	4.0 million SNPs	318	2
Perlegen Sciences	24 European-American	1.6 million SNPs	297	3
Gene-specific sequencing				
SeattleSNPs PGA: Panel 1	23 unrelated CEPH	203 genes	39	4
SeattleSNPs PGA: Panel 2	23 unrelated CEU	105 genes	3	4
Innate Immunity SNPs PGA	23 unrelated European	83 genes	21	5
NIEHS SNPs Panel 1	22 unrelated European	201 genes	9	6

N genes/SNPs based on human coding genes (release 29 in genome build 36.3); N NF-κB from among $n = 319$ listed in Supplemental Table 1; CEU, samples collected in 1980 from U.S. residents with northern and western European ancestry by the Centre d'Etude du Polymorphisme Humain (CEPH); References: 1. <http://www.1000genomes.org/>; accessed April 2010; 2. <http://www.hapmap.org/>; accessed November 2008; 3. <http://genome.perlegen.com/>; accessed November 2008; 4. <http://pga.mbt.washington.edu/>; accessed November 2008; 5. <http://www.innateimmunity.net/>; accessed November 2008; 6. <http://egp.gs.washington.edu/>; accessed November 2008