

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/262108602>

Transient dynamics of A β contribute to toxicity in Alzheimer's disease

ARTICLE in CELLULAR AND MOLECULAR LIFE SCIENCES CMLS · MAY 2014

Impact Factor: 5.81 · DOI: 10.1007/s00018-014-1634-z · Source: PubMed

CITATIONS

14

READS

67

4 AUTHORS:



[Ellen Hubin](#)

Vrije Universiteit Brussel

8 PUBLICATIONS 24 CITATIONS

[SEE PROFILE](#)



[Nico A J Van Nuland](#)

Vrije Universiteit Brussel

97 PUBLICATIONS 2,575 CITATIONS

[SEE PROFILE](#)



[Kerensa Broersen](#)

University of Twente

35 PUBLICATIONS 844 CITATIONS

[SEE PROFILE](#)



[Kris Pauwels](#)

Vrije Universiteit Brussel

24 PUBLICATIONS 250 CITATIONS

[SEE PROFILE](#)

Transient dynamics of A β contribute to toxicity in Alzheimer's disease

E. Hubin · N. A. J. van Nuland · K. Broersen ·
K. Pauwels

Received: 18 March 2014 / Revised: 15 April 2014 / Accepted: 22 April 2014
© The Author(s) 2014. This article is published with open access at Springerlink.com

Abstract The aggregation and deposition of the amyloid- β peptide (A β) in the brain has been linked with neuronal death, which progresses in the diagnostic and pathological signs of Alzheimer's disease (AD). The transition of an unstructured monomeric peptide into self-assembled and more structured aggregates is the crucial conversion from what appears to be a harmless polypeptide into a malignant form that causes synaptotoxicity and neuronal cell death. Despite efforts to identify the toxic form of A β , the development of effective treatments for AD is still limited by the highly transient and dynamic nature of interconverting forms of A β . The variability within the in vivo "pool" of different A β peptides is another complicating factor. Here we review the dynamical interplay between various components that influence the heterogeneous A β system, from intramolecular A β flexibility to intermolecular dynamics between various A β alloforms and external factors. The complex dynamics of A β contributes to the causative role of A β in the pathogenesis of AD.

Keywords Alzheimer's disease · Amyloid- β peptide · A β dynamics · Intrinsically disordered peptide · Aggregation

Introduction

Experimental studies and clinical trials are ongoing in the search for an effective prevention or treatment of Alzheimer's disease (AD) [1–3]. These studies and trials often target the amyloid-beta peptide (A β), which plays a major role in AD pathogenesis [4]. Effective drug development has remained without success and this is thought to originate from the fact that A β can appear in many different shapes that can interconvert within a dynamical interplay. This finding triggered a vast exploration of the many conformations the peptide can adopt, as well as the aim to precisely pinpoint which of these conformations can be claimed as "the toxic species", such that specific drug targeting can be employed. To complicate matters even more, a heterogeneous pool of monomeric A β varying in length from 37 to 49 amino acids is produced by proteolytic cleavage from the transmembrane amyloid precursor protein (APP) by β - and γ -secretases [5, 6] (Fig. 1). Most research effort has been focused on the most abundant form A β_{1-40} , which comprises 40 amino acids. The longer and less abundant A β_{1-42} , C-terminally extended by two residues, has been found to be more aggregation-prone [7]. Nonetheless, it has recently been discovered by us [8–10] and other groups [11–14] that the co-occurrence of peptides varying in length can affect the neurotoxic and aggregation potential of the total A β pool. It was also recognized that particularly small aggregated forms of A β are potentially toxic, rather than the mature amyloid fibrils as observed in the brain of AD patients. Therefore, a lot of research has aimed at understanding the A β aggregation mechanism and identifying the intermediate species that occur along the aggregation pathway [15, 16]. The current amyloid cascade hypothesis suggests that AD-related synapto- and neurotoxicity might be mediated by soluble A β oligomers [17, 18],

E. Hubin · K. Broersen
Nanobiophysics Group, MIRA Institute for Biomedical
Technology and Technical Medicine, Faculty of Science
and Technology, University of Twente, 7500 AE Enschede,
The Netherlands

E. Hubin · N. A. J. van Nuland · K. Pauwels
Structural Biology Brussels, Department of Biotechnology
(DBIT), Vrije Universiteit Brussel (VUB), Pleinlaan 2,
1050 Brussels, Belgium

E. Hubin · N. A. J. van Nuland · K. Pauwels (✉)
Structural Biology Research Center, VIB, Pleinlaan 2,
1050 Brussels, Belgium
e-mail: krpauwel@vub.ac.be




Fig. 1 Heterogeneity in the Aβ peptide pool. Sequential proteolytic events by the β- and γ-secretase of the amyloid precursor protein (APP) give rise to the carboxy-terminal fragment (CTF), APP intracellular domain (AICD), and the amyloid-β peptide (Aβ). The heterogeneity in the Aβ pool originates from the proteolysis by the

γ-secretase, but also post-translational modifications contribute to the formation of various Aβ alloforms. Mutations in Aβ and other exogenous factors can influence the dynamics that are observed within the Aβ system

which have proven notoriously difficult to study in detail in vivo with the currently available technology. The dynamics, stability, and transient lifetime of potentially toxic species further hamper the possibility to precisely pinpoint the toxic structural aspects of Aβ aggregates. Moreover, the dynamic behavior of aggregation intermediates may actually provide an important source for toxicity of Aβ as isolated Aβ oligomers are only toxic in the presence of Aβ monomers that provide a source for continued growth of oligomers into fibrillar species [13, 19].

This review discusses how Aβ peptide dynamics can influence and contribute to Aβ-induced toxicity. Aβ dynamics is mainly considered on two levels. First, we define *intramolecular dynamics* of Aβ as the intrinsic disorder or polypeptide backbone flexibility that is present in isolated Aβ monomeric peptides or aggregation states. Second, we define *intermolecular dynamics* as (1) the interplay between different Aβ alloforms present in the in vivo Aβ pool and (2) the dynamical equilibrium that exists between different Aβ species. With the term *alloform*, we refer to a distinct form of the Aβ peptide that is commonly treated as a single kind of peptide species, like Aβ length variants or side chain modifications. Finally, several external factors and interaction partners that can influence Aβ dynamics are addressed. The potential importance of Aβ dynamics in understanding AD pathology is highlighted with the aim of shaping new research orientations for AD treatment.

Intramolecular dynamics

The Aβ monomer has a high tendency to self-assemble into large aggregates and fibrils. It is increasingly recognized that despite the highly packed and ordered state of these higher-order aggregates, they often do contain a significant portion of flexible and intrinsically disordered regions [20]. The intrinsically disordered nature of the Aβ monomer is fairly well documented, but revealing the structural disorder in oligomers and fibrils has proven more challenging due to the difficulties in studying this phenomenon. In this section, we discuss the intrinsic structural disorder that is present in every Aβ aggregation state, and we illustrate how it contributes to Aβ-induced toxicity.

The intrinsically disordered Aβ monomer

Although the pathological hallmark of AD comprises insoluble Aβ deposits in neuritic plaques in the brain of AD patients, monomeric Aβ peptides have also been purified and characterized from brain tissue [21–24]. Size exclusion chromatography (SEC) experiments suggested that the freshly dissolved peptide eluted as a single low molecular weight species, consistent with a monomer or dimer [25–27]. These low molecular weight Aβ species were competent to deposit onto pre-existing amyloid in preparations of AD cortex, with a first-order kinetic dependence




Fig. 2 Various structures of A β that correspond to different experimental conditions and phases in the aggregation landscape. **a** Four representatives of the structural ensemble of monomeric A β_{1-42} under aqueous conditions as derived from a combined molecular dynamics/NMR approach [38]. Extended as well as collapsed coil conformations with secondary structural elements can be observed. **b** A β_{1-40}

in presence of 50 mM NaCl at 15 °C [33] and A β_{1-42} in presence of 30 % hexafluoroisopropanol [32] contain an α -helical segment. **c** Fibril polymorphism illustrated by fibrillar A β_{1-42} [53], D23N A β_{1-40} [74] and **d** the ultrastructure of A β_{1-40} [83], and brain-derived A β_{1-40} [89]

on soluble A β concentration [26]. Translational diffusion measurements by nuclear magnetic resonance (NMR) techniques conclusively demonstrated that the form of the peptide active in plaque deposition is a monomer [26]. Further NMR data revealed that monomeric A β exists in solution as disordered coils that lack regular α -helical or β -stranded structure [28–30]. Despite the challenging task because of its unstructured and amyloidogenic nature, the A β monomer is now well recognized as an intrinsically disordered peptide (IDP). This implies that the monomeric A β peptide does not display a unique fold, as would be the case for a typical well-folded protein, but rather comprises a mixture of rapidly interconverting conformations whereby the polypeptide backbone can sample the conformational space without any stable and well-defined conformational ensemble (Fig. 2). Yet, it is possible to bias the ensemble toward distinct secondary structure elements by changing solution conditions and/or the oxidation state of Met₃₅ [30–33].

Some experimental studies suggested that A β is not entirely a “random coil”. Ion mobility mass spectrometry (MS) combined with theoretical modeling showed that A β_{1-42} in aqueous solution adopts both extended chain as well as collapsed-coil structures [34]. Limited proteolysis successfully identified structured and disordered regions within A β [35]. This approach revealed a proteolytically

resistant decapeptide, Ala₂₁–Ala₃₀, that was found in NMR studies to form a turn-like structure [30]. When the dynamics of monomeric A β_{1-40} in solution was studied using ¹⁵N-relaxation experiments, it revealed structural propensities that correlate well with the secondary structure segments of the peptide that are present in the fibrils, and with the α -helical structure in membrane-mimicking systems [32, 36]. NMR studies further revealed subtle differences between A β_{1-40} and A β_{1-42} monomers whereby a modest increase in C-terminal rigidity has been observed in A β_{1-42} versus A β_{1-40} [37]. Various molecular dynamics simulations also hinted that distinct intramolecular interaction patterns occur in A β_{1-42} [28, 38, 39]. Such subtle differences between A β_{1-40} and A β_{1-42} were confirmed by molecular dynamics simulations [40, 41]. Experimental results in combination with computational simulations have thus proven very powerful to shed light on the conformational landscape of IDPs. The emerging picture of A β comprises an IDP that can adapt a variety of collapsed and extended monomeric conformations and transiently samples long-range intramolecular interactions without exclusively stabilizing a specific globular fold.

Even though the physiological function of A β remains obscure, the intrinsic structural flexibility offers certain advantages: high specificity and low affinity in the case of

binding-induced folding IDPs (mostly exploited in signaling pathways), and high binding promiscuity that is frequently used by hub proteins in large interaction networks [42]. So its IDP nature facilitates the interaction of the peptide with many different binding partners (see “[Other players in the game](#)”), including identical peptides and other A β alloforms. In addition, the high intramolecular flexibility of A β also simplifies post-translational modifications because the involved side chains are readily accessible (see “[The in vivo A \$\beta\$ pool: a cocktail of different interacting species](#)”).

There is a well-established link between intrinsic polypeptide disorder and functional promiscuity. Protein moonlighting, the phenomenon of proteins exhibiting more than one unique biological function, is typically mediated by intrinsically disordered regions in polypeptides [43]. As IDPs can play a role in numerous biological processes, it is not surprising to find some of them involved in human diseases.

Intrinsic fibril flexibility can underlie disease progression and phenotype

A β fibrils contain high order and rigidity compared to A β monomers, but still retain a considerable amount of disorder in the N-terminal segment [44–47] and they are often polymorphous. The inherent disorder of A β fibrils and the associated fibril polymorphism could underlie time-dependent structural changes during aging in AD and differences in disease progression and phenotype.

The molecular dynamic nature of A β fibrils

Even though the amyloid fibril state of A β has traditionally been viewed as a rigid or semi-rigid state with the typical cross- β X-ray fiber diffraction pattern [48, 49], part of the peptide in this conformation is also flexible. This flexibility has been illustrated first by solid-state and solution NMR [50–54], electron paramagnetic resonance (EPR) [44], site-directed mutagenesis [55], limited proteolysis and hydrogen-deuterium exchange (HDX) evaluated by MS [45–47, 56], and even X-ray crystallography [57]. These studies suggested a hairpin-like arrangement of each A β monomer stacked within the fibril, consisting of two semi-rigidly organized β -strands linked by a flexible connecting region (Fig. 2). The hydrophobic C-terminus of A β_{1-42} in the fibril is highly resistant to HDX and forms the fibril core [53, 54]. In contrast, the C-terminus of A β_{1-40} in the fibril contains slightly more disorder [52, 56, 58–61]. The N-terminal segment, which can range from 10 to 19 residues depending on the study, remains intrinsically disordered for both A β_{1-40} and A β_{1-42} fibrils (Table 1). This relatively hydrophilic part of the polypeptide chain is excluded from the H-bonded β -sheet fibril core and remains exposed to the solvent [44–47, 52–54, 58–61]. Recently, differential scanning calorimetry suggested that thermal denaturation of amyloid fibrils can take place and that this process can be considered as a reversible equilibrium under certain experimental conditions, highlighting the dynamic nature of fibrils [62]. These

Table 1 Secondary structure assignments of A β fibrils and structures deposited in the PDB

Peptide	Flexible regions (solvent-exposed)	β -structured regions (non-exposed)	Method	References
A β_{1-40}	N-terminus (Asp ₁ -Phe ₁₉) C-terminus (Met ₃₅ -Val ₄₀)	Phe ₂₀ -Leu ₃₄	HDX-MS coupled with online proteolysis	[191]
A β_{1-40}	N-terminus (Asp ₁ -His ₁₄) C-terminus (Gly ₃₇ -Val ₃₉) Turn? (Ser ₂₆ -Asn ₂₇)	Gln ₁₅ -Asp ₂₃ Lys ₂₈ -Met ₃₅	HDX-solution NMR	[52]
A β_{1-40}	N-terminus (Asp ₁ -His ₁₄) C-terminus (Gly ₃₇ -Val ₄₀) Turns (Glu ₂₂ -Asp ₂₃ , Gly ₂₉ -Ala ₃₀)	Gln ₁₅ -Ala ₂₁ Val ₂₄ -Lys ₂₈ Ile ₃₁ -Val ₃₆	Scanning proline mutagenesis	[58]
A β_{1-40}	N-terminus (Asp ₁ -Tyr ₁₀) Bend (Gly ₂₅ -Gly ₂₉)	Val ₁₂ -Val ₂₄ C-terminus (Ala ₃₀ -Val ₄₀)	Solid-state NMR	[50]
A β_{1-40}	N-terminus (Asp ₁ -Gly ₉) Bend/loop (Asp ₂₃ -Gly ₂₉)	Tyr ₁₀ -Glu ₂₂ C-terminus (Ala ₃₀ -Val ₄₀)	Solid-state NMR	[59]
A β_{1-40} A β_{1-42}	N-terminus (Asp ₁ -Tyr ₁₀) C-terminus (Val ₄₀ -Ala ₄₂ ?) Turn/bend? (Asp ₂₃ -Gly ₂₉)	His ₁₄ -Gly ₃₈	Site-directed spin labeling-EPR	[44]
A β_{1-42}	N-terminus (Asp ₁ -Leu ₁₇) Turn (Asn ₂₇ -Ala ₃₀)	Val ₁₈ -Ser ₂₆ C-terminus (Ile ₃₁ -Ala ₄₂)	HDX-solution NMR	[53]
A β_{1-42}	N-terminus (Asp ₁ -Tyr ₁₀) Bend region? (Ser ₂₆ -Asn ₂₇)	Glu ₁₁ -Gly ₂₅ C-terminus (Lys ₂₈ -Ala ₄₂)	HDX-solution NMR	[54]

Fibril structures deposited in the PDB: synthetic A β_{1-40} (2LMN, 2LMO, 2LMP, 2LMQ), brain-derived A β_{1-40} (2M4J), synthetic D23N A β_{1-40} (2LNQ), recombinant A β_{1-42} (2BEG)

observations illustrate the impact of the various dynamics within the A β system.

The inherent flexibility of A β fibrils also allows the internal fibril structure to evolve in time. Multidimensional infrared spectroscopy revealed that fresh and 4-year-old fibrils were structurally heterogeneous due to trapped water molecules that perturbed the H-bonding pattern in time [63]. Recently, Nilsson and coworkers [64] revealed conformational rearrangements during aging in plaques in the brains of AD mouse models using different luminescent conjugated polythiophenes.

Although ignored for a long time, structural disorder in fibrils seems to occur in various amyloidogenic proteins (e.g. α -synuclein, tau, and multiple prions) (reviewed in [20]). Structural disorder in fibrils has been suggested to stabilize fibril formation by accommodating destabilizing residues and by limiting the unfavorable entropy associated with the formation of the highly ordered cross- β spine.

A β fibrils are polymorphic entities

Overall fibril topology has been studied using cryo-electron microscopy and 3D reconstruction. In general, A β fibrils exhibit multiple distinct morphologies that can differ in fibril symmetry, width, twist period, and curvature [65, 66]. This structural diversity is not limited to A β fibrils, but appears to be a fundamental property of the amyloid state [67–69]. *Inter-sample polymorphism* commonly occurs in vitro in different fibril growth conditions and is subject to pH, temperature, agitation, and salt conditions [70, 71]. A Darwinian-type “survival of the fittest” competition allows the type of fibril that is kinetically the most accessible in a given environment to be the most populated [72]. However, A β_{1-40} can also form at least 12 structurally distinct morphotypes under the same solution conditions (*intra-sample polymorphism*) indicating that this polymorphism arises from an intrinsic variability [73]. Interconversion between fibril polymorphs coexisting in solution can occur, resulting in the thermodynamically more stable polymorph, as was monitored by solid-state NMR over a period of several weeks for A β_{1-40} [74, 75].

Amyloid polymorphism can have several molecular origins that are not mutually exclusive [76–79]. First, mass-per-length values obtained from scanning TEM indicate that fibrils can be composed of one to five protofilaments (the minimal fibrillar entities) [80, 81]. Second, distinct orientations and modes of lateral association of protofilaments by different patterns of inter-residue interactions determine if protofilaments are oriented side-by-side [50, 82], offset from one another [76, 77], or wound around a hollow core [79]. Third, solid-state NMR demonstrated that agitated (striated) and quiescent (twisted) fibrils differ in the residues participating in the β -strands and such variations in

the underlying protofilament substructure can contribute to polymorphism [59, 83]. Surprisingly, the Iowa mutant (D23N A β_{1-40}) was recently found to form metastable fibrils with an antiparallel cross- β spine, indicating that a familial disease-related mutation can have profound effects on fibril structure [74]. Although the cross- β spine of A β fibers is a common feature, fibrils show a great variety of structural complexity that appears inherent to the dynamic nature of the peptide.

Fibril polymorphism could lead to different pathological outcomes

Fibrils can initiate inflammation in brain tissues and cell-cultured microglia and astrocytes. Fibril-induced inflammation then leads to the secretion of pro-inflammatory cytokines and the production of free radicals causing oxidative damage [84, 85]. Substantial evidence provided that different fibril morphologies exert different toxicities in vitro, although toxic activity of oligomeric A β was reported to exceed that of the fibrillar form multiple times [53, 59, 86–88]. For example, oligomeric A β correlated more strongly to cognitive impairment as compared to fibrillar A β of amyloid plaques [86, 87].

Fibril polymorphism could explain the weak correlation between plaque load and cognitive impairment. If plaques are comprised of different fibril polymorphs, different levels of toxicity could be associated to these amyloid deposits. In this case, the structural diversity of fibrils may account for differences in disease progression and phenotype as has been suggested by Tycko and coworkers [89]. They reported that A β fibrils seeded from human brain extracts differed between patients with different clinical history and neuropathology [89]. Moreover, fibril polymorphism has been linked previously to different phenotypes for hereditary transthyretin amyloidosis [90]. In this regard, the different architectures of wild-type A β and Iowa D23N fibrils, comprising respectively parallel and antiparallel β -sheet orientations, could underlie the different pathological outcomes: sporadic AD versus early onset AD associated with cerebral amyloid angiopathy (CAA).

A β oligomers: a mishmash of conformations and sizes

Since the A β plaque load and AD severity could not be correlated [86, 87], growing evidence has revealed that soluble oligomers, either on- or off-pathway to fibrils (see “[The in vivo A \$\beta\$ pool: a cocktail of different interacting species](#)”), play a primary role in AD. Soluble oligomers have commonly been associated with disease severity, the loss of synapses and neuronal damage (reviewed in [18]). The low abundance, heterogeneity, low solubility, and transient nature of A β oligomers have hindered structural studies. It

now becomes clear that A β oligomers exist in a broad range of interconverting assemblies varying in size, conformation, and associated toxicity (reviewed in [91, 92]).

A β oligomers can cause toxicity by a variety of mechanisms (reviewed in [93]). To enable drug design, it is essential to establish the key determinants of oligomer toxicity. Several studies report that neurotoxic activity varies with A β oligomer size with small oligomers ($n < 14$) being most toxic [94, 95]. However, oligomer size is not sufficient to define toxicity as A β oligomers with similar size have been shown to exert different toxicities [96–98]. The underlying peptide conformation also needs to be taken into account as the interplay between A β oligomer size and conformation plays an important role in toxicity (reviewed in [92]). The design of a well-controlled study to investigate size and conformational impact on toxicity is notoriously difficult as different oligomer conformations and sizes are in continuous exchange. However, studies in which different conformations or sizes have been enriched or stabilized by means of crosslinking have been performed and careful conclusions can be drawn from such studies. For example, different A β oligomer conformations have been shown to induce neurotoxicity by distinct mechanisms in human cortical neurons [99]. One possibility to classify oligomers according to their underlying structure is based on recognition by conformation-dependent antibodies [100–103]. Surprisingly, soluble oligomers of a wide variety of amyloidogenic polypeptides (A β , α -synuclein, islet amyloid polypeptide, polyglutamine, lysozyme, human insulin and prion peptide) react with the oligomer-specific A11 antibody developed in the laboratory of Charles Glabe, suggesting that there has to be a common denominator to their toxic origin. Interestingly, pre-incubation of mouse hippocampal neurons with the A11 antibody, before treatment with A β , rescues them from the neurotoxic effects induced by A β [8]. It has been suggested that A11 positive oligomers are composed of antiparallel β -sheets, based on Fourier transform infrared (FTIR) spectroscopy. This antiparallel signature might represent a critical step in perturbation or permeabilization of cell membranes leading to cell toxicity [104]. Later studies using FTIR, EPR, and X-ray crystallography have confirmed that oligomeric species can be characterized by an antiparallel β -sheet orientation, while most fibrils consist of in-register, parallel β -sheets [105–109]. Moreover, antiparallel oligomers displayed a lower content in secondary structure and faster HDX kinetics compared to fibrils, suggesting a higher intrinsic flexibility [104].

Apart from size and peptide conformation, this intrinsic flexibility of the A β oligomer can also be a key determinant of A β -induced toxicity. Several studies have shown that the N-terminus retains a degree of flexibility upon oligomerization and is exposed to the solvent [41, 109–111]. Ahmed and coworkers reported solution NMR measurements of

A β_{1-42} pentamers [111]. The authors found that the loosely packed N-terminal segment of A β was defined by HDX ratios approaching 1 for residues Asp₁-Gly₉, indicating high solvent accessibility and nearly complete exchange within the acquisition time (<1.5 h). In contrast, Val₄₀-Ala₄₂ were less solvent accessible and most likely buried within the center of the oligomer. Similar results were obtained for packing of the A β peptide within A β_{1-42} dodecamers. Site-directed spin labeling of A β_{1-42} combined with EPR spectroscopy showed that the N-terminus was loosely packed within the dodecamer, while residues Ile₃₂-Val₄₀ formed a tight core [109]. Increased structural disorder and solvent exposure of hydrophobic segments of the oligomer have been suggested to be a common feature of highly toxic, soluble aggregates [96–98, 112]. Recent work has shown that the most cytotoxic, oligomeric species of the E22G (arctic) variant of A β_{1-42} interacted more strongly with 1-anilinoaphthalene 8-sulfonate (ANS), a dye sensitive to exposed hydrophobic patches [112]. A higher degree of solvent-exposed, hydrophobic regions was further shown to lead to a disturbed cellular calcium homeostasis, likely due to disruption of the cell membrane [98]. Moreover, oligomers have been shown to bind with higher affinity and cause more disruption of synthetic membranes as compared to the higher-ordered fibrils [113]. These data emphasize the importance of intrinsic disorder and molecular flexibility of A β oligomers for the toxicity mechanism.

In conclusion, a re-evaluation of the oligomer cascade hypothesis is needed (reviewed in [114]). Whereas earlier hypotheses held one single oligomer of a predefined size responsible for toxicity [23, 115], it is obvious that a diverse “A β oligomeric soup” exists, consisting of a large variety of rapidly exchangeable polymorphs that differ in size, conformation, hydrophobicity, solvent exposure, intrinsic disorder (or internal flexibility), and toxicity. The oligomer cascade hypothesis should take into account that it is likely that the entire dynamic A β oligomeric soup contributes to the heterogeneity of AD progression and phenotype, via various toxic mechanisms.

Intermolecular dynamics

As the in vivo A β pool is a mix of species influencing one another, one must also consider the dynamics between different A β species when regarding A β -related toxicity. First, A β peptides of various lengths are produced due to the heterogeneous cleavage pattern of APP by γ -secretase [5, 6]. This gives rise to the production of A β_{1-40} , smaller amounts of A β_{1-42} , and trace amounts of peptides ranging in length from 37 to 49 amino acids [116–118]. Second, a dynamical equilibrium exists between different aggregation states during A β aggregation. Studying the behavior of A β

peptide mixtures and revealing the dynamics of interconversion among different aggregate species will be crucial in understanding the AD-related toxic effects of A β .

The in vivo A β pool: a cocktail of different interacting species

The large majority of biophysical and cell biological studies investigating the role of A β in AD have focused either on pure A β_{1-40} or on pure A β_{1-42} , the two predominant A β alloforms present in the brain [7, 119]. The in vivo A β pool not only contains different A β peptide lengths but also comprises post-translationally modified A β [120] (Fig. 1). A β peptides can undergo racemization [121, 122], isomerization [123], phosphorylation [124, 125], oxidation [126, 127], non-enzymatic glycation [128], and pyroglutamylation [129]. Post-translational oxidation of Met₃₅ affects fibril flexibility within A β plaques [127]. Met₃₅ oxidation also has been shown to impede the rate of A β aggregation in vitro [30], possibly by decreasing the β -strand content of the C-terminal region [130]. Furthermore, proteins can become modified by non-enzymatic glycation upon aging. Advanced glycation end products (AGEs), found in A β plaques and in neurons, and their receptor RAGE play an important role in AD by contributing to oxidative stress and by triggering inflammation signaling pathways [128, 131, 132]. For other modifications, it remains largely unknown how they can affect A β aggregation dynamics.

Various forms of A β co-exist and co-deposit in amyloid fibrils and plaques [23, 128]. It has become clear that biologically relevant mixtures of A β alloforms behave in a more complex manner in vitro than anticipated from their behavior in isolation, in terms of aggregation properties and toxicity [8–12]. For example, A β_{1-38} and A β_{1-40} exerted little toxicity in isolation, but were highly toxic to a neuroblastoma cell line when tested in a mixture, whereas addition of A β_{1-38} to A β_{1-42} had a protective effect [10].

Recently, it has been demonstrated that minor shifts in the A β_{1-42} :A β_{1-40} ratio can modulate neurotoxicity [8]. The aggregation of samples of A β lengths in various compositions were monitored by NMR allowing simultaneous investigation of both A β_{1-42} and A β_{1-40} in the same sample by combining ¹⁵N-isotope-labeling of one A β alloform with ¹⁵N-edited filter experiments [9]. It was revealed that A β_{1-42} and A β_{1-40} directly interact and influence oligomer formation and aggregation kinetics. Moreover, cross-seeding data revealed structural differences between the different ratios at the level of the oligomeric state. A subtle change in the A β_{1-42} :A β_{1-40} ratio was suggested to induce differences in conformational plasticity of the oligomeric peptide mixtures [9]. High molecular weight (HMW) mass spectra further showed that a continuous range of

oligomeric intermediates were formed upon incubation of A β through a monomer addition process for the time frame within which toxicity exists [8, 9]. This observation is in agreement with the “coalescence and reorganization model of amyloid formation” [133], but also with the principle of a template-dependent dock-and-lock-and-block mechanism whereby the locking of a peptide cannot efficiently occur unless the previously loaded peptide has assembled into the correct position [134]. This can be envisaged in the following way: intrinsically disordered A β monomers diffuse freely and can attach individually to each other, to a pre-existing oligomer or to the fiber surface, especially through the distal ends. The crucial step occurs when the incoming monomer collides with the docking surface. In the case of a productive association, a permanent attachment can then take place, perhaps accompanied by a minor structural rearrangement. The conformational constraints of the monomers will therefore influence the efficiency and kinetics of the aggregation as well as the architecture of A β fibrils [135] (see “A β fibrils are polymorphic entities”). Alloform differences of the monomeric conformation are essential at this point to interpret productive or non-productive interactions [38, 40, 41], particularly in the complex in vivo pool of peptides. A β_{1-42} has the tendency to sample more fibril-like conformations compared to A β_{1-40} and as such can simply dock to fibril-like oligomers leading to highly productive (on-pathway) interactions. The more rigid and less flexible C-terminus of A β_{1-42} was suggested to enable the formation of a larger number of intramolecular contacts than A β_{1-40} [37, 40] and therefore provide a more extensive hydrophobic surface for intermolecular interactions. Experiments using amino acid substitutions in the C-terminal part of A β_{1-40} and A β_{1-42} confirmed that (i) the stability of the β -hairpin structure was increased by reducing the backbone flexibility and strengthening the hydrophobic interactions between the putative β -strands, (ii) destabilizing mutations in the C-terminal part of A β_{1-42} lead to a more A β_{1-40} -like behavior, and (iii) stabilizing mutations in the C-terminus of A β_{1-40} lead to a more A β_{1-42} -like behavior [143]. The conformational search of the incoming peptide for binding on the docking surface and for the proper orientation to lock-in-place could explain the complex aggregation behavior of A β alloform mixtures. The balance between productive and non-productive interactions in the transient encounter states is essential to guide the kinetics of aggregation, which in turn will define the time window within which the toxic species exist. Now that it becomes evident from independent research groups that the pattern of oligomer formation is mainly influenced by (patho)physiologically relevant A β_{1-42} :A β_{1-40} ratios [136], it is also important to realize that independent (on- and off-) pathways exist for oligomerization and fibrillization of A β [137, 138].

Experimental approaches to obtain insight into complex A β dynamics

It seems logical that the assembly and disassembly of toxic species is a dynamic and continuous process, at least in the initial stages, that is directed by the A β pool composition. However, the possibility that toxicity is present over a series of conformers or sizes should not be disregarded [91, 92, 94, 139, 140]. The question is thus how biophysical parameters influence this process in vivo and affect the relative distribution of A β species over toxic and non-toxic conformations over time. Given the complexity of the biophysical environment in which A β aggregation occurs in vivo, such a question is extremely difficult to address. Nevertheless, it is possible to analyze the dynamic features of this process in simplified and controlled conditions in vitro, and to evaluate the effect of the relative concentrations of A β_{1-40} and A β_{1-42} (and other alloforms) to the generation of neurotoxic species over time.

The combination of high-resolution NMR and HMW MS is perfectly suited to investigate the individual aggregation behavior of the diverse A β alloforms in complex and heterogeneous sample compositions. This can yield a comprehensive *aggregation fingerprint* that allows us to understand how the different compositions of the A β peptide pool influence the overall aggregation behavior. This aggregation fingerprint can be related to cytotoxicity, membrane integrity, apoptotic responses, and functional readouts such as microelectrode arrays (MEA), in which synaptic activities at different timeframes and under various conditions are monitored in response to A β [8, 141]. Such a fingerprint also opens perspectives to the diagnostics and therapeutics field when it can be correlated to biomarkers. Patient-specific treatment (personalized medicine) could be based on the detailed characterization of the composition of the A β pool. It will be essential to correlate the aggregation fingerprint of such compositions with disease severity and the (ir)reversibility of the disease “progress”. It is also important to cover the overall dynamics in these pools rather than focusing on particular “toxic” intermediates that are only transient in the aggregation process. This will allow tackling the source of toxicity and limiting the time frame in which the toxic assemblies can exist. Aggregation fingerprints will thus be essential to better understand the A β -induced pathogenesis of AD and the biophysical processes that underlie the cell biological responses.

The interaction between different species present during A β aggregation

NMR relaxation measurements showed that monomers are constantly binding to and being released from oligomers in vitro [142, 143]. Estimates showed that approximately

3 % of the peptide within the oligomer undergoes exchange with free monomer in pseudo-equilibrium conditions, suggesting that exchange occurs predominantly from the oligomer surface. A large part of the hydrophobic C-terminal region is involved in the association of monomer onto the oligomer surface [142]. In a next aggregation phase, protofibrils are formed that are also in constant exchange with monomers through the same surface region [144]. An elegant combination of ^{19}F -NMR and other biophysical techniques revealed a heterogeneous mixture of small A β oligomers that exist in pseudo-equilibrium with protofibrils and fibrils during the early stages of aggregation [145].

Protofibrils self-associate and give rise to mature fibrils that can thermodynamically be considered as the most stable aggregation state due to the high density of intermolecular hydrogen bonding and steric zipper interactions [146]. However, fibrils are not static and irreversible end species, as was the traditional view, but were shown to continuously dissociate and reassociate through both fibril ends [147]. A β_{1-40} fibrils recycle to a greater extent than A β_{1-42} fibrils, which could be attributed to a difference in fibril dissolution rate. These findings are consistent with a dynamical model for interpreting plaque morphology, in which aggregation and disaggregation were proposed to be in steady-state equilibrium [148]. The species involved in the fibril recycling process are still a matter of debate. Differential solution NMR isotope labeling experiments revealed that A β_{1-40} monomers can replace A β_{1-42} on A β_{1-42} aggregates, recycling A β_{1-42} monomers back into solution [14]. Later reports confirmed the constant recycling of A β_{1-40} and A β_{1-42} monomers and competition of binding for the ends of protofibrillar and fibrillar aggregates [13]. Alternatively, the accumulation of fibrils could be associated with the generation of diffusible lower molecular weight aggregates. This idea is consistent with the observation of a halo of oligomeric A β surrounding senile plaques when analyzed by array tomography [149]. Recently, Knowles and coworkers demonstrated that the secondary nucleation pathway can be a major source of oligomers once the critical concentration of amyloid fibrils (in the order of 10 nM) has formed [150]. Hereby, the surfaces of existing fibrils catalyze the nucleation of new aggregates from the monomeric state, with a rate dependent on both the concentration of the monomers and that of the existing fibrils. As the critical fibril concentration is lower than the aggregate loads present in brains of AD patients, this pathway is likely to be active in the brain [150].

The dynamical equilibrium potentially contributes to A β -associated toxicity

The co-existence of different A β aggregate species should be taken into account when analyzing A β toxicity studies.

For example, fibrils act as a reservoir of soluble aggregates that can diffuse and induce toxic effects. The halo of oligomers surrounding senile plaques co-localizes with loss of excitatory synapses and spine collapse [149] and the disruption of dendritic spines in the vicinity of plaques is dependent on their distance from these plaques [151]. Moreover, fibrils can be destabilized by brain lipids and reverted into neurotoxic soluble protofibrils [139]. Amyloid fibrils can thus be toxic per se (see “Fibril polymorphism could lead to different pathological outcomes”) or can function as a potential source of neurotoxic oligomeric species [152, 153]. It has also been suggested that the ongoing polymerization process, rather than the formation of one stable aggregate, is responsible for A β -related toxicity [19, 154]. In accordance with this hypothesis, crude A β_{1-42} preparations containing a monomeric and heterogeneous mixture of A β_{1-42} oligomers and protofibrils were more toxic than purified monomeric, protofibrillar fractions or fibrils. The toxicity of protofibrils was directly linked with their interactions with monomeric A β_{1-42} and strongly dependent on their ability to convert into fibrils. Moreover, the ongoing A β aggregation process, rather than distinct aggregation states, elicited alterations in astrocyte metabolic phenotypes [19]. Therefore, insight into the dynamic equilibrium is required to fully understand A β toxicity.

Other players in the game

The modulation of A β production, aggregation, and degradation by environmental factors [155–157], genetic risk factors [158–161], post-translational modifications [127], and an individual’s lifestyle [162–169] has been extensively reviewed before and does not lie in the scope of this review. Only a few reports discuss the influence of these factors on A β dynamics.

Metals have been shown to affect A β intramolecular dynamics. Binding of zinc to the N-terminus of the A β monomer leads to a decrease in the intrinsic mobility of this region and the formation of a turn-like conformation in residues Val24-Lys28 promoting aggregation, as shown by ^{15}N relaxation measurements [170]. Copper can also bind to the N-terminus, causing a structural ordering in this region [171], but slowing down aggregation [110].

There is evidence that membrane composition and properties, in turn, play a critical role in A β cytotoxicity associated with its conformational changes and aggregation into oligomers and fibrils ([172–174], reviewed in [175]). Moreover, interaction with lipid membranes can modulate A β peptide conformation and aggregation properties (reviewed in [175, 176]).

Genetic evidence suggested a role for chaperones in AD [177] and abundant chaperone levels block formation of

A β aggregates as was demonstrated in a *Caenorhabditis elegans* disease model [178]. In vitro results indicated a role for heat shock proteins in the early aggregation events by interfering with the dynamical aggregation process [179]. The BRICHOS domain, a chaperone-like domain found in lung surfactant protein C, is reported to be a potent in vitro inhibitor of A β aggregation [180]. The contribution of chaperones in the context of AD is reviewed in [181].

Interactions of A β with small molecules designed to target A β toxicity and/or A β aggregation have also been extensively studied. These ligands are not only interesting in light of drug development, but also provide a tool for addressing the modulation of A β dynamics upon ligand interaction [182–184].

As the A β monomer concentration affects the dynamical equilibrium between monomers, oligomers, protofibrils, and fibrillar A β , it is also worthwhile to consider factors that modulate A β metabolism. Aluminium is known to increase the A β brain burden in experimental animals and this might be due to a direct influence upon A β anabolism or to direct or indirect effects on A β catabolism [185]. Holtzman and coworkers reported that human cerebrospinal fluid A β levels undergo diurnal fluctuations and that this cycle is disturbed following plaque formation before the appearance of any cognitive symptoms [186]. A β fluctuations were affected by perturbation of the orexin signaling pathway and the sleep-wake cycle and this suggested that sleep abnormalities in earlier life might predispose an individual to AD [187]. Cholesterol has been suggested to provide stability to membrane-adjacent lipid rafts and therefore facilitate the A β cleavage from APP [168]. Recent evidence showed that the γ -secretase subunit composition defines the A β profile and affects the ratio between alloforms [6]. This implies that external factors influencing the γ -secretase subunit composition will have a profound effect on A β toxicity.

Conclusions

Understanding the intrinsic molecular flexibility, dynamics of interactions, and the structural behavior of the various A β peptides is crucial to comprehend the molecular mechanisms underlying the pathophysiology of Alzheimer’s disease. This will allow a more rational design of therapeutic intervention strategies to halt the disease progress and neutralize the malignant action of A β aggregation. To gain understanding of these events is difficult if not impossible to follow in real-time in the human brain. Therefore, these events are often mimicked in the test tube in research laboratories where information on A β behavior can be followed in molecular detail using advanced biophysical and

biochemical assays in the course of seconds to hours or days, which happen in patients over a range of years.

The intrinsically disordered nature of A β raises the question of whether this peptide may act as signaling peptide, which is known to require a high degree of flexibility. It is striking to observe that many proteins involved in human diseases are in fact classified as IDPs (alpha-synuclein, tau, multiple prions) [188, 189]. This raises the question as to whether protein flexibility may act as a disease-contributing factor as opposed to the generally accepted idea that specific sizes or conformations of oligomeric forms of these peptides induce pathogenesis. In this review we state that different types of dynamics can be distinguished varying from inter- to intramolecular factors as well as external factors and that recent observations strongly indicate that indeed the contribution of dynamics to pathogenesis warrants further investigation. As the dynamic nature of A β and its ability to undergo conformational changes and aggregation has hampered its study, promising new experimental approaches and chemical tools [182] are being developed to address A β dynamics, having the major advantage that they can be used directly without the need for modification of A β with additional amino acids or fluorophores [110, 190]. While a lot has been learned in the past from the behavior of the A β system, it is clear that the picture is still incomplete and extremely complex. Variability in terms of space (intra- and extracellular space, brain compartments, patient-to-patient differences, etc.) and time (circadian rhythm, aging, lifestyle, etc.) imposes additional dynamical factors, emphasizing the importance to better understand the fluctuating microenvironment. Therefore, it is opportune to compare the A β system to a complex ecosystem or society, where minor perturbations might have profound effects that can result in cataclysmic events. Various A β alloforms interact and mutually influence each other's behavior, but they also interact with the complex biological cell surface where they might exert a toxic effect by interfering with its normal functionality. Therefore, a holistic view of the dynamical A β ecosystem would enable us to initiate a successful ecosystem management strategy to prevent or remediate the AD pathobiology.

We summarized the evidence supporting the role of structural flexibility and in particular of the intrinsic protein disorder in the A β system to AD pathogenesis. A more systematic approach to the study of molecular flexibility in the A β system is required. This knowledge should then be integrated into future research efforts to optimize the clinical outcomes of drug trials.

Acknowledgments EH is supported by a FWO doctoral fellowship. NvN is supported by the VIB and the Flemish Hercules Foundation. KB is supported by a grant from the Internationale Stichting Alzheimer Onderzoek (ISAO), an Odysseus II award from FWO and a UTWIST fellowship. KP is the recipient of a FWO Pegasus long-term

postdoctoral fellowship and is supported by the Stichting Alzheimer Onderzoek (SAO-FRA).

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Mangialasche F, Solomon A, Winblad B, Mecocci P, Kivipelto M (2010) Alzheimer's disease: clinical trials and drug development. *Lancet Neurol* 9(7):702–716
- Huang Y, Mucke L (2012) Alzheimer mechanisms and therapeutic strategies. *Cell* 148(6):1204–1222
- Hamaguchi T, Ono K, Yamada M (2006) Anti-amyloidogenic therapies: strategies for prevention and treatment of Alzheimer's disease. *Cell Mol Life Sci* 63(13):1538–1552
- Hardy J, Allsop D (1991) Amyloid deposition as the central event in the aetiology of Alzheimer's disease. *Trends Pharmacol Sci* 12(10):383–388
- Takami M et al (2009) Gamma-secretase: successive tripeptide and tetrapeptide release from the transmembrane domain of beta-carboxyl terminal fragment. *J Neurosci* 29(41):13042–13052
- Acx H et al (2013) Signature A β profiles are produced by different γ -secretase complexes. *J Biol Chem* 289(7):4346–4355
- Finder VH, Vodopivec I, Nitsch RM, Glockshuber R (2010) The recombinant amyloid-beta peptide Abeta1-42 aggregates faster and is more neurotoxic than synthetic Abeta1-42. *J Mol Biol* 396(1):9–18
- Kuperstein I et al (2010) Neurotoxicity of Alzheimer's disease A β peptides is induced by small changes in the A β 42 to A β 40 ratio. *EMBO J* 29(19):3408–3420
- Pauwels K et al (2012) Structural basis for increased toxicity of pathological a β 42:a β 40 ratios in Alzheimer disease. *J Biol Chem* 287(8):5650–5660
- Vandersteen A et al (2012) Molecular plasticity regulates oligomerization and cytotoxicity of the multipetide-length amyloid- β peptide pool. *J Biol Chem* 287(44):36732–36743
- Yoshiike Y, Chui DH, Akagi T, Tanaka N, Takashima A (2003) Specific compositions of amyloid-beta peptides as the determinant of toxic beta-aggregation. *J Biol Chem* 278(26):23648–23655
- Snyder SW et al (1994) Amyloid-beta aggregation: selective inhibition of aggregation in mixtures of amyloid with different chain lengths. *Biophys J* 67(3):1216–1228
- Jan A, Gokce O, Luthi-Carter R, Lashuel HA (2008) The ratio of monomeric to aggregated forms of Abeta40 and Abeta42 is an important determinant of amyloid-beta aggregation, fibrillogenesis, and toxicity. *J Biol Chem* 283(42):28176–28189
- Yan Y, Wang C (2007) Abeta40 protects non-toxic Abeta42 monomer from aggregation. *J Mol Biol* 369(4):909–916
- Morris AM, Watzky MA, Finke RG (2009) Protein aggregation kinetics, mechanism, and curve-fitting: a review of the literature. *Biochim Biophys Acta* 1794(3):375–397
- Finder VH, Glockshuber R (2007) Amyloid-beta aggregation. *Neurodegener Dis* 4(1):13–27
- Karran E, Mercken M, De Strooper B (2011) The amyloid cascade hypothesis for Alzheimer's disease: an appraisal for the development of therapeutics. *Nat Rev Drug Discov* 10(9):698–712
- Haass C, Selkoe DJ (2007) Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's amyloid beta-peptide. *Nat Rev Mol Cell Biol* 8(2):101–112

19. Jan A et al (2011) Abeta42 neurotoxicity is mediated by ongoing nucleated polymerization process rather than by discrete Abeta42 species. *J Biol Chem* 286(10):8585–8596
20. Tompa P (2009) Structural disorder in amyloid fibrils: its implication in dynamic interactions of proteins. *FEBS J* 276(19):5406–5415
21. Glenner GG, Wong CW (1984) Alzheimer's disease: initial report of the purification and characterization of a novel cerebrovascular amyloid protein. *Biochem Biophys Res Commun* 120:885–890
22. Klyubin I et al (2008) Amyloid beta protein dimer-containing human CSF disrupts synaptic plasticity: prevention by systemic passive immunization. *J Neurosci* 28(16):4231–4237
23. Shankar GM et al (2008) Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. *Nat Med* 14(8):837–842
24. Jin M et al (2011) Soluble amyloid beta-protein dimers isolated from Alzheimer cortex directly induce Tau hyperphosphorylation and neuritic degeneration. *Proc Natl Acad Sci USA* 108(14):5819–5824
25. Hilbich C, Kisters-Woike B, Reed J, Masters CL, Beyreuther K (1991) Aggregation and secondary structure of synthetic amyloid beta A4 peptides of Alzheimer's disease. *J Mol Biol* 218(1):149–163
26. Tseng BP et al (1999) Deposition of monomeric, not oligomeric, Abeta mediates growth of Alzheimer's disease amyloid plaques in human brain preparations. *Biochemistry* 38(32):10424–10431
27. Jan A, Hartley DM, Lashuel HA (2010) Preparation and characterization of toxic Abeta aggregates for structural and functional studies in Alzheimer's disease research. *Nat Protoc* 5(6):1186–1209
28. Zhang S et al (2000) The Alzheimer's peptide a beta adopts a collapsed coil structure in water. *J Struct Biol* 130(2–3):130–141
29. Riek R, Güntert P, Döbeli H, Wipf B, Wüthrich K (2001) NMR studies in aqueous solution fail to identify significant conformational differences between the monomeric forms of two Alzheimer peptides with widely different plaque-competence, A beta(1–40)(ox) and A beta(1–42)(ox). *Eur J Biochem* 268(22):5930–5936
30. Hou L et al (2004) Solution NMR studies of the A beta(1–40) and A beta(1–42) peptides establish that the Met35 oxidation state affects the mechanism of amyloid formation. *J Am Chem Soc* 126(7):1992–2005
31. Shao H, Jao S, Ma K, Zagorski MG (1999) Solution structures of micelle-bound amyloid beta-(1–40) and beta-(1–42) peptides of Alzheimer's disease. *J Mol Biol* 285(2):755–773
32. Tomaselli S et al (2006) The alpha-to-beta conformational transition of Alzheimer's Abeta-(1–42) peptide in aqueous media is reversible: a step-by-step conformational analysis suggests the location of beta conformation seeding. *Chem Bio Chem* 7(2):257–267
33. Vivekanandan S, Brender JR, Lee SY, Ramamoorthy A (2011) A partially folded structure of amyloid-beta(1–40) in an aqueous environment. *Biochem Biophys Res Commun* 411(2):312–316
34. Baumketner A et al (2006) Structure of the 21–30 fragment of amyloid beta-protein. *Protein Sci* 15(6):1239–1247
35. Lazo ND, Grant MA, Condrón MC, Rigby AC, Teplow DB (2005) On the nucleation of amyloid beta-protein monomer folding. *Protein Sci* 14(6):1581–1596
36. Danielsson J, Andersson A, Jarvet J, Gräslund A (2006) 15N relaxation study of the amyloid beta-peptide: structural propensities and persistence length. *Magn Reson Chem* 44 Spec No:S114–121
37. Yan Y, Wang C (2006) Abeta42 is more rigid than Abeta40 at the C terminus: implications for Abeta aggregation and toxicity. *J Mol Biol* 364(5):853–862
38. Sgourakis NG, Yan Y, McCallum SA, Wang C, Garcia AE (2007) The Alzheimer's peptides Abeta40 and 42 adopt distinct conformations in water: a combined MD/NMR study. *J Mol Biol* 368(5):1448–1457
39. Sgourakis NG et al (2011) Atomic-level characterization of the ensemble of the A $\beta_{(1-42)}$ monomer in water using unbiased molecular dynamics simulations and spectral algorithms. *J Mol Biol* 405(2):570–583
40. Yang M, Teplow DB (2008) Amyloid beta-protein monomer folding: free-energy surfaces reveal alloform-specific differences. *J Mol Biol* 384(2):450–464
41. Rosenman DJ, Connors CR, Chen W, Wang C, García AE (2013) A β monomers transiently sample oligomer and fibril-like configurations: ensemble characterization using a combined MD/NMR approach. *J Mol Biol* 425(18):3338–3359
42. Wright PE, Dyson HJ (2009) Linking folding and binding. *Curr Opin Struct Biol* 19(1):31–38
43. Tompa P, Szász C, Buday L (2005) Structural disorder throws new light on moonlighting. *Trends Biochem Sci* 30(9):484–489
44. Török M et al (2002) Structural and dynamic features of Alzheimer's Abeta peptide in amyloid fibrils studied by site-directed spin labeling. *J Biol Chem* 277(43):40810–40815
45. Kheterpal I, Zhou S, Cook KD, Wetzel R (2000) Abeta amyloid fibrils possess a core structure highly resistant to hydrogen exchange. *Proc Natl Acad Sci USA* 97(25):13597–13601
46. Kheterpal I, Williams A, Murphy C, Bledsoe B, Wetzel R (2001) Structural features of the Abeta amyloid fibril elucidated by limited proteolysis. *Biochemistry* 40(39):11757–11767
47. Wang SS, Tobler SA, Good TA, Fernandez EJ (2003) Hydrogen exchange-mass spectrometry analysis of beta-amyloid peptide structure. *Biochemistry* 42(31):9507–9514
48. Sunde M et al (1997) Common core structure of amyloid fibrils by synchrotron X-ray diffraction. *J Mol Biol* 273(3):729–739
49. Jahn TR et al (2010) The common architecture of cross-beta amyloid. *J Mol Biol* 395(4):717–727
50. Petkova AT et al (2002) A structural model for Alzheimer's beta-amyloid fibrils based on experimental constraints from solid state NMR. *Proc Natl Acad Sci USA* 99(26):16742–16747
51. Bertini I, Gonnelli L, Luchinat C, Mao J, Nesi A (2011) A new structural model of A β 40 fibrils. *J Am Chem Soc* 133(40):16013–16022
52. Whittemore NA et al (2005) Hydrogen-deuterium (H/D) exchange mapping of Abeta(1–40) amyloid fibril secondary structure using nuclear magnetic resonance spectroscopy. *Biochemistry* 44(11):4434–4441
53. Luhurs T et al (2005) 3D structure of Alzheimer's amyloid-beta(1–42) fibrils. *Proc Natl Acad Sci USA* 102(48):17342–17347
54. Olofsson A, Sauer-Eriksson AE, Ohman A (2006) The solvent protection of Alzheimer amyloid-beta-(1–42) fibrils as determined by solution NMR spectroscopy. *J Biol Chem* 281(1):477–483
55. Morimoto A et al (2004) Analysis of the secondary structure of beta-amyloid (Abeta42) fibrils by systematic proline replacement. *J Biol Chem* 279(50):52781–52788
56. Kheterpal I, Chen M, Cook KD, Wetzel R (2006) Structural differences in Abeta amyloid protofibrils and fibrils mapped by hydrogen exchange-mass spectrometry with on-line proteolytic fragmentation. *J Mol Biol* 361(4):785–795
57. Sawaya MR et al (2007) Atomic structures of amyloid cross-beta spines reveal varied steric zippers. *Nature* 447(7143):453–457

58. Williams AD et al (2004) Mapping abeta amyloid fibril secondary structure using scanning proline mutagenesis. *J Mol Biol* 335(3):833–842
59. Petkova AT et al (2005) Self-propagating, molecular-level polymorphism in Alzheimer's beta-amyloid fibrils. *Science* 307(5707):262–265
60. Scheidt HA, Morgado I, Rothmund S, Huster D, Fändrich M (2011) Solid-state NMR spectroscopic investigation of A β protofibrils: implication of a β -sheet remodeling upon maturation into terminal amyloid fibrils. *Angew Chem Int Ed Engl* 50(12):2837–2840
61. Scheidt HA, Morgado I, Rothmund S, Huster D (2012) Dynamics of amyloid β fibrils revealed by solid-state NMR. *J Biol Chem* 287(3):2017–2021
62. Morel B, Varela L, Conejero-Lara F (2010) The thermodynamic stability of amyloid fibrils studied by differential scanning calorimetry. *J Phys Chem B* 114(11):4010–4019
63. Ma J et al (2013) Intrinsic structural heterogeneity and long-term maturation of amyloid β peptide fibrils. *ACS Chem Neurosci* 4(8):1236–1243
64. Nyström S et al (2013) Evidence for age-dependent in vivo conformational rearrangement within A β amyloid deposits. *ACS Chem Biol* 8(6):1128–1133
65. Fändrich M, Meinhardt J, Grigorieff N (2009) Structural polymorphism of Alzheimer Abeta and other amyloid fibrils. *Prion* 3(2):89–93
66. Tycko R, Wickner RB (2013) Molecular structures of amyloid and prion fibrils: consensus versus controversy. *Acc Chem Res* 46(7):1487–1496
67. Crowther RA, Goedert M (2000) Abnormal tau-containing filaments in neurodegenerative diseases. *J Struct Biol* 130(2–3):271–279
68. Jiménez JL, Tennent G, Pepys M, Saibil HR (2001) Structural diversity of ex vivo amyloid fibrils studied by cryo-electron microscopy. *J Mol Biol* 311(2):241–247
69. Bousset L et al (2013) Structural and functional characterization of two alpha-synuclein strains. *Nat Commun* 4:2575
70. Kodali R, Williams AD, Chemuru S, Wetzel R (2010) Abeta(1-40) forms five distinct amyloid structures whose beta-sheet contents and fibril stabilities are correlated. *J Mol Biol* 401(3):503–517
71. Klement K et al (2007) Effect of different salt ions on the propensity of aggregation and on the structure of Alzheimer's abeta(1-40) amyloid fibrils. *J Mol Biol* 373(5):1321–1333
72. Pedersen JS, Otzen DE (2008) Amyloid-a state in many guises: survival of the fittest fibril fold. *Protein Sci* 17(1):2–10
73. Meinhardt J, Sachse C, Hortschansky P, Grigorieff N, Fändrich M (2009) Abeta(1-40) fibril polymorphism implies diverse interaction patterns in amyloid fibrils. *J Mol Biol* 386(3):869–877
74. Qiang W, Yau WM, Luo Y, Mattson MP, Tycko R (2012) Antiparallel β -sheet architecture in Iowa-mutant β -amyloid fibrils. *Proc Natl Acad Sci USA* 109(12):4443–4448
75. Qiang W, Kelley K, Tycko R (2013) Polymorph-specific kinetics and thermodynamics of β -amyloid fibril growth. *J Am Chem Soc* 135(18):6860–6871
76. Sachse C et al (2006) Quaternary structure of a mature amyloid fibril from Alzheimer's Abeta(1-40) peptide. *J Mol Biol* 362(2):347–354
77. Sachse C, Fändrich M, Grigorieff N (2008) Paired beta-sheet structure of an Abeta(1-40) amyloid fibril revealed by electron microscopy. *Proc Natl Acad Sci USA* 105(21):7462–7466
78. Schmidt M et al (2009) Comparison of Alzheimer Abeta(1-40) and Abeta(1-42) amyloid fibrils reveals similar protofilament structures. *Proc Natl Acad Sci USA* 106(47):19813–19818
79. Zhang R et al (2009) Interprotofilament interactions between Alzheimer's Abeta1-42 peptides in amyloid fibrils revealed by cryoEM. *Proc Natl Acad Sci USA* 106(12):4653–4658
80. Goldsbury CS et al (2000) Studies on the in vitro assembly of a beta 1-40: implications for the search for a beta fibril formation inhibitors. *J Struct Biol* 130(2–3):217–231
81. Goldsbury C, Frey P, Olivieri V, Aepli U, Müller SA (2005) Multiple assembly pathways underlie amyloid-beta fibril polymorphisms. *J Mol Biol* 352(2):282–298
82. Petkova AT, Yau WM, Tycko R (2006) Experimental constraints on quaternary structure in Alzheimer's beta-amyloid fibrils. *Biochemistry* 45(2):498–512
83. Paravastu AK, Leapman RD, Yau WM, Tycko R (2008) Molecular structural basis for polymorphism in Alzheimer's beta-amyloid fibrils. *Proc Natl Acad Sci USA* 105(47):18349–18354
84. Cameron B, Landreth GE (2010) Inflammation, microglia, and Alzheimer's disease. *Neurobiol Dis* 37(3):503–509
85. Ill-Raga G et al (2010) Amyloid- β peptide fibrils induce nitro-oxidative stress in neuronal cells. *J Alzheimers Dis* 22(2):641–652
86. McKee AC, Kosik KS, Kowall NW (1991) Neuritic pathology and dementia in Alzheimer's disease. *Ann Neurol* 30(2):156–165
87. Berg L et al (1998) Clinicopathologic studies in cognitively healthy aging and Alzheimer's disease: relation of histologic markers to dementia severity, age, sex, and apolipoprotein E genotype. *Arch Neurol* 55(3):326–335
88. Seilheimer B et al (1997) The toxicity of the Alzheimer's beta-amyloid peptide correlates with a distinct fiber morphology. *J Struct Biol* 119(1):59–71
89. Lu JX et al (2013) Molecular structure of β -amyloid fibrils in Alzheimer's disease brain tissue. *Cell* 154(6):1257–1268
90. Ihse E et al (2008) Amyloid fibril composition is related to the phenotype of hereditary transthyretin V30M amyloidosis. *J Pathol* 216(2):253–261
91. Benilova I, Karran E, De Strooper B (2012) The toxic A β oligomer and Alzheimer's disease: an emperor in need of clothes. *Nat Neurosci* 15(3):349–357
92. Broersen K, Rousseau F, Schymkowitz J (2010) The culprit behind amyloid beta peptide related neurotoxicity in Alzheimer's disease: oligomer size or conformation? *Alzheimers Res Ther* 2(4):12
93. Kaye R, Lasagna-Reeves CA (2013) Molecular mechanisms of amyloid oligomers toxicity. *J Alzheimers Dis* 33(Suppl 1):S67–S78
94. Ono K, Condron MM, Teplow DB (2009) Structure-neurotoxicity relationships of amyloid beta-protein oligomers. *Proc Natl Acad Sci USA* 106(35):14745–14750
95. Cizas P et al (2010) Size-dependent neurotoxicity of beta-amyloid oligomers. *Arch Biochem Biophys* 496(2):84–92
96. Campioni S et al (2010) A causative link between the structure of aberrant protein oligomers and their toxicity. *Nat Chem Biol* 6(2):140–147
97. Nekooki-Machida Y et al (2009) Distinct conformations of in vitro and in vivo amyloids of huntingtin-exon1 show different cytotoxicity. *Proc Natl Acad Sci USA* 106(24):9679–9684
98. Ladiwala AR et al (2012) Conformational differences between two amyloid β oligomers of similar size and dissimilar toxicity. *J Biol Chem* 287(29):24765–24773
99. Deshpande A, Mina E, Glabe C, Busciglio J (2006) Different conformations of amyloid beta induce neurotoxicity by distinct mechanisms in human cortical neurons. *J Neurosci* 26(22):6011–6018
100. Kaye R et al (2003) Common structure of soluble amyloid oligomers implies common mechanism of pathogenesis. *Science* 300(5618):486–489
101. Glabe CG (2008) Structural classification of toxic amyloid oligomers. *J Biol Chem* 283(44):29639–29643
102. Kaye R et al (2009) Annular protofibrils are a structurally and functionally distinct type of amyloid oligomer. *J Biol Chem* 284(7):4230–4237

103. Kayed R et al (2010) Conformation-dependent monoclonal antibodies distinguish different replicating strains or conformers of prefibrillar A β oligomers. *Mol Neurodegener* 5:57
104. Cerf E et al (2009) Antiparallel beta-sheet: a signature structure of the oligomeric amyloid beta-peptide. *Biochem J* 421(3):415–423
105. Sarroukh R et al (2011) Transformation of amyloid β (1–40) oligomers into fibrils is characterized by a major change in secondary structure. *Cell Mol Life Sci* 68(8):1429–1438
106. Celej MS et al (2012) Toxic prefibrillar α -synuclein amyloid oligomers adopt a distinctive antiparallel β -sheet structure. *Biochem J* 443(3):719–726
107. Vandersteen A et al (2012) A comparative analysis of the aggregation behavior of amyloid- β peptide variants. *FEBS Lett* 586(23):4088–4093
108. Laganowsky A et al (2012) Atomic view of a toxic amyloid small oligomer. *Science* 335(6073):1228–1231
109. Gu L, Liu C, Guo Z (2013) Structural insights into A β 42 oligomers using site-directed spin labeling. *J Biol Chem* 288(26):18673–18683
110. Zhang Y et al (2013) Pulsed hydrogen-deuterium exchange mass spectrometry probes conformational changes in amyloid beta (A β) peptide aggregation. *Proc Natl Acad Sci USA* 110(36):14604–14609
111. Ahmed M et al (2010) Structural conversion of neurotoxic amyloid-beta(1–42) oligomers to fibrils. *Nat Struct Mol Biol* 17(5):561–567
112. Bolognesi B et al (2010) ANS binding reveals common features of cytotoxic amyloid species. *ACS Chem Biol* 5(8):735–740
113. Williams TL et al (2011) A β 42 oligomers, but not fibrils, simultaneously bind to and cause damage to ganglioside-containing lipid membranes. *Biochem J* 439(1):67–77
114. Teplow DB (2013) On the subject of rigor in the study of amyloid β -protein assembly. *Alzheimers Res Ther* 5(4):39
115. Lesné S et al (2006) A specific amyloid-beta protein assembly in the brain impairs memory. *Nature* 440(7082):352–357
116. Qi-Takahara Y et al (2005) Longer forms of amyloid beta protein: implications for the mechanism of intramembrane cleavage by gamma-secretase. *J Neurosci* 25(2):436–445
117. Wiltfang J et al (2002) Highly conserved and disease-specific patterns of carboxyterminally truncated Abeta peptides 1–37/38/39 in addition to 1–40/42 in Alzheimer's disease and in patients with chronic neuroinflammation. *J Neurochem* 81(3):481–496
118. Vigo-Pelfrey C, Lee D, Keim P, Lieberburg I, Schenk DB (1993) Characterization of beta-amyloid peptide from human cerebrospinal fluid. *J Neurochem* 61(5):1965–1968
119. Masters CL, Selkoe DJ (2012) Biochemistry of amyloid β -protein and amyloid deposits in Alzheimer disease. *Cold Spring Harb Perspect Med* 2(6):a006262
120. Moro ML, Collins MJ, Cappellini E (2010) Alzheimer's disease and amyloid beta-peptide deposition in the brain: a matter of 'aging'? *Biochem Soc Trans* 38(2):539–544
121. Mori H et al (1994) Racemization: its biological significance on neuropathogenesis of Alzheimer's disease. *Tohoku J Exp Med* 174(3):251–262
122. Kubo T, Kumagai Y, Miller CA, Kaneko I (2003) Beta-amyloid racemized at the Ser26 residue in the brains of patients with Alzheimer disease: implications in the pathogenesis of Alzheimer disease. *J Neuropathol Exp Neurol* 62(3):248–259
123. Kuo YM, Webster S, Emmerling MR, De Lima N, Roher AE (1998) Irreversible dimerization/tetramerization and post-translational modifications inhibit proteolytic degradation of A beta peptides of Alzheimer's disease. *Biochim Biophys Acta* 1406(3):291–298
124. Milton NG (2001) Phosphorylation of amyloid-beta at the serine 26 residue by human cdc2 kinase. *Neuroreport* 12(17):3839–3844
125. Kumar S et al (2011) Extracellular phosphorylation of the amyloid β -peptide promotes formation of toxic aggregates during the pathogenesis of Alzheimer's disease. *EMBO J* 30(11):2255–2265
126. Dong J et al (2003) Metal binding and oxidation of amyloid-beta within isolated senile plaque cores: Raman microscopic evidence. *Biochemistry* 42(10):2768–2773
127. Hou L et al (2013) Modification of amyloid- β 1–42 fibril structure by methionine-35 oxidation. *J Alzheimers Dis* 37(1):9–18
128. Vitek MP et al (1994) Advanced glycation end products contribute to amyloidosis in Alzheimer disease. *Proc Natl Acad Sci USA* 91(11):4766–4770
129. Jawhar S, Wirths O, Bayer TA (2011) Pyroglutamate amyloid- β (A β): a hatchet man in Alzheimer disease. *J Biol Chem* 286(45):38825–38832
130. Brown AM, Lemkul JA, Schaum N, Bevan DR (2014) Simulations of monomeric amyloid β -peptide (1–40) with varying solution conditions and oxidation state of Met35: implications for aggregation. *Arch Biochem Biophys* 545:44–52
131. Srikanth V et al (2011) Advanced glycation endproducts and their receptor RAGE in Alzheimer's disease. *Neurobiol Aging* 32(5):763–777
132. Gu L, Guo Z (2013) Alzheimer's A β 42 and A β 40 peptides form interlaced amyloid fibrils. *J Neurochem* 126(3):305–311
133. Serio TR et al (2000) Nucleated conformational conversion and the replication of conformational information by a prion determinant. *Science* 289(5483):1317–1321
134. Esler WP et al (2000) Alzheimer's disease amyloid propagation by a template-dependent dock-lock mechanism. *Biochemistry* 39(21):6288–6295
135. Brännström K, Ohman A, Olofsson A (2011) A β peptide fibrillar architectures controlled by conformational constraints of the monomer. *PLoS One* 6(9):e25157
136. Johnson RD et al (2013) Single-molecule imaging reveals a β 42:a β 40 ratio-dependent oligomer growth on neuronal processes. *Biophys J* 104(4):894–903
137. Necula M, Kaye R, Milton S, Glabe CG (2007) Small molecule inhibitors of aggregation indicate that amyloid beta oligomerization and fibrillization pathways are independent and distinct. *J Biol Chem* 282(14):10311–10324
138. Necula M et al (2007) Methylene blue inhibits amyloid Abeta oligomerization by promoting fibrillization. *Biochemistry* 46(30):8850–8860
139. Martins IC et al (2008) Lipids revert inert Abeta amyloid fibrils to neurotoxic protofibrils that affect learning in mice. *EMBO J* 27(1):224–233
140. Hepler RW et al (2006) Solution state characterization of amyloid beta-derived diffusible ligands. *Biochemistry* 45(51):15157–15167
141. Chong SA et al (2011) Synaptic dysfunction in hippocampus of transgenic mouse models of Alzheimer's disease: a multi-electrode array study. *Neurobiol Dis* 44(3):284–291
142. Fawzi NL, Ying J, Torchia DA, Clore GM (2010) Kinetics of amyloid beta monomer-to-oligomer exchange by NMR relaxation. *J Am Chem Soc* 132(29):9948–9951
143. Krishnamoorthy J, Brender JR, Vivekanandan S, Jahr N, Ramamoorthy A (2012) Side-chain dynamics reveals transient association of A β (1–40) monomers with amyloid fibers. *J Phys Chem B* 116(46):13618–13623
144. Fawzi NL, Ying J, Ghirlardo R, Torchia DA, Clore GM (2011) Atomic-resolution dynamics on the surface of amyloid- β protofibrils probed by solution NMR. *Nature* 480(7376):268–272
145. Suzuki Y et al (2013) Resolution of oligomeric species during the aggregation of A β 1–40 using (19)F NMR. *Biochemistry* 52(11):1903–1912

146. Hartl FU, Hayer-Hartl M (2009) Converging concepts of protein folding in vitro and in vivo. *Nat Struct Mol Biol* 16(6):574–581
147. Carulla N, Zhou M, Giralte E, Robinson CV, Dobson CM (2010) Structure and intermolecular dynamics of aggregates populated during amyloid fibril formation studied by hydrogen/deuterium exchange. *Acc Chem Res* 43(8):1072–1079
148. Cruz L et al (1997) Aggregation and disaggregation of senile plaques in Alzheimer disease. *Proc Natl Acad Sci USA* 94(14):7612–7616
149. Koffie RM et al (2009) Oligomeric amyloid beta associates with postsynaptic densities and correlates with excitatory synapse loss near senile plaques. *Proc Natl Acad Sci USA* 106(10):4012–4017
150. Cohen SI et al (2013) Proliferation of amyloid- β 42 aggregates occurs through a secondary nucleation mechanism. *Proc Natl Acad Sci USA* 110(24):9758–9763
151. Spires-Jones TL et al (2009) Passive immunotherapy rapidly increases structural plasticity in a mouse model of Alzheimer disease. *Neurobiol Dis* 33(2):213–220
152. Xue WF, Hellewell AL, Hewitt EW, Radford SE (2010) Fibril fragmentation in amyloid assembly and cytotoxicity: when size matters. *Prion* 4(1):20–25
153. Cremades N et al (2012) Direct observation of the interconversion of normal and toxic forms of α -synuclein. *Cell* 149(5):1048–1059
154. Wogulis M et al (2005) Nucleation-dependent polymerization is an essential component of amyloid-mediated neuronal cell death. *J Neurosci* 25(5):1071–1080
155. Tiiman A, Palumaa P, Tõugu V (2013) The missing link in the amyloid cascade of Alzheimer's disease-metal ions. *Neurochem Int* 62(4):367–378
156. Pithadia AS, Lim MH (2012) Metal-associated amyloid- β species in Alzheimer's disease. *Curr Opin Chem Biol* 16(1–2):67–73
157. Watt AD, Villemagne VL, Barnham KJ (2013) Metals, membranes, and amyloid- β oligomers: key pieces in the Alzheimer's disease puzzle? *J Alzheimers Dis* 33(Suppl 1):S283–S293
158. Bu G (2009) Apolipoprotein E and its receptors in Alzheimer's disease: pathways, pathogenesis and therapy. *Nat Rev Neurosci* 10(5):333–344
159. Holtzman DM, Herz J, Bu G (2012) Apolipoprotein E and apolipoprotein E receptors: normal biology and roles in Alzheimer disease. *Cold Spring Harb Perspect Med* 2(3):a006312
160. Mahley RW, Weisgraber KH, Huang Y (2006) Apolipoprotein E4: a causative factor and therapeutic target in neuropathology, including Alzheimer's disease. *Proc Natl Acad Sci USA* 103(15):5644–5651
161. Kim J, Basak JM, Holtzman DM (2009) The role of apolipoprotein E in Alzheimer's disease. *Neuron* 63(3):287–303
162. Ramesh BN, Rao TS, Prakasam A, Sambamurti K, Rao KS (2010) Neuronutrition and Alzheimer's disease. *J Alzheimers Dis* 19(4):1123–1139
163. Smid SD, Maag JL, Musgrave IF (2012) Dietary polyphenol-derived protection against neurotoxic β -amyloid protein: from molecular to clinical. *Food Funct* 3(12):1242–1250
164. Radak Z et al (2010) Exercise plays a preventive role against Alzheimer's disease. *J Alzheimers Dis* 20(3):777–783
165. Farina N, Rusted J, Tabet N (2013) The effect of exercise interventions on cognitive outcome in Alzheimer's disease: a systematic review. *Int Psychogeriatr* 1–10
166. Hoffmann K et al (2013) preserving cognition, quality of life, physical health and functional ability in Alzheimer's disease: the effect of physical exercise (ADEX trial): rationale and design. *Neuroepidemiology* 41(3–4):198–207
167. Li S et al (2013) Environmental novelty activates β 2-adrenergic signaling to prevent the impairment of hippocampal LTP by A β oligomers. *Neuron* 77(5):929–941
168. Silva T, Teixeira J, Remião F, Borges F (2013) Alzheimer's disease, cholesterol, and statins: the junctions of important metabolic pathways. *Angew Chem Int Ed Engl* 52(4):1110–1121
169. Ju YE, Lucey BP, Holtzman DM (2013) Sleep and Alzheimer disease pathology—a bidirectional relationship. *Nat Rev Neurol*
170. Rezaei-Ghaleh N, Giller K, Becker S, Zweckstetter M (2011) Effect of zinc binding on β -amyloid structure and dynamics: implications for A β aggregation. *Biophys J* 101(5):1202–1211
171. Syme CD, Nadal RC, Rigby SE, Viles JH (2004) Copper binding to the amyloid-beta (A β) peptide associated with Alzheimer's disease: folding, coordination geometry, pH dependence, stoichiometry, and affinity of A β -(1–28): insights from a range of complementary spectroscopic techniques. *J Biol Chem* 279(18):18169–18177
172. Peters I et al (2009) The interaction of beta-amyloid protein with cellular membranes stimulates its own production. *Biochim Biophys Acta* 1788(5):964–972
173. Williams TL, Serpell LC (2011) Membrane and surface interactions of Alzheimer's A β peptide—insights into the mechanism of cytotoxicity. *FEBS J* 278(20):3905–3917
174. Sciacca MF et al (2012) Two-step mechanism of membrane disruption by A β through membrane fragmentation and pore formation. *Biophys J* 103(4):702–710
175. Kotler SA, Walsh P, Brender JR, Ramamoorthy A (2014) Differences between amyloid- β aggregation in solution and on the membrane: insights into elucidation of the mechanistic details of Alzheimer's disease. *Chem Soc Rev*. doi:10.1039/C3CS60431D
176. Burke KA, Yates EA, Legleiter J (2013) Biophysical insights into how surfaces, including lipid membranes, modulate protein aggregation related to neurodegeneration. *Front Neurol* 4:17
177. Magrané J, Smith RC, Walsh K, Querfurth HW (2004) Heat shock protein 70 participates in the neuroprotective response to intracellularly expressed beta-amyloid in neurons. *J Neurosci* 24(7):1700–1706
178. Fonte V et al (2002) Interaction of intracellular beta amyloid peptide with chaperone proteins. *Proc Natl Acad Sci USA* 99(14):9439–9444
179. Evans CG, Wisén S, Gestwicki JE (2006) Heat shock proteins 70 and 90 inhibit early stages of amyloid beta-(1–42) aggregation in vitro. *J Biol Chem* 281(44):33182–33191
180. Knight SD, Presto J, Linse S, Johansson J (2013) The BRI-CHOS domain, amyloid fibril formation, and their relationship. *Biochemistry* 52(43):7523–7531
181. Koren J et al (2009) Chaperone signalling complexes in Alzheimer's disease. *J Cell Mol Med* 13(4):619–630
182. Lee S et al (2014) Rational design of a structural framework with potential use to develop chemical reagents that target and modulate multiple facets of Alzheimer's disease. *J Am Chem Soc* 136(1):299–310
183. Ramamoorthy A, Lim MH (2013) Structural characterization and inhibition of toxic amyloid- β oligomeric intermediates. *Biophys J* 105(2):287–288
184. DeToma AS, Salamekh S, Ramamoorthy A, Lim MH (2012) Misfolded proteins in Alzheimer's disease and type II diabetes. *Chem Soc Rev* 41(2):608–621
185. Exley C (2005) The aluminium-amyloid cascade hypothesis and Alzheimer's disease. *Subcell Biochem* 38:225–234
186. Roh JH et al (2012) Disruption of the sleep-wake cycle and diurnal fluctuation of β -amyloid in mice with Alzheimer's disease pathology. *Sci Transl Med* 4(150):150ra122
187. Kang JE et al (2009) Amyloid-beta dynamics are regulated by orexin and the sleep-wake cycle. *Science* 326(5955):1005–1007
188. Deleersnijder A, Gerard M, Debyser Z, Baekelandt V (2013) The remarkable conformational plasticity of alpha-synuclein: blessing or curse? *Trends Mol Med* 19(6):368–377

189. Uversky VN (2014) The triple power of D³: protein intrinsic disorder in degenerative diseases. *Front Biosci (Landmark Ed)* 19:181–258
190. Santi S et al (2013) Real-time amyloid aggregation monitoring with a photonic crystal-based approach. *Chem Phys Chem* 14(15):3476–3482
191. Kheterpal I, Wetzel R (2006) Hydrogen/deuterium exchange mass spectrometry: a window into amyloid structure. *Acc Chem Res* 39(9):584–593