

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/8934840>

X-Ray Crystal Structure of CutA from *Thermotoga maritima* at 1.4 Å Resolution

ARTICLE in PROTEINS STRUCTURE FUNCTION AND BIOINFORMATICS · NOVEMBER 2003

Impact Factor: 2.63 · DOI: 10.1002/prot.10585 · Source: PubMed

CITATIONS

14

READS

15

9 AUTHORS, INCLUDING:



Alex Savtchenko

Sanford Burnham Prebys Medical Discovery I...

157 PUBLICATIONS 4,147 CITATIONS

SEE PROFILE



Roman Aleksander Laskowski

European Molecular Biology Laboratory

99 PUBLICATIONS 30,908 CITATIONS

SEE PROFILE



Andrzej Joachimiak

Argonne National Laboratory

400 PUBLICATIONS 16,455 CITATIONS

SEE PROFILE



Rongguang Zhang

131 PUBLICATIONS 8,937 CITATIONS

SEE PROFILE

Published in final edited form as:

Proteins. 2004 January 1; 54(1): 162–165. doi:10.1002/prot.10585.

X-Ray Crystal Structure of CutA From *Thermotoga maritima* at 1.4 Å Resolution

Alexei Savchenko¹, Tatiana Skarina¹, Elena Evdokimova¹, James D. Watson⁴, Roman Laskowski⁴, Cheryl H. Arrowsmith^{1,3}, Aled M. Edwards^{1,2,*}, Andrzej Joachimiak^{5,*}, and Rong-guang Zhang⁵

¹Ontario Center for Structural Proteomics, University Health Network, Toronto, Ontario, Canada

²Banting and Best Department of Medical Research, Toronto, Ontario, Canada

³Ontario Cancer Institute and Department of Medical Biophysics, University of Toronto, Toronto, Ontario, Canada

⁴The Department of Crystallography, Birkbeck College, London

⁵Biosciences Division and Structural Biology Center, Argonne National Laboratory, Argonne, Illinois

Introduction

The structure of the CutA protein from *Thermotoga maritima* (tmCutA) was determined at 1.4 Å resolution using the Se-Met multiwavelength anomalous diffraction (MAD) technique. This protein (TIGR annotation - TM1056, DNA bases 1,069,580–1,069,885) is conserved in numerous bacteria, archaea and eucarya, including plants and mammals (COG1324, Fig. 1). The CutA *Escherichia coli* homolog—CutA1 (35% ID) is involved in divalent cation homeostasis,¹ while the mammalian homolog—mCutA (40% ID) was found to be associated with cell surface acetylcholinesterase.² However, the biological function of the CutA proteins is yet to be determined.

The tmCutA monomer assumes a two-layer α/β sandwich fold ($\beta 1\beta 2\alpha 1\beta 3\beta 4\alpha 3\beta 5\beta 6\alpha 5$) shared by a large number of proteins (CATH number 3.30.70.120). Two short ($\beta 2\beta 5$ and two long ($\beta 3\beta 4$) β -strands arranged in an antiparallel fashion $\beta 5\uparrow\beta 2\downarrow\beta 3\uparrow\beta 4\downarrow$ form a curved sheet. Three α -helices cover one side of the molecule (Fig. 2a). Each monomer is elongated with overall dimensions approximately $25 \times 30 \times 60$ Å (Fig. 2a). The monomers are assembled into a trimer with dimensions $55 \times 55 \times 35$ Å through interaction of the edges of three β -strands ($\beta 3$ on the top and $\beta 5\beta 6$ on the bottom) (Fig. 2b). In the trimer, the subunits are related by crystallographic three fold axes. The same oligomeric state was observed for tmCutA in solution by size exclusion experiments (see Methods) and appears to be functionally relevant. In the trimer, there is a cavity that is accessible from outside via three solvent accessible channels. The inner surface of the cavity is lined with several conserved residues including Cys35, which may serve as a potential metal binding site.

Structural homologues of tmCutA were identified using Dali search³ and SSAP structure comparison program 4. The closest structural matches, with a Dali score of 9.4 Z, were the *E.*

coli signal transducer proteins GlnB (PII)⁵ and GlnK⁶ (Fig. 2c). These two proteins are 67% identical to each other but have insignificant sequence similarity (less than 15% ID) to tmCutA. GlnB and GlnK, which are also trimeric, are involved in maintaining nitrogen homeostasis^{7–9} and have been shown to cooperatively bind ATP and α -ketoglutarate.^{6,10} The ATP binding sites of GlnB and GlnK are located in clefts between the monomers,¹¹ but most of the conserved amino acids that mediate ATP binding in PII proteins are not found in tmCutA. The cavity on the side of tmCutA is formed by a number of conserved aromatic (Tyr45, Trp47, Tyr81, and Trp101) and charged residues (Asp54, Glu56, Glu82) (Figs. 1 and 2c), which makes this cleft a strong candidate for a conserved function.

In PII-type proteins, the two central β -strands β 3 and β 4 are joined by a flexible loop called the T-loop, which is composed of 17 amino acids. The T-loop plays a key role in protein-protein interactions with downstream effector proteins.¹² The β 3 and β 4 strands in tmCutA, while longer than the corresponding strands in the PII proteins (12 residues in the β 3 strand of tmCutA compared with 9 in the β 2 strand of PII proteins), are joined by only a two-amino acid (Lys48Gly49) turn and form a hairpin. Although the residues in the C-terminal extension of β 3-strand in tmCutA proteins (Tyr45, Trp46, and Trp47, Fig. 1) are conserved among tmCutA homologs, the structure of this region in tmCutA is different than in PII proteins and thus this region of tmCutA may not participate in protein-protein interactions.

CutA is annotated as being involved in metal homeostasis. Several other structural homologs of tmCutA, including the PII proteins, metallochaperone Atx1 (5.9 Z score) and the metal-binding domain of the Menkes copper-transporting ATPase (4.5 Z score) have metal-binding properties.^{13,14} These proteins coordinate the metal through a CXXC sequence motif,^{15,16} but this motif is not found in CutA. Therefore, tmCutA may represent a new branch of this family that differs from PII and metal binding proteins. The biochemical experiments are under way to define possible ligands and functional partners of CutA proteins.

Methods

The tmCutA gene was subcloned, expressed and its product purified and screened for crystallization as described previously.¹⁷ Crystals for X-ray diffraction data collection were obtained from hanging drop vapor diffusion conditions containing 2 μ l of Se-Met derivative of the protein plus 2 μ l of 2 M ammonium sulphate, 0.1 M Tris HCl at pH 8.5 and 2% glycerol, over 2–5 days at 21°C. The crystals were flash frozen with crystallization buffer completed with 25% glycerol.

Diffraction data (Table I) were collected at beamline 19ID of the Advanced Photon Source, Argonne National Laboratory following the approach described earlier.¹⁸ The three-wavelength inverted beam MAD data up to 1.4 Å were collected from one Se-Met labeled protein crystal at 100 K with 3 seconds exposure/1°/frame using a 100 mm crystal-to-detector distance. The total oscillation range was 90°, as predicted using strategy module within HKL2000 suite.¹⁹ All data were processed and scaled with HKL2000 (Table I). The structure was determined by MAD phasing using CNS.²⁰ The initial model was built automatically using wARP,²¹ and refined to 1.4 Å using CNS against the peak data. The final R factor was 0.218, and the free R was 0.241 (Table II). Final refinement permitted us to define the coordinates of all 101 amino acids of tmCutA, seven (ALYFMGH) N-terminal amino acids belonging to His6-tag and two (GS) C-terminal amino acids, which resulted from the cloning artifact.

HPLC size-exclusion chromatography was performed on a Superdex-75 and Superdex-200 columns (10 \times 300 mm) pre-equilibrated with 10 mM HEPES pH 7.5, 0.5 M NaCl. The column was calibrated with cytochrome C (12.4 kDa), carbonic anhydrase (29 kDa), bovine serum albumin (66 kDa), alcohol dehydrogenase (150 kDa), β -amylase (200 kDa), and Blue Dextran

(2,000 kDa). A 25- μ L of tmCutA protein sample at a 2 mg ml⁻¹ concentration or premixed with standard proteins was centrifuged at 14,000 rpm for 10 min before being injected into the column through a 20 μ L injection loop. Filtration was carried out at 20°C at a flow rate of 1 ml min⁻¹. The eluted proteins were detected by measuring the absorbance at 280 nm.

Acknowledgments

Atomic coordinates have been deposited in the Protein Data Bank (PDB) with PDB-ID 1KR4 and accession number RCSB015255. We wish to thank all members of the Structural Biology Center at Argonne National Laboratory for their help in conducting these experiments and S. Beasley for help in preparation of this manuscript. CHA is a CIHR Investigator.


The submitted manuscript has been created by the University of Chicago as Operator of Argonne National Laboratory ("Argonne") under Contract No. W-31-109-ENG-38 with the U.S. Department of Energy. The U.S. Government retains for itself, and others acting on its behalf, a paid-up, nonexclusive, irrevocable worldwide license in said article to reproduce, prepare derivative works, distribute copies to the public, and perform publicly and display publicly, by or on behalf of the Government.

Grant sponsor: National Institutes of Health; Grant Number: GM62414; Grant sponsor: U.S. Department of Energy, Office of Biological and Environmental Research, under contract W-31-109-Eng-38; Grant sponsor: the Ontario Research and Development Challenge Fund; Grant sponsor: National Institutes of Health; Grant Number: GM26290.

REFERENCE

1. Fong ST, Camakaris J, Lee BT. Molecular genetics of a chromosomal locus involved in copper tolerance in *Escherichia coli* K-12. *Mol Microbiol* 1995;15:1127–1137. [PubMed: 7623666]
2. Perrier AL, Cousin X, Boschetti N, Haas R, Chatel JM, Bon S, Roberts WL, Pickett SR, Massoulie J, Rosenberry TL, Krejci E. Two distinct proteins are associated with tetrameric acetylcholinesterase on the cell surface. *J Biol Chem* 2000;275:34260–34265. [PubMed: 10954708]
3. Holm L, Sander C. Dali: a network tool for protein structure comparison. *Trends Biochem Sci* 1995;20:478–480. [PubMed: 8578593]
4. Orengo CA, Taylor WR. SSAP: sequential structure alignment program for protein structure comparison. *Methods Enzymol* 1996;266:617–635. [PubMed: 8743709]
5. Cheah E, Carr PD, Suffolk PM, Vasudevan SG, Dixon NE, Ollis DL. Structure of the *Escherichia coli* signal transducing protein PII. *Structure* 1994;2:981–990. [PubMed: 7866749]
6. Xu Y, Cheah E, Carr PD, van Heeswijk WC, Westerhoff HV, Vasudevan SG, Ollis DL. GlnK, a PII-homologue: structure reveals ATP binding site and indicates how the T-loops may be involved in molecular recognition. *J Mol Biol* 1998;282:149–165. [PubMed: 9733647]
7. Magasanik B. The regulation of nitrogen utilization in enteric bacteria. *J Cell Biochem* 1993;51:34–40. [PubMed: 8094391]
8. Ninfa AJ, Atkinson MR. PII signal transduction proteins. *Trends Microbiol* 2000;8:172–179. [PubMed: 10754576]
9. Arcondeguy T, Jack R, Merrick M. P(II) signal transduction proteins, pivotal players in microbial nitrogen control. *Microbiol Mol Biol Rev* 2001;65:80–105. [PubMed: 11238986]
10. Kamberov ES, Atkinson MR, Ninfa AJ. The *Escherichia coli* PII signal transduction protein is activated upon binding 2-ketoglutarate and ATP. *J Biol Chem* 1995;270:17797–17807. [PubMed: 7629080]
11. Xu Y, Carr PD, Huber T, Vasudevan SG, Ollis DL. The structure of the PII-ATP complex. *Eur J Biochem* 2001;268:2028–2037. [PubMed: 11277925]
12. Martinez-Argudo I, Contreras A. PII T-loop mutations affecting signal transduction to NtrB also abolish yeast two-hybrid interactions. *J Bacteriol* 2002;184:3746–3748. [PubMed: 12057972]
13. Gitschier J, Moffat B, Reilly D, Wood WI, Fairbrother WJ. Solution structure of the fourth metal-binding domain from the Menkes copper-transporting ATPase. *Nat Struct Biol* 1998;5:47–54. [PubMed: 9437429]

14. Rosenzweig AC, Huffman DL, Hou MY, Wernimont AK, Pufahl RA, O'Halloran TV. Crystal structure of the Atx1 metallochaperone protein at 1.02 Å resolution. *Structure Fold Des* 1999;7:605–617. [PubMed: 10404590]
15. DiDonato M, Narindrasorasak S, Forbes JR, Cox DW, Sarkar B. Expression, purification, and metal binding properties of the N-terminal domain from the wilson disease putative copper-transporting ATPase (ATP7B). *J Biol Chem* 1997;272:33279–33282. [PubMed: 9407118]
16. Lutsenko S, Petrukhin K, Cooper MJ, Gilliam CT, Kaplan JH. N-terminal domains of human copper-transporting adenosine triphosphatases (the Wilson's and Menkes disease proteins) bind copper selectively in vivo and in vitro with stoichiometry of one copper per metal-binding repeat. *J Biol Chem* 1997;272:18939–18944. [PubMed: 9228074]
17. Zhang RG, Skarina T, Katz JE, Beasley S, Khachatryan A, Vyas S, Arrowsmith CH, Clarke S, Edwards A, Joachimiak A, Savchenko A. Structure of *Thermotoga maritima* stationary phase survival protein SurE: a novel acid phosphatase. *Structure (Camb)* 2001;9:1095–1106. [PubMed: 11709173]
18. Walsh MA, Evans G, Sanishvili R, Dementieva I, Joachimiak A. MAD data collection — current trends. *Acta Crystallogr D Biol Crystallogr* 1999;55:1726–1732. [PubMed: 10531522]
19. Otwinowski Z, Minor W. Processing of X-ray diffraction data collected in oscillation mode. *Methods Enzymol* 1997;307–326.
20. Brunger AT, Adams PD, Clore GM, DeLano WL, Gros P, Grosse-Kunstleve RW, Jiang JS, Kuszewski J, Nilges M, Pannu NS, Read RJ, Rice LM, Simonson T, Warren GL. Crystallography & NMR system: a new software suite for macromolecular structure determination. *Acta Crystallogr D Biol Crystallogr* 1998;54:905–921. [PubMed: 9757107]
21. Perrakis A, Harkiolaki M, Wilson KS, Lamzin VS. ARP/wARP and molecular replacement. *Acta Crystallogr D Biol Crystallogr* 2001;57:1445–1450. [PubMed: 11567158]

**Fig. 1.**

Multiple sequence alignment of CutA from *Thermotoga maritima* (Thema) against other periplasmic divalent cation tolerance proteins from bacteria (Aquae = *Aquifex aeolicus*, Deira = *Deinococcus radiodurans*) and archaea (Pyrho = *Pyrococcus horikoshii*, Arcfu = *Archaeoglobus fulgidus*, Thevo = *Thermoplasma volcanium*), together with one from Mouse and a brain acetylcholinesterase putative membrane anchor from Human. Sequence conservation is color coded from red for invariant residues to orange to grey to none. The secondary structure of the *T. maritima* CutA is schematically shown in purple above the alignment.




Fig. 2. Ribbon representations of the tmCutA monomer (**a**), trimer (**b**) and superimposed structures of tmCutA and GlnK' complex with ATP (**c**). **a:** The α -helices and β -strains are colored in red and blue, respectively. **b:** Each monomer in the tmCutA trimer is differentiated by color. **c:** The tmCutA is shown on green, GlnK is shown in blue and ATP molecules are colored in red.

TABLE I

Summary of Crystal and MAD Data

| | | | |
|---------------------------|---|--------|---------|
| Unit cell | a = b = 52.237 Å, c = 33.846 Å $\alpha = \beta = 90^\circ$ $\gamma = 120^\circ$ | | |
| Space group | P3 | | |
| MW (Da) (108 residues) | 12420 | | |
| Mol (AU) | 1 | | |
| SeMet (AU) | 3 | | |
| | | | |
| | | | |
| MAD data collection | | | |
| | Edge | Peak | Remote |
| Wavelength (Å) | 0.9795 | 0.9793 | 0.95354 |
| Resolution range (Å) | 1.4 | 1.4 | 1.4 |
| No. of unique reflections | 19614 | 19651 | 19451 |
| Completeness (%) | 95.9 | 96.1 | 95.1 |
| R merge (%) | 6.7 | 7.1 | 5.6 |

TABLE II

Crystallographic Statistics

| Resolution range (Å) | Phasing | | | |
|---|---------|---------------|----------|---------------|
| | Centric | | Acentric | |
| | FOM | Phasing power | FOM | Phasing power |
| 30.0–1.4 | 0 | 0 | 0.5555 | 2.3552 |
| Density modification | | | | 0.80826 |
| | | | | 2.3552 |
| Refinement | | | | |
| Resolution range (Å) | | | | 40–1.4 |
| No. of reflections | | | | 19651 |
| σ cutoff | | | | 0.0 |
| R-value (%) | | | | 20.94 |
| Free R-value (%) | | | | 24.08 (1571) |
| Rms deviations from ideal geometry | | | | |
| bond length (1–2) (Å) | | | | 0.006 |
| angle (°) | | | | 1.2 |
| dihedral (°) | | | | 21.8 |
| improper (°) | | | | 0.66 |
| Number of atoms | | | | |
| protein | | | | 930 |
| water | | | | 124 |
| Mean B-factor (Å ²) for all atoms | | | | 15.3 |
| Ramachandran plot statistics (%) | | | | |
| Residues in most favored regions | | | | 99.2 |
| Residues in additional allowed regions | | | | 0.8 |
| Residues in disallowed region | | | | 0.0 |