

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/227612592>

# A superfamily of metalloenzymes unifies phosphopentomutase and cofactor-independent phosphoglycerate mutase with alkaline phosphatase and sulfatases

ARTICLE *in* PROTEIN SCIENCE · AUGUST 1998

Impact Factor: 2.85 · DOI: 10.1002/pro.5560070819 · Source: PubMed

---

CITATIONS

98

---

READS

25

## 3 AUTHORS:



**Michael Galperin**

National Institutes of Health

220 PUBLICATIONS 12,776 CITATIONS

SEE PROFILE



**Eugene V Koonin**

National Institutes of Health

579 PUBLICATIONS 46,711 CITATIONS

SEE PROFILE



**Amos Bairoch**

University of Geneva

244 PUBLICATIONS 44,630 CITATIONS

SEE PROFILE

## FOR THE RECORD

# A superfamily of metalloenzymes unifies phosphopentomutase and cofactor-independent phosphoglycerate mutase with alkaline phosphatases and sulfatases

MICHAEL Y. GALPERIN,<sup>1</sup> AMOS BAIROCH,<sup>2</sup> AND EUGENE V. KOONIN<sup>1</sup>

<sup>1</sup>National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Bethesda, Maryland 20894

<sup>2</sup>Département de Biochimie Médicale, Université de Genève, 1, rue Michel Servet CH-1211, Genève 4, Switzerland

(RECEIVED December 22, 1997; ACCEPTED May 4, 1998)

**Abstract:** Sequence analysis of the probable archaeal phosphoglycerate mutase resulted in the identification of a superfamily of metalloenzymes with similar metal-binding sites and predicted conserved structural fold. This superfamily unites alkaline phosphatase, N-acetylgalactosamine-4-sulfatase, and cerebroside sulfatase, enzymes with known three-dimensional structures, with phosphopentomutase, 2,3-bisphosphoglycerate-independent phosphoglycerate mutase, phosphoglycerol transferase, phosphonate monoesterase, streptomycin-6-phosphate phosphatase, alkaline phosphodiesterase/nucleotide pyrophosphatase PC-1, and several closely related sulfatases. In addition to the metal-binding motifs, all these enzymes contain a set of conserved amino acid residues that are likely to be required for the enzymatic activity. Mutational changes in the vicinity of these residues in several sulfatases cause mucopolysaccharidosis (Hunter, Maroteaux-Lamy, Morquio, and Sanfilippo syndromes) and metachromatic leucodystrophy.

**Keywords:** alkaline phosphatase; autotaxin; inherited disease; mucopolysaccharidosis; nucleotide pyrophosphatase PC-1; phosphoglycerate mutase; phosphopentomutase; sulfatase deficiency

Phosphoglycerate mutase (EC 5.4.2.1), a key glycolytic enzyme, is found in two forms, which differ in their requirement for 2,3-bisphosphoglycerate and show no detectable sequence similarity to one another (Grana et al., 1992, 1995). Although the 2,3-bisphosphoglycerate-dependent enzyme, found in bacterial, yeast, and animal cells, is relatively well studied (reviewed in Fothergill-Gilmore & Watson, 1989), the information about the structure or catalytic mechanism of the 2,3-bisphosphoglycerate-independent

form (iPGM) is limited (Singh & Setlow, 1979a; Blattler & Knowles, 1980; Huang & Dennis, 1995). iPGM has been found in bacteria (Singh & Setlow, 1978; Watabe & Freese, 1979), archaea (Yu et al., 1994), plants (Leadlay et al., 1977; Botha & Dennis, 1986), and in some invertebrates (Carreras et al., 1982). Several bacterial and plant iPGM genes have been sequenced (Grana et al., 1992, 1995; Leyva-Vazquez & Setlow, 1994; Morris et al., 1995). Sequence analysis of the maize iPGM showed significant similarity to alkaline phosphatase (AP), including conservation of several metal-binding residues of the AP active center, and a similar catalytic mechanism for the two enzymes, namely the formation of a phosphoserine intermediate stabilized by divalent cations, has been suggested (Grana et al., 1992). This conclusion, however, has been disputed (Huang et al., 1993) because (1) activity of iPGM from castor bean appeared to be metal independent (Botha & Dennis, 1986), and (2) sequence alignment of several plant iPGMs did not reveal the conserved Asp-Ser-Ala triad, like the one in the AP active center.

When the first complete genome of an archaeon, *Methanococcus jannaschii*, was sequenced, genes for all the enzymes of the lower (tri-carbon) portion of the glycolytic pathway, except PGM, were easily identified (Bult et al., 1996). The PGM-coding gene has been reported missing (Selkov et al., 1997), even though iPGM activity had been experimentally demonstrated in closely related *Methanococcus maripaludis* (Yu et al., 1994). Likewise, no PGM-encoding gene was recognized in the recently sequenced genomes of two other archaea, *Methanobacterium thermoautotrophicum* (Smith et al., 1997) and *Archaeoglobus fulgidus* (Klenk et al., 1997). On the other hand, when the set of *Methanococcus jannaschii* proteins was searched for the closest homolog of known bacterial and eukaryotic iPGMs, a candidate protein, MJ1612 (originally annotated as phosphonopyruvate decarboxylase), has been identified and predicted to possess iPGM activity (Koonin et al., 1997). However, MJ1612 appeared to be related also to phosphopentomutases (phosphodeoxyribomutases) and several other en-

Reprint requests to: Michael Y. Galperin, National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Bethesda, Maryland 20894; e-mail: galperin@ncbi.nlm.nih.gov.

zymes. This prompted us to investigate the possible relationships between iPGMs and phosphopentomutases.

Here, we show a pattern of sequence conservation between phosphopentomutases, iPGMs, and APs, which suggests conservation of the structural fold and similar reaction mechanisms. In accordance with the recent structural studies (Bond et al., 1997; Lukatela et al., 1998), similar conserved motifs were found in N-acetylgalactosamine-4-sulfatase, cerebroside sulfatase, and several related sulfatases. These findings define a new superfamily of proteins, which we refer to as the alkaline phosphatase superfamily.

The nonredundant protein sequence database at the National Center for Biotechnology Information (Bethesda, Maryland) was searched using the PSI-BLAST (Position-Specific Iterative BLAST) program, which converts local gapped alignment produced by BLASTP into position-specific weight matrices that are then used for iterative database scanning (Altschul et al., 1997). The multiple

alignment was constructed using the alignment (-m4) option of PSI-BLAST with subsequent manual refinement on the basis of the structural alignment of AP and sulfatases, which was generated using Dali (Holm & Sander, 1998).

Sequence analysis of MJ1612 showed high similarity to another *M. jannaschii* protein, MJ0010. Corresponding pairs of paralogs were found in genomes of two other archaea, *M. thermoautotrophicum* (Smith et al., 1997) and *A. fulgidus* (Klenk et al., 1997). Sequence database searches confirmed similarity of each of these proteins to a putative phosphonopyruvate decarboxylase from *Streptomyces hygroscopicus* (Lee et al., 1995); they also revealed a highly statistically significant ( $P < 10^{-8}$ ) similarity between all these proteins and iPGMs (Fig. 1). Iterative searches using the PSI-BLAST program resulted in identification of similar conserved regions in phosphopentomutases, APs, and related enzymes, and in several previously uncharacterized proteins (Fig. 1).



**Fig. 1.** Multiple alignment of the alkaline phosphatase superfamily. The proteins are listed under their unique SWISS-PROT (left column) and GenBank (right column) identifiers; 1160616, human autotaxin (Clair et al., 1997); 927036,  $\text{Ca}^{2+}$ -ATPase from *Flavobacterium odoratum* (Desrosiers et al., 1996; Peiffer et al., 1996); 1196755, phosphonoacetate hydrolase from *Pseudomonas fluorescens* (Kulakova et al., 1997); 1177864, phosphonate monoesterase from *Burkholderia caryophylli* (Dotson et al., 1996). The numbers indicate distances to the ends of each protein and the sizes of the gaps between aligned segments. Red and blue shading indicate conserved amino acid residues that are involved in metal binding in alkaline phosphatase (IALK) and sulfatases (IFSU and IAUk); their positions in mature enzymes are indicated above such residues. Conserved residues identified in this work are colored red and magenta. Black shading indicates the residues that were found mutated in patients with genetic disorders (intermediate or severe forms of hypophosphatasia, mucopolysaccharidosis, or metachromatic leucodystrophy). The references for particular mutations can be found in SWISS-PROT database (Bairoch & Apweiler, 1997). Yellow shading indicates uncharged amino acid residues (A, I, L, V, M, F, Y, or W) with a propensity to form a  $\beta$ -strand. Conserved small residues (G, A, or S) are shown in green, the residues conserved among several protein families are in bold. The consensus includes amino acid residues conserved in all sequences (upper case) and those conserved in the majority of the sequences (lower case). U stands for a bulky hydrophobic residue (I, L, V, M, F, Y, W), O stands for a small residue (G, A, S), - stands for D or E, \$ indicates any charged residue (D, E, K, R, N, Q), and dot stands for any residue. In the structure line,  $\alpha$  indicates  $\alpha$ -helix and  $\beta$  indicates  $\beta$ -strand.

In each case, PSI-BLAST searches using iPGMs or phosphopentomutases as the query produced highly significant sequence alignments ( $P < 10^{-6}$ ) with each other in the second or third iteration, and with APs and sulfatases in the fifth or sixth iteration. Inspection of the multiple alignment of all these proteins showed conservation of the core structural elements of APs and sulfatases (Fig. 1), suggesting that they belong to a distinct superfamily with a common structural fold.

This superfamily includes enzymes with substantially different activities (isomerases, hydrolases, and a putative lyase), which, however, all act on similar phosphocarbohydrate (or sulfocarbohydrate) substrates (Table 1). Remarkably, AP is known to have phosphotransferase activity (Coleman, 1992, and references therein), while iPGM can also function as a phosphatase (Breathnach & Knowles, 1977). Indeed, the conserved region in all these proteins (Fig. 1) contains the amino acid residues that are known to be involved in phosphate binding in AP (Kim & Wyckoff, 1991) and sulfate binding in sulfatases (Bond et al., 1997; Lukatela et al., 1998).

The alignment on Figure 1 demonstrates that all the amino acid residues that interact with Zn1 (Asp-327, His-331, and His-412) and Zn2 (Asp-51, Asp-369, and His-370) in AP (Kim & Wyckoff, 1991) are absolutely conserved in phosphocarbohydrate-binding proteins of the AP superfamily (Fig. 2). On the other hand, Mg binding residues of AP are much less conserved, as Glu-322 is substituted by Asn in phosphopentomutases and iPGMs, while Asp-153 and Thr-155 (Fig. 2) do not seem to be conserved at all. As noted earlier (Bond et al., 1997; Lukatela et al., 1998), the residues that coordinate Zn2 in AP are also conserved in sulfatases (Fig. 1).

The strong conservation of metal-binding residues in both phosphopentomutase and iPGM indicates that both these enzymes are metal dependent. Indeed, phosphopentomutase from *Escherichia coli* requires  $Mn^{2+}$ ,  $Ni^{2+}$ , or  $Co^{2+}$  for activity, binding two metal atoms per enzyme molecule (Hammer-Jespersen & Munch-Petersen, 1970; Hammer-Jespersen, 1983). Similar data were reported for the rat liver enzyme (Barsky & Hoffee, 1983).

**Table 1.** Properties of the enzymes of the alkaline phosphatase superfamily

Enzyme (EC No.)	SWISS-PROT symbol	pH optimum	Metal requirements	Human disease caused by a mutation	Refs.
Phosphoglycerate mutase (EC 5.4.2.1)	PMGL_ECOLI	7.7–9.5	Mn	— <sup>a</sup>	1, 2 <sup>b</sup>
Phosphopentomutase (EC 5.4.2.7)	DEOB_ECOLI	8.0–8.5	Mn or Co	Not known	3
Alkaline phosphatase (EC 3.1.3.1)	PPB_ECOLI PPBT_HUMAN	8.0–10.5	Zn + Mg	Hypophosphatasia	4, 5
Streptomycin-6-phosphatase (EC 3.1.3.39)	STRK_STRGR	n.d. <sup>c</sup>	n.d.	—	6
Alkaline phosphodiesterase/nucleotide pyrophosphatase (EC 3.1.4.1/3.6.1.9)	PC1_HUMAN	8.0–9.0	Mn, Mg, or Ca	Not known; increased activity in type II diabetes	7
Phosphoglycerol transferase (EC 2.7.8.20)	MDOB_ECOLI	8.9	Mn	—	8
Ca <sup>2+</sup> -ATPase	—	8.0	Ca	—	20
Phosphonopyruvate decarboxylase <sup>d</sup>	BCPC_STRHY	n.d.	Mg	—	9
Phosphonoacetate hydrolase (EC 3.11.1.1)	—	7.7–9.0	Zn, Mn, or Co	—	10
Phosphonate monoesterase	—	8.5–9.0	Mn	—	11
Arylsulfatase (EC 3.1.6.1)	ASLA_ECOLI ARSE_HUMAN	n.d.	n.d.	Chondrodysplasia punctata	12
Steroid sulfatase (EC 3.1.6.2)	STS_HUMAN	6.0–7.5	n.d.	X-linked ichthyosis	13
N-acetylgalactosamine 6-sulfatase (EC 3.1.6.4)	GA6S_HUMAN	3.5–4.0	n.d.	MPS <sup>e</sup> type IVA (Morquio A syndrome)	14
Cerebroside sulfatase (EC 3.1.6.8)	ARSA_HUMAN	4.8–6.2	Mg	Metachromatic leucodystrophy	14
N-acetylgalactosamine 4-sulfatase (EC 3.1.6.12)	ARSB_HUMAN	n.d.	Ca?	MPS type VI (Maroteaux-Lamy syndrome)	16
Iduronate 2-sulfatase (EC 3.1.6.13)	IDS_HUMAN	4.0–5.7	n.d.	MPS type II (Hunter syndrome)	17
N-acetylglucosamine 6-sulfatase (EC 3.1.6.14)	GL6S_HUMAN	3.9–5.7	n.d.	MPS type IIID (Sanfilippo D syndrome)	18
N-sulfoglucosamine sulfatase (EC 3.10.1.1)	SPHM_HUMAN	3.9–4.1	n.d.	MPS type IIIA (Sanfilippo A syndrome)	19

<sup>a</sup>Absence of 2,3-bisphosphoglycerate-dependent PGM in humans causes myopathies.


<sup>b</sup>References: 1, Singh and Setlow (1979a); 2, Carreras et al. (1982); 3, Hammer-Jespersen and Munch-Petersen (1970); 4, Henthorn et al. (1992); 5, Murphy et al. (1995); 6, Mansouri and Piepersberg (1991); 7, Oda et al. (1993); 8, Jackson and Kennedy (1983); 9, Nakashita et al. (1997); 10, McGrath et al. (1995); 11, Dotson et al. (1996); 12, Parenti et al. (1997); 13, Alperin and Shapiro (1997); 14, Bielicki and Hopwood (1991); 15, Lukatela et al. (1998); 16, Bond et al. (1997); 17, Bielicki et al. (1990); 18, Freeman and Hopwood (1987); 19, Freeman and Hopwood (1992); 20, Desrosiers et al. (1996).

<sup>c</sup>Not determined.

<sup>d</sup>Phosphonopyruvate decarboxylase activity of BCPC\_STRHY has not been demonstrated experimentally and could have been encoded by a different gene; actual function of this protein remains unidentified.

<sup>e</sup>MPS, Mucopolysaccharidosis.





**Fig. 2.** Conserved residues in the enzymes of the alkaline phosphatase superfamily. The scheme of the active center of the *E. coli* alkaline phosphatase (modified from Kim & Wyckoff, 1991); amino acid residues that are conserved in phosphopentomutases and phosphoglyceromutases (Fig. 1) are labeled in bold; nonconserved amino acid residues are labeled in italic, W indicates water molecules.

Bacterial iPGMs also require  $Mn^{2+}$  for activity (Singh & Setlow, 1978, 1979b; Watabe & Freese, 1979; Kuhn et al., 1993). Recently, a detailed study of  $Mn^{2+}$  binding by iPGM from *Bacillus megaterium* demonstrated a cooperativity in Mn-dependent activation of iPGM with a Hill coefficient of  $2.1 \pm 0.1$ , indicating that two Mn atoms bind per iPGM molecule (Kuhn et al., 1995). Thus, phosphopentomutase and bacterial iPGM each require two Mn atoms for activity.

The metal requirements of the plant iPGM have been a subject of some controversy. It was first reported that iPGM from wheat germ was inhibited by EDTA (Leadlay et al., 1977; Smith & Hass, 1985) and required  $Mn^{2+}$  or  $Co^{2+}$  for reactivation after denaturation (Smith et al., 1986). In contrast to these data, the activity of castor bean iPGM was reported unaffected by passing the enzyme solution through Chelex resin (Botha & Dennis, 1986). No data were presented, however, and the experimental protocol used has not been shown to completely remove trace metals from the reaction mixture. Based on the high level of sequence similarity between the plant and bacterial iPGMs (Grana et al., 1995; Fig. 1), it would be reasonable to suggest that plant enzymes are also metal dependent.

Sequence analysis shows that alkaline phosphodiesterase/nucleotide pyrophosphatase PC-1, a cell surface enzyme, implicated in pathogenesis of cancer and diabetes (Maddux et al., 1995), is also a member of the AP superfamily. Comparison of PC-1 with AP (not shown) shows that Thr-204 of PC1\_MOUSE aligns with the active site Ser-102 of AP (Fig. 2); it is similarly phosphorylated during the catalytic cycle of PC-1 (Belli et al., 1995). Autotaxin, a human tumor motility-stimulating protein, very similar to PC-1 (Clair et al., 1997; Fig. 1) has the same conserved region around Thr-208, indicating that it could also be phosphorylated. Divalent cations ( $Ca^{2+}$ ,  $Mg^{2+}$ , or  $Mn^{2+}$ ) are required for the activity of PC-1 (Oda et al., 1993); they also improve its thermal stability (Belli et al., 1994). Nucleotide pyrophosphatase activity has also been found in *Haemophilus influenzae* (Kahn & Anderson, 1986); it could belong to one of the previously uncharacterized *H. influenzae* proteins shown in Figure 1.  $Mn^{2+}$  was also shown to stim-

ulate the activity of three additional members of the AP superfamily (Fig. 1; Table 1), phosphoglycerol transferase (Jackson & Kennedy, 1983), phosphonoacetate hydrolase (McGrath et al., 1995), and phosphonate monoesterase from a glyphosate-degrading bacterium (Dotson et al., 1996). Another unusual member of the AP superfamily is the  $Ca^{2+}$ -dependent ATPase that requires two Ca atoms for activity (Desrosiers et al., 1996; Peiffer et al., 1996). Finally, an outer membrane protein YHBX\_ECOLI, associated with the adherence of enteropathogenic *E. coli* O157:H7 to human epithelial cells (Zhao et al., 1996), is also a member of this superfamily. The conservation of the predicted catalytic residues (Fig. 1) suggests that this protein possesses phosphatase activity that may be important for pathogenicity.

Several amino acid residues that form the active center of AP (Kim & Wyckoff, 1991) or sulfatases (Bond et al., 1997; Lukatela et al., 1998) are not conserved in phosphopentomutase and iPGM. The structure of the AP active center (Fig. 2) shows that one of these missing amino acid residues, Arg-166, binds the two remaining O atoms of the phosphate group (Kim & Wyckoff, 1991; Coleman, 1992) and thus assists in loosening the bond between the P atom and Zn1-bound O atom of the leaving  $RO^-$  group. Neither could we identify a counterpart of the phosphorylated Ser-102 of AP in phosphopentomutase or iPGM, even though iPGM has been suggested to form a phosphoenzyme intermediate (Blattler & Knowles, 1980).

The balance between phosphotransferase and phosphatase reaction may be affected by the difference in the metal specificity between AP, on one hand, and phosphopentomutase and iPGM, on the other hand. Even though AP is maximally active with  $Zn^{2+}$  ions, substitution of  $Mn^{2+}$  or  $Co^{2+}$  for  $Zn^{2+}$  still produced an enzyme with detectable activity. The decreased activity of such enzymes was largely due to the lower rate of the enzyme dephosphorylation, caused by a tighter binding of phosphate (Applebury et al., 1970; Coleman, 1992). Such an arrangement favors phosphotransferase reaction, which could be a reason for the  $Mn^{2+}$  dependence of phosphopentomutase and iPGM.

The alignment in Figure 1 also shows several highly conserved amino acid residues that have no known role in enzyme activity. These include Asp-346, Thr-367, Thr-413, Asp-437, and Thr-441 of AP and additional Thr residues in phosphopento- and phosphoglyceromutases, which have no counterparts in other enzymes. Examination of the three-dimensional structures of AP and sulfatases shows that in phosphopentomutases and iPGMs, only the equivalents of Thr-367, Thr-413, and Gln-410 of AP are likely to be positioned close enough to the phosphate-binding site to participate in binding of the carbohydrate moiety of the phosphocarbohydrate substrate (Fig. 1).

Asp-346 and Asp-437 of AP, located at ca. 20 Å from the metal-binding site in both AP and sulfatases, could be involved in the maintenance of the structural integrity of these enzymes. It is also possible that they participate in a relay that directs substrates to the active sites of these enzymes. In any case, the absence of activity in D255H and D335V mutants of human cerebroside sulfatase (Hess et al., 1996; Lissens et al., 1996) demonstrates that these conserved Asp residues are required for sulfatase activity, and suggests that they might be important in other enzymes as well. Analysis of genetic disorders that result from mutations in human genes coding for known enzymes may offer additional insight into the organization of their molecules. Several inherited disorders, such as hypophosphatasia, chondrodysplasia, metachromatic leukodystrophy, and various mucopolysaccharidoses are caused by

**Table 2.** Phylogenetic distribution of the enzymes of the alkaline phosphatase superfamily

Enzyme	Phylogenetic distribution									
	Bacteria					Archaea				
	Ec <sup>a</sup>	Hi	Hp	Sy	Bs	Mg	Mj	Mt	Af	Sc
Cofactor-independent phosphoglycerate mutase <sup>b,c</sup>	yibO	— <sup>d</sup>	HP0974	slr1945	<b>pgm</b>	MG430	MJ1612	MTH1591	AF1751	—
Phosphopentomutase <sup>b</sup>	<b>deoB</b> yhfW	—	HP1179	—	drm	—	MJ0010	MTH418	AF1425	—
Alkaline phosphatase	<b>phoA</b>	—	—	—	<b>phoA</b> <b>phoB</b>	—	—	—	—	<b>YDR481c</b>
Alkaline phosphodiesterase/ nucleotide pyrophosphatase	—	HI0841	—	—	—	—	—	—	—	—
Phosphoglycerol transferase	<b>mdoB</b>	HI1246	HP0578	—	yqgS	—	—	—	—	—
Sulfatases	aslA yidJ bl498	—	—	—	—	—	—	—	—	—
Uncharacterized	yhbX ybiP yhiW yidB yefM yijP yigX	HI0275 HI1005 HI1064	HP0022	—	yfiE yfiI yvgJ	—	—	—	AF0247	YJL062w YKL165c YLL031c
										1256283 1017806 1125842
										10 or more proteins
										10 or more enzymes

<sup>a</sup> Ec, *Escherichia coli*; Hi, *Haemophilus influenzae*; Hp, *Helicobacter pylori*; Sy, *Synechocystis* sp.; Bs, *Bacillus subtilis*; Mg, *Mycobacterium genitalium*; Mj, *Methanococcus jannaschii*; Af, *Archaeoglobus fulgidus*; Sc, *Saccharomyces cerevisiae*; Ce, *Caenorhabditis elegans*; Hs, *Homo sapiens*. The proteins are listed under their original authors' gene designations or their GenBank identifiers. Human phosphopentomutase fragment is from the expressed sequence tags (EST) database.

<sup>b</sup> The proteins with experimentally demonstrated enzymatic activity are shown in bold type; other proteins are predicted to possess indicated activities on the basis of sequence similarity to previously characterized proteins (Fig. 1).

<sup>c</sup> Ec, Hi, Sy, Bs, Sc, and Hs have genes coding for 2,3-bisphosphate-dependent phosphoglycerate mutase.

<sup>d</sup> No corresponding gene is present in the complete genome.

missense mutations in the genes for sulfatases that belong to the AP superfamily. Some of these mutations result in single amino acid substitutions in the conserved motifs shown on Figure 1 and abolish the enzymatic activity (reviewed in Henthorn et al., 1992; Parenti et al., 1997).

The phylogenetic distribution of the AP superfamily enzymes is unusual. While some bacteria, such as *E. coli* or *Bacillus subtilis*, encode both iPGMs and APs, archaea and eukaryotes usually have only one of these enzymes (Table 2). Thus, iPGM activity has not been found in vertebrates (Carreras et al., 1982); the respective gene is also absent from the yeast genome. Instead, fungi and vertebrates have a different, 2,3-bisphosphoglycerate-dependent form of phosphoglycerate mutase (Fothergill-Gilmore & Watson, 1989). In algae, iPGM is encoded in the chloroplast, while in higher plants it is nuclear encoded and absent from the chloroplast genome. Plant iPGMs thus appear to have chloroplast origin. On the other hand, AP, found in yeast and animal cells, so far has not been described in plants. Conceivably, AP and iPGM could have evolved from a common ancestral enzyme, with selective loss of one of these enzymes in various eukaryotic branches. The pairs of paralogous archaeal proteins (e.g., MJ0010 and MJ1612) have shown significant sequence similarity to all the enzymes of the AP superfamily and may resemble the ancestral phosphomutases.

The distribution of the two classes of PGMs in bacteria is also remarkable. While organisms with larger genomes, such as *E. coli*, *B. subtilis*, and *Synechocystis* sp. have genes coding for both classes of this enzyme (Table 2), the organisms with smaller genomes code for only one of them. Thus, iPGM is the only form of this enzyme that is encoded in the genomes of such human pathogens as *Mycoplasma genitalium*, *Mycoplasma pneumoniae*, and *Helicobacter pylori*, the causative agents of nongonococcal male urethritis, atypical pneumonia, and gastric ulcer, respectively. The importance of iPGM for the metabolism of these bacteria and its apparent absence in vertebrates (Carreras et al., 1982) suggest that iPGM may be a plausible target for new, specialized antibacterial drugs. The 3D structure of iPGM, once determined, will facilitate the development of such drugs and will help resolve the remaining questions about its catalytic mechanism.

## References

- Alperin ES, Shapiro LJ. 1997. Characterization of point mutations in patients with X-linked ichthyosis. Effects on the structure and function of the steroid sulfatase protein. *J Biol Chem* 272:20756–20763.
- Altschul SF, Madden TL, Schaffer AA, Zhang J, Zheng Z, Miller W, Lipman DJ. 1997. Gapped BLAST and PSI-BLAST—A new generation of protein database search programs. *Nucleic Acids Res* 25:3389–3402.
- Applebury ML, Johnson BP, Coleman JE. 1970. Phosphate binding to alkaline phosphatase. Metal ion dependence. *J Biol Chem* 245:4968–4976.
- Bairoch A, Apweiler R. 1997. The SWISS-PROT protein sequence data bank and its supplement TrEMBL. *Nucleic Acids Res* 25:31–36.
- Barsky DL, Hoffee PA. 1983. Purification and characterization of phosphoenolpyruvate carboxylase from rat liver. *Biochim Biophys Acta* 743:162–171.
- Belli SI, Mercuri FA, Sali A, Goding JW. 1995. Autophosphorylation of PC-1 (alkaline phosphodiesterase 1/nucleotide pyrophosphatase) and analysis of the active site. *Eur J Biochem* 228:669–676.
- Belli SI, Sali A, Goding JW. 1994. Divalent cations stabilize the conformation of plasma cell membrane glycoprotein PC-1 (alkaline phosphodiesterase 1). *Biochem J* 304:75–80.
- Bielicki J, Freeman C, Clements PR, Hopwood JJ. 1990. Human liver iduronate-2-sulfatase. Purification, characterization and catalytic properties. *Biochem J* 271:75–86.
- Bielicki J, Hopwood JJ. 1991. Human liver N-acetylgalactosamine 6-sulphatase. Purification and characterization. *Biochem J* 279:515–520.
- Blattler WA, Knowles JR. 1980. Phosphoglycerate mutases: Stereochemical course of the phosphoryl group transfers catalyzed by the cofactor-dependent enzyme from rabbit muscle and the cofactor-independent enzyme from wheat germ. *Biochemistry* 19:738–743.
- Bond CS, Clements PR, Ashby SJ, Collyer CA, Harrop SJ, Hopwood JJ, Guss JM. 1997. Structure of a human lysosomal sulfatase. *Structure* 5:277–289.
- Botha FC, Dennis DT. 1986. Isozymes of phosphoglyceromutase from the developing endosperm of *Ricinus communis*: Isolation and kinetic properties. *Arch Biochem Biophys* 245:96–103.
- Breathnach R, Knowles JR. 1977. Phosphoglycerate mutase from wheat germ: Studies with <sup>18</sup>O-labeled substrate, investigations of the phosphatase and phosphoryl transfer activities, and evidence for a phosphoryl-enzyme intermediate. *Biochemistry* 16:3054–3060.
- Bult CJ, White O, Olsen GJ, Zhou L, Fleischmann RD, Sutton GG, Blake JA, FitzGerald LM, Clayton RA, Gocayne JD, Kerlavage AR, Dougherty BA, Tomb J-F, Adams MD, Reich CI, Overbeek R, Kirkness EF, Weinstock KG, Merrick JM, Glodek A, Scott JL, Geoghegan NSM, Weidman JF, Fuhrmann JL, Nguyen D, Utterback TR, Kelley JM, Peterson JD, Sadow PW, Hanna MC, Cotton MD, Roberts KM, Hurst MA, Kaine BP, Borodovsky M, Klenk H-P, Fraser CM, Smith HO, Woese CR, Venter JC. 1996. Complete genome sequence of the methanogenic archaeon, *Methanococcus jannaschii*. *Science* 273:1058–1073.
- Carreras J, Mezquita J, Bosch J, Bartrons R, Pons G. 1982. Phylogeny and ontogeny of the phosphoglycerate mutases—IV. Distribution of glycerate-2,3-P2 dependent and independent phosphoglycerate mutases in algae, fungi, plants and animals. *Comp Biochem Physiol [B]* 71:591–597.
- Clair T, Lee HY, Liotta LA, Stracke ML. 1997. Autotaxin is an exoenzyme possessing 5'-nucleotide phosphodiesterase/ATP pyrophosphatase and ATPase activities. *J Biol Chem* 272:996–1001.
- Coleman JE. 1992. Structure and mechanism of alkaline phosphatase. *Annu Rev Biophys* 21:441–483.
- Desrosiers MG, Gately LJ, Gambel AM, Menick DR. 1996. Purification and characterization of Ca<sup>2+</sup>-ATPase of *Flavobacterium odoratum*. *J Biol Chem* 271:3945–3951.
- Dotson SB, Smith CE, Ling CS, Barry GF, Kishore GM. 1996. Identification, characterization, and cloning of a phosphonate monoester hydrolase from *Burkholderia caryophylli* PG2982. *J Biol Chem* 271:25754–25761.
- Fothergill-Gilmore LA, Watson HC. 1989. The phosphoglycerate mutases. *Adv Enzymol* 62:227–313.
- Freeman C, Hopwood JJ. 1987. Human liver N-acetylglucosamine-6-sulphate sulphatase. Catalytic properties. *Biochem J* 246:355–365.
- Freeman C, Hopwood JJ. 1992. Human glucosamine-6-sulphatase deficiency. Diagnostic enzymology towards heparin-derived trisaccharide substrates. *Biochem J* 282:605–614.
- Grana X, de Lecea L, el-Maghrabi MR, Urena JM, Caellas C, Carreras J, Puigdomenech P, Pilakis SJ, Climent F. 1992. Cloning and sequencing of a cDNA encoding 2,3-bisphosphoglycerate-independent phosphoglycerate mutase from maize. Possible relationship to the alkaline phosphatase family. *J Biol Chem* 267:12797–12803.
- Grana X, Perez de la Ossa P, Broceno C, Stocker M, Garriga J, Puigdomenech P, Climent F. 1995. 2,3-Bisphosphoglycerate-independent phosphoglycerate mutase is conserved among different phylogenetic kingdoms. *Comp Biochem Physiol B Biochem Mol Biol* 112:287–293.
- Hammer-Jespersen K. 1983. Nucleoside catabolism. In: Munch-Petersen A, ed. *Metabolism of nucleotides, nucleosides and nucleobases in microorganisms*. London: Academic Press. pp 203–258.
- Hammer-Jespersen K, Munch-Petersen A. 1970. Phosphodeoxyribomutase from *Escherichia coli*. Purification and some properties. *Eur J Biochem* 17:397–407.
- Henthorn PS, Raducha M, Fedde KN, Lafferty MA, Whyte MP. 1992. Different missense mutations at the tissue-nonspecific alkaline phosphatase gene locus in autosomal recessively inherited forms of mild and severe hypophosphatasia. *Proc Natl Acad Sci USA* 89:9924–9928.
- Hess B, Kafert S, Heinisch U, Wenger DA, Zlotogora J, Gieselmann V. 1996. Characterization of two arylsulfatase A missense mutations D335V and T274M causing late infantile metachromatic leukodystrophy. *Hum Mutat* 7:311–317.
- Holm L, Sander C. 1998. Touring protein fold space with Dali/FSSP. *Nucleic Acids Res* 26:318–321.
- Huang Y, Blakeley SD, McAleese SM, Fothergill-Gilmore LA, Dennis DT. 1993. Higher-plant cofactor-independent phosphoglyceromutase: Purification, molecular characterization and expression. *Plant Mol Biol* 23:1039–1053.
- Huang Y, Dennis DT. 1995. Histidine residues 139, 363 and 500 are essential for catalytic activity of cofactor-independent phosphoglyceromutase from developing endosperm of the castor plant. *Eur J Biochem* 229:395–402.
- Jackson BJ, Kennedy EP. 1983. The biosynthesis of membrane-derived oligosaccharides. A membrane-bound phosphoglycerol transferase. *J Biol Chem* 258:2394–2398.

- Kahn DW, Anderson BM. 1986. Characterization of *Haemophilus influenzae* nucleotide pyrophosphatase. An enzyme of critical importance for growth of the organism. *J Biol Chem* 261:6016–6025.
- Kim EE, Wyckoff HW. 1991. Reaction mechanism of alkaline phosphatase based on crystal structures. Two-metal ion catalysis. *J Mol Biol* 218:449–464.
- Klenk HP, Clayton RA, Tomb J-F, White O, Nelson KE, Ketchum KA, Dodson RJ, Gwinn M, Hickey EK, Peterson JD, Richardson DL, Kerlavage AR, Graham DE, Kyrpides NC, Fleischmann RD, Quackenbush J, Lee NH, Sutton GG, Gill S, Kirkness EF, Dougherty BA, McKenney K, Adams MD, Loftus B, Peterson JD, Reich CI, McNeil LK, Badger JH, Glodek A, Zhou L, Overbeek R, Gocayne JD, Weidman JF, McDonald L, Utterback TR, Cotton MD, Spriggs T, Artach P, Kaine BP, Sykes SM, Sadow PW, D'Andrea KP, Bowman C, Fujii C, Garland SA, Mason TM, Olsen GJ, Fraser CM, Smith HO, Woese CR, Venter JC. 1997. The complete genome sequence of the hyperthermophilic, sulphate-reducing archaeon *Archaeoglobus fulgidus*. *Nature* 390:364–370.
- Koonin EV, Mushegian AR, Galperin MY, Walker DR. 1997. Comparison of archaeal and bacterial genomes: Computer analysis of protein sequences predicts novel functions and suggests a chimeric origin for the archaea. *Mol Microbiol* 25:619–637.
- Kuhn NJ, Setlow B, Setlow P. 1993. Manganese(II) activation of 3-phosphoglycerate mutase of *Bacillus megaterium*: pH-sensitive interconversion of active and inactive forms. *Arch Biochem Biophys* 306:342–349.
- Kuhn NJ, Setlow B, Setlow P, Cammack R, Williams R. 1995. Cooperative manganese (II) activation of 3-phosphoglycerate mutase of *Bacillus megaterium*: A biological pH-sensing mechanism in bacterial spore formation and germination. *Arch Biochem Biophys* 320:35–42.
- Kulakova AN, Kulakov LA, Quinn JP. 1997. Cloning of the phosphonoacetate hydrolase gene from *Pseudomonas fluorescens* 23F encoding a new type of carbon-phosphorus bond cleaving enzyme and its expression in *Escherichia coli* and *Pseudomonas putida*. *Gene* 195:49–53.
- Leadlay PF, Breathnach R, Gatehouse JA, Johnson PE, Knowles JR. 1977. Appendix: Phosphoglycerate mutase from wheat germ: Isolation, crystallization, and properties. *Biochemistry* 16:3045–3053.
- Lee SH, Hidaka T, Nakashita H, Seto H. 1995. The carboxyphosphonoenolpyruvate synthase-encoding gene from the bialaphos-producing organism *Streptomyces hygroscopicus*. *Gene* 153:143–144.
- Leyva-Vazquez MA, Setlow P. 1994. Cloning and nucleotide sequences of the genes encoding triose phosphate isomerase, phosphoglycerate mutase, and enolase from *Bacillus subtilis*. *J Bacteriol* 176:3903–3910.
- Lissens W, Vervoort R, Van Regemorter N, Van Bogaert P, Freund M, Verellen-Dumoulin C, Seneca S, Liebaers I. 1996. A D255H substitution in the arylsulphatase A gene of two unrelated Belgian patients with late-infantile metachromatic leukodystrophy. *J Inher Metab Dis* 19:782–786.
- Lukatela G, Krauss N, Theis K, Selmer T, Gieselmann V, von Figura K, Saenger W. 1998. Crystal structure of human arylsulphatase A: The aldehyde function and the metal ion at the active site suggest a novel mechanism for sulfate ester hydrolysis. *Biochemistry* 37:3654–3664.
- Maddux BA, Sbraccia P, Kumakura S, Sasson S, Youngren J, Fisher A, Spencer S, Grupe A, Henzel W, Stewart TA, et al. 1995. Membrane glycoprotein PC-1 and insulin resistance in non-insulin-dependent diabetes mellitus. *Nature* 373:448–451.
- Mansouri K, Piepersberg W. 1991. Genetics of streptomycin production in *Streptomyces griseus*: Nucleotide sequence of five genes, strFGHIK, including a phosphatase gene. *Mol Gen Genet* 228:459–469.
- McGrath JW, Wisdom GB, McMullan G, Larkin MJ, Quinn JP. 1995. The purification and properties of phosphonoacetate hydrolase, a novel carbon-phosphorus bond-cleavage enzyme from *Pseudomonas fluorescens* 23F. *Eur J Biochem* 234:225–230.
- Morris VL, Jackson DP, Grattan M, Ainsworth T, Cuppels DA. 1995. Isolation and sequence analysis of the *Pseudomonas syringae* pv. *tomato* gene encoding a 2,3-diphosphoglycerate-independent phosphoglyceromutase. *J Bacteriol* 177:1727–1733.
- Murphy JE, Tibbitts TT, Kantrowitz ER. 1995. Mutations at positions 153 and 328 in *Escherichia coli* alkaline phosphatase provide insight towards the structure and function of mammalian and yeast alkaline phosphatases. *J Mol Biol* 253:604–617.
- Nakashita H, Watanabe K, Hara O, Hidaka T, Seto H. 1997. Studies on the biosynthesis of bialaphos. Biochemical mechanism of C–P bond formation: Discovery of phosphonopyruvate decarboxylase which catalyzes the formation of phosphonoacetaldehyde from phosphonopyruvate. *J Antibiot* 50:212–219.
- Oda Y, Kuo MD, Huang SS, Huang JS. 1993. The major acidic fibroblast growth factor (aFGF)-stimulated phosphoprotein from bovine liver plasma membranes has aFGF-stimulated kinase, autoadenylation, and alkaline nucleotide phosphodiesterase activities. *J Biol Chem* 268:27318–27326.
- Parenti G, Meroni G, Ballabio A. 1997. The sulfatase gene family. *Curr Opin Genet Dev* 7:386–391.
- Peiffer WE, Desrosiers MG, Menick DR. 1996. Cloning and expression of the unique  $\text{Ca}^{2+}$ -ATPase from *Flavobacterium odoratum*. *J Biol Chem* 271:5095–5100.
- Selkov E, Maltsev N, Olsen GJ, Overbeek R, Whitman WB. 1997. A reconstruction of the metabolism of *Methanococcus jannaschii* from sequence data. *Gene* 197:GC11–GC26.
- Singh RP, Setlow P. 1978. Phosphoglycerate mutase in developing forespores of *Bacillus megaterium* may be regulated by the intrasporal level of free manganous ion. *Biochem Biophys Res Commun* 82:1–5.
- Singh RP, Setlow P. 1979a. Purification and properties of phosphoglycerate phosphomutase from spores and cells of *Bacillus megaterium*. *J Bacteriol* 137:1024–1027.
- Singh RP, Setlow P. 1979b. Regulation of phosphoglycerate phosphomutase in developing forespores and dormant and germinated spores of *Bacillus megaterium* by the level of free manganous ions. *J Bacteriol* 139:889–898.
- Smith DR, Doucette-Stamm LA, Deloughery C, Lee H, Dubois J, Aldredge T, Bashirzadeh R, Blakely D, Cook R, Gilbert K, Harrison D, Hoang L, Keagle P, Lumm W, Pothier B, Qiu D, Spadafora R, Vicaire R, Wang Y, Wierzbowski J, Gibson R, Jiwani N, Caruso A, Bush D, Safer H, Patwell D, Prabhakar S, McDougall S, Shimer G, Goyal A, Pietrovsky S, Church GM, Daniels CJ, Mao J-I, Rice P, Nolling J, Reeve JN. 1997. Complete genome sequence of *Methanobacterium thermoautotrophicum* deltaH: Functional analysis and comparative genomics. *J Bacteriol* 179:7135–7155.
- Smith GC, Hass LF. 1985. Wheat germ phosphoglycerate mutase: Purification, polymorphism, and inhibition. *Biochem Biophys Res Commun* 131:743–749.
- Smith GC, McWilliams AD, Hass LF. 1986. Wheat germ phosphoglycerate mutase: Evidence for a metalloenzyme. *Biochem Biophys Res Commun* 136:336–340.
- Watabe K, Freese E. 1979. Purification and properties of the manganese-dependent phosphoglycerate mutase of *Bacillus subtilis*. *J Bacteriol* 137:773–778.
- Yu JP, Ladapo J, Whitman WB. 1994. Pathway of glycogen metabolism in *Methanococcus maripaludis*. *J Bacteriol* 176:325–332.
- Zhao S, Meng J, Doyle MP, Meinersman R, Wang G, Zhao P. 1996. A low molecular weight outer-membrane protein of *Escherichia coli* O157:H7 associated with adherence to INT407 cells and chicken caeca. *J Med Microbiol* 45:90–96.