

Confinement, Quartet Mechanism and Cluster Property.

L. BONORA, P. PASTI and M. TONIN

Istituto di Fisica dell'Università - Padova

Istituto Nazionale di Fisica Nucleare - Sezione di Padova

(ricevuto il 9 Febbraio 1981)

Confinement implies a violation of the cluster property ⁽¹⁾. STROCCHI ⁽²⁾ has shown that non-Abelian gauge QFT are the only theories, among renormalizable QFT, in which a violation of the cluster property can arise. The violation of the cluster property typically arises in a theory where some two-point Green's functions in the momentum space has a double-pole singularity at $q^2 = 0$ (massless dipole propagators) ^(*).

From another point of view KUGO ⁽⁴⁾ proposed a confinement criterion based on the so-called quartet mechanism.

One can ask whether the two approaches are related, *i.e.* whether the Kugo mechanism leads to a violation of the cluster property.

Here we shall give a heuristic argument supporting an affirmative answer. This argument is suggested by the superfield formulation of the gauge theories proposed in ref. ⁽⁵⁾.

Let us consider the $(4+2)$ -dimensional superspace M obtained adding two Grassmann variables θ and $\bar{\theta}$ to the Minkowski space. In M we define the one-form \hat{a} :

$$(1) \quad \hat{a} = \hat{U}^+(A_\mu dx^\mu) \hat{U} + \hat{U}^+ d\hat{U},$$

where

$$(2) \quad \hat{U}(x, \theta, \bar{\theta}) = \exp [\theta \bar{c}(x) + \bar{\theta} c(x) + \theta \bar{\theta} (B(x) + \frac{1}{2} (c\bar{c} + \bar{c}c))]]$$

and

$$A_\mu(x) = A_\mu^\alpha(x) \tau^\alpha, \quad c = c^\alpha \tau^\alpha, \quad \bar{c} = \bar{c}^\alpha \tau^\alpha, \quad B = B^\alpha \tau^\alpha;$$

A_μ^α, B^α are real, θ, c^α are Hermitian and $\bar{\theta}, \bar{c}^\alpha$ are anti-Hermitian.

⁽¹⁾ H. ARAKI, K. HEPP and D. RUELE: *Helv. Phys. Acta*, **35**, 164 (1962).

⁽²⁾ F. STROCCHI: *Phys. Lett. B*, **62**, 60 (1976); *Phys. Rev. D*, **17**, 2010 (1978).

^(*) A confined model with dipole gluon fields has been studied in ref. ⁽³⁾.

⁽³⁾ E. D'EMILIO and M. MINTCHEV: Pisa IFUP preprint (1980).

⁽⁴⁾ T. KUGO: *Phys. Lett. B*, **83**, 93 (1979).

⁽⁵⁾ L. BONORA and M. TONIN: *Phys. Lett. B*, **98**, 48 (1981).

τ^α are the anti-Hermitian representative matrices which generate the Lie algebra of the gauge group G . $A_\mu^\alpha(x)$ are the gauge fields, $c^\alpha(x)$ and $\bar{c}^\alpha(x)$ the anticommuting FP fields and $B^\alpha(x)$ the auxiliary fields.

From eq. (1) one can write

$$(3) \quad \hat{\alpha} = \Phi_\mu(x, \theta, \bar{\theta}) dx^\mu + \eta(x, \theta, \bar{\theta}) d\bar{\theta} + \bar{\eta}(x, \theta, \bar{\theta}) d\theta,$$

where the superfields Φ_μ^α , η^α , $\bar{\eta}^\alpha$ are given by

$$(4) \quad \begin{cases} \Phi_\mu^\alpha(x, \theta, \bar{\theta}) = A_\mu^\alpha(x) + \theta D_\mu \bar{c}^\alpha(x) + \bar{\theta} D_\mu c^\alpha(x) + \theta \bar{\theta} (D_\mu B(x) + D_\mu c(x) \times \bar{c}(x))^\alpha, \\ \eta^\alpha(x, \theta, \bar{\theta}) = c^\alpha(x) + \theta \bar{B}^\alpha(x) - \frac{\bar{\theta}}{2} (c(x) \times c(x))^\alpha + \theta \bar{\theta} (\bar{B}(x) \times c(x))^\alpha, \\ \bar{\eta}^\alpha(x, \theta, \bar{\theta}) = \bar{c}^\alpha(x) - \frac{\theta}{2} (\bar{c}(x) \times \bar{c}(x))^\alpha + \bar{\theta} B^\alpha(x) + \theta \bar{\theta} (c(x) \times B(x))^\alpha \end{cases}$$

with

$$B^\alpha(x) + \bar{B}^\alpha(x) + (c(x) \times \bar{c}(x))^\alpha = 0.$$

The curvature two-form $\hat{\varrho}$

$$\hat{\varrho} = d\hat{\alpha} + \frac{1}{2} [\hat{\alpha}, \hat{\alpha}]$$

is

$$\hat{\varrho} = \hat{U}^+ (\mathbf{F}_{\mu\nu} dx^\mu \wedge dx^\nu) \hat{U} = \hat{\mathbf{F}}_{\mu\nu}(x, \theta, \bar{\theta}) dx^\mu \wedge dx^\nu,$$

where

$$\mathbf{F}_{\mu\nu} = \partial_\mu \mathbf{A}_\nu - \partial_\nu \mathbf{A}_\mu + [\mathbf{A}_\mu, \mathbf{A}_\nu].$$

In this notation the Yang-Mills Lagrangian density equipped with a gauge-fixing term and the corresponding FP term is

$$(5) \quad \mathcal{L} = \text{Tr} \left\{ -\frac{1}{4g^2} (\hat{\mathbf{F}}_{\mu\nu} \hat{\mathbf{F}}^{\mu\nu}) + \frac{d}{d\theta} \frac{d}{d\bar{\theta}} (\Phi^\mu \Phi_\mu) + \xi \left(\frac{d\eta}{d\theta} \right)^2 \right\}.$$

This Lagrangian density is manifestly invariant under translations in $\bar{\theta}$ and θ . These nilpotent translations are just the BRS ⁽⁶⁾ and the anti-BRS ⁽⁷⁻⁹⁾ transformations, generated by the nilpotent charges Q_B and \bar{Q}_B .

The Fock space \mathcal{V} of the QFT defined by the Lagrangian density (5) has indefinite metric; the S -matrix invariant, semi-definite positive, physical subspace \mathcal{V}_{ph} is defined

⁽⁶⁾ C. BECCHI, A. ROUET and R. STORA: *Commun. Math. Phys.*, **42**, 127 (1975); *Ann. Phys. (N. Y.)*, **98**, 287 (1976).

⁽⁷⁾ G. CURCI and R. FERRARI: *Nuovo Cimento A*, **32**, 151 (1976); *Phys. Lett. B*, **63**, 91 (1976); *Nuovo Cimento A*, **35**, 273 (1976).

⁽⁸⁾ M. QUIROS and F. J. DE URRIES: preprint IEM (1980), Madrid; M. QUIROS, F. J. DE URRIES, J. HOYOS, M. L. MAZON and E. RODRIGUEZ: preprint IEM (1980), Madrid.

⁽⁹⁾ I. OJIMA: *Prog. Theor. Phys.*, **64**, 625 (1980).

by the condition ^(5,9)

$$(6) \quad \mathcal{V}_{\text{ph}} \equiv \{|\alpha\rangle \in \mathcal{V}; Q_{\text{B}}|\alpha\rangle = 0; \bar{Q}_{\text{B}}|\alpha\rangle = 0\}$$

a strong version of the physical condition of ref. ^(7,10). Since \mathcal{V} is nondegenerate, to each $|\alpha, k\rangle \in \mathcal{V}$ with ghost number k , will correspond a *conjugate* state $|\alpha, -k\rangle$ with opposite ghost number such that

$$\langle \alpha, -k | \alpha, k \rangle \neq 0.$$

Let us recall briefly the quartet confinement mechanism purposed by KUGO ⁽⁴⁾.

If $\varrho(x)$ is the interpolating field of a particle of given mass and spin and $\varrho^{\text{as}}(x)$ is the corresponding asymptotic field, we shall consider the BRS transform of $\varrho(x)$

$$(7) \quad [iQ_{\text{B}}, \varrho(x)]_{\pm} = \sigma(x).$$

Let us suppose that, in the channel of the (composite) operator $\sigma(x)$, a (bound) one-particle state is present with the same mass and spin of $\varrho(x)$. We shall call $\varrho^{\text{as}}(x)$ the asymptotic field of this (bound) state. Then $\varrho^{\text{as}}(x)$, $\sigma^{\text{as}}(x)$, together with their conjugate fields, form a quartet and the corresponding particles appear in the physical space \mathcal{V}_{ph} only in zero-norm combinations, so that they are confined.

For instance, in the case of the gluons

$$(8) \quad [iQ_{\text{B}}, A_{\mu}^{\alpha}] = \partial_{\mu} c^{\alpha} + (A_{\mu} x c)^{\alpha}.$$

If the operators $(A_{\mu} x c)^{\alpha}$ develop zero-mass bound states described by the asymptotic fields $C_{\mu}^{\text{as}\alpha}$, the gluons are confined.

Let us consider eq. (8) together with

$$(9) \quad [i\bar{Q}_{\text{B}}, A_{\mu}^{\alpha}] = (D_{\mu} \bar{c})^{\alpha},$$

$$(10) \quad [i\bar{Q}_{\text{B}}, D_{\mu} c^{\alpha}] = Y_{\mu}^{\alpha} = -[Q_{\text{B}}, D_{\mu} \bar{c}^{\alpha}],$$

where

$$Y_{\mu}^{\alpha} = D_{\mu} B^{\alpha} + (D_{\mu} c \times \bar{c})^{\alpha}.$$

We shall assume that $(D_{\mu} \bar{c})^{\alpha}$ and Y_{μ}^{α} too develop zero-mass bound states with asymptotic fields $\bar{C}_{\mu}^{\text{as}\alpha}$ and $B_{\mu}^{\text{as}\alpha}$. Then C_{μ}^{as} , $\bar{C}_{\mu}^{\text{as}}$, B_{μ}^{as} and the gluons asymptotic fields A_{μ}^{as} form a quadruplet. In superfield notation ⁽⁵⁾ we have

$$(11) \quad \Phi_{\mu}^{\text{as}}(x, \theta, \bar{\theta}) = A_{\mu}^{\text{as}}(x) + \bar{\theta}(c_{\mu}^{\text{as}}(x) + \partial_{\mu} c_{\mu}^{\text{as}}(x)) + \theta(\bar{c}_{\mu}^{\text{as}}(x) + \partial_{\mu} \bar{c}_{\mu}^{\text{as}}(x)) + \theta\bar{\theta}(B_{\mu}^{\text{as}}(x) + \partial_{\mu} B_{\mu}^{\text{as}}(x))$$

in eq. (11) we have omitted the renormalization constants in front of the asymptotic fields, *i.e.* our asymptotic fields are not normalized.

⁽¹⁰⁾ T. KUGO and I. OJIMA: *Phys. Lett. B*, **73**, 459 (1978); *Prog. Theor. Phys. Suppl.*, **66**, 31 (1979).

Let us write down the effective Lagrangian \mathcal{L}_{eff} for these asymptotic fields. \mathcal{L}_{eff} should satisfy the following requirements.

First of all it must be invariant under BRS and anti-BRS transformations.

Secondly it is quadratic in the asymptotic fields, since they are free fields. Moreover, the truncated effective Lagrangian $\hat{\mathcal{L}}_{\text{eff}}$,

$$\hat{\mathcal{L}}_{\text{eff}} = \mathcal{L}_{\text{eff}} - \mathcal{L}_{\text{g.f.}},$$

where $\mathcal{L}_{\text{g.f.}}$ is the gauge-fixing term, must be independent of the auxiliary fields B^α ($B^\alpha = B^{\alpha \text{as}}$).

Finally we remark that the original N -parameter gauge group G becomes the direct product of N Abelian U_1 groups.

The key point is that, if the bound states described previously exist, the gauge invariance of a Lagrangian term no longer ensures its BRS and anti-BRS invariance.

So, the term

$$\mathcal{L}_0 = -\frac{1}{4} \text{Tr}(\mathbf{F}_{\mu\nu}^{\text{as}} \mathbf{F}^{\mu\nu \text{as}}),$$

where

$$\mathbf{F}_{\mu\nu}^{\text{as}} = \partial_\mu \mathbf{A}_\nu^{\text{as}} - \partial_\nu \mathbf{A}_\mu^{\text{as}}$$

is gauge invariant, but breaks the BRS and anti-BRS symmetry and therefore cannot appear in \mathcal{L}_{eff} .

The simplest quadratic, gauge-invariant as well as BRS and anti-BRS invariant effective Lagrangian is

$$(12) \quad \mathcal{L}_{\text{eff}} = -\text{Tr} \left(\frac{d}{d\theta} \frac{d}{d\bar{\theta}} \left[\frac{1}{4} \Phi_{\mu\nu}^{\text{as}} \Phi^{\mu\nu \text{as}} + \zeta \Phi_{\theta\mu}^{\text{as}} \Phi^{\bar{\theta}\mu \text{as}} \right] \right),$$

where

$$(13) \quad \left\{ \begin{array}{l} \Phi_{\mu\nu}^{\text{as}} = \partial_\mu \Phi_\nu^{\text{as}} - \partial_\nu \Phi_\mu^{\text{as}}, \\ \Phi_{\theta\mu}^{\text{as}} = \frac{\partial}{\partial\theta} \Phi_\mu^{\text{as}} - \partial_\mu \bar{\eta}^{\text{as}}, \\ \Phi_{\bar{\theta}\mu}^{\text{as}} = \frac{\partial}{\partial\bar{\theta}} \Phi_\mu^{\text{as}} - \partial_\mu \eta^{\text{as}}, \end{array} \right.$$

and

$$\begin{aligned} \eta^{\text{as}} &= \mathbf{c}^{\text{as}} - \theta \mathbf{B}^{\text{as}}, \\ \bar{\eta}^{\text{as}} &= \bar{\mathbf{c}}^{\text{as}} + \bar{\theta} \mathbf{B}^{\text{as}}. \end{aligned}$$

In terms of the component fields, eq. (12) becomes

$$(14) \quad \mathcal{L}_{\text{eff}} = \text{Tr} \left(\mathbf{F}_{\mu\nu}^{\text{as}} \partial^\mu \mathbf{B}^{\nu \text{as}} + \zeta \mathbf{B}_\mu^{\text{as}} \mathbf{B}^{\mu \text{as}} + \frac{1}{2} (\partial_\mu \bar{\mathbf{C}}_\nu^{\text{as}} - \partial_\nu \bar{\mathbf{C}}_\mu^{\text{as}}) (\partial_\mu \mathbf{C}_\nu^{\text{as}} - \partial_\nu \mathbf{C}_\mu^{\text{as}}) \right).$$

We shall not consider a term proportional to $\text{Tr}((\partial^2/\partial\theta\partial\bar{\theta}) \Phi_\mu^{\text{as}} \Phi^{\mu \text{as}})$ in \mathcal{L}_{eff} , which would give a mass to the vector particles.

With a Gaussian integration over B_μ^{as} the effective Lagrangian (14) becomes

$$(15) \quad \mathcal{L}_{\text{eff}} \Rightarrow -\text{Tr} \left(\frac{1}{4\xi} (\partial_\mu \mathbf{F}_{\mu\nu}^{as})^2 + \frac{1}{2} (\partial_\mu \bar{\mathbf{C}}_\nu^{as} - \partial_\nu \bar{\mathbf{C}}_\mu^{as}) (\partial_\mu \mathbf{C}_\nu^{as} - \partial_\nu \mathbf{C}_\mu^{as}) \right).$$

The Lagrangian (15) implies that the propagator of the gluon fields in the momentum space is

$$(16) \quad \Delta_{\mu\nu}(q) = \frac{1}{q^4} \left(g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \right) + \dots$$

This dipole propagator leads of course to a violation of the cluster property.

* * *

We are grateful to Drs. S. BRIGUGLIO, U. GASPARINI and P. F. MARCHETTI for discussions.

One of us (LB) acknowledges the Physics Department of University of Massachusetts at Boston for the kind hospitality.