

## Cytotoxic lipid esters from *Peucedanum ledebourielloides*

Xing Zheng · Jiang Du · Yunlong Xu ·  
Duanfang Liao · George R. Pettit

Received: 11 November 2008 / Accepted: 17 March 2009 / Published online: 23 April 2009  
© Birkhäuser Boston 2009

**Abstract** Two cancer cell growth inhibitory esters, 1,2-dipalmitoyl-3-glucosyl glycerol (**1**) and 1,6-dihydroxy-hexane-bis-palmitoyl ester (**2**), together with arachidic acid-2-hydroxy-glycerol ester, daucosterol, and oleanolic acid, were isolated from the roots of *Peucedanum ledebourielloides* (Apiaceae family). The structures were determined by spectroscopic analyses. The esters **1** and **2** displayed significant activity against the SGC-7901, HT-29, and HL-60 cancer cell lines.

**Keywords** Apiaceae family · *Peucedanum ledebourielloides* · Ester · Anticancer

### Introduction

Terrestrial plants continue to be a well-established source of medically useful drugs, and the currently important anticancer drugs provide ample illustration (Newman

---

X. Zheng (✉) · D. Liao  
Institute of Pharmacy and Pharmacology, Nanhua University, Hengyang 421001,  
People's Republic of China  
e-mail: zhengxing5018@yahoo.com

X. Zheng · G. R. Pettit  
Department of Chemistry and Biochemistry, Arizona State University, Tempe, AZ 85287, USA

G. R. Pettit  
e-mail: bpettit@asu.edu

J. Du  
Department of Chemistry, Colorado State University, Fort Collins, CO 80526, USA

Y. Xu  
State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650204, People's Republic of China

*et al.*, 2007; Jiangsu 1991). In the present study, we investigated *Peucedanum ledebourielloides* (Apiaceae family) as a potential source of cancer cell growth inhibitors. *P. ledebourielloides* is a shrub that grows in Huaying County (Shaanxi Province), People's Republic of China. The root of this plant is a well-known traditional medicine in China and has essentially replaced *Saposhnikovia divaricata* to treat arthritis pain as well as other uses (Cragg *et al.*, 2005). Various compounds, including coumarins, flavones, lignans, and volatile oils, have been isolated from many species of *Peucedanum* (Cisowski 1983; Duh *et al.*, 1991; Kong *et al.*, 2003; Li *et al.* 2003; Rao *et al.*, 1997; Zhang *et al.*, 2003). Some have been reported to show anticancer activity (Duh *et al.*, 1991; Schillaci *et al.*, 2003; Zhang *et al.*, 2003). However, the chemical constituents of *P. ledebourielloides* have not been previously investigated.

Continuation of our studies concerning new sources of naturally occurring antineoplastic agents (Pettit *et al.*, 2007; Zheng *et al.*, 2003a, 2003b), two cancer cell growth inhibitors, 1,2-dipalmitoyl-3-glucosyl glycerol (**1**), and 1,6-dihydroxy-hexane-bis-palmitoyl ester (**2**), as well as arachidic acid-2-hydroxy-glycerol ester, daucosterol, and oleanolic acid, were isolated from *Peucedanum ledebourielloides*. Esters from many higher plants exhibit a wide variety of biological activities, including antineoplastic (Boeryd *et al.*, 1986; Graca *et al.*, 2002). We report the isolation, structure, and cancer cell growth inhibitory activities of two esters, 1,2-dipalmitoyl-3-glucosyl glycerol (**1**) and 1,6-dihydroxy-hexane-bis-palmitoyl ester (**2**).

## Materials and methods

### General

Melting points were measured using a Kofler-type melting point apparatus and are uncorrected. IR spectra were recorded with a PE-577 spectrometer (in  $\text{cm}^{-1}$ ; KBr pellets). The  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR spectra were recorded using a Bruker Am-400 unit with TMS as an internal standard (chemical shifts ( $\delta$ ) in ppm). The mass spectra were recorded by using a VG Austospec-3000 spectrometer; in  $m/z$  (rel. int. in % of the base peak). Column chromatography (CC) was performed with silica gel (200–300 mesh, Qingdao Marine Chemical Co., China). Fractions were monitored by TLC, and spots were visualized by heating silica gel plates sprayed with 10%  $\text{H}_2\text{SO}_4$  in ethanol.

### Plant material

The roots of *P. ledebourielloides* were collected in the Huaying County of Shaanxi Province, People's Republic of China, and were identified by Prof. L. R. Xu. A voucher specimen was deposited at the Xibei Institute of Botany (No. K. T. Fu 17188).

### Extraction and isolation

The dry powdered roots of *P. ledebourilloide* (1 kg) were extracted with MeOH ( $3 \times 4$  L) at room temperature for a week. Removal of the organic solvent under


reduced pressure provided a combined MeOH extract (58 g), which was subjected to separation by column chromatography (2 kg silica gel,  $5 \times 100$  cm) using gradient elution (10–60%) with a n-hexane-ethyl acetate. Five major fractions (I–V) were obtained on the basis of TLC monitoring of the individual fractions. Fraction III (0.4 g) was further separated by column chromatography over silica gel ( $4 \times 75$  cm, 300 g) eluted with n-hexane-ethyl acetate [2:98 (4 L), 5:95 (4 L), 10:90 (5 L), 15:85 (5 L), V:V] to obtain four subfractions (*Fr.* 3.1, *Fr.* 3.2, *Fr.* 3.3, *Fr.* 3.4). *Fr.* 3.2 was further purified by repeated column chromatography over silica gel ( $4 \times 75$  cm, 300 g) and elution with n-hexane-ethyl acetate [2:98 (4 L), 5:95 (4 L), 10:90 (4 L), V:V] to yield diester **2** (18 mg), arachidic acid-2-hydroxy-glycerol ester (12 mg), daucosterol (17 mg), and oleanolic acid (13 mg). *Fr.* 3.4 provided glycerine ester **1** (22 mg) following purification chromatography over silica gel ( $3 \times 45$  cm, 160 g) and with n-hexane-ethyl acetate [10:90 (5 L), V:V].

### Identification of compounds

Glucoside **1** was obtained as amorphous colorless powder that gave an  $[M-H]^+$  ion in the HRFABMS at  $m/z$  729.5510, corresponding to molecular formula  $C_{41}H_{78}O_{10}$ . The IR absorption at  $3100\text{--}3700\text{ cm}^{-1}$  and  $1735\text{ cm}^{-1}$  indicated the presence of one or more hydroxyl groups and ester carboxyls. The  $^{13}\text{C}$ -NMR and DEPT spectral data suggested two long-chain aliphatic ester groups corresponding to  $\delta$  173.6 (s), 173.9 (s), 14.3 (q), 23.0 (t), 25.3 (t), 29.9 (t, strong), 32.2 (t), and 34.4 (t). Furthermore, the height of every peak between 14–35 ppm was nearly twice that of the others, which served to confirm the presence of two long-chain esters. The existence of 1- $\beta$ -D-glucoside also was confirmed by  $^{13}\text{C}$ - and DEPT spectra signals at  $\delta$  100.4(d), 75.2(d), 75.1(d), 69.8(d), and 63.6(t). Those assignments left only three unaccounted for signals:  $\delta$  66.7(t), 71.1(d), and 55(t) generated by a glycerol ester. Thus, the structure of glycerol ester **1** was deduced to be 1,2-dipalmitoyl-3-glucosyl glycerol and further suggested by results of EIMS spectroscopy (Fig. 1).

Diester **2** also was obtained as a colorless powder, which yielded a molecular ion in its HRFABMS spectra at  $[M-H]^+$  at  $m/z$  593.5583, pointing to molecular formula  $C_{38}H_{74}O_4$ . The  $^1\text{H}$ -NMR spectrum of **2** showed one methyl at  $\delta$  0.86 (t,  $J = 6.8$  Hz), a methane bonded to a oxygen at  $\delta$  4.03 (t,  $J = 6.8$  Hz), another methane connected to a carbonyl at  $\delta$  2.26 (t,  $J = 6.8$  Hz), and a number of methane signals at 1–1.96 on the  $^{13}\text{C}$ -NMR spectrum of diester **2** indicated long aliphatic ester chains. Further analysis of the NMR data and molecular formula gave results consistent with a symmetrical compound. The prominent palmityl mass fragment in the EIMS spectrum completed the spectral data necessary to assign the new diester **2** as 1,6- dihydroxy-hexane-bis-palmityl ester (Fig. 2).


Arachidic acid-2-hydroxy-glycerol ester, daucosterol, and oleanolic acid were identified by agreement between their spectroscopic data and that published in the literature (Giron *et al.*, 1992; Rawat *et al.*, 1988; Wandji *et al.*, 2003).



**Fig. 1** Mass special fragmentation of glucoside **1**

## Results and discussion

The active compounds were identified as 1,2-dipalmitoyl-3-glucosyl glycerol and 1,6-dihydroxy-hexane-bis-palmitoyl ester. The structures of the compounds are shown in Figs. [1](#) and [2](#).



**Fig. 2** Diester **2** mass special fragmentation

**Table 1** Human cancer cell line growth inhibition values (IC<sub>50</sub> µg/ml) for 1,2-dipalmitoyl-3-glucosyl glycerol (**1**) and 1,6-dihydroxy-hexane-bis-palmitoyl ester (**2**)<sup>a</sup>

Cancer cell lines <sup>b</sup>			
	SGC-7901	HT-29	HL-60
Compound			
<b>1</b>	4.9	0.21	27.7
<b>2</b>	1.6	34.2	0.49
<b>5-FU</b> <sup>c</sup>	0.58	10.26	25.4

<sup>a</sup> In DMSO

<sup>b</sup> Cancer type: SGC-7901(gastric carcinoma), HT-29 (colon), HL-60 (promyelocytic leukemia)

<sup>c</sup> 5-fluorouracil (99%) from Sigma Aldrich Co., USA as positive control

### 1,2-Dipalmitoyl-3-glucosyl glycerol (**1**)

Colorless powder amorphous, mp 203–206°C; IR  $\nu_{\max}$  (log  $\epsilon$ ) 3700–3100, 2920, 2860, 1735, 1455, 1060, and 1040 cm<sup>-1</sup>; <sup>1</sup>H-NMR (C<sub>5</sub>D<sub>5</sub>N, 400 MHz)  $\delta$  4.73 (d,  $J$  = 7.6 Hz, H-1''), 3.41 (m, H-2''), 3.44 (m, H-3''), 3.39 (m, H-4''), 3.43 (m, H-5''), 3.69 (m, H-6''). <sup>13</sup>C-NMR (C<sub>5</sub>D<sub>5</sub>N, 100 MHz)  $\delta$  173.9 (C-16), 173.6 (C-16'), 100.4 (C-1''), 75.2 (C-3''), 75.1 (C-2''), 71.1 (C-18), 69.8 (C-4''), 66.7 (C-17), 63.6 (C-6''), 55.0 (C-19), 32.2 (C-15'), 23.0 (C-2 and C-2'), 14.3 (C-1 and C-1'), 25.1, 29.9 (C-3-C-14 and C-3'-C-14'); and EIMS  $m/z$  (%): 730 (3), 551 (45), 311 (95), 239 (100).

### Diester **2**

Colorless powder (amorphous), mp 65–68°C. <sup>1</sup>H-NMR (C<sub>5</sub>D<sub>5</sub>N, 400 MHz)  $\delta$  4.03 (4H, t,  $J$  = 6.8 Hz, H-17 and H-17'), 2.26 (4H, t,  $J$  = 6.8 Hz, H-15 and H-15'), 0.86 (6H, t,  $J$  = 6.8 Hz, H-1 and H-1'); <sup>13</sup>C-NMR (C<sub>5</sub>D<sub>5</sub>N, 100 MHz)  $\delta$  173.9 (C-16 and C-16'), 64.4 (C-17 and C-17'), 34.5 (C-15 and C-15'), 22.7 (C-2 and C-2'), 14.1 (C-1 and C-1'), 25.1, 29.3 (C3-C14 and C-3'-C14'); EIMS  $m/z$  (%): 594 (12), 257 (82), 239 (29).

## Cytotoxicity assay

Aliphatic ester **1** and **2** were evaluated for in vitro anticancer activity against the SGC-7901, HT-29, and HL-60 cell lines by MTT-based assay. These experiments were performed in 96-well plates essentially as described by Mosmann (1983). The IC<sub>50</sub> concentrations listed represent the concentration that results in a 50% decrease in cell growth during 6-day incubation. Both 1,2-dipalmitoyl-3-glucosyl glycerol (**1**) and 1,6-dihydroxy-hexane-bis-palmitoyl ester (**2**) exhibited significant cancer cell growth inhibitory activities against human gastric carcinoma SGC-7901, human colon cancer HT-29, and human promyelocytic leukemia HL-60 cell lines (Table 1).

**Acknowledgement** The authors thank the Department of Education of Hunan (07B065) and the Department Science and Technology of Hunan (2007FJ4155) for financial support. Special thanks are due to the Analytical Center, State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany of Chinese Academy of Sciences, for recording IR, MS, and NMR spectra. The authors also thank Prof. Langran Xu for plant identification. Two authors (GRP and JD) are pleased to acknowledge the financial support of CA90441-5 and CA90441-06A1 awarded by the Division of Cancer Treatment and Diagnosis, National Cancer Institute, DHHS, the Arizona Biomedical Research Commission, and the assistance of Professor Fiona Hogan.

## References

- Boeryd B, Hallgren B (1986) Tumor transplantation resistance in mice of various ages neonatally fed diets having different fat compositions. *Cancer J* 1:72–76
- Cisowski W (1983) Flavonoid compounds of the herb *Peucedanum japonicum* Thunb. *Pol J Chem* 57(10–12):1283–1286
- Cragg GM, Newman DJ (2005) Plants as source of anticancer agents. *J Ethnopharmacol* 100:72–79. doi:10.1016/j.jep.2005.05.011
- Duh CY, Wang SK, Wu YC (1991) Cytotoxic pyranocoumarins from *Peucedanum japonicum*. *Phytochem* 30:2812–2814. doi:10.1016/0031-9422(91)85156-T
- Giron D, Link R, Bouissel S (1992) Analysis of mono-, di- and triglycerides in pharmaceutical excipients by capillary supercritical fluid chromatography. *J Pharm Biomed* 10:821–830. doi:10.1016/0731-7085(91)80087-P
- Graca J, Schreiber L, Rodrigues J, Pereira H (2002) Glycerol and glyceryl esters of omega-hydroxyacids in cutins. *Phytochemistry* 61:205–215
- Jiangsu Institute of Botany (1991) Outline of new China herbal. Shanghai Science and Technology Press, Shanghai, p 339
- Kong LY, Li Y, Niwa M (2003) A new pyranocoumarin from *Peucedanum praeruptorum*. *Heterocycle* 60:1915–1919
- Li FW, Lou M, Zhu XX, Li RS, Shen JF, Zhang YJ (2003) Study on the preparation of  $\beta$ -cyclodextrin inclusion compound of volatile oils in Jiketing Granula. *Zhongguo Yaoke Daxue Xuebao* 34:29–31
- Mosmann T (1983) Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* 65:55–63. doi:10.1016/0022-1759(83)90303-4
- Newman DJ, Cragg GM (2007) Natural products as sources of new drugs over the last 25 years. *J Nat Prod* 70:461–477. doi:10.1021/np068054v
- Pettit GR, Tan R, Pettit RK, Smith TH, Feng S, Doubek DL, Richert L, Hamblin J, Weber C, Chapuis JC (2007) Antineoplastic agents. 560. Isolation and structure of kitastatin 1 from an Alaskan *Kitasatospora* sp. *J Nat Prod* 70:1069–1072. doi:10.1021/np068072c
- Rao GX, Liu QX, Dai WS, Sun HD (1997) Chemical constituents of *Peucedanum turgeniifolium*. *Tianran Chanwu Yanjiu Yu Kaifa* 9:9–11
- Rawat MSM, Negi DS, Panwar MS, Pant G (1988) Chemical examination of *Rheum moorcroftianum* (rhizomes), a rare high altitudinal species of Garhwal. *Fitoterapia* 59:248–249

- Schillaci D, Venturella F, Venuti F, Plescia F (2003) Antimicrobial and antiproliferative activity of *Peucedanum nebrodense*. J Ethnopharmacol 87:99–101. doi:[10.1016/S0378-8741\(03\)00116-8](https://doi.org/10.1016/S0378-8741(03)00116-8)
- Wandji J, Tillequin F, Mulholland DA, Shirri JC, Tsabang N, Seguin E, Verite P, Libot F, Fomum ZT (2003) Pentacyclic triterpenoid and saponins from *Gambeya boukokoensis*. Phytochemistry 64:845–849. doi:[10.1016/S0031-9422\(03\)00495-3](https://doi.org/10.1016/S0031-9422(03)00495-3)
- Zhang JX, Fong WF, Wu JY, Yang M, Cheung HY (2003) Pyranocoumarins isolated from *Peucedanum praeruptorum* as differentiation inducers in human leukemic HL-60 cells. Planta Med 69:223–229. doi:[10.1055/s-2003-38490](https://doi.org/10.1055/s-2003-38490)
- Zheng X, Cao JG, Meng WD, Qing FL (2003a) Synthesis and anticancer effect of B-Ring trifluoromethylated flavonoids. Bioorg Med Chem Lett 13:3423–3427. doi:[10.1016/S0960-894X\(03\)00752-2](https://doi.org/10.1016/S0960-894X(03)00752-2)
- Zheng X, Meng WD, Xu YY, Cao JG, Qing FL (2003b) Synthesis and anticancer effect of chrysin derivatives. Bioorg Med Chem Lett 13:881–884. doi:[10.1016/S0960-894X\(02\)01081-8](https://doi.org/10.1016/S0960-894X(02)01081-8)