

# Long-term effects of L- and N-type calcium channel blocker on uric acid levels and left atrial volume in hypertensive patients

Mitsuru Masaki<sup>1,2,3</sup> · Toshiaki Mano<sup>1</sup> · Akiyo Eguchi<sup>1,2,3</sup> · Shohei Fujiwara<sup>1</sup> · Masataka Sugahara<sup>1</sup> · Shinichi Hirotsu<sup>1</sup> · Takeshi Tsujino<sup>4</sup> · Kazuo Komamura<sup>1,5</sup> · Masahiro Koshihara<sup>2,3</sup> · Tohru Masuyama<sup>1</sup>

Received: 17 August 2015 / Accepted: 15 January 2016  
© Springer Japan 2016

**Abstract** Left ventricular (LV) diastolic dysfunction is associated with hypertension and hyperuricemia. However, it is not clear whether the L- and N-type calcium channel blocker will improve LV diastolic dysfunction through the reduction of uric acid. The aim of this study was to investigate the effects of anti-hypertensive therapy, the L- and N-type calcium channel blocker, cilnidipine or the L-type calcium channel blocker, amlodipine, on left atrial reverse remodeling and uric acid in hypertensive patients. We studied 62 patients with untreated hypertension, randomly assigned to cilnidipine or amlodipine for 48 weeks. LV diastolic function was assessed with the left atrial volume index (LAVI), mitral early diastolic wave ( $E$ ), tissue Doppler early diastolic velocity ( $E'$ ) and the ratio ( $E/E'$ ). Serum uric acid levels were measured before and after treatment. After treatment, systolic and diastolic blood pressures equally dropped in both groups. LAVI,  $E/E'$ , heart rate and uric acid levels decreased at 48 weeks in the cilnidipine group but not in the amlodipine group. The % change from baseline to 48 weeks in LAVI,  $E$

wave,  $E/E'$  and uric acid levels were significantly lower in the cilnidipine group than in the amlodipine group. Larger %-drop in uric acid levels were associated with larger %-reduction of LAVI ( $p < 0.01$ ). L- and N-type calcium channel blocker but not L-type calcium channel blocker may improve LV diastolic function in hypertensive patients, at least partially through the decrease in uric acid levels.

**Keywords** Hypertension · Uric acid · L- and N-type calcium channel blocking drugs · Diastolic dysfunction

## Abbreviations

CCB	Calcium channel blocker
UA	Uric acid
LAV	Left atrial volume
LV	Left ventricular
RAS	Renin–angiotensin system
SBP	Systolic blood pressure
DBP	Diastolic blood pressure
LAVI	Left atrial volume index
LVMI	Left ventricular mass index
RWT	Relative wall thickness
$E$	Peak velocities of early diastolic phase
$A$	Late diastolic phase of mitral inflow
$E'$	Mitral annulus velocities
HR	Heart rate
LAD	Left atrial diameter
LVEDD	Left ventricular end-diastolic diameter
LVESD	Left ventricular end-systolic diameter
LVEF	Left ventricular ejection fraction
LVMI	Left ventricular mass index
DT	Deceleration time

✉ Toshiaki Mano  
mano@hyo-med.ac.jp

<sup>1</sup> Cardiovascular Division, Department of Internal Medicine, Hyogo College of Medicine, 1-1 Mukogawa-cho, Nishinomiya 663-8501, Japan

<sup>2</sup> Division of Clinical Laboratory Medicine, Hyogo College of Medicine, Nishinomiya, Japan

<sup>3</sup> Center for Health Evaluation and Promotion, Hyogo College of Medicine Hospital, Nishinomiya, Japan

<sup>4</sup> Department of Pharmacy, School of Pharmacy, Hyogo University of Health Sciences, Kobe, Japan

<sup>5</sup> Osaka Clinic, Takeda Pharmaceutical Company Limited, Osaka, Japan

## Introduction

Hypertension leads to diastolic dysfunction with left atrial (LA) enlargement and left ventricular hypertrophy [1, 2]. Diastolic dysfunction is associated with increased morbidity and mortality [3] due to enhanced activity of the sympathetic nervous system and the oxidative stress [4–6]. Recent animal studies have shown that L- and N-type calcium channel blocker compared with L-type calcium channel blocker improved the left ventricular diastolic dysfunction and cardiac oxidative stress in Dahl salt-sensitive rats [7]. Moreover, L- and N-type calcium channel blocker has greater inhibition of sympathetic nervous system in hypertensive patients [8–10].

Hyperuricemia play an important role in development of hypertension, cardiovascular disease, chronic kidney disease and metabolic syndrome [11, 12]. L- and N-type calcium channel blocker, cilnidipine decreased serum uric acid levels by increasing the nitrogen monoxide excretion [13].

However, it is unclear whether the L- and N-type calcium channel blocker, cilnidipine or the L-type calcium channel blocker, amlodipine could decrease serum uric acid level and LA volume in patients with untreated hypertension. Thus, the aim of this study was to investigate the impact of decrease in uric acid as a contributor to regression of LA volume index in hypertensive patients following long-term treatment with cilnidipine or amlodipine.

## Materials and methods

### Subjects

We enrolled 62 patients with untreated hypertension who presented to outpatient clinic of Hyogo College of Medicine between January 2010 and November 2015. Patients were included in the study if they met the following criteria: 20 years of age or older, systolic blood pressure (SBP) and diastolic blood pressure (DBP) of 140 and 90 mmHg or over. Patients were excluded if they had secondary hypertension of any cause, angina pectoris or acute coronary artery disease, current or recent history of congestive heart failure, valvular heart diseases, cardiac arrhythmias, renal dysfunction (serum creatinine level over 2.0 mg/dL). Informed consent was obtained from all patients, and the study was designed to comply with the ethical principles of our institution. Eligible patients were randomly assigned in a 1:1 ratio to receive either amlodipine 2.5 mg daily (amlodipine group) or cilnidipine 10 mg daily (cilnidipine group). Target SBP and DBP were below 140 and 90 mmHg, respectively. If blood pressure reduction did not achieve the target level after 4 weeks, the daily dose was

increased to 10 mg amlodipine or 20 mg cilnidipine. The third and fourth steps of treatment included the addition of a thiazide and/or alpha-blocker.

### Blood chemistry

Among 62 enrolled subjects, we could collect blood samples for uric acid measurement from 32 patients in cilnidipine and 27 patients in amlodipine groups. 3 subjects could not be measured because of missing blood samples. Blood samples ( $n = 59$ ) were taken between 0900 and 1100 hours and were immediately placed on ice and centrifuged within 1 h. The specimens were stored at  $-80^{\circ}\text{C}$  until analysis. Uric acid was measured by an automatic biochemical analysis systems Hitachi LABOSPECT.

### Echocardiographic studies

Transthoracic echocardiography was performed at baseline and 48 weeks after treatment in all patients. Echocardiography was recorded with iE33 (Philips Medical Systems, Bothell, Washington). A standard, comprehensive, M-mode, 2-dimensional echocardiography and Doppler study were conducted according to the guideline of the American Society of Echocardiography [14]. LA volume was calculated with a formula using an ellipsoid model and was indexed to the body surface area, i.e., LA volume index (LAVI) [15]. LV mass index (LVMI) and relative wall thickness (RWT) were measured by the method described previously [16]. Peak velocities of early diastolic phase ( $E$ ) and late diastolic phase ( $A$ ) of mitral inflow, and the  $E/A$  ratio were measured by pulsed-wave Doppler echocardiography with the sample volume between mitral leaflet tips. Mitral annulus velocities ( $E'$ ) and  $E/E'$  ratio were measured at the septal annulus by tissue Doppler imaging.

### Ethics

This study was conducted in accordance with the principles outlined in the Declaration of Helsinki after receiving approval from the institutional review board of Hyogo College of Medicine. All subjects provided written informed consent prior to participation.

### Statistical analysis

The primary outcomes included changes in uric acid and LA volume in hypertensive patients following long-term treatment with amlodipine or cilnidipine. Continuous data were presented as mean and standard error of the mean ( $\pm\text{SEM}$ ). We compared values at baseline and after treatment using the Wilcoxon signed ranks test. Spearman's rank correlation coefficient was used to examine

the correlations between the change in serum uric acid and LAVI, SBP, heart rate and  $E/E'$ .  $p$  values  $<0.05$  were considered significant. Statistical computations were performed with JMP version 10.0.1 (SAS Institute, Inc, Cary, North Carolina).

## Results

### Patient characteristics

We analyzed the data of 35 men and 27 women with mean age of 61 years. The final dose of amlodipine was  $5.8 \pm 0.4$  mg/day and that of cilnidipine was  $10.0 \pm 0.9$  mg/day. Thiazide indapamide 1 mg was added to the basal

medications in four patients of amlodipine group and three patients of cilnidipine group and, whereas alpha blocker doxazosin 1 mg was added in three patients of amlodipine group and in no patients of cilnidipine group. Baseline characteristics were similar in both groups of patients (Table 1).

### Changes in blood pressure and echocardiographic parameters and serum uric acid


SBP, DBP and LVMI decreased similarly in the cilnidipine and amlodipine groups after 48 weeks of treatment. The serum uric acid, LAVI,  $E/E'$  and heart rate significantly decreased in the cilnidipine group but not in the amlodipine group.  $E$  wave significantly increased in the amlodipine group but not in the cilnidipine group (Table 1).

**Table 1** Changes in hemodynamic and echocardiographic parameters, and uric acid levels in amlodipine and cilnidipine groups

	Amlodipine		Cilnidipine	
	Before treatment	48 weeks	Before treatment	48 weeks
Age (years)	60 $\pm$ 2		62 $\pm$ 2	
Sex (female/male)	11/19, 30		16/16, 32	
Hight (cm)	161 $\pm$ 2		160 $\pm$ 2	
Body weight (kg)	65 $\pm$ 2	65 $\pm$ 2	61 $\pm$ 2	61 $\pm$ 2
SBP (mmHg)	165 $\pm$ 3	132 $\pm$ 3***	162 $\pm$ 3	132 $\pm$ 2***
DBP (mmHg)	94 $\pm$ 2	81 $\pm$ 2***	93 $\pm$ 2	77 $\pm$ 2***
HR (bpm)	74 $\pm$ 2	74 $\pm$ 2	71 $\pm$ 2	65 $\pm$ 2*
LAD (mm)	37 $\pm$ 1	37 $\pm$ 1	38 $\pm$ 1	37 $\pm$ 1
LAVI (ml/m <sup>2</sup> )	21 $\pm$ 1	24 $\pm$ 1*	24 $\pm$ 1	22 $\pm$ 1**
LVEDD (mm)	48 $\pm$ 1	48 $\pm$ 1	48 $\pm$ 1	48 $\pm$ 1
LVESD (mm)	29 $\pm$ 1	30 $\pm$ 1	31 $\pm$ 1	30 $\pm$ 1
RWT	0.42 $\pm$ 0.06	0.38 $\pm$ 0.01**	0.41 $\pm$ 0.02	0.36 $\pm$ 0.01*
LVEF (%)	67 $\pm$ 1	67 $\pm$ 1	66 $\pm$ 2	69 $\pm$ 1
LVMI (g/m <sup>2</sup> )	104 $\pm$ 3	93 $\pm$ 3***	100 $\pm$ 6	86 $\pm$ 4***
$E$ (m/s)	58 $\pm$ 3	68 $\pm$ 3**	62 $\pm$ 2	63 $\pm$ 2
$A$ (m/s)	72 $\pm$ 3	78 $\pm$ 3	73 $\pm$ 3	78 $\pm$ 3
$E/A$	0.9 $\pm$ 0.1	0.9 $\pm$ 0.1	0.9 $\pm$ 0.1	0.9 $\pm$ 0.1
DT (ms)	202 $\pm$ 8	202 $\pm$ 9	206 $\pm$ 8	215 $\pm$ 10
$E'$ (cm/s)	5.7 $\pm$ 0.3	6.1 $\pm$ 0.3**	5.2 $\pm$ 0.2	6.1 $\pm$ 0.3***
$E/E'$	11 $\pm$ 1	12 $\pm$ 1	12 $\pm$ 1	11 $\pm$ 1**
Serum uric acid (mg/dl)	5.2 $\pm$ 0.2	5.5 $\pm$ 1.3	5.6 $\pm$ 0.2	5.4 $\pm$ 0.2*
Comorbidity				
Dyslipidemia (%)	4 (13)		7 (22)	
Diabetes mellitus (%)	3 (10)		4 (13)	
Smoking (%)	2 (6)		3 (9)	
Baseline Medication				
Statin $n$ (%)	4 (13)		4 (13)	
Sulfonylurea $n$ (%)	3 (10)		4 (13)	

SBP systolic blood pressure, DBP diastolic blood pressure, HR heart rate, LAD left atrial diameter, LAVI left atrial volume index, LVEDD left ventricular end-diastolic diameter, LVESD left ventricular end-systolic diameter, RWT relative wall thickness, LVEF left ventricular ejection fraction, LVMI left ventricular mass index, DT deceleration time

\*  $p < 0.05$  versus baseline; \*\*  $p < 0.01$  versus baseline; \*\*\*  $p < 0.001$  versus baseline



**Fig. 1** Percent change in serum uric acid (a), LAVI (b), trans mitral *E* wave peak velocity (*E*) (c), and ratio of *E* to mitral annular *E'* velocity (*E/E'*) (d)

### Effects of cilnidipine and amlodipine on LAVI, *E/E'* and serum uric acid

Percent changes from the baseline in LAVI, *E* wave, *E/E'* and serum uric acid levels were lower in the cilnidipine group than in the amlodipine group ( $-9 \pm 3$  vs.  $26 \pm 10$  %,  $p < 0.01$ ,  $3 \pm 3$  vs.  $20 \pm 5$  %,  $p < 0.01$ ,  $-12 \pm 4$  vs.  $12 \pm 6$  %,  $p < 0.01$ ,  $-3 \pm 3$  vs.  $8 \pm 5$  %,  $p < 0.05$  as percentage reduction from the values before treatment) (Fig. 1a–d).

### Correlation between serum uric acid and LAVI

Larger percent-drop in serum uric acid were associated with larger percent-reduction of LAVI in the cilnidipine group ( $\rho = 0.49$ ,  $p < 0.01$ ) but not in the amlodipine group (Fig. 2). No association was showed between the percent changes in serum uric acid and the changes in SBP, heart rate and *E/E'* in both groups.


### Discussion

Main findings in the present study were: (1) While blood pressure reduction over 48 weeks was similar between cilnidipine and amlodipine groups, serum uric acid, LAVI, *E/E'* and heart rate decreased only in cilnidipine group; (2) In cilnidipine group, percent decrease in LAVI was correlated with percent decrease in serum uric acid.

These data suggest that cilnidipine decreased LV filling pressure and LA volume at least partially through the reduction of uric acid levels in hypertensive patients.

### Effect of cilnidipine on hemodynamics and left atrial volume

Enlargement of LA has adverse cardiovascular outcomes, including heart failure [17–19], atrial fibrillation [20], stroke [21], and mortality [2]. LA dilatation has been



**Fig. 2** Correlation between percent change in serum uric acid and percent change in LAVI. *Left* amlodipine, *right* cilnidipine

associated with LV diastolic dysfunction, independent of age, gender and LV ejection fraction [1, 22]. Since LA dilatation can be attributed to a sustained elevation of LV filling pressure and an atrial volume overload due to diastolic dysfunction in hypertensive patients [23, 24], the reduction of LA volume might be associated with improvement in cardiovascular events. In this study, LAVI,  $E/E'$  and heart rate decreased only in cilnidipine group, not in amlodipine group. L- and N-type calcium channel blocker, cilnidipine has improved diastolic dysfunction or cardiac fibrosis in hypertensive patients and hypertensive rats [7, 25, 26]. Moreover, cilnidipine has anti-sympathetic nerve activity, anti-inflammation effect, anti-oxidation effect, anti-renin-angiotensin-aldosterone system, in addition to its antihypertensive effect [10, 27–31].

Takatsu et al. [7] described that cilnidipine reduced left ventricular diastolic dysfunction compared with amlodipine in Dahl salt-sensitive rats. Soeki et al. [10] reported that cilnidipine inhibited the oxidative stress in hypertensive patients. Konda et al. [27] found that cilnidipine, compared with amlodipine, inhibited renin-angiotensin system due to the anti-sympathetic nerve activity in hypertensive model rats. Aritomi et al. [52] demonstrated that cilnidipine suppressed the aldosterone synthase (CYP11B2) expression in human adrenocortical cells. The improvement in diastolic function with cilnidipine might be at least partially attributed to the inhibition of cardiac oxidative stress, inflammation and renin-angiotensin-aldosterone gene expression.

On the other hand, L-type calcium channel blocker, amlodipine, might activate renin-angiotensin-aldosterone system due to increase in sympathetic tone via a rapid and large reduction in blood pressure [28, 32, 33]. Nishimura

et al. [34] showed increment in LV end-diastolic pressure as well as prolongation of the time constant of relaxation by calcium channel blocker. In this study, LAVI and  $E$  were increased in the amlodipine group. These data suggested that L-type calcium channel blocker, amlodipine may not improve LV preload through activated sympathetic nerve as compared to L- and N-type calcium channel blocker, cilnidipine. These data suggest cilnidipine may have a beneficial effect on anti-sympathetic nerve activity, LV filling pressure and LA volume as well.

However, it has not been fully clarified whether the LA volume reduction directly results in the improvement in cardiovascular outcomes. Further research intervention studies are required to establish whether the LA volume reduction directly results in the improvement in cardiovascular mortality.

#### **Effect of cilnidipine on serum uric acid and left atrial reverse remodeling**

There are several possible mechanisms that might explain how cilnidipine could decrease serum uric acid level. Hamada et al. reported that cilnidipine significantly decreased the serum uric acid level and increased uric nitrogen monoxide synthesis in hypertensive patients. Similarly, Uchida et al. observed that cilnidipine improved uric acid metabolism in hypertensive patients with chronic kidney disease. This mechanism of the serum uric acid reduction by cilnidipine inhibited the production of uric acid precursor, hypoxanthine, in the skeletal muscle [13, 51].

The present study, the larger decrease in serum uric acid was associated with the larger reduction of LAVI in



cilnidipine group. Previous studies demonstrated that hyperuricemia was associated with the dilatation of left atrial size or elevated left atrial pressures [35, 36]. In addition, hyperuricemia promote hypertension, metabolic syndrome, cardiac fibrosis, cardiac oxidative stress, LV hypertrophy, LV diastolic dysfunction and endothelial dysfunction [6, 11, 12, 38–42]. The pro-oxidant activity of serum uric acid increased the inflammatory cytokines such as monocyte chemo attractant protein-1 (MCP-1) and cyclooxygenase (COX-2), and the oxidant species such as peroxynitrite [43–45]. Randomized trial showed that allopurinol, a xanthine oxidase inhibitors, decreased plasma renin activity and blood pressure in patients with hypertension [37]. This study suggested that decreasing serum uric acid may be associated with afterload reduction and improving renin–angiotensin–aldosterone system. Therefore, it is speculated that the lowering of uric acid would be accompanied by the left atrial size reduction.

However, serum uric acid is not only pro-oxidant effects [43–45] but also anti-oxidant effects [48, 49]. Administration of uric acid has been reported to improve endothelial function [49, 50]. Sugihara et al. [47] found that very low serum uric acid (below 0.8 mg/dL) was associated with endothelial dysfunction by increasing oxidative stress. The correlation between serum uric acid levels and cardiovascular events were J-shaped in PIUMA study [46]. Although raising serum uric acid levels protected against oxidative stress, these studies investigate the effect of acute uric acid administration, younger population and very low serum uric acid concentrations. Middle aged hypertensive patients were included in the present study. Moreover, there were no patients with very low serum uric acid levels in the present study. Therefore, serum uric acid in this study may be recognized as a marker of pro-oxidant rather than an anti-oxidant.

### Limitations

There are several limitations in this study. First, the small number of patients is enrolled in this study. Therefore, a large number of patients is needed to confirm our results. Second, neither invasive hemodynamic evaluation nor the activation of oxidative stress, renin–angiotensin–aldosterone and hypoxanthine were available in the present study. In future, we need to elaborate the detailed relations between invasive hemodynamic monitoring and cardiac structure. Third, we used only one calcium channel blocker for each hypertensive patient group (cilnidipine and amlodipine). Therefore the evaluation of other calcium channel blockers may be necessary in the future study.

### Perspectives

The cilnidipine seems to decrease serum uric acid and LAVI, differently from amlodipine. There were significant

correlations between percent decrease in serum uric acid and percent decreases in LAVI. These data suggest that suppression of serum uric acid by the L- and N-type calcium channel blocker contributed to reverse remodeling of LA in hypertensive patients. The L- and N-type calcium channel blocker, which improve LV diastolic dysfunction, may be useful in heart failure with preserved ejection fraction.

**Acknowledgments** We thank Ms. M. Tanaka, S. Makihara and C. Misumi for their excellent technical assistance in the acquisition of echocardiographic tracings. This work was supported by a Grant-in-Aid for Researchers, Hyogo College of Medicine, 2014.

### Compliance with ethical standards

**Conflict of interest** The authors declare no conflict of interest.

### References

1. Tsang TS, Barnes ME, Gersh BJ, Bailey KR, Seward JB (2002) Left atrial volume as a morphophysiologic expression of left ventricular diastolic dysfunction and relation to cardiovascular risk burden. *Am J Cardiol* 90:1284–1289
2. Pritchett AM, Mahoney DW, Jacobsen SJ, Rodeheffer RJ, Karon BL, Redfield MM (2005) Diastolic dysfunction and left atrial volume: a population-based study. *J Am Coll Cardiol* 45:87–92
3. Dahlöf B, Pennert K, Hansson L (1992) Reversal of left ventricular hypertrophy in hypertensive patients. A metaanalysis of 109 treatment studies. *Am J Hypertens* 5:95–110
4. Grossman W (1991) Diastolic dysfunction in congestive heart failure. *N Engl J Med* 325:1557–1564
5. Zile MR, Brutsaert DL (2002) New concepts in diastolic dysfunction and diastolic heart failure: part II: causal mechanisms and treatment. *Circulation* 105:1503–1508
6. Bergamini C, Ciccoira M, Rossi A, Vassanelli C (2009) Oxidative stress and hyperuricaemia: pathophysiology, clinical relevance, and therapeutic implications in chronic heart failure. *Eur J Heart Fail* 11:444–452
7. Takatsu M, Hattori T, Murase T, Ohtake M, Kato M, Nashima K, Nakashima C, Takahashi K, Ito H, Niinuma K, Aritomi S, Murohara T, Nagata K (2012) Comparison of the effects of cilnidipine and amlodipine on cardiac remodeling and diastolic dysfunction in Dahl salt-sensitive rats. *J Hypertens* 30:1845–1855
8. Sakata K, Shirotani M, Yoshida H, Nawada R, Obayashi K, Togi K, Miho N (1999) Effects of amlodipine and cilnidipine on cardiac sympathetic nervous system and neurohormonal status in essential hypertension. *Hypertension* 33:1447–1452
9. Sakata K, Yoshida H, Tamekiyo H, Obayashi K, Nawada R, Doi O, Mori N (2003) Comparative effect of cilnidipine and quinapril on left ventricular mass in mild essential hypertension. *Drugs Exp Clin Res* 29:117–123
10. Soeki T, Kitani M, Kusunose K, Yagi S, Taketani Y, Koshiba K, Wakatsuki T, Orino S, Kawano K, Sata M (2012) Renoprotective and antioxidant effects of cilnidipine in hypertensive patients. *Hypertens Res* 35:1058–1062
11. Oda E (2014) Serum uric acid is an independent predictor of metabolic syndrome in a Japanese health screening population. *Heart Vessels* 29:496–503
12. Kawada T, Andou T, Fukumitsu M (2015) Metabolic syndrome showed significant relationship with carotid atherosclerosis. *Heart Vessels*. doi:10.1007/s00380-015-0668-y

13. Hamada T, Yamada K, Mizuta E, Watanabe A, Osaki T, Ishida K, Hasegawa A, Sakata S, Mishima M, Ogino K, Nosaka Y, Miyazaki S, Ohtahara A, Ninomiya H, Kato M, Yoshida A, Taniguchi S, Yamamoto K, Hisatome I (2012) Effects of cilnidipine on serum uric acid level and urinary nitrogen monoxide excretion in patients with hypertension. *Clin Exp Hypertens* 34:470–473
14. Schiller NB, Shah PM, Crawford M, DeMaria A, Devereux R, Feigenbaum H, Gutgesell H, Reichek N, Sahn D, Schnittger I, Silverman N, Tajik A (1989) Recommendations for quantitation of the left ventricle by two-dimensional echocardiography. American Society of Echocardiography Committee on Standards, Subcommittee on Quantitation of Two-Dimensional Echocardiograms. *J Am Soc Echocardiogr* 2:358–367
15. Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, Picard MH, Roman MJ, Seward J, Shanewise JS, Solomon SD, Spencer KT, Sutton MS, Stewart WJ (2005) Recommendations for chamber quantification: a report from the American Society of Echocardiography's Guidelines and Standards Committee and the Chamber Quantification Writing Group, developed in conjunction with the European Association of Echocardiography, a branch of the European Society of Cardiology. *J Am Soc Echocardiogr* 18:1440–1463
16. Yoshida C, Goda A, Naito Y, Nakaboh A, Matsumoto M, Otsuka M, Ohyanagi M, Hirotani S, Lee-Kawabata M, Tsujino T, Masuyama T (2011) Role of plasma aldosterone concentration in regression of left-ventricular mass following antihypertensive medication. *J Hypertens* 29:357–363
17. Takemoto Y, Barnes ME, Seward JB, Lester SJ, Appleton CA, Gersh BJ, Bailey KR, Tsang TS (2005) Usefulness of left atrial volume in predicting first congestive heart failure in patients  $\geq 65$  years of age with well-preserved left ventricular systolic function. *Am J Cardiol* 96:832–836
18. Gottdiener JS, Kitzman DW, Aurigemma GP, Arnold AM, Manolio TA (2006) Left atrial volume, geometry, and function in systolic and diastolic heart failure of persons  $\geq 65$  years of age (the cardiovascular health study). *Am J Cardiol* 97:83–89
19. Zile MR, Gottdiener JS, Hetzel SJ, McMurray JJ, Komajda M, McKelvie R, Baicu CF, Massie BM, Carson PE (2011) Prevalence and significance of alterations in cardiac structure and function in patients with heart failure and a preserved ejection fraction. *Circulation* 124:2491–2501
20. Vaziri SM, Larson MG, Benjamin EJ, Levy D (1994) Echocardiographic predictors of nonrheumatic atrial fibrillation. The Framingham Heart Study. *Circulation* 89:724–730
21. Benjamin EJ, D'Agostino RB, Belanger AJ, Wolf PA, Levy D (1995) Left atrial size and the risk of stroke and death. The Framingham Heart Study. *Circulation* 92:835–841
22. Tsang MY, Barnes ME, Tsang TS (2012) Left atrial volume: clinical value revisited. *Curr Cardiol Rep* 14:374–380
23. Douglas PS (2003) The left atrium: a biomarker of chronic diastolic dysfunction and cardiovascular disease risk. *J Am Coll Cardiol* 42:1206–1207
24. Raman SV (2010) The hypertensive heart. An integrated understanding informed by imaging. *J Am Coll Cardiol* 55:91–96
25. Kosaka T, Nakagawa M, Ishida M, Iino K, Watanabe H, Hasegawa H, Ito H (2009) Cardioprotective effect of an L/N-type calcium channel blocker in patients with hypertensive heart disease. *J Cardiol* 54:262–272
26. Kobayashi N, Mori Y, Mita S, Nakano S, Kobayashi T, Tsubokou Y, Matsuoka H (2001) Effects of cilnidipine on nitric oxide and endothelin-1 expression and extracellular signal-regulated kinase in hypertensive rats. *Eur J Pharmacol* 422:149–157
27. Konda T, Enomoto A, Aritomi S, Niinuma K, Koganei H, Ogawa T, Nitta K (2009) Different effects of L/N-type and L-type calcium channel blockers on the renin–angiotensin–aldosterone system in SHR/Izm. *Am J Nephrol* 30:155–161
28. Takahara A, Nakamura Y, Wagatsuma H, Aritomi S, Nakayama A, Satoh Y, Akie Y, Sugiyama A (2009) Long-term blockade of L/N-type  $\text{Ca}^{2+}$  channels by cilnidipine ameliorates repolarization abnormality of the canine hypertrophied heart. *Br J Pharmacol* 158:1366–1374
29. Fan YY, Kohno M, Nakano D, Ohsaki H, Kobori H, Suwarni D, Ohashi N, Hitomi H, Asanuma K, Noma T, Tomino Y, Fujita T, Nishiyama A (2010) Cilnidipine suppresses podocyte injury and proteinuria in metabolic syndrome rats: possible involvement of N-type calcium channel in podocyte. *J Hypertens* 28:1034–1043
30. Aritomi S, Konda T, Yoshimura M (2012) L/N-type calcium channel blocker suppresses reflex aldosterone production induced by antihypertensive action. *Heart Vessels* 27:419–423
31. Nishida M, Ishikawa T, Saiki S, Sunggip C, Aritomi S, Harada E, Kuwahara K, Hirano K, Mori Y, Kim-Mitsuyama S (2013) Voltage-dependent N-type  $\text{Ca}^{2+}$  channels in endothelial cells contribute to oxidative stress-related endothelial dysfunction induced by angiotensin II in mice. *Biochem Biophys Res Commun* 434:210–216
32. Lefrandt JD, Heitmann J, Sevré K, Castellano M, Hausberg M, Fallon M, Fluckiger L, Urbigkeit A, Rostrup M, Agabiti-Rosei E, Rahn KH, Murphy M, Zannad F, de Kam PJ, van Roon AM, Smit AJ (2001) The effects of dihydropyridine and phenylalkylamine calcium antagonist classes on autonomic function in hypertension: the VAMPHYRE study. *Am J Hypertens* 14:1083–1089
33. Struck J, Muck P, Trubger D, Handrock R, Weidinger G, Dendorfer A, Dodt C (2002) Effects of selective angiotensin II receptor blockade on sympathetic nerve activity in primary hypertensive subjects. *J Hypertens* 20:1143–1149
34. Nishimura RA, Schwartz RS, Holmes DR Jr, Tajik AJ (1993) Failure of calcium channel blockers to improve ventricular relaxation in humans. *J Am Coll Cardiol* 21:182–188
35. Chao TF, Hung CL, Chen SJ, Wang KL, Chen TJ, Lin YJ, Chang SL, Lo LW, Hu YF, Tuan TC, Chen SA (2013) The association between hyperuricemia, left atrial size and new-onset atrial fibrillation. *Int J Cardiol* 168:4027–4032
36. Hooper MM, Hohlfeld JM, Fabel H (1999) Hyperuricaemia in patients with right or left heart failure. *Eur Respir J* 13:682–685
37. Feig DI, Soletsky B, Johnson RJ (2008) Effect of allopurinol on blood pressure of adolescents with newly diagnosed essential hypertension: a randomized trial. *JAMA* 300:924–932
38. Culleton BF, Larson MG, Kannel WB, Levy D (1999) Serum uric acid and risk for cardiovascular disease and death: the Framingham Heart Study. *Ann Intern Med* 131:7–13
39. Khosla UM, Zharikov S, Finch JL, Nakagawa T, Roncal C, Mu W, Krotova K, Block ER, Prabhakar S, Johnson RJ (2005) Hyperuricemia induces endothelial dysfunction. *Kidney Int* 67:1739–1742
40. Kanbay M, Segal M, Afsar B, Kang DH, Rodriguez-Iturbe B, Johnson RJ (2013) The role of uric acid in the pathogenesis of human cardiovascular disease. *Heart* 99:759–766
41. Pan KL, Lin JC, Lin CL, Chen MC, Chang PJ, Hsiao JF, Chang ST, Chung CM (2014) The effects of gout on left atrial volume remodelling: a prospective echocardiographic study. *Rheumatology (Oxford)* 53:867–874
42. Nishino M, Mori N, Yoshimura T, Nakamura D, Lee Y, Taniike M, Makino N, Kato H, Egami Y, Shutta R, Tanouchi J, Yamada Y (2014) Higher serum uric acid and lipoprotein(a) are correlated with coronary spasm. *Heart Vessels* 29:186–190
43. Kanellis J, Watanabe S, Li JH, Kang DH, Li P, Nakagawa T, Wamsley A, Sheikh-Hamad D, Lan HY, Feng L, Johnson RJ (2003) Uric acid stimulates monocyte chemoattractant protein-1 production in vascular smooth muscle cells via mitogen-activated protein kinase and cyclooxygenase-2. *Hypertension* 41:1287–1293

44. Santos CX, Anjos EI, Augusto O (1999) Uric acid oxidation by peroxynitrite: multiple reactions, free radical formation, and amplification of lipid oxidation. *Arch Biochem Biophys* 372:285–294
45. Abuja PM (1999) Ascorbate prevents prooxidant effects of urate in oxidation of human low density lipoprotein. *FEBS Lett* 446:305–308
46. Verdecchia P, Schillaci G, Reboldi G, Santeusano F, Porcellati C, Brunetti P (2000) Relation between serum uric acid and risk of cardiovascular disease in essential hypertension. The PIUMA study. *Hypertension* 36:1072–1078
47. Sugihara S, Hisatome I, Kuwabara M, Niwa K, Maharani N, Kato M, Ogino K, Hamada T, Ninomiya H, Higashi Y, Ichida K, Yamamoto K (2015) Depletion of uric acid due to SLC22A12 (URAT1) loss-of-function mutation causes endothelial dysfunction in hypouricemia. *Circ J* 79:1125–1132
48. Ames BN, Cathcart R, Schwiers E, Hochstein P (1981) Uric acid provides an antioxidant defense in humans against oxidant- and radical-caused aging and cancer: a hypothesis. *Proc Natl Acad Sci USA* 78:6858–6862
49. Waring WS, Webb DJ, Maxwell SR (2001) Systemic uric acid administration increases serum antioxidant capacity in healthy volunteers. *J Cardiovasc Pharmacol* 38:365–371
50. Waring WS, McKnight JA, Webb DJ, Maxwell SR (2006) Uric acid restores endothelial function in patients with type 1 diabetes and regular smokers. *Diabetes* 55:3127–3132
51. Uchida S, Takahashi M, Sugawara M, Saito T, Nakai K, Fujita M, Mochizuki K, Shin I, Morita T, Hikita T, Itakura H, Takahashi Y, Mizuno S, Ohno Y, Ito K, Ito T, Soma M (2014) Effects of the N/L-type calcium channel blocker cilnidipine on nephropathy and uric acid metabolism in hypertensive patients with chronic kidney disease (J-CIRCLE study). *J Clin Hypertens (Greenwich)* 16:746–753
52. Aritomi S, Wagatsuma H, Numata T, Uriu Y, Nogi Y, Mitsui A, Konda T, Mori Y, Yoshimura M (2011) Expression of N-type calcium channels in human adrenocortical cells and their contribution to corticosteroid synthesis. *Hypertens Res* 34:193–201