

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/226821475>

# Biochemical characterization of glutamine synthetase from the diazotrophic cyanobacterium, *Anabaena doliolum*

ARTICLE *in* CURRENT MICROBIOLOGY · JULY 1992

Impact Factor: 1.42 · DOI: 10.1007/BF01570962

---

CITATION

1

---

READS

43

2 AUTHORS, INCLUDING:



Prakash S Bisen

Jiwaji University

359 PUBLICATIONS 1,925 CITATIONS

SEE PROFILE

## Biochemical Characterization of Glutamine Synthetase from the Diazotrophic Cyanobacterium, *Anabaena doliolum*

P.S. Bisen and S. Shanthi


Department of Microbiology, Barkatullah University, Bhopal, India

**Abstract.** In cyanobacteria, the glutamine synthetase–L-glutamine-2-oxoglutarate aminotransferase (GS-GOGAT) pathway is the major ammonia-assimilating route. The GS of *Anabaena doliolum* was synthesized more under N<sub>2</sub>-fixing conditions, followed by ammonium, nitrate, and nitrite as nitrogen sources. The activities of both the glutamine synthetase, Mg<sup>2+</sup>-dependent biosynthetic and Mn<sup>2+</sup>-dependent  $\gamma$ -glutamyl transferase were optimum at pH 7. The active site of the enzyme bears sulfhydryl (-SH) groups; this was confirmed with the -SH group inhibitors, para-chloromercuribenzoate (pCMB) and N-ethylmaleimide (NEM). The biosynthetic and  $\gamma$ -glutamyl transferase activities showed specificity for the divalent cations, Mg<sup>2+</sup> and Mn<sup>2+</sup>, respectively. The other divalent cations Co<sup>2+</sup>, Cu<sup>2+</sup>, and Ni<sup>2+</sup> were poor substitutes. This enzyme also required these divalent cations to stabilize its structure and function under extreme conditions such as high and low temperatures and urea denaturation. The glutamate analog L-methionine-D,L-sulfoximine, inactivated the enzyme, whereas the GOGAT inhibitor, azaserine, had no effect on the enzyme activity. The GS enzyme required de novo protein synthesis.


Inorganic nitrogen is assimilated via the glutamine synthetase/glutamate synthase pathway under conditions of a limited supply of ammonia [2–6, 8, 9, 11, 13, 14, 18, 24, 26, 29]. NH<sub>4</sub><sup>+</sup> incorporation into glutamine is catalyzed by glutamine synthetase (GS) followed by the reductive reaction of glutamine with  $\alpha$ -oxoglutarate to yield two molecules of glutamate, the reaction catalyzed by glutamate synthase (GOGAT) [2–6, 9, 11, 13, 14, 18, 24, 26, 29]. In cyanobacteria, this pathway has been shown to be the major ammonia-assimilating route [5, 8, 13, 18, 24, 26, 29].

Glutamine synthetase catalyzes a number of reactions [15, 17, 25] forming the basis of a number of different assays.

(1) Biosynthetic reaction:



(2) Transferase reaction:



Biologically, the most significant reaction catalyzed by GS is the biosynthetic reaction involving the conversion of L-glutamate to L-glutamine, utilizing the available ammonia.

The work reported in this paper represents isolation and biochemical characterization of glutamine synthetase biosynthetic and  $\gamma$ -glutamyl transferase activities from the diazotrophic, filamentous cyanobacterium *Anabaena doliolum*.

### Materials and Methods

**Organism and culture conditions.** Cells of *Anabaena doliolum* were grown in eight times diluted Allen and Arnon medium [1], without any nitrogen source and containing different nitrogen sources, viz., 0.1 mM ammonium chloride, 20 mM potassium nitrate, 1 mM sodium nitrite, and 2 mM glutamine; buffered with 50 mM 4-(hydroxyethyl)-1-piperazine ethanesulfonic acid (HEPES)/NaOH to pH 7.0. The cultures were grown at 28°C under illumination of 10 W/m<sup>2</sup> provided by white fluorescent tubes. Cyanobacterial cells were collected in the exponential phase and used for the preparation of the enzyme extract [20].

**Extraction and assay of glutamine synthetase enzyme.** Two hundred milliliters of cyanobacterial cultures in the exponential phase were centrifuged and washed with 30 mM Tris buffer (pH 8.0) containing 10 mM MgCl<sub>2</sub> · 6H<sub>2</sub>O, 1 mM MnCl<sub>2</sub> · 4H<sub>2</sub>O and resuspended in the same buffer [20]. The cells were broken by passage

Table 1. Effect of nitrogen source on the activity of glutamine synthetase biosynthetic and  $\gamma$ -glutamyl transferase activity

Nitrogen source	Biosynthetic activity <sup>a</sup>	$\gamma$ -Glutamyl transferase activity <sup>a</sup>
N <sub>2</sub> -fixing	4.7	25.9
NH <sub>4</sub> <sup>+</sup> grown	2.4	13.3
NO <sub>3</sub> <sup>-</sup> grown	1.4	7.7
NO <sub>2</sub> <sup>-</sup> grown	0.8	4.6
Glutamine grown	0.03	20.9

<sup>a</sup> An average of three independent readings was taken into consideration and the activity expressed as  $\mu$ moles product formed/mg protein per min.

through a chilled French Pressure Cell twice at 20,000 lb/in<sup>2</sup> [20]. The homogenate was centrifuged at 6000 *g* for 20 min, and the supernatant thus obtained was used for the estimation of glutamine synthetase biosynthetic and  $\gamma$ -glutamyl transferase activities.

The Mn<sup>2+</sup>-dependent  $\gamma$ -glutamyl transferase was estimated by determining the rate of  $\gamma$ -glutamylhydroxamate formed [21]. The Mg<sup>2+</sup>-dependent biosynthetic activity of glutamine synthetase was determined from the rate of release of Pi [21].

**Analytical methods.** Protein contents in the extracts were mea-

sured with bovine serum albumin (BSA) as the standard protein [12].

Various inhibitors were first dissolved in the minimum possible volumes of ethanol diluted with distilled water or only in distilled water to obtain stock solutions. Stock solutions were filter-sterilized before being added to the cultures.

**Chemicals.** All chemicals at their analytical grades were obtained from Sigma Chemical Co. (USA), BDH, and SRL (India).

## Results

### Influence of different nitrogen sources on GS activity.

$\gamma$ -Glutamyl transferase activity was maximum in extracts obtained from N<sub>2</sub>-fixing cultures, followed by glutamine, ammonium, nitrate, and nitrite (Table 1). The GS biosynthetic activity was nearly five to six times lower than the transferase activity with representative response of nitrogen sources similar to the transferase activity.

### Effect of pH on the GS $\gamma$ -glutamyl transferase and biosynthetic activity.

Enzyme activities were optimum at pH 7. The biosynthetic activity showed a gradual decline in activity in the alkaline range. The transferase activity, on the other hand, had a




Fig. 1. Effect of pH on glutamine synthetase biosynthetic and transferase activities.




Fig. 2. (a) Rate of glutamine synthetase biosynthetic and transferase activities at varying temperatures of the enzyme. (b) Rate of glutamine synthetase biosynthetic and transferase activities at varying temperatures of the enzyme and reaction mixture.

much sharper drop on either side of the optimal pH (Fig. 1).

**Effect of temperature on the enzyme activity.** The enzyme activity was assayed over a wide range of temperature (10°–70°C). In one set, the enzyme was

incubated at these temperatures for a suitable time period before being added to the reaction mixture, and the GS activity was assayed (Fig. 2a). In the second set, both the enzyme and the reaction mixture were incubated at different temperatures and the activity estimated (Fig. 2b). In both sets, the

Table 2. Effect of various divalent cations on glutamine synthetase biosynthetic and  $\gamma$ -glutamyl transferase activities

Metal ion	Biosynthetic activity <sup>a</sup> metal ion (mM) <sup>b</sup>	Transferase activity <sup>a</sup> metal ion (mM) <sup>c</sup>
Mg <sup>2+</sup>	11.3	2.17
Mn <sup>2+</sup>	1.1	25.9
Co <sup>2+</sup>	0.013	2.19
Cu <sup>2+</sup>	0.0	2.58
Ca <sup>2+</sup>	0.0	0.0
Ni <sup>2+</sup>	0.0	0.0

<sup>a</sup> An average of three independent readings was taken into consideration in terms of  $\mu$ M product formed/mg protein per min.

<sup>b</sup> 125 mM of other divalent cations was added in place of Mg<sup>2+</sup> in the reaction mixture, and the enzyme activity was determined.

<sup>c</sup> 0.3 mM of other divalent cations was added in place of Mn<sup>2+</sup> in the reaction mixture, and the enzyme activity was determined.

temperature required for optimal  $\gamma$ -glutamyl transferase and biosynthetic activity was 30°C.

**Divalent cation specificity.** The relative effectiveness of various divalent cations in catalyzing the biosynthetic and  $\gamma$ -glutamyl transferase reaction is shown in Table 2. Maximum transferase activity was recorded at 0.3 mM Mn<sup>2+</sup>, whereas Mg<sup>2+</sup>, Co<sup>2+</sup>, and Cu<sup>2+</sup> expressed low activity at the same concentration. Ca<sup>2+</sup> and Ni<sup>2+</sup> did not support the enzyme activity.

Biosynthetic activity was maximum with 125 mM Mg<sup>2+</sup>, Mn<sup>2+</sup> and Co<sup>2+</sup> showed very little biosynthetic activity at the same concentration. Cu<sup>2+</sup>, Ca<sup>2+</sup>, and Ni<sup>2+</sup> showed no activity (Table 2).

**Effect of urea on GS activity.** Urea with increasing concentration was very effective in denaturing the native enzyme, resulting in loss of transferase as well as biosynthetic activity (Fig. 3a and b). Both Mg<sup>2+</sup> and Mn<sup>2+</sup> were effective in partially reducing the denaturation caused by urea. Mg<sup>2+</sup> gave 40% protection to the biosynthetic enzyme with 5 mM urea, while Mn<sup>2+</sup> was much more effective and gave 60% protection with same urea concentration.

**Effect of inhibitors on GS activity.** Inhibition of GS activity by L-methionine-D,L-sulfoximine (MSX) was concentration dependent from 1 to 4  $\mu$ M. Azaserine, an inhibitor of GOGAT, did not inhibit the glutamine synthetase activity. The enzyme activity was arrested by the protein synthesis inhibitor, chloramphenicol. The -SH group blocker, para-chloromercuribenzoate (pCMB), and the thiol reagent, N-ethylmaleimide, blocked the enzyme activity, in-

Table 3. Effect of various inhibitors on the activity of glutamine synthetase enzyme

Additions	% residual activity
Control	100
MSX (1 $\mu$ M)	2.4 (97.6) <sup>a</sup>
MSX (2 $\mu$ M)	0.8 (99.2)
MSX (4 $\mu$ M)	0.1 (99.1)
Azaserine (4 $\mu$ M)	98.9 (1.1)
Chloramphenicol (200 $\mu$ g/ml)	2.1 (97.9)
pCMB (0.1 mM)	1.6 (98.4)
NEM (2.5 mM)	6.8 (93.2)

<sup>a</sup> Percent inhibition given in parentheses.

dicating that -SH groups are present at the active site of the enzyme.

## Discussion

The data presented here are on partial characterization of the biochemical parameters of glutamine synthetase from *Anabaena doliolum*. A variety of effectors in the cellular milieu have been proposed to have a regulatory role.

Glutamine synthetase (GS) and glutamate synthase are the key ammonia-assimilating enzymes in cyanobacteria [8, 18, 19, 26, 29]. The enzyme was synthesized more under N<sub>2</sub>-fixing conditions, followed by ammonium, nitrate, and nitrite. Similar inhibitory response of GS synthesis to the different nitrogen sources was also observed in *Anabaena variabilis* [19], *Anabaena cylindrica* [18], and *Nostoc muscorum* [22].

The pH optimum of 7.0 for both Mg<sup>2+</sup>-dependent biosynthetic and Mn<sup>2+</sup>-dependent transferase activities resembled that for the enzyme from the cyanobacterium *Anacystis nidulans* [7] and from animal, plant, and bacterial sources [16, 28], but differed slightly from that in the cyanobacteria *A. cylindrica* [20] and *Anabaena* L-31 [27], where the pH optimum was around 7.8.

GS in *Anabaena doliolum* has been found to be much more stable in the presence of divalent cations, Mg<sup>2+</sup> and Mn<sup>2+</sup>. This was evident when the activities were compared in the presence and absence of divalent cations at high and low temperatures with urea treatment. It was suggested that under these denaturing conditions, the -SH groups on the enzyme become more exposed after the removal of the divalent cations, resulting in the conversion of the active enzyme to a relaxed state. The divalent cat-




Fig. 3. (a) Percentage inhibition of the  $\gamma$ -glutamyl transferase activity by urea and the alleviation of the inhibition by manganese. (b) Percentage inhibition of the glutamine synthetase biosynthetic activity by urea and its alleviation by magnesium.

ions,  $Mg^{2+}$  and  $Mn^{2+}$ , decrease the denaturation owing to their capacity to induce aggregation of the enzyme molecules, causing less exposure of the vulnerable portions of the enzyme to the external influences by reducing the surface area [27, 28]. These observations are in agreement with the results for the GS from *E. coli* [10, 21], *A. cylindrica* [20], and *Anabaena* L-31 [27].

The important role of the -SH groups present at the active site of the enzyme is further confirmed by pCMB and NEM, which block the -SH groups and inhibit the enzyme activity. *Anabaena* L-31 GS also showed inhibition by pCMB [27].

$Mg^{2+}$  and  $Mn^{2+}$  stabilize the enzyme structure and are specific for biosynthetic and  $\gamma$ -glutamyl transferase activities respectively [20, 23, 27]. This

was further confirmed with other divalent cations,  $\text{Co}^{2+}$ ,  $\text{Cu}^{2+}$ ,  $\text{Ca}^{2+}$ , and  $\text{Ni}^{2+}$ . These cations proved poor substitutes, possibly because they cannot regulate the enzyme structure and, hence, the catalytic function of the enzyme. A similar role of divalent cations has been observed in *Anabaena* CA [23], *A. cylindrica* [20], and *Anabaena* L-31 [27].

The analog of glutamate, MSX, competes with it for a common binding site on the catalytic part of the enzyme. Once firmly and irreversibly bound, it causes irreversible inactivation of the enzyme activity [8, 22, 24]. In contrast to MSX, which decreased the intracellular glutamine pool, azaserine, a specific glutamate synthase inhibitor, increased the glutamine pool and decreased the glutamate pool with no effect on GS activity [8, 18]. Lastly, the inhibition of GS activity with chloramphenicol suggested that the enzyme system was synthesized de novo on the ribosomes [18, 27].

Thus, studies on the regulation of glutamine synthetase in *A. doliolum*, a major inorganic nitrogen assimilatory enzyme, were crucial to be able to understand how this enzyme regulated the production of a diverse array of nitrogenous compounds essential for the maintenance and survival of living systems.

#### ACKNOWLEDGMENT

We express our gratitude to the Council of Scientific and Industrial Research, New Delhi, for awarding the Senior Research Fellowship to S. Shanthi.

#### Literature Cited

- Allen MB, Arnon DI (1955) Studies on nitrogen-fixing blue-green algae I. Growth and nitrogen fixation by *Anabaena cylindrica*. *Lemm Plant Physiol* 30:366–372
- Bender RA, Janssen KA, Resnick AD, Blumenberg M, Foor F, Magasanik B (1977) Biochemical parameters of glutamine synthetase from *Klebsiella aerogenes*. *J Bacteriol* 129:1001–1009
- Brown CM, MacDonald-Brown DS, Meers JL (1974) Physiological aspects of microbial inorganic nitrogen metabolism. *Adv Microb Physiol* 11:1–52
- Dharmawardene MWN, Stewart WDP, Stanley SO (1972) Nitrogenase activity, amino acid pool patterns and amination in blue-green algae. *Planta* 108:133–145
- Dharmawardene MWN, Haystead A, Stewart WDP (1973) Glutamine synthetase of the nitrogen-fixing alga *Anabaena cylindrica*. *Arch Mikrobiol* 90:281–295
- Deuel TF, Prusiner S (1974) Regulation of glutamine synthetase from *Bacillus subtilis* by divalent cations, feedback inhibitors and L-glutamine. *J Biol Chem* 249:257–264
- Edmond D, Rondeau N, Cedergreen RJ (1979) Distinctive properties of glutamine synthetase from the cyanobacterium *Anacystis nidulans*. *Can J Biochem* 57:843–851
- Flores E, Ramos JL, Herrero A, Guerrero MG (1983) Nitrate assimilation by cyanobacteria. In: Papageorgiou GC, Packer L (eds) *Photosynthetic prokaryotes: cell differentiation and function*, pp. 363–384
- Haystead A, Dharmawardene MWN, Stewart WDP (1973) Ammonia assimilation in a nitrogen-fixing blue-green alga. *Pl Sci Lett* 1:439–445
- Kingdom HS, Shapiro BM, Stadtman ER (1967) Regulation of glutamine synthetase VIII ATP: glutamine synthetase adenyltransferase, an enzyme that catalyzes alterations in the regulatory properties of glutamine synthetase. *Proc Natl Acad Sci USA* 58:1703–1710
- Lea PJ, Mifflin BJ (1975) Glutamine (amide); 2-oxoglutarate aminotransferase in blue-green algae. *Biochem Soc Trans* 3:381–384
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ (1951) Protein measurements by Folin phenol reagent. *J Biol Chem* 193:266–275
- Meeks JC, Wolk CP, Thomas J, Lockau W, Shaffer PW, Austin SM, Chien WS, Galonsky A (1977) The pathways of assimilation of  $\text{NH}_4^+$  by the cyanobacterium *Anabaena cylindrica*. *J Biol Chem* 252:7894–7900
- Mifflin BJ, Lea PJ (1976) The pathway of nitrogen assimilation in plants. *Phytochemistry* 15:873–885
- Miller RE, Shelton E, Stadtman ER (1974) Zinc-induced paracrystalline aggregation of glutamine synthetase. *Arch Biochem Biophys* 163:155–171
- O'Neal D, Jay KW (1973) Glutamine synthetase of pea leaves. I. Purification, stabilisation and pH optima. *Arch Biochem Biophys* 159:113–122
- Orr J, Haselkorn R (1981) Kinetic and inhibition studies of glutamine synthetase from the cyanobacterium *Anabaena* 7120. *J Biol Chem* 256:13099–13104
- Rowell P, Enticott S, Stewart WDP (1977) Glutamine synthetase and nitrogenase activity in the blue-green alga *Anabaena cylindrica*. *New Phytologist* 79:41–54
- Sakhurieva GN, Polukhina LE, Shestakov SV (1981) Glutamine synthetase in *Anabaena variabilis* mutants with derepressed nitrogenase. *Mikrobiologiya* 51:258–262
- Sawhney SK, Nicholas DJD (1978) Effects of amino acids, adenine nucleotides and inorganic pyrophosphate on glutamine synthetase from *Anabaena cylindrica*. *Biochim Biophys Acta* 527:485–496
- Shapiro BM, Stadtman ER (1970) Glutamine synthetase (*Escherichia coli*). *Methods Enzymol* 17A:910–922
- Singh HN, Singh RK, Sharma R (1983) An L-methionine-D,L-sulfoximine-resistant mutant of the cyanobacterium *Nostoc muscorum* showing inhibitor resistant  $\gamma$ -glutamyl transferase, defective glutamine synthetase and producing extracellular ammonia during  $\text{N}_2$  fixation. *FEBS Lett* 154:10–14
- Stacey G, Tabita FR, Van Baalen C (1977) Nitrogen and ammonia assimilation in the cyanobacteria. Purification of glutamine synthetase from *Anabaena* sp. strain CA. *J Bacteriol* 132:596–603
- Stewart WDP, Rowell P (1975) Effects of L-methionine-D,L-sulfoximine on the assimilation of newly fixed  $\text{NH}_3$ , acetylene reduction and heterocyst production in *Anabaena cylindrica*. *Biochem Biophys Res Commun* 65:846–856
- Tempest DW, Meers JL, Brown CM (1970) Influence of environment on the content and composition of microbial free amino acid pools. *J Gen Microbiol* 64:171–185

26. Thomas J, Wolk CP, Shaffer PW, Austin SM, Galonsky A (1975) The initial organic products of fixation of  $^{15}\text{N}$ -labeled nitrogen gas by the blue-green alga *Anabaena cylindrica*. *Biochem Biophys Res Commun* 67:501–507
27. Tuli R, Thomas J (1981) In vivo regulation of glutamine synthetase by ammonium in the cyanobacterium *Anabaena* L-31. *Arch Biochem Biophys* 206:181–189
28. Wedler FC, Hoffmann FM (1974) Glutamine synthetase of *Bacillus stearothermophilus*. I. Purification and basic properties. *Biochemistry* 13:3207–3214
29. Wolk CP, Thomas J, Shaffer PW, Austin SM, Galonsky A (1976) Pathway of nitrogen metabolism after fixation of  $^{15}\text{N}$ -labeled nitrogen gas by the cyanobacterium, *Anabaena cylindrica*. *J Biol Chem* 251:5027–5034