

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/243588181>

Ab-initio study of singlet and triplet excitation energies in oligothiophenes [phys. stat. sol. (c)1, No. 3, 539-542 (2004)]

ARTICLE *in* PHYSICA STATUS SOLIDI (C) · NOVEMBER 2004

DOI: 10.1002/pssc.200404034

READS

15

3 AUTHORS, INCLUDING:



Eduardo Fabiano

Italian National Research Council

74 PUBLICATIONS 1,220 CITATIONS

SEE PROFILE



Fabio Della Sala

Italian National Research Council

182 PUBLICATIONS 5,057 CITATIONS

SEE PROFILE

Erratum

Ab-initio study of singlet and triplet excitation energies in oligothiophenes

[phys. stat. sol. (c) 1, No. 3, 539–542 (2004)]

E. Fabiano, F. Della Sala*, and R. Cingolani

National Nanotechnology Laboratory, INFN-Dipartimento di Ingegneria dell'Innovazione,
Università di Lecce, Via per Arnesano, 73100 Lecce, Italy

Received 4 November 2004

Published online 22 November 2004

PACS 71.15.Mb, 78.40.Me, 79.60.Fr, 99.10.Cd

The authors report an error in the fit for the 3A state shown in Fig. 1 of their article and present a new corrected figure.

© 2004 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim

In Fig. 1 we reported linear fits of the calculated excitation energies for T1, T2, T3, and T4 and compared them with experimental results. We have now found that the fit for the 3A state is not valid because the 3A state of the monomer has a different character as compared to its oligomers. Here we report in Fig. 1 the new linear fits for all the states considering only excitation energies for T2, T3, and T4. While similar results are obtained for the 1B and 3B states, a qualitatively different behavior is found for 3A state: the second triplet excited-state lies energetically below the first singlet excited-state for long oligomers. In order to verify this trend, we have calculated the excitation energies of pentathiophene (T5) and we found that the 3A state lies indeed 0.26 eV below the 1B state. This new finding shows a discrepancy between the theoretical and the experimental evolution with the number of monomers of the 3A state. However all the other results and conclusions of the article are still valid due to the very small energetical difference between the 1B and 3A states.




Fig. 1 Theoretical (lines) and experimental (points) energy of the 1B , 3B and 3A excited states as a function of inverse of oligomer size $1/n$.

* Corresponding author: e-mail: fabio.dellasala@unile.it