

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/234142703>

Introduction of histidine units using benzotriazolide activation

ARTICLE *in* JOURNAL OF PEPTIDE SCIENCE · FEBRUARY 2013

Impact Factor: 1.55 · DOI: 10.1002/psc.2483 · Source: PubMed

CITATIONS

4

READS

58

5 AUTHORS, INCLUDING:



[Kiran Bajaj](#)

Birla Institute of Technology and Science Pi...

29 PUBLICATIONS 301 CITATIONS

SEE PROFILE



[Siva S. Panda](#)

Georgia Regents University

82 PUBLICATIONS 315 CITATIONS

SEE PROFILE



[Mohamed Ibrahim](#)

University of Florida

16 PUBLICATIONS 51 CITATIONS

SEE PROFILE



Introduction of histidine units using benzotriazole activation

Kiran Bajaj,^a Siva S. Panda,^a Mohamed A. Ibrahim,^{a,b,c} Said A. El-Feky^b and Alan R. Katritzky^{a,d*}

***N^α*-Boc-*N^{im}*-(4-toluenesulfonyl-L-histidylbenzotriazole) enables convenient acylation of *N*-, *O*-, *S*-, and *C*-nucleophiles with no detectable racemization. We report efficient syntheses of novel histidine-containing di-, tri-, and tetra-peptides and models for the preparation of potentially biologically active histidine *N*-, *O*-, *S*-, and *C*-conjugates. Copyright © 2013 European Peptide Society and John Wiley & Sons, Ltd.**

Supporting information can be found in the online version of this article.

Keywords: histidine; conjugates; benzotriazole; acylation

Introduction

Histidine is of prime importance as an essential component of the active site of many enzymes [1], which is widely distributed in nature, and its derivatives exhibit diverse biological activities [2,3]: for example, naphthalene-substituted histidine derivatives act as useful antihypertensive agents [4], a histidine-estrone conjugate has pronounced anti-tumor activity [5], and histidine-tocopherol conjugates are antioxidants and prodrug delivery vehicles [6]. Carnosine (β -alanyl-L-histidine) and carbinine (β -alanyl-histamine) occur in the non-protein fraction of mammalian tissues [7]. In proteins, histidine residues play an important role in interactions with metal ions because of the presence of three coordinating groups. Histidine coordinates strongly and forms six membered rings in which a metal is attached to an imidazole nitrogen atom and the α -amino group [8]. For example, the N-terminus of the protein bovine serum albumin (Asp-Thr-His-Lys-OH) specifically binds the Cu(II) ion because of the histidine residue [9]. Histidine-containing peptides act as catalysts in Michael addition of nitroalkanes [10], direct asymmetric aldol reactions [11], and asymmetric synthesis of cyanohydrins [12].

Histidine residues are often involved in neutral hydrolytic metalloenzyme active centers and have the potential to bind DNA. The unprotonated imidazole (pH > 7) is nucleophilic, whereas the protonated form (pH < 7) serves as a proton donor.

Histidine is often susceptible to racemization during solid-phase synthesis (2.1% racemization detected for the L-His-containing peptide and 1.9% detected for the D-His-containing peptide) [13]. Ishiguro *et al.* reported unexpected chain-terminating side reactions caused by histidine in solid-phase peptide synthesis [14]. Racemization during the synthesis of polypeptides could be a serious problem as a high degree of steric homogeneity is usually required for biological or physical studies. The detection of small amounts of diastereoisomeric impurities and their removal from synthetic polypeptide preparations are often difficult; thus, synthetic procedures that minimize the risk of racemization are commonly used. In a histidine unit, the presence of the imidazole

ring and an amino group has complicated the development of synthetic methods to prepare histidine C-terminus derivatives. The nucleophilic imidazole ring of histidine easily undergoes *N*-acylation in absence of appropriate protecting groups, which can result in side reactions such as acyl transfer. In particular, histidine derivatives are known to be prone to racemization during activating and coupling processes involving the π -nitrogen of the imidazole ring [15]. Thus, several protecting groups as well as activating reagents have been introduced in attempts to solve problems such as the regioselective protection of the π -nitrogen, and this enables avoidance of racemization. For this purpose, the *N^{tr}*-benzyloxymethyl (Bom) group has been widely employed in Boc chemistry [16], and the *N^{tr}*-Trt protecting group [17], used in combination with *N^{tr}*-Fmoc protection [17,18]. Although the steric hindrance of the Trt group may help reduce racemization to a certain extent, *N^{tr}*-protecting groups cannot inherently prevent racemization. Other *N^{tr}*-protecting groups including *N^{tr}*-*t*-butoxymethylhistidine [His(Bum)] [19], *N^{tr}*-1-adamantyloxymethylhistidine [His(1-Adom)] [20], and 4-methoxybenzyloxymethyl (MBom) [21,22] have been introduced. The use of several activating agents has been reported in literature including (i) solution phase synthesis of a histidine-containing dipeptide derivative from Z-Phe-OH and H-His-OMe using isobutyl chloroformate [12], (ii) preparation of the histidine dipeptide (Boc-His(Tos)-Leu-OH) from Boc-His(Tos)-

* Correspondence to: Alan R. Katritzky, Center for Heterocyclic Compounds, Department of Chemistry, University of Florida, Gainesville, FL 32611-7200, USA. E-mail: katritzky@chem.ufl.edu

a Center for Heterocyclic Compounds, Department of Chemistry, University of Florida, Gainesville, FL 32611-7200, USA

b Department of Pharmaceutical Organic Chemistry, Faculty of Pharmacy, Zagazig University, Zagazig 44519, Egypt

c Department of Organic Chemistry, College of Pharmacy, Misr University for Science and Technology, Al-Motamayez District, P.O. Box 77, Egypt

d Chemistry Department, King Abdulaziz University, Jeddah 21589, Saudi Arabia

OSu and H-Leu-OH [23], (iii) dipeptide (Boc-Ala-His(Tos)-OMe) from protected histidine and alanine in the presence of EDCI and Et₃N [24], and (iv) dipeptide (Boc-His(MBS)-Gly-OBzl) utilizing *p*-TsOH, and diethyl phosphorocyanidate as a coupling agent [25]. Literature routes to C-terminus histidine conjugates include the use of the coupling reagents (i) DCC with DMAP [26,27], (ii) EDCI-HOBt [28], and (iii) isobutyl chloroformate [29]. These methods showed good yields, but products needed column purification. Additionally, some reports indicate that histidine derivatives racemize on activation with DCC [30].

We have extensively applied *N*-acylbenzotriazoles for *N*-acylation, *C*-acylation, *O*-acylation, and *S*-acylation [31]. In these reactions, *N*-(*Z*-aminoacyl)benzotriazoles coupled with unprotected amino acids to produce peptides in 85–95% yield with minimal racemization [32]. *N*-(*Z*- α -aminoacyl)benzotriazoles prepared from *N*-protected α -amino acids with unfunctionalized side chains such as Ala, Val, and Phe coupled successfully with unprotected amino acids (Ala, Val, Phe, Ser, Trp) in the presence of Et₃N in partially aqueous solution (CH₃CN/H₂O). We have also prepared *N*-(*Z*- or Fmoc- α -aminoacyl)benzotriazoles derived from Tyr, Trp, Cys, Met, and Gln containing unprotected side-chain functionality [33], and successfully coupled them with α -amino acids (L-Ala, L-Phe) in partially aqueous medium to form the expected peptides. NMR and HPLC analysis supported the preservation of the original chirality. These findings were further confirmed by NMR and HPLC comparisons of the diastereomeric dipeptides prepared by coupling *N*-(*Z*-Tyr, Trp, Cys, Met, Gln and Fmoc-Trp, Met acyl)benzotriazole with H-DL-Ala-OH, H-DL-Phe-OH [33]. We have also reported successful benzotriazole activation in racemization free tri- and tetra-peptides [31]. Herein, we report another method to introduce histidine residues at the N-terminus in solution phase peptide synthesis by utilization of benzotriazole-activated histidine intermediate **3a** and **3b** (Scheme 1). Compound **3a** was also used to synthesize novel histidine conjugates by *N*-, *O*-, *S*-, and *C*-acylations. There is a previous reference [23] for the preparation of **5c** (Scheme 2), but no experimental details are given.

Materials and Methods

Melting points were determined on a capillary point apparatus equipped with a digital thermometer. NMR spectra were recorded in CDCl₃, DMSO-*d*₆, or CD₃OD-*d*₄ on Mercury, Gemini, or Varian NMR spectrometers operating at 300 MHz for ¹H (with TMS as an internal standard) and 75 MHz for ¹³C NMR. Elemental analyses were performed on a Carlo Erba-EA1108 instrument. All microwave-assisted reactions were carried out with a single mode cavity Discover Microwave Synthesizer (CEM Corporation, NC, USA). The reaction mixtures were transferred into a 10 mL glass pressure microwave tube equipped with a magnetic stirrer bar. The tube was closed with a silicon septum, and the reaction mixture was subjected to microwave irradiation (Discover mode; run time: 60 s; PowerMax-cooling mode).

Preparation of *N*^α-Boc-*N*^{im}-protected-L-histidylbenzotriazole (**3a,b**)

Compounds **3a,b** were synthesized by irradiating an equimolar amount of *N*^α-Boc-*N*^{im}-protected-L-histidine with 1-(methylsulfonyl)-1*H*-benzo[d][1,2,3]triazole (BtSO₂Me) in the presence of 1.0 eq. of Et₃N in anhydrous DMF for 30 min at 50 °C and 20 W

irradiation power. Completion of the reaction was monitored by TLC. After completion, the reaction mixture was quenched with water. The precipitate obtained was washed with saturated solution of Na₂CO₃ to afford the desired products, which were further crystallized from methylene chloride/hexane.

(*S*)-*tert*-Butyl-(1-(1*H*-benzo[d][1,2,3]triazol-1-yl)-1-oxo-3-(1-tosyl-1*H*-imidazol-4-yl)propan-2-yl)carbamate (Boc-His(Tos)-Bt, **3a**)

White microcrystals (65%); mp 150–152 °C; ¹H NMR (CDCl₃) δ 8.18 (d, *J* = 8.4 Hz, 1H), 8.13 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J* = 1.5 Hz, 1H), 7.75 (d, *J* = 8.4 Hz, 2H), 7.66 (t, *J* = 7.5 Hz, 1H), 7.53 (t, *J* = 7.5 Hz, 1H), 7.34 (d, *J* = 8.4 Hz, 2H), 7.01 (s, 1H), 6.12–6.06 (m, 1H), 5.96–5.84 (m, 1H), 3.40–3.22 (m, 2H), 2.44 (s, 3H), 1.44 (s, 9H); ¹³C NMR (CDCl₃) δ 170.7, 155.6, 146.5, 146.1, 139.6, 136.8, 134.9, 131.3, 130.8, 130.6, 127.5, 126.6, 120.5, 115.1, 114.6, 80.6, 54.3, 30.8, 28.5, 22.0. HRMS (+ESI-TOF): *m/z* for C₂₄H₂₆N₆O₅S [M + H]⁺ calcd 511.1758, found 511.1749.

(*S*)-*tert*-Butyl-1-(1*H*-benzo[d][1,2,3]triazol-1-yl)-3-(1-benzyl-1*H*-imidazol-4-yl)-1-oxopropan-2-ylcarbamate (Boc-His(Bzl)-Bt, **3b**)

White microcrystals (62%); mp 85–87 °C; ¹H NMR (CDCl₃) δ 8.16 (d, *J* = 8.2 Hz, 1H), 8.07 (d, *J* = 8.2 Hz, 1H), 7.59 (t, *J* = 7.9 Hz, 1H), 7.50–7.45 (m, 2H), 7.32–7.26 (m, 3H), 7.05–7.03 (m, 2H), 6.63 (d, *J* = 7.7 Hz, 1H), 6.55 (s, 1H), 5.94–5.87 (m, 1H), 4.98 (s, 2H), 3.48–3.27 (m, 2H), 1.44 (s, 9H); ¹³C NMR (CDCl₃) δ 170.9, 155.6, 145.7, 137.4, 137.0, 135.9, 131.1, 130.4, 128.8, 128.1, 127.1, 126.2, 120.0, 117.2, 114.3, 79.9, 54.8, 50.7, 30.4, 28.3.

General Procedure for the Preparation of Histidine-containing di-, tri-, and tetra-peptides (**5a–f**)

To a solution of compound **3** (1.0 eq.) in acetonitrile, a solution of amino acid, dipeptide or tripeptide (1.0 eq.), and Et₃N (1.2 eq.) in water was added dropwise, and the reaction mixture was stirred at room temperature for 3–4 h. After complete consumption of compound **3** (TLC), the solution was concentrated under reduced pressure. The solution was neutralized by 4N HCl and extracted with ethyl acetate, dried over anhydrous MgSO₄, and evaporated under reduced pressure to afford a white solid, which was either crystallized from diethyl ether or purified by flash column chromatography with 30% ethyl acetate–hexane solution as eluent.

(*S*)-2-((*S*)-2-((*tert*-Butoxycarbonyl)amino)-3-(1-tosyl-1*H*-imidazol-4-yl)propanamido)-3-phenylpropanoic acid (Boc-L-His(Tos)-L-Phe-OH, **5a**)

White microcrystals (79%); mp 133–134 °C; ¹H NMR (CDCl₃) δ 7.98 (s, 1H), 7.81 (d, *J* = 6.9 Hz, 2H), 7.38–7.29 (m, 3H), 7.26–7.18 (m, 5H), 7.09 (s, 1H), 5.39 (d, *J* = 8.7 Hz, 1H), 4.76–4.68 (m, 1H), 4.45 (t, *J* = 9.0 Hz, 1H), 3.20–2.92 (m, 3H), 2.53–2.41 (m, 1H), 2.42 (s, 3H), 1.35 (s, 9H); ¹³C NMR (CDCl₃) δ 174.3, 170.9, 155.2, 146.9, 139.9, 136.5, 134.6, 130.7, 129.9, 128.5, 127.7, 127.1, 115.6, 79.6, 54.2, 53.7, 38.7, 32.8, 28.4, 22.0; Anal. calcd for C₂₇H₃₂N₄O₇S: C, 58.26; H, 5.79; N, 10.07; found: C, 58.15; H, 6.13; N, 10.09.

2-((S)-2-((tert-Butoxycarbonyl)amino)-3-(1-tosyl-1H-imidazol-4-yl)propanamido)-3-phenylpropanoic acid (Boc-His(Tos)-DL-Phe-OH, 5a+5a')

White microcrystals (72%); mp 175–177 °C; ¹H NMR (CDCl₃) δ 8.97–8.96 (d, *J* = 4.4 Hz, 1H), 8.16–8.07 (m, 1H), 7.50–7.47 (m, 2H), 7.27–6.98 (m, 8H), 4.51–4.43 (m, 1H), 4.32–4.25 (m, 1H), 3.19–2.64 (m, 4H), 2.29 (s, 3H), 1.32 (s, 9H). ¹³C NMR (CDCl₃) δ 172.6, 170.3, 170.1, 155.1, 145.5, 137.7, 137.3, 133.7, 129.5, 129.5, 129.2, 128.2, 128.1, 126.5, 125.5, 116.8, 116.6, 78.5, 53.3, 53.0, 36.8, 36.6, 28.1, 27.1, 20.8; Anal. calcd for C₂₇H₃₂N₄O₇S: C, 58.36; H, 5.79; N, 10.07; found: C, 57.87; H, 6.02; N, 9.83.

(S)-2-((S)-3-(1-Benzyl-1H-imidazol-4-yl)-2-(tert-butoxycarbonylamino)propanamido)-3-methylbutanoic acid (Boc-His(Tos)-L-Val-OH, 5b)

Oil (72%); ¹H NMR (CDCl₃) δ 8.04 (s, 1H), 7.82 (d, *J* = 8.1 Hz, 2H), 7.35 (d, *J* = 8.1 Hz, 1H), 7.11 (s, 1H), 5.52 (d, *J* = 8.7 Hz, 1H), 4.58–4.49 (m, 2H), 3.13 (d, *J* = 14.1 Hz, 1H), 2.61–2.50 (m, 1H), 2.43 (s, 3H), 2.25–2.19 (m, 1H), 1.31 (s, 9H), 0.98–0.92 (m, 6H). ¹³C NMR (CDCl₃) δ 174.4, 171.5, 155.4, 146.9, 139.9, 136.6, 134.6, 130.8, 130.7, 127.8, 115.6, 79.7, 57.8, 53.9, 32.4, 31.8, 28.4, 22.0, 18.8, 18.1. HRMS (+ESI-TOF): *m/z* for C₂₃H₃₂N₄O₅ [M + H]⁺ calcd 507.1919, found 507.1926.

2-((S)-3-(1-Benzyl-1H-imidazol-4-yl)-2-(tert-butoxycarbonylamino)propanamido)-3-methylbutanoic acid (Boc-His(Tos)-DL-Val-OH, 5b+5b')

Oil (70%); ¹H NMR (CDCl₃) δ 8.03 (s, 1H), 7.83 (d, *J* = 8.1 Hz, 2H), 7.36 (d, *J* = 8.1 Hz, 2H), 7.21–7.14 (m, 1H), 5.86–5.84 (m, 1H), 5.65–5.62 (m, 1H), 4.68–4.50 (m, 2H), 3.18–3.05 (m, 1H), 2.74–2.64 (m, 1H), 2.43 (s, 3H), 2.28–2.19 (m, 1H), 1.31 (d, *J* = 12.0 Hz, 9H), 0.96–0.90 (m, 6H). ¹³C NMR (CDCl₃) δ 174.6, 174.3, 171.6, 171.4, 155.6, 155.4, 146.8, 139.9, 136.5, 134.6, 130.7, 127.7, 115.6, 79.8, 79.7, 66.0, 57.7, 57.3, 53.9, 31.7, 31.6, 28.4, 21.9, 18.9, 18.8, 18.0, 17.9, 15.4. HRMS (+ESI-TOF): *m/z* for C₂₃H₃₂N₄O₅ [M + H]⁺ calcd 507.1919, found 507.1932.

(S)-2-((S)-2-((tert-Butoxycarbonyl)amino)-3-(1-tosyl-1H-imidazol-4-yl)propanamido)-4-methylpentanoic acid (Boc-L-His(Tos)-L-Leu-OH, 5c)

White microcrystals (89%); mp 121–123 °C [lit. mp 146–148 °C]¹⁶; ¹H NMR (CDCl₃) δ 8.02 (s, 1H), 7.83 (d, *J* = 8.4 Hz, 2H), 7.75 (d, *J* = 7.5 Hz, 1H), 7.36 (d, *J* = 8.4 Hz, 2H), 7.15 (s, 1H), 5.60 (d, *J* = 9.0 Hz, 1H), 4.64–4.52 (m, 2H), 3.14–3.05 (m, 1H), 2.68–2.56 (m, 1H), 2.43 (s, 3H), 1.78–1.56 (m, 3H), 1.32–1.22 (m, 9H), 0.96 (d, *J* = 6.3 Hz, 3H), 0.94 (d, *J* = 6.6 Hz, 3H); ¹³C NMR (CDCl₃) δ 175.6, 171.4, 155.4, 146.8, 139.9, 136.5, 134.6, 130.7, 127.7, 115.7, 79.7, 53.7, 51.5, 42.6, 32.3, 31.8, 28.3, 25.0, 22.8, 22.6, 21.9, 14.3.

2-((S)-2-((tert-Butoxycarbonylamino)-3-(1-tosyl-1H-imidazol-4-yl)propanamido)propanoic acid (Boc-L-His(Tos)-L-Ala-OH, 5d)

White microcrystals (92%); mp 151–153 °C; ¹H NMR (CDCl₃) δ 8.02 (s, 1H), 7.84–7.82 (d, *J* = 8.1 Hz, 2H), 7.38–7.35 (d, *J* = 8.1 Hz, 2H), 7.12 (s, 1H), 5.66–5.63 (d, *J* = 8.2 Hz, 1H), 5.46–5.44 (d, *J* = 7.9 Hz, 1H), 4.63–4.50 (m, 1H), 3.19–3.10 (m, 1H), 2.70–2.48 (m, 2H), 2.44 (s, 3H), 1.51–1.27 (m, 12H); ¹³C NMR (CDCl₃) δ 157.9, 171.3, 155.3, 146.9, 139.8, 136.5, 134.5, 130.8, 127.8, 115.7, 79.7, 53.6, 49.2, 48.8, 32.5, 28.3, 22.0, 18.9. HRMS (+ESI-TOF): *m/z* for C₂₁H₂₈N₄O₇S [M + Na]⁺ calcd 503.1571, found 503.1571.

(6S,12S)-12-Isopropyl-2,2-dimethyl-4,7,10-trioxo-6-((1-tosyl-1H-imidazol-4-yl)methyl)-3-oxa-5,8,11-triazatridecan-13-oic acid (Boc-L-His(Tos)-Gly-L-Val-OH, 5e)

White microcrystals (68%); mp 113–115 °C; ¹H NMR (CDCl₃) δ 8.02–8.00 (m, 1H), 7.87–7.74 (m, 3H), 7.45–7.14 (m, 5H), 5.95–5.87 (m, 1H), 4.49–4.39 (m, 1H), 4.11–3.78 (m, 2H), 3.08–2.93 (m, 2H), 2.42 (s, 3H), 2.42–2.13 (m, 1H), 1.33 (s, 9H), 0.94–0.90 (m, 6H); ¹³C NMR (CDCl₃) δ 174.7, 172.6, 169.8, 155.8, 146.8, 139.9, 136.8, 134.6, 130.7, 127.7, 126.0, 115.4, 80.3, 58.3, 58.1, 54.2, 43.4, 30.7, 28.4, 21.9, 19.4, 18.1. Anal. calcd for C₂₅H₃₅N₅O₈S: C, 53.08; H, 6.24; N, 12.38; found: C, 52.68; H, 6.43; N, 12.32.

(6S,15S)-15-((S)-sec-Butyl)-2,2-dimethyl-4,7,10,13-tetraoxo-6-((1-tosyl-1H-imidazol-4-yl)methyl)-3-oxa-5,8,11,14-tetraaza-hexadecan-16-oic acid (Boc-L-His(Tos)-Gly-Gly-L-Ile-OH, 5f)

White microcrystals (60%); mp 140–142 °C; ¹H NMR (DMSO-*d*₆) δ 12.62 (s, 1H), 8.28 (s, 1H), 8.22–8.10 (m, 2H), 7.92 (d, *J* = 8.1 Hz, 2H), 7.49 (d, *J* = 8.4 Hz, 2H), 7.37 (s, 1H), 6.88 (d, *J* = 8.1 Hz, 1H), 4.23–4.16 (m, 2H), 3.79 (t, *J* = 5.7 Hz, 2H), 3.66 (d, *J* = 5.4 Hz, 2H), 2.92–2.80 (m, 1H), 2.75–2.65 (m, 1H), 2.40 (s, 3H), 1.82–1.70 (m, 1H), 1.30 (s, 9H), 1.24–1.11 (m, 2H), 0.85–0.78 (m, 6H); ¹³C NMR (DMSO-*d*₆) δ 172.8, 171.5, 168.9, 168.6, 155.1, 146.3, 140.7, 136.7, 134.4, 130.6, 127.2, 114.7, 78.3, 56.2, 53.5, 42.2, 41.6, 36.4, 30.3, 28.1, 24.6, 21.2, 15.5, 11.3. Anal. calcd for C₂₈H₄₀N₆O₉S: C, 52.82; H, 6.33; N, 13.20; found: C, 52.88; H, 6.70; N, 13.09.

Procedure for Preparation of Compound 7

To a solution of compound **3b** (1.0 eq.) in acetonitrile, a solution of D-H-Phe-OMe (1.0 eq.) and Et₃N (1.2 eq.) in water was added dropwise, and the reaction mixture was stirred at room temperature for 3 h. After complete consumption of compound **3b** (TLC), the solution was concentrated under reduced pressure. The residue was extracted with ethyl acetate, dried over anhydrous MgSO₄, and evaporated under reduced pressure to afford the desired product as an oil.

Methyl 2-((S)-3-(1-benzyl-1H-imidazol-4-yl)-2-(tert-butoxycarbonylamino)propanamido)-3-phenylpropanoate (Boc-His(Bzl)-Phe-OMe, 7)

Oil (69%); ¹H NMR (CDCl₃) δ 7.42–7.38 (m, 1H), 7.34–7.16 (m, 7H), 7.13–7.06 (m, 3H), 6.96–6.94 (m, 1H), 6.71–6.63 (m, 1H), 6.39–6.27 (m, 1H), 4.97 (s, 2H), 4.79–4.75 (m, 1H), 3.69 (s, 1H), 3.63 (s, 1H), 3.59 (s, 1H), 3.06–2.84 (m, 4H), 1.40 (d, *J* = 4.0 Hz, 9H). ¹³C NMR (CDCl₃) δ 175.4, 171.6, 171.5, 171.4, 155.6, 138.5, 138.2, 136.7, 136.0, 135.8, 129.2, 128.9, 128.5, 128.4, 128.3, 128.2, 127.4, 127.3, 126.9, 126.8. HRMS (+ESI-TOF): *m/z* for C₂₈H₃₄N₄O₅ [M + H]⁺ calcd 507.2602, found 507.2604.

Procedure for Preparation of Compound 8a

N-(3-Aminopropyl)-imidazole (1.0 eq.) was dissolved in THF and Et₃N (1.5 eq.). *N*²-Boc-*N*^{1m}-4-toluenesulfonyl-histidylbenzotriazole **3a** (1.0 eq.) was added to the solution, and the mixture was stirred at room temperature for 3 h. The mixture was acidified with 4N HCl, concentrated and then diluted with ethyl acetate. The organic layer was washed with 10% Na₂CO₃ solution, water and dried over anhydrous MgSO₄, filtered, and evaporated to give a white solid, which was further washed with diethyl ether to give the desired compound.

(S)-tert-Butyl-1-((3-(1H-imidazol-1-yl)propyl)amino)-1-oxo-3-(1-tosyl-1H-imidazol-4-yl) propan-2-yl)carbamate (8a)

White microcrystals (92%); mp 64–66 °C; ^1H NMR (CDCl_3) δ 7.93 (s, 1H), 7.80 (d, J = 8.4 Hz, 2H), 7.46 (s, 1H), 7.35 (d, J = 8.1 Hz, 2H), 7.10 (s, 1H), 7.04 (s, 1H), 6.91 (br s, 2H), 5.93 (d, J = 7.2 Hz, 1H), 4.40–4.33 (m, 1H), 3.87 (t, J = 6.9 Hz, 2H), 3.14 (q, J = 6.3 Hz, 2H), 3.05 (dd, J = 14.7, 5.4 Hz, 1H), 2.90 (dd, J = 14.7, 5.4 Hz, 1H), 2.43 (s, 3H), 1.92–1.84 (m, 2H), 1.42 (s, 9H); ^{13}C NMR (CDCl_3) δ 171.7, 155.9, 146.7, 140.7, 137.3, 136.3, 134.8, 130.6, 129.7, 127.6, 119.0, 115.1, 80.5, 54.4, 44.2, 36.4, 31.2, 30.3, 28.5, 21.9; HRMS (+ESI-TOF): m/z for $\text{C}_{24}\text{H}_{33}\text{N}_6\text{O}_5\text{S}$ $[\text{M} + \text{H}]^+$ calcd 517.2228, found 517.2229.

Procedure for Preparation of Compound 8b

A dried heavy-walled Pyrex tube containing a small stir bar was mixed with compound **3a** (0.3 g, 0.587 mmol) and 2-aminopyridine (0.056 g, 0.587 mmol) in dry DMF (5 mL) for 30 min at 50 °C and 20 W irradiation power. After completion (TLC), the reaction mixture was quenched with water. The precipitate obtained was washed with saturated Na_2CO_3 solution and water to afford a solid, which was flash chromatographed with 30% ethyl acetate/hexane solution to give compound **8b**.

(S)-tert-Butyl-1-oxo-1-(pyridin-2-ylamino)-3-(1-tosyl-1H-imidazol-4-yl)propan-2-yl)carbamate (8b)

White microcrystals (91%); mp 87–88 °C; ^1H NMR (CDCl_3) δ 9.00 (br s, 1H), 8.24 (d, J = 4.5 Hz, 1H), 8.14 (d, J = 8.4 Hz, 1H), 7.94 (s, 1H), 7.73 (d, J = 8.1 Hz, 2H), 7.67 (t, J = 8.0 Hz, 1H), 7.25 (d, J = 8.4 Hz, 2H), 7.10 (s, 1H), 7.08–7.00 (m, 1H), 6.18–6.04 (m, 1H), 4.62–4.54 (m, 1H), 3.22–3.12 (m, 1H), 2.99 (dd, J = 15.0, 5.1 Hz, 1H), 2.39 (s, 3H), 1.44 (s, 9H); ^{13}C NMR (CDCl_3) δ 170.3, 155.4, 151.1, 147.9, 146.4, 140.4, 138.5, 136.6, 134.9, 130.5, 127.4, 120.0, 115.1, 114.2, 80.7, 55.1, 30.2, 28.4, 21.9; HRMS (+ESI-TOF): m/z for $\text{C}_{23}\text{H}_{28}\text{N}_5\text{O}_5\text{S}$ for $[\text{M} + \text{H}]^+$ calcd 486.1806, found 486.1796.

General Procedure for Preparation of Compound 8c,d

The *O*-nucleophile (0.783 mmol) was added to a solution of compound **3a** and DMAP (0.05 g, 0.392 mmol) in dry THF (2 mL). The mixture was irradiated in microwave at 60 °C for 60 min and 100 W. The solvent was evaporated, and the residue was purified over flash chromatography to give **8c** or **8d**.

(S)-4-Ethylphenyl 2-(tert-butoxycarbonylamino)-3-(1-tosyl-1H-imidazol-4-yl)propanoate (8c)

Oil (50%), ^1H NMR (CDCl_3) δ 7.96 (d, J = 1.3 Hz, 1H), 7.79 (d, J = 8.4 Hz, 2H), 7.32 (d, J = 8.1 Hz, 2H), 7.13 (d, J = 8.1 Hz, 3H), 6.87 (d, J = 8.4 Hz, 2H), 5.77 (d, J = 8.1 Hz, 1H), 4.77 (br s, 1H), 3.26–3.07 (m, 2H), 2.62 (q, J = 7.6 Hz, 2H), 2.42 (s, 3H), 1.43 (s, 9H), 1.22 (t, J = 7.5 Hz, 3H); ^{13}C NMR (CDCl_3) δ 170.6, 155.6, 148.6, 146.4, 142.0, 140.2, 136.8, 135.0, 130.6, 128.8, 127.5, 121.2, 115.0, 80.1, 53.2, 30.5, 30.4, 28.4, 21.9, 15.7. HRMS (+ESI-TOF) m/z for $\text{C}_{26}\text{H}_{32}\text{N}_3\text{O}_6\text{S}$ $[\text{M} + \text{H}]^+$ calcd 514.2006, found 514.2025.

(S)-Naphthalen-2-yl 2-(tert-butoxycarbonylamino)-3-(1-tosyl-1H-imidazol-4-yl)propanoate (8d)

Oil (55%). ^1H NMR (CDCl_3) δ 8.00 (d, J = 1.2 Hz, 1H), 7.84–7.76 (m, 5H), 7.51–7.46 (m, 3H), 7.27 (d, J = 7.8 Hz, 2H), 7.18 (s, 1H), 7.12 (dd, J = 8.9, 2.3 Hz, 1H), 5.78 (d, J = 8.1 Hz, 1H), 4.86–4.83 (m, 1H), 3.33–3.12 (m, 2H), 2.37 (s, 3H), 1.45 (s, 9H); ^{13}C NMR (CDCl_3) δ 170.6, 155.6, 148.3, 146.5, 140.2, 136.9, 135.0, 133.8, 131.7, 130.6, 129.6, 127.9, 127.5, 126.8, 126.0, 121.0, 118.6, 115.2, 80.3, 53.4, 30.5, 28.5, 21.9. HRMS (+ESI-TOF) m/z for $\text{C}_{28}\text{H}_{29}\text{N}_3\text{O}_6\text{S}$ $[\text{M} + \text{H}]^+$ calcd 536.1850, found 536.1850.

General Procedure for Preparation of Compound 8e,f

The mercapto nucleophile (1.0 eq.) was dissolved in THF and Et_3N (1.5 eq.). N^Z -Boc- N^{im} -4-toluenesulfonyl-L-histidinylbenzotriazole **3a** (1.0 eq.) was added to the solution, and the mixture was stirred at room temperature for 2 h. The mixture was concentrated under reduced pressure, then diluted with ethyl acetate. The organic layer was washed with 10% Na_2CO_3 solution, water and dried over anhydrous MgSO_4 , filtered, and evaporated to give the desired compound.

(S)-S-Benzyl 2-((tert-butoxycarbonylamino)-3-(1-tosyl-1H-imidazol-4-yl)propanethioate (8e)

White microcrystals (80%); mp 60–61 °C; ^1H NMR (CDCl_3) δ 7.91 (s, 1H), 7.78 (d, J = 8.4 Hz, 2H), 7.32 (d, J = 7.8 Hz, 2H), 7.28–7.13 (m, 5H), 7.05 (s, 1H), 6.10 (d, J = 8.7 Hz, 1H), 4.70–4.57 (m, 1H), 3.97 (s, 2H), 3.09 (dd, J = 15.0, 6.0 Hz, 1H), 2.95 (dd, J = 13.2, 5.1 Hz, 1H), 2.40 (s, 3H), 1.42 (s, 9H); ^{13}C NMR (CDCl_3) δ 201.0, 155.3, 146.4, 139.9, 136.8, 136.5, 134.8, 130.5, 128.8, 128.6, 127.4, 127.3, 115.0, 80.3, 59.7, 33.3, 30.0, 28.4, 21.8; Anal. calcd for $\text{C}_{25}\text{H}_{29}\text{N}_3\text{O}_5\text{S}_2$: C, 58.23; H, 5.67; N, 8.15; found: C, 58.11; H, 6.00; N, 8.54.

(S)-Methyl 2-((2-((tert-butoxycarbonylamino)-3-(1-tosyl-1H-imidazol-4-yl)propanoyl) thio)acetate (8f)

White microcrystals (78%); mp 110–111 °C; ^1H NMR (CDCl_3) δ 7.94 (d, J = 1.2 Hz, 1H), 7.81 (d, J = 8.1 Hz, 2H), 7.36 (dd, J = 8.1, 0.6 Hz, 2H), 7.10 (s, 1H), 6.10 (d, J = 8.7 Hz, 1H), 4.68–4.58 (m, 1H), 3.70 (s, 3H), 3.58 (d, J = 10.5 Hz, A part of AB system, 1H), 3.53 (d, J = 10.5 Hz, B part of AB system, 1H), 3.09 (dd, J = 14.7, 6.2 Hz, 1H), 2.97 (dd, J = 14.7, 4.5 Hz, 1H), 2.44 (s, 3H), 1.44 (s, 9H); ^{13}C NMR (CDCl_3) δ 200.3, 169.1, 155.4, 155.3, 146.5, 139.6, 136.5, 134.9, 130.5, 127.5, 115.3, 80.5, 59.7, 52.8, 31.2, 29.9, 28.4, 21.8; Anal. calcd for $\text{C}_{21}\text{H}_{27}\text{N}_3\text{O}_7\text{S}_2$: C, 50.69; H, 5.47; N, 8.44; found: C, 51.04; H, 5.89; N, 8.08.

Procedure for Preparation of Compound 8g

To a solution of **3a** (0.07 g, 0.137 mmol) and malononitrile (0.02 g, 0.274 mmol) in dry THF (1 mL), diisopropylethylamine (0.026 mL, 0.151 mmol) was added. The mixture was irradiated in microwave at 60 °C for 30 min and 100 W. The solvent was evaporated, and the residue was purified by flash chromatography to give **8g**.

(S)-tert-Butyl 4,4-dicyano-3-oxo-1-(1-tosyl-1H-imidazol-4-yl)butan-2-ylcarbamate (8g)


Yellow microcrystal (52%), mp 75–77 °C. ^1H NMR ($\text{CD}_3\text{OD}-d_4$) δ 8.05 (s, 1H), 7.84 (d, $J=8.4$ Hz, 2H), 7.39 (d, $J=8.2$ Hz, 2H), 7.23 (s, 1H), 4.61 (br s, 1H), 3.27–3.24 (m, 1H), 2.86–2.66 (m, 2H), 2.38 (s, 3H), 1.31 (s, 9H); ^{13}C NMR (CD_3OD) δ 194.2, 157.2, 148.1, 141.8, 138.0, 136.2, 131.8, 128.7, 121.7, 120.5, 116.5, 80.5, 55.5, 32.7, 28.8, 21.8. HRMS (+ESI-TOF) m/z for $\text{C}_{21}\text{H}_{23}\text{N}_5\text{O}_5\text{S}$ $[\text{M} + \text{H}]^+$ calcd 480.1320, found: 480.1312.

Procedure for Preparation of Compound 8h


Thiophenol (1.0 eq.) was dissolved in THF and Et_3N (1.5 eq.). N^{α} -Boc- N^{im} -4-benzyl-L-histidylbenzotriazole **3b** (1.0 eq.) was added to the solution, and the mixture was stirred at room temperature for 2 h. The mixture was concentrated under reduced pressure and then diluted with ethyl acetate. The organic layer was washed with 10% Na_2CO_3 solution, water and dried over anhydrous MgSO_4 , filtered, and evaporated to give the desired compound.

(S)-S-Phenyl 3-(1-benzyl-1H-imidazol-4-yl)-2-(tert-butoxycarbonylamino)propanethioate (8h)

Yellow microcrystals (60%); mp 130–132 °C; ^1H NMR (CDCl_3) δ 7.55–7.51 (br s, 1H), 7.42–7.31 (m, 10H), 7.00 (s, 1H), 5.34 (br s, 2H), 4.74–4.62 (m, 1H), 4.02 (d, $J=6.3$ Hz, 1H), 3.37–3.34 (m, 2H), 1.39 (s, 9H). ^{13}C NMR (CDCl_3) δ 199.2, 169.7, 155.7, 134.8, 134.7, 133.2, 131.3, 129.6, 129.1, 128.7, 127.4, 118.6, 80.4, 73.9, 53.0, 28.3, 18.9. HRMS (+ESI-TOF): m/z for $\text{C}_{24}\text{H}_{27}\text{N}_3\text{O}_3\text{S}$ $[\text{M} + \text{H}]^+$ calcd 438.1846, found 438.1850.



Scheme 1. Synthesis of N^{α} -Boc- N^{im} -protected-L-histidylbenzotriazole **3a,b**.



Scheme 2. (a) Syntheses of histidine-containing di-, tri- and tetra-peptides **5a-f**. (b) Syntheses of histidine-containing dipeptides **5a+5a'**, **5b+5b'**.

Table 1. Syntheses of histidine containing di-, tri-, and tetra-peptides **5a–f**, **5a+5a'**, **5b+5b'**

Product	Reactant 4	Yield (%)	mp (°C) (lit. mp)	$[\alpha]_D^{23}$
5a	H-L-Phe-OH 4a	79	133–134	–11
5a+5a'	H-DL-Phe-OH 4a'	72	175–177	Racemic
5b	H-L-Val-OH 4c	72	Oil	–44
5b+5b'	H-DL-Val-OH 4c'	70	Oil	Racemic
5c	H-L-Leu-OH 4d	89	121–123 (146–148) [21]	–15
5d	H-L-Ala-OH 4d	92	151–153	–8
5e	H-Gly-L-Val-OH 4e	68	113–115	–12
5f	H-Gly-Gly-L-Ile-OH 4f	60	140–142	–9






Figure 1. (a) ^{13}C NMR spectra of compound **5b**. (b) ^{13}C NMR spectra of compound **5b+5b'**.

**Scheme 3.** Syntheses of histidine-containing dipeptide **7**.**Scheme 4.** *N*-, *O*-, *S*-, and *C*-acylations with *N*^α-Boc-*N*^{im}-protected-L-histidyl-benzotriazole **3a,b**.**Table 2.** *N*-, *O*-, *S*-, and *C*-acylation with *N*^α-Boc-*N*^{im}-protected-L-histidyl-benzotriazole **8a–h**

Product	Reactant	Nucleophile	Conditions	Yield >(%)	Mp (°C)
8a	3a	<i>N</i> -(3-Aminopropyl) imidazole	A	92	64–66
8b	3a	2-Aminopyridine	B	91	87–88
8c	3a	Ethylphenol	C	50	Oil
8d	3a	β-Napthol	C	55	Oil
8e	3a	Benzylmercaptan	A	80	60–61
8f	3a	Methyl mercaptoacetate	A	78	110–111
8g	3a	Malononitrile	D	52	75–77
8h	3b	Thiophenol	A	60	130–132

A: THF, TEA, 2 h, 20 °C; B: DMF, 0.5 h, MW (50 °C, 20 W); C: THF, DMAP, 1 h MW (60 °C, 100 W); D: THF, DIPEA, 0.5 h, MW (60 °C, 100 W).

corresponding pure derivatives **8a–h** in good yields (Scheme 4, Table 2). HPLC analysis of these conjugates shows racemization in case of **8h**, which again indicates that benzyl-protected histidine is more prone to racemization than tosyl-protected histidine.

Conclusion

In conclusion, *N*^α-Boc-*N*^{im}-4-toluenesulfonyl-L-histidyl-benzotriazole **3a** is a novel, crystalline coupling reagent that (i) is sufficiently reactive to form amide and ester bonds at ambient temperature, (ii) provides good to excellent yields without detectable racemization, and (iii) is inexpensive to prepare. Hence, *N*^α-Boc-*N*^{im}-4-toluenesulfonyl-L-histidyl-benzotriazole **3a** offers an efficient preparative route to prepare *N*-protected histidine-containing peptides and conjugates with *N*-, *O*-, *S*-, and *C*-nucleophiles in synthetically useful yields without racemization.

Acknowledgements

We thank the University of Florida, the Kenan Foundation, King Abdulaziz University, Jeddah, Saudi Arabia. We also thank Dr Hall for English checking.

References

- Hunt GRA. The design of integrated inorganic experiments. An example from organo-transition-element chemistry. *J. Chem. Edu.* 1976; **53**: 53–57.
- Wright D. Pharmaceuticals containing analgesic histidine derivatives. PCT Int. Appl. WO 2010115130 A1 20101007, 2010.
- Nakamura K, Nakazawa Y, Kawamura M, Higashiura K, Miura T. Preparation of novel histidine derivatives as analgesics. PCT Int. Appl. WO 2007086354 A1 20070802, 2007.
- Nakano K, Fujikura T, Hara R, Ichihara M, Fukunaga Y, Shibasaki M. Histidine derivatives as superior renin inhibitors. Eur. Pat. Appl. US 281316 A2 19880907, 1988.
- Zhang Z, Shi X, Zhang C, Zhang H, Ma Y. Estrone amino acid ester compounds with antitumor activity and synthetic method. Faming Zhuanli Shenqing CN 102079771 A 20110601, 2011.
- Sabliov CM, Astete CE, Spivak D. Vitamin E conjugates, and their uses as antioxidants and prodrug delivery vehicles. PCT Int. Appl. WO 2011146589 A2 20111124, 2011.
- Babizhayev MA, Seguin M-C, Gueyene J, Evstigneeva RP, Ageyeva EA, Zheltukhina GA. L-Carnosine (β-alanyl-L-histidine) and carnosine (β-alanylhistamine) act as natural antioxidants with hydroxyl-radical-scavenging and lipid-peroxidase activities. *Biochem. J.* 1994; **304**: 509–516.
- Kumar G. Synthesis and characterization of mixed ligand copper (II) complexes of histidine and hippuric acid. *Chemistry* 2011; **20**: 1–8.
- Quagraine EK, Kraatz H-B, Reid RS. Peptides mimicking the N-terminal Cu(II)-binding site of bovine serum albumin: synthesis, characterization and coordination with Cu(II) ions. *J. Inorg. Biochem.* 2001; **85**: 23–32.
- Linton BR, Reutershan MH, Aderman CM, Richardson EA, Brownell KR, Ashley CW, Evans CA, Miller SJ. Asymmetric Michael addition of α-nitro-ketones using catalytic peptides. *Tetrahedron Lett.* 2007; **48**: 1993–1997.
- Tsogoeva SB, Wei S. (S)-Histidine-based dipeptides as organic catalyst for direct asymmetric aldol reactions. *Tetrahedron: Asymmetry* 2005; **16**: 1947–1951.
- Tanaka K, Mori A, Inoue S. The cyclic dipeptide *cyclo* [(S)-phenylalanyl-(S)-histidyl] as a catalyst for asymmetric addition of hydrogen cyanide to aldehydes. *J. Org. Chem.* 1990; **55**: 181–185.
- Angeletti RH, Bibbs L, Bonewald LF, Fields GB, Kelly JW, McMurray JS, Moore WT, Weintraub ST. Analysis of racemization during “standard” solid phase peptide synthesis: a multicenter study. *Techniques in Protein Chem. VIII* 1997; 875–890.
- Ishiguro T, Eguchi C. Unexpected chain-terminating side reaction caused by histidine and acetic anhydride in solid-phase peptide synthesis. *Chem. Pharm. Bull.* 1989; **37**: 506–508.
- Isidro-Llobet A, Álvarez M, Albericio F. Amino acid-protecting groups. *Chem. Rev.* 2009; **109**: 2455–2504.
- (a) Jones JH, Ramage WJ. An approach to the prevention of racemization in the synthesis of histidine-containing peptides. *J. Chem. Soc. Chem. Comm.* 1978, 472–473; (b) Jones JH, Stachulski AV. The use of the *N*(π)-phenacyl group for the protection of the histidine side chain in peptide synthesis. *J. Chem. Soc., Perkin Trans. 1* 1979, 2261–2267.
- Palasek SA, Cox ZJ, Collins JM. Limiting racemization and aspartimide formation in microwave-enhanced Fmoc solid phase peptide synthesis. *J. Pept. Sci.* 2007; **13**: 143–148.
- Barlos K, Papaioannou D, Theodoropoulos D. Efficient “one-pot” synthesis of *N*-trityl-amino acids. *J. Org. Chem.* 1982; **47**: 1324–1326.
- Colombo R, Colombo F, Jones JH. Acid-labile histidine side-chain protection: the *N*^{tert}-butoxymethyl group. *J. Chem. Soc. Chem. Comm.* 1984; 292–293.

- 20 Okada Y, Wang J, Yamamoto T, Mu Y, Yokoi T. Amino acids and peptides. Part 45. Development of a new N^{ϵ} -protecting group of histidine, N^{ϵ} -(1-adamantyloxymethyl)histidine, and its evaluation for peptide synthesis. *J. Chem. Soc., Perkin Trans. 1*. 1996; **17**: 2139–2143.
- 21 Hibino H, Nishiuchi Y. 4-Methoxybenzyloxymethyl group as an N^{ϵ} -protecting group for histidine to eliminate side-chain-induced racemization in the Fmoc strategy. *Tetrahedron Lett*. 2011; **52**: 4947–4949.
- 22 Hibino H, Miki Y, Nishiuchi Y. Synthesis and application of N^{ϵ} -Fmoc- N^{ϵ} -4-methoxybenzyloxymethylhistidine in solid phase peptide synthesis. *J. Pept. Sci.* 2012; **18**: 763–769.
- 23 Fukushima Y. Stable β -sheet formation and enhanced hydrolytic catalysis of a sequential alternating amphiphilic polypeptide containing catalytic triads in a serine protease. *Bull. Chem. Soc. Jpn.* 1996; **69**: 431–439.
- 24 Algen J, Laurence E, Jeanne B, Goeldner P, Christian H. Synthesis and properties of a 2-diazohistidine derivative: a new photoactivatable aromatic amino acid analog. *Helv. Chim. Acta* 1990; **73**: 86–96.
- 25 Ueda J-I, Hanaki A, Hatano K, Nakajima T. Autoxidation of ascorbic acid catalyzed by the copper(II) bound to L-histidine oligopeptides, (His) $_i$ -Gly and acetyl-(His) $_i$ -Gly ($i = 9, 19, 29$). Relationship between catalytic activity and coordination mode. *Chem. Pharm. Bull.* 2000; **48**: 908–913.
- 26 Kvasnica M, Budesinsky M, Swaczynova J, Pouzar V, Kohout L. Platinum (II) complexes with steroidal esters of L-methionine and L-histidine: synthesis, characterization and cytotoxic activity. *Bioorg. Med. Chem.* 2008; **16**: 3704–3713.
- 27 Zheng M, Zhang X, Zhao M, Chang HW, Wang W, Wang Y, Peng S. (3S)- N -(L-Aminoacyl)-1,2,3,4-tetrahydroisoquinolines, a class of novel antithrombotic agents: synthesis, bioassay, 3D QSAR, and ADME analysis. *Bioorg. Med. Chem.* 2008; **16**: 9574–9587.
- 28 Morris T, Sandham D, Caddick S. A microwave enhanced cross-metathesis approach to peptidomimetics. *Org. Biomol. Chem.* 2007; **5**: 1025–1027.
- 29 Histidine derivatives. Brit. GB 1352414 A 19740508, 1974.
- 30 Jones H, Ramage WJ, Witty MJ. Mechanism of racemization of histidine derivatives in peptides synthesis. *Int. J. Pept. Protein Res.* 1980; **15**: 301–303.
- 31 Bajaj K, Panda SS, El-Nachef C, Katritzky AR. Syntheses of chiral N -(protected) tri- and tetrapeptide conjugates. *Chem. Biol. Drug Des.* 2012; **80**: 17–26.
- 32 Katritzky AR, Suzuki K, Singh SK. Highly diastereoselective peptide chain extensions of unprotected amino acids with N -(Z- α -aminoacyl) benzotriazoles. *Synthesis* 2004; 2645–2652.
- 33 Katritzky AR, Angrish P, Huer D, Suzuki K. N -(Z- and Fmoc- α -aminoacyl) benzotriazoles: stable derivatives enabling peptide coupling of Tyr, Trp, Cys, Met, and Gln with free amino acids in aqueous media with complete retention of chirality. *Synthesis* 2005; 397–402.