

See discussions, stats, and author profiles for this publication at:
<https://www.researchgate.net/publication/262486396>

Hairy Root Culture: An Alternative Terpenoid Expression Platform

CHAPTER · JANUARY 2013

DOI: 10.1007/978-3-642-22144-6_136

CITATION

1

READS

50

1 AUTHOR:



Jocelyne Trémouillaux-Guiller

University F.Rabelais,Tours, F...

39 PUBLICATIONS 551 CITATIONS

SEE PROFILE

Jocelyne Trémouillaux-Guiller

Contents

| | | |
|----|---|------|
| 1 | Introduction | 2942 |
| 2 | Terpenoid Biosynthesis in Plants | 2944 |
| 3 | <i>Agrobacterium rhizogenes</i> : A Natural Engineer of Genetic Transformation | 2948 |
| 4 | Emergence and Development of Hairy Roots | 2950 |
| 5 | Ri-Plants Taken as Bioreactors | 2950 |
| 6 | Transgenic Plant Devoid of Hairy-Root Phenotype | 2952 |
| 7 | Novel Genes Revealed by Ri-T-DNA Activation Tagging and Transcriptome Analysis | 2953 |
| 8 | Engineering of Terpene Metabolism Pathways | 2954 |
| 9 | Hairy Roots: Fascinating Routes to Produce Terpenoids | 2955 |
| 10 | Elicitation Treatment: A Booster System of Secondary Products | 2956 |
| 11 | Hairy Root Cultures on a Large Scale: A True Challenge | 2958 |
| 12 | Therapeutic Merits of Plant Terpene/Terpenoids | 2959 |
| 13 | Conclusion | 2961 |
| | References | 2962 |

Abstract

A fantastic source of natural products encompassing the vast group of the terpenes, alkaloids, and phenolic compounds is provided by the plants. About 25,000 of these secondary metabolites hold a strong potential in a large domain of industrial and pharmaceutical applications. Plant families, such as *Lamiaceae*, *Asteraceae*, and *Taxaceae*, synthesize several terpenoid classes with a great economic value. Hairy roots, resulting from *Agrobacterium rhizogenes*-mediated transformation of plant cells, can be harnessed for terpenoid production on a large scale. *A. rhizogenes* over other transformation systems makes possible an

J. Trémouillaux-Guiller
University F. Rabelais, Tours, France
e-mail: j.tremouillaux@sfr.fr

easy regeneration of transgenic plants. Based on the deletion of *rol* genes located on the wild-type Ri-T-DNA, disarmed versions are created in order to overcome abnormalities of Ri-plants. Terpenoid accumulation from hairy roots can be optimized by elicitation treatment with jasmonates. Ri-T-DNA activation tagging system and transcriptome analyses are powerful tools to isolate new function genes involved in the terpenoid biosynthesis. Such genes encoding enzymes involved in the secondary metabolism can be inserted between the T-DNA borders. Lastly, some pharmacological properties of these plant terpenes/terpenoids involved in the therapy of more or less severe pathologies are reported.

Keywords

Activation tagging • *Agrobacterium rhizogenes* • Elicitation • Hairy roots • Metabolic engineering • Pharmacological properties • Ri-plants • Ri-T-DNA • Secondary metabolite production • Terpenes • Terpenoids

Abbreviations

| | |
|--------|---|
| 2, 4-D | 2, 4-Dichlorophenoxyacetic acid |
| Ac-MVA | Acetate mevalonate pathway |
| BABA | β -aminobutyric acid |
| BAP | 6-Benzylaminopurine |
| CPPU | 4- <i>N</i> -(2-chloro-4 pyridyl) – <i>N'</i> -phenylurea |
| DMAPP | Dimethylallyl pyrophosphate |
| DXP | Deoxyxylulose 5-phosphate |
| DXS | 1-Deoxy-D-xylulose 5-phosphate synthase |
| HMGR | 3-Hydroxy-3-methylglutaryl CoA reductase |
| IAA | Indole 3 acetic acid |
| IPP | Isopentenyl pyrophosphate |
| Kin | Kinetin |
| MeJA | Methyl jasmonate |
| MEP | 2-C-methyl-D-erythritol 4-phosphate pathway |

1 Introduction

Plants represent most structurally vast and varied source of natural products, although less than half of the estimated 200,000+ secondary metabolites have been identified at the present time [1]. In nature, a plethora of phytochemicals have specialized functions in ecological interactions and are frequently produced in response to pathogen or abiotic stresses [2]. These natural secondary compounds can play a beneficial or deleterious role in the plant-insect, plant-pathogen, and plant-plant relations [3]. Owing to their large number and great diversity, the secondary metabolites hold an immense potential in industrial and/or medicinal application fields [4], notably as cosmetics, food additives, perfumes, and biopharmaceuticals for the human and

animal healthcare [5]. Based on their biosynthetic origins, these secondary products can be divided into three major groups including the alkaloids ($\pm 12,000$ types), phenolic compounds ($\pm 8,000$ types), and terpenes ($\pm 25,000$ types) which do not seem to be all directly involved in the development and growth of plants [4, 6]. The terpenes that are hydrocarbons exclusively composed of carbon and hydrogen and the terpene-derived terpenoids (or isoprenoids) by oxidation or some alteration of the carbon skeleton encompass more than 40,000 structures and constitute the largest and mostly diverse class of all known natural molecules [7]. Among them, a great chemical diversity is composed of primary metabolites (chlorophylls, carotenoids, cytokinins, gibberellins) and more than 25,000 secondary or specialized metabolites have harmful or beneficial effects on other organisms [3, 6]. Specialized terpenes/terpenoids have a myriad of important industrial applications notably, as flavoring agents, perfumes, insecticides, and biopharmaceuticals [8]. Higher plants biosynthesize commonly terpenes via two IPP (isopentenyl pyrophosphate) sources generated from two independent pathways: acetate-mevalonate (Ac-MEV) (in the cytosol) and 2-C-methyl-D-erythritol 4-phosphate (MEP) or deoxyxylulose 5-phosphate (DXP) (in the plastids) pathways (Fig. 95.1), present in vegetative tissues, flowers, and more occasionally in roots [9]. The early steps of terpene/terpenoid biosynthesis in the plants start by the formation and assembly of C5 isoprene units (1 isoprene unit = a monomer form) that result from two separate pathways. Polymerization of C5 isoprene units leads, for example, to monoterpenes C₁₀ (2 isoprene units), sesquiterpenes C₁₅ (3 isoprene units), and diterpenes C₂₀ (4 isoprene units) [9] (Fig. 95.1). Indeed, terpene biosynthesis occurs, in the cytosol or the plastids, through the condensation of the five-carbon precursor isopentenyl pyrophosphate (IPP) (active isoprene unit) and its allylic isomer dimethylallyl pyrophosphate (DMAPP). The sequential head-to-tail addition of IPP units to DMAPP leads to the prenyl diphosphate geranyl pyrophosphate (GPP), farnesyl pyrophosphate (FPP), and geranylgeranyl pyrophosphate (GGPP). These three previous components serve respectively as precursors for sesquiterpenes, triterpenes (in the cytosol) and monoterpenes, diterpenes and tetraterpenes (in the plastids) (Fig. 95.1). Several terpenoids of a great industrial/medicinal interest are accumulated in the plant roots [10].

In the past three decades, hairy root cultures have been extensively investigated *in vitro* as a fascinating and alternative research tool for producing secondary metabolites at levels comparable to those found in the wild roots [11–20] and/or for revealing novel biomolecules [21, 22]. Hairy roots offer many advantages over cellular suspension cultures, notably their high growth rate, genetic stability, and hormonal independence [23]. Compared with transformed roots, undifferentiated cell cultures show most often a permanent instability responsible for low levels of secondary products and/or a rapid loss of the accumulation potential [24–27].

This chapter focuses on the terpenoid production from plant species, hairy root cultures, and Ri-plants; the terpenoid increase by hairy root elicitation; and recent strategies as T-DNA activation tagging/transcriptome analyses for isolating novel genes and hairy root engineering in order to improve terpene metabolism pathways. Lastly, therapeutic properties of some plant-derived terpenoids are reported.




Fig. 95.1 *Terpene biosynthesis pathways and their subcellular localization in the plants.* Different classes of terpenes are respectively formed in the cytosol or the plastid by two independent pathways in the plants, that is, acetate-mevalonate pathway (MEV) (cytosol) and methylerythritol 4-phosphate (MEP) or deoxyxylulose 5-phosphate pathway (DXP) (plastid). Monoterpenes, diterpenes, and tetraterpenes are derived from IPP and DMAPP from the plastidial MEP or DXP pathway. Sesquiterpenes and triterpenes are biosynthesized from IPP and DMAPP from the cytosol pathway. *Black square* with a white question mark suggests a possible transport of IPP (isopentenylpyrophosphate) from the plastid to the cytosol. Other metabolites involved in the different steps are *DMAPP* dimethylallylpyrophosphate, *FPP* farnesylpyrophosphate, *GA3P* D- glyceraldehyde- 3-phosphate, *GPP* geranylpyrophosphate, *GGPP* geranylgeranylpyrophosphate. *TPSs* in the circle correspond to terpene synthases. *Broken arrows* show several enzymatic steps (Adapted from Aharoni et al. [8] and Sallaud et al. [154])

2 Terpenoid Biosynthesis in Plants

Plant species are capable of manufacturing different types of secondary products which can be harnessed by humans for their beneficial properties in a large domain of industrial or medicinal applications [4]. World Health Organization (WHO) estimates that up to 80% of people rely mainly on traditional herbs as remedies for their medicines [28, 29]. Extracted from entire plants, secondary products are used by food and pharmaceutical industries, although most often numerous natural plant-derived molecules remain undiscovered or unexplored for their pharmacological properties [1]. Nowadays, the distribution of terpenes/terpenoids in the nature is extensively studding [30] and numerous terpenoids, presenting tremendous industrial and therapeutic properties, have already been identified in perennial herbs or woody plant species (Table 95.1). A large number of highly efficient terpenes, including

Table 95.1 Different terpene/terpenoid classes synthesized via the Ac Mevalonate/MEV pathway + (in the cytosol) and via the MEP or DXP pathway ° (in the plastid) in some plant species that are known for their therapeutic properties

| Terpene class | Plant species | References |
|---|--|----------------|
| <i>Monoterpenes (C₁₀H₁₆)/monoterpenoids</i> | | |
| Menthol | <i>Mentha piperita</i> L. | [9] |
| Menthol-geraniol | <i>Anethum graveolens</i> L. | [124] |
| Monoterpene alcohols (Linalool, geraniol, nerol. . .) | <i>Vitis vinifera</i> L. | [32] |
| Essential oils (E.O) | <i>Centaureum erythraea</i> Rafn | [99] |
| Oxygenated monoterpenoids (E.O.) | <i>Pelargonium graveolens</i> L'Her Ex Ait | [153] |
| Volatile terpenoids (Cavacrol, p-cymene) | <i>Ocimum basilicum</i> L. | [36] |
| Thymol; para-cymene; α, γ terpinene | <i>Ocimum gratissimum</i> L. | [35] |
| Volatile oils | <i>Panax ginseng</i> CA Meyer, <i>Prunus japonica</i> | [47] |
| Monoterpenoids | <i>Salvia sclarea</i> L. | [40] |
| <i>Sesquiterpenes (C₁₅H₂₄)/Sesquiterpenoids +</i> | | |
| Sesquiterpenoids | <i>Solanum tuberosum</i> L. | [97] |
| | <i>Solanum truncatula</i> L. | [93] |
| | <i>Solanum habrochaites</i> | [154] |
| Sesquiterpene lactone/bilobalide | <i>Ginkgo biloba</i> L. | [155] |
| | <i>Salvia sclarea</i> L. | [140] |
| Sesquiterpene volatiles | <i>Vitis vinifera</i> L. | [31, 32] |
| Sesquiterpene lactone, with an endoperoxide bridge) | <i>Artemisia annua</i> L. | [129] |
| (Artemisinin) | <i>Artemisia indica</i> Willd | [42, 54] |
| Terpene aldehyde – sesquiterpenoid (gossypol) | <i>Gossypium barbadense</i> | [135] |
| <i>Diterpenes (C₂₀ H₃₂)/diterpenoids °</i> | | |
| Ginkgolides/diterpene trilactones | <i>Ginkgo biloba</i> L. | [73, 155] |
| Abietane diterpenes – Diterpenoid: tanshinones | <i>Salvia scalrea</i> L. | [40] |
| | <i>Salvia miltiorrhiza</i> Bunge | [41, 119] |
| Complex diterpenoid: diterpene-alkaloid | <i>Taxus cuspidata</i> Sieb. &Zucc. | [57] |
| | <i>Taxus media</i> Hicksii | [58] |
| | <i>Taxus baccata</i> L. | [59] |
| | <i>Taxus brevifolia</i> Nutt Sieb | [56] |
| Diterpenoids: | <i>Panax ginseng</i> CA Meyer | [44] |
| <i>Triterpenes (C₃₀ H₄₈)/Triterpenoids +</i> | | |
| Ginseng triterpenoid saponins – Ginsenosides | <i>Panax ginseng</i> CA Meyer | [49, 120, 158] |
| Saponins | <i>Salvia sclarea</i> L. | [40] |
| Saponins-glycyrrhizin | <i>Glycyrrhiza glabra</i> L. | [157, 161] |
| | <i>Glycyrrhiza radix</i> L. | |
| Dammarane-type saponin glycosides | <i>Gynostemma pentaphyllum</i> (Thunb.) | [50, 51] |
| Aralia saponins V | <i>Aralia elata</i> Miq | [103] |

(continued)

Table 95.1 (continued)

| Terpene class | Plant species | References |
|--|--------------------------------------|------------|
| C-13-norisoprenoid – Lancemaside A | <i>Codonopsis lanceolata</i> Trautv. | [104, 160] |
| Triterpenoid glycosides (saponins) Escin | <i>Aesculus hippocastanum</i> L. | [43] |
| | <i>Aesculus indica</i> L. | [151] |
| <i>Tetraterpenes (C40 H64)/Tetraterpenoids</i> ° | | |
| Carotenoids | <i>Aesculus hippocastanum</i> L. | [42, 43] |

monoterpenes and sesquiterpenes, play a significant role in the interactions of plants with other environmental organisms [8]; notably terpene volatiles, carrying out important information, permit the pollinators to locate flowers and protect reproductive tissues against diverse attacks [31]. Monoterpenoids, sesquiterpenes, and norisoprenoids of low molecular weights, which have commonly been identified as volatile compounds or volatile oils emitted from flowers, fruits, and leaves of numerous plants, possess important values, such as flavor, fragrance, and aroma compounds. Likewise, other terpenoid volatiles, mainly monoterpene alcohols, for example, linalool, geraniol, nerol, and terpineol, have been considered as flavor and aroma products of grapevine berries and vine [32]. The composition of floral volatiles, for example, pollen volatiles from several grapevine varieties serve as attractants for pollinators, but also may function to deter herbivore attacks and defend the flowers against the pathogen, desiccation, or UV-light damages. Likewise, sesquiterpene volatiles at the highest levels were detected in this floral organ of *Vitis vinifera* L. cv. Cabernet Sauvignon immediately before bloom and at bloom [31]. Aromatic herbs, such as *Ocimum basilicum* L. (sweet basil), *Thymus vulgaris* L., and *Ocimum gratissimum* L., of the family Lamiaceae are largely utilized because of multiple properties as distinctive aroma additives to food or as medicinal herbs [9, 33–35]. These aromatic oils, stored in the leaves, are composed of aroma monoterpenes (eugenol, thymol, carvacrol, and linalool) and exhibit potent antioxidant properties [36]. Essential oils, that is, green pesticides are an excellent alternative to synthetic pesticides in order to reduce negative impacts to human health and environment [37].

Plant species, belonging to different genera and synthesizing several under classes or types of terpenes/terpenoids, are succinctly described below (Table 95.1). The first genus *Salvia*, the largest genus of the family Lamiaceae (or Labiatae), distributed throughout the world, offers a great and cosmopolitan group of about 236 genera and 900–1,000 species [38] and is phytochemically characterized by their diterpene contents [39, 40]. *Salvia* species that are of widespread interest owing to their large spectrum of medicinal activities can be cultivated in several countries of Europe [41]. Turkey is one of the major centers of diversity for genus *Salvia*, with 93 species of which 54 % are endemic [38]. Two subgenera *Salvia* and *Sclarea* possess abietane diterpenes, whereas *Calosphaea* has clerodane diterpenes and *Leonia*, abietane and clerodane diterpenes. Considered as compounds specific to this species the diterpenoid tanshinones of *Salvia miltiorrhiza* (Chinese sage), contained in the rhizomes, have received more attention [41]. The aerial parts of *Salvia sclarea* contain mostly monoterpenoids and

sesquiterpenoids, whereas twelve diterpenoids and two triterpenoids have been identified in the roots of this species. Most of the diterpenoids, abietane or arranged abietane-type, show various biological activities [40]. Genus *Aesculus* belonging to Hippocastanaceae or Horse-chestnut family regroups 100 species or varieties. *Aesculus* sp. are deciduous trees and shrubs among which several species, such as *Aesculus hippocastanum*, *Aesculus chinensis*, *Abronia turbinata*, and *Acalypha indica*, have been cultured as pharmaceutical crops for providing Standardized Therapeutic Extracts (STEs) [42]. More than 210 biomolecules have been isolated from the genus *Aesculus* and identified as belonging to different terpenoid classes, triterpenoids, triterpenoid glycosides (saponins), and tetraterpenoids (carotenoids) [42, 43] (Table 95.1). Two Eurasian species, of which aescin and individual compounds have been identified and other *Aesculus* sp. exhibit potent antitumor, antiviral, antioxidative, anti-inflammatory effects. The root of *Panax ginseng* CA Meyer is largely used as traditional medicine since ancient time due to its stimulating and tonic properties [44]. Genus *Panax* (Araliaceae family), regrouping about 11 slow-growing species, that is, perennial plants with strong roots, has been used for therapeutic aspects for centuries. Among them *Panax japonicus* (Japanese ginseng), *Panax quinquefolius* (American ginseng), and *Panax ginseng* (Korea or Asia ginseng) [45, 46] are cultured in Japan, China, Korea, but also in the United States and Canada [47]. *P. ginseng* is the most required species for its restorative properties [48], and its roots contain active principles detected as ginseng triterpenoid saponins and ginsenosides (Rx), that is, saponin glycosides, the most prominent constituents of ginseng [49]. The chemical compounds of different *Panax* species are relatively similar and approximately 38 types of ginsenosides have been identified with variable levels according to the *Panax* species [46]. *Panax ginseng* is close to *Gynostemma pentaphyllum* because of the secondary metabolite production and pharmacological properties that its chemical entities confer on this plant species. *Gynostemma pentaphyllum* Thunb. or jiaogulan (Chinese name), a member of the family Cucurbitaceae and genus *Gynostemma*, is an oriental medicinal herb biosynthesizing numerous Dammarane-type saponin glycosides, and about 90 compounds known as gypenosides that have been isolated and considered as responsible for the pharmacological properties recently attributed to this species [50, 51] (Table 95.1).

Lastly, two other important genera regroup precious species biosynthesizing phytochemicals involved in the treatment of mortal pathologies. In first, the genus *Artemisia* is one of the largest of the family Asteraceae with more than 500 species and includes a large number of aromatic plants. Genus *Artemisia* is mostly composed of perennial plants and distributed worldwide, mainly across the temperate zones of the Northern Hemisphere, but a few species can also be observed on the Southern Hemisphere [52]. Overall *Artemisia* species are considered as herbs (*Artemisia annua* L., *Artemisia vulgaris* L.), subshrubs (*A. changaica* Krasch, *A. crithmifolia* L.), or shrubs (*A. tridentate* Nutt.) with strong aroma [52, 53]. Artemisinin, a sesquiterpene lactone with an endoperoxide bridge [42, 54], has been isolated from species *Artemisia annua* L., a model plant owing to its small genome, which is widely used to treat the malaria disorders [42] (Table 95.1).

Another important genus *Taxus*, composed of small coniferous trees and shrubs, belongs to the family Taxaceae including many species, such as *Taxus baccata*, *Taxus brevifolia*, *Taxus chinensis*, *Taxus Canadensis* or *Taxus cuspidata* or still *Taxus media*. These coniferous trees, relatively slow-growing, can reach between 1 and 40 m in length and produce red berry-like fruits. Known commonly as yews, these species are poisonous because most of the trees produce taxol. Taxol or Paclitaxel (its generic name) that was isolated in 1971 [55] from the bark of the yew tree (*Taxus brevifolia* Nutt) is a complex diterpenoid (diterpene alkaloid) possessing very strong antineoplastic efficiency due to its unique mode of action on the cell microtubules [56–58]. Considered as an alternative product to taxol, taxotere or its generic name “docetaxel” is a chemical entity, semi-synthetically via the conversion of non-cytotoxic molecules: baccatin III and 10-deacetylbaccatin III precursors of taxol extracted from the needles of *Taxus baccata* L., the European yew [59] which has provided an interesting and renewable source of taxanes with the similar action mode for treating the same variety of cancer pathologies.

In the last decades, numerous biotechnological investigations have made possible the establishment of hairy root cultures from the plant species, previously described, in order to produce terpene/terpenoids of pharmaceutically great interest.

3 *Agrobacterium rhizogenes*: A Natural Engineer of Genetic Transformation

Phytopathogen gram-negative soil bacterium, *Agrobacterium rhizogenes*, is able to infect wounded plant cells [60] for inducing a tumorous pathology, referred to as “hairy root syndrome” [61, 62] and revealed by the emergence of adventitious roots at or near to the inoculation site of the agrobacteria [63] (Fig. 95.2). Pathogenicity of *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes* depends respectively on a large and extrachromosomal Ti- (for tumor inducing) or Ri- (for root-inducing) plasmid carrying a DNA region termed Ti-T-DNA or Ri-T-DNA respectively, [64] that can be horizontally transferred and stably integrated into the nuclear genome of higher plants [62], as dicotyledon, gymnosperm, and some monocotyledon host species [65]. Ri-T-DNA harbors between its specific border sequences (T_R (R =right border) and T_L (L =left border)) of 25-bp repeats, notably the T_L -DNA region carrying the root loci (*rol*) genes responsible for the hairy root phenotype. Ri plasmids, such as *pRiA4* or *pRi1855*, belonging to the hypervirulent agropine-type *Agrobacterium rhizogenes* strains, possess a T-DNA formed of two independent regions [66] termed T_R -DNA and T_L -DNA. The T_R -DNA region carries the *aux1-2* genes that encode enzymes involved in auxin synthesis and play an ancillary role in the transgenic root formation [67]. Eighteen open reading frames (*Orfs*) have been identified from the T_L -DNA region, among them the 10, 11, 12, and 15 *Orfs* correspond respectively to the *A*, *B*, *C*, and *D rol* genes [68]. Other genes, termed *ops* genes, are equally located in the T_R -DNA and code for opine synthesis, that is, particular amino acid derivatives serving for specific food to the bacteria [69–72]. Genetic transformation of recalcitrant species, as




Fig. 95.2 Establishment, culture on a large scale and terpenoid production from hairy roots. (a) Both partners involved in the genetic transformation: *Agrobacterium rhizogenes* (a pathogen gram-negative soil bacterium) harbors a large Ri-plasmid carrying the wild-type T-DNA (black arrows) or an engineered T-DNA that can be transferred into the plant genome (black arrow) and one wounded leaf. (b) After 48–72 h of co-culture with bacterium/plant cells, emergence of adventitious primary roots at or near the infected site. (c) Each primary root excised on the leaf explant and placed on solid medium grows vigorously for generating a true clone of hairy root (HR). (d) A crucial step corresponds to the transfer of the root tip segments of HR into the agitated liquid medium. (e) One well-established HR clone grows rapidly and shows the characteristics of the HR phenotype, that is, plagiotropic growth, high branching in hormone-free medium. (f) Scale-up of hairy root culture in specific bioreactor for commercial exploitation of valuable terpenoids. (g) Separation and identification of terpene/terpenoids by liquid chromatography/mass spectrometry (LC/MS) or by high performance liquid chromatography (HPLC)

Ginkgo biloba, required the use of hypervirulent agropine-type strains [73]. On the contrary some Ri plasmids, as *Ri8196* or *pRi2659*, have a simple T_L -DNA capable of initiating hairy roots and producing different opine types (mannopine, mikimopine, or cucumopine) according to the *A. rhizogenes* strain [74]. Fundamental studies have improved our understanding of the different molecular events resulting in the T-DNA transfer through the host cytoplasm up to nuclear plant genome. Nevertheless, the final steps of T-DNA insertion into the plant genome, involving host factors, are still unknown. The Ti/Ri plasmids possess *vir* genes (*vir* for virulence) that encode Vir proteins responsible for the plant-bacterium attachment, formation, and protection of the simple and mobile copy of the T-DNA for its translocation from bacterium to the host nucleus [75–77]. Most *vir*

genes, carried by the Ti/Ri plasmids and involved in the T-DNA secretion system, are similar. However, the Ri plasmid possesses GALL genes coding for two full-length GALLS (-FL and -CT) proteins, instead of the *virE2/virE1* genes present in *A. tumefaciens* [78, 79].

4 Emergence and Development of Hairy Roots

In the previous decades, the strong power of *A. rhizogenes* to transform various plants and induce hairy roots has been largely demonstrated [71]. About 200 species of higher plants (at least 30 families), mostly dicotyledonous species, could be genetically transformed with *A. rhizogenes* and led to the development of adventitious roots [80]. This efficient mode of bacterial transformation is depending on several parameters, including the strain type, bacterial concentration, co-culture time of the bacterium/explant, temperature, light/darkness, as well as the plant species and explant type [81]. Different surface-sterilized explant types (i.e., leaf discs, leaves, stem segments, petioles, roots, embryos, or plantlets) are wounded and then inoculated with *Agrobacteria*. After 2–4 weeks primary adventitious roots, coming from the differentiation of one single plant cell transformed by one bacterium [62], appear at, or near, wound site of plant cells infected by *A. rhizogenes* [82] (Fig. 95.2). Roots, reaching about 1–1.5 cm in length, are excised and placed separately on solid or semi-solid hormone-free medium for initiating a true clone. Hairy root phenotype is characterized by a fast hormone-free growth, an unusual ageotropism, a profusion of root hairs with abundant lateral branching [83–85] (Fig. 95.2). Hairy root cultures can present more and less than root thickness, root length, number of root hairs and root branching, according to plant species and *A. rhizogenes* strain used for hairy root induction [86]. Sometimes, Ri-roots can become brown and stop their growth in liquid medium, because the most crucial step results from the transit of the hairy root clones into agitated liquid medium [87] (Fig. 95.2). But the morphology of *A. rhizogenes*–induced hairy roots is similar in structure to wild-type roots with a few notable exceptions [66]. A recent paper underlined that the *A. rhizogenes*–mediated transformation has made possible the establishment of hairy root cultures from 116 plant species [88].

5 Ri-Plants Taken as Bioreactors

A. rhizogenes–mediated transformation can be exploited for producing terpenoids using plants as bioreactors. Advantages of transformation with *A. rhizogenes* over other methods encompass adventitious organs/roots and rapid shoot regeneration reducing the risk of somaclonal variation [89]. Moreover, this transformation is that it enables the remarkable development of Ri-plants via a marker-free selection through the observation of the hairy root morphology, as the primary indicator of

genetic transfer [90]. Lastly, another fundamental advantage from the wild-type *A. rhizogenes* strains, compared with the *A. tumefaciens* ones, is that the hairy root creation can naturally lead to the plantlets. Thereby such plants are free from the legal controls of GMOs in Japan [91]. One pioneered plant regeneration, performed successfully from *Brassica napus* hairy roots, displayed that fertile Ri-plants could sexually transmit the hairy root phenotype to their progenies [83]. Such a heredity of Mendelian type has recently been confirmed by Crane et al. [92] from Ri-plants of *Medicago truncatula*, an interesting legume, that emits a large spectrum of volatile terpenoids, including monoterpenes, sesquiterpenes, and tetranor-terpenoids, induced by herbivore attacks [93]. Studies reported that transgenic plants were regenerated from hairy roots of 62 different taxa, representing 53 plant species from 24 families [89] or still up to 79 plant species [94].

Besides, Ri-plants potentially can equally offer promising horticultural/floricultural applications. Classical breeding in floriculture serves to create new plant varieties, although the available gene pool for introducing novel agronomic or aesthetic features is genetically limited to the parental background [95]. Appeared after the *rol* gene insertion into the plant genome, cytological abnormalities make possible the creation of the novel floricultural varieties with exciting characteristics. For instance, Ri-regenerants of *Kalanchoe blossfeldiana*, an economically important plant in Europe, exhibited a dwarf and compact growth habit compared to an elongated growth habit, commercially disadvantageous, of the control plants [90].

An overview of the literature reveals that plant regeneration from hairy root segments can take, spontaneously, directly or indirectly (after a callusing phase before shoot regeneration) several developmental ways. Thereby, buds were initiated from adventitious Ri-root parts with or without exogenous growth regulators [66], for example in *Aesculus hippocastanum*, a plant species producing triterpenic glycosides [43], buds emerged from hairy roots placed on the solid medium devoid of exogenous hormones [96]. Nevertheless, the number of *A. hippocastanum* buds was found superior when the Ri-roots were placed on the medium supplemented with BAP. Rooting of these shoots was initiated more rapidly with IBA (in 80% of explants) than IAA (in 40% of explants) [96]. Another regeneration model was reported from genus *Solanum* which is one of the largest genus of family Solanaceae comprising about 84 genera and 3,000 species and producing numerous flavonoids and terpenoids, such as sesquiterpenoids in *Solanum tuberosum* [97] (Table 95.1). *Solanum nigrum* hairy roots, maintained on a hormone-free medium, developed spontaneously plantlets capable of producing in vitro flowers and fruits, whereas control plantlets remained in the vegetative state [89]. *Pelargonium graveolens* hairy roots, coming from Ri-T-DNA-mediated transformation of leaves and internodal segments, expressed their shoot regenerative capacity from both explant types [98]. *Centaureum erythraea* Rafn (*Gentianaceae*) is a traditional medicinal plant of Croatia that accumulates monoterpenoids in the aerial parts [99] (Table 95.1). *C. erythraea* hairy roots, placed in the medium added with IAA and BAP, formed shoots which were rooted at 68–97% without any hormonal

supplement [100]. The Ri-plants regeneration could be associated to the formation of embryogenic calli [100]. *Panax ginseng* CA Meyer possesses numerous pharmacological activities, mostly attributed to one type of its constituents, namely, ginsenosides belonging to triterpenoid saponins [46] (Table 95.1). Placed on the medium without any growth regulator and in the dark, shoots and yellow calli in *Panax ginseng* appeared simultaneously from hairy root clones. Embryogenic calli, transferred to light, gave rise to somatic embryos leading to shoots/plantlets [101]. Transformed callus-derived embryogenic cell lines of *Panax ginseng* have been able to develop embryos and plantlets arising from only *A. rhizogenes* *rol* C-gene expression [102]. Root segment-derived hairy root lines of *Aralia elata*, a wood shrub producing pharmaceutically important metabolites as triterpenoids (*Aralia* saponins V) [103], and cultured on nutritive medium supplemented with 2, 4-D could generate embryogenic calli able to develop somatic embryos up to the cotyledonary stage without any phytohormone. Embryo germination led to plantlets and 100% of them were successfully acclimatized under greenhouse conditions [103]. Embryogenic calli, cotyledonary embryos and plantlets were developed from hairy root segments of *Codonopsis lanceolata* (Campanulaceae family) cultured in medium supplemented with 2, 4-D and accumulated, in the stems, higher contents of three major triterpenoids (lancemaside A, foetidissimoside A, and aster saponin Hb) than wild-type plants [104].

The third organogenesis type requires an intermediate callus phase before the formation of shoots and plantlets. Thereby, hairy root-derived calluses of *Mentha piperita* proliferated abundantly in the presence of 1 μ M NAA, 10 μ M CPPU and 10% coconut powder in the culture medium. Placed onto solid hormone-free medium *M. piperita* calli gave rise to shoots and roots [105].

6 Transgenic Plant Devoid of Hairy-Root Phenotype

Ri-Regenerants, arising from genetic transformation by the T-DNA insertion, exhibit frequently typical abnormalities due to the *rol* gene expression as wrinkled leaves, short internodes, dwarfness, numerous lateral shoots, flowers, and plagiotropic roots with an extensive branching [94, 106, 107]. Nevertheless, normal morphology of the leaves, a good stem elongation, and an extensive root system could be observed from Ri-regenerants of *Alhagi pseudoalhagi*, and similarly of *Hypericum perforatum* [108]. In contrast, transgenic somatic embryo-derived plantlets of *Aralia elata* exhibited an aberrant hairy-root phenotype compared to the mother plants [103]. This particular aspect of Ri-plants can be discouraging for initiating certain programs of applied research, notably, in the agronomy fields. Easy protocols have been established to overcome undesirable abnormalities of the Ri-plants by creating disarmed versions, as it has been achieved with *A. tumefaciens* [66]. The abnormal hairy-root phenotype can be suppressed after deletion of the oncogenes, that is, *rol* genes, located on

the wild-type Ri-T-DNA and responsible for these morphological singularities. This deletion is performed without any consequence for the T-DNA insertion, since with exception of the right and left borders none of the other T-DNA regions are indispensable to the horizontal transfer of foreign genes into the host genome [109]. Therefore, the *Rol* genes carried by the wild-type Ri-T-DNA may be replaced by additional genes overexpressing secondary metabolites/recombinant proteins or encoding the resistance to herbicides, insects, and various pathogens [110]. Two vector types, co-integrative and binary vectors, can be used as genetic transformation systems [111]. A binary vector system contains, for example, the Ti plasmid carrying only virulence genes, whereas its T-DNA has been removed [112] and the second plasmid carries the foreign gene inserted between the border sequences of the Ri-T-DNA with or without deleting the *rol* genes [113]. Plants coming from leaf explants of *Centaurea erythraea* Raf co-cultured with the *A. rhizogenes* strain LBA 9402, carrying a binary vector, were reported by Piatczak et al. [100]. Using binary vectors, deleted oncogenes, the problem of the surprising hairy root phenotype may be bypassed.

7 Novel Genes Revealed by Ri-T-DNA Activation Tagging and Transcriptome Analysis

Hairy roots, coming from the genetic horizontal transfer of the wild or recombined Ri-T-DNA vectors in the host plant genome, represent an ideal engineering platform for exploiting new biotechnology tools (Fig. 95.2). DNA activation tagging [114] and RNA silencing processes followed by transcriptome analyses are recent powerful tools capable of revealing novel genes, notably those involved in the secondary metabolism, but also, of identifying gene function [114]. Activation tagging process permits to generate gain-of-function mutations, whereas gene silencing leads to loss-of-function approaches [115]. RNA silencing corresponds to a natural mechanism of genetic control taking place in virus resistance, genome maintenance, and developmental control in plants [116]. The classical loss-of-function process can be limiting because of the absence of visible phenotypes which restrict the identification of mutants [16]. Thus, one of the most frequently applied approaches is DNA-activation tagging insertional mutagenesis [117] which can circumvent many disadvantages of loss of function [44]. The *Agrobacterium* spp. T-DNA vector, carrier of multimerized transcriptional enhancers of the cauliflower mosaic virus (CaMV) 35 S promoter, is considered as a powerful mutagen [44] capable of discovering novel dominant or semi-dominant mutations observed in primary transformants [117]. Activation tagging system leads to a random genomic insertion, in the plant genome, of Ri-T-DNA constructs harboring constitutive enhancer sequences positioned next to the right- or left-hand T-DNA border, which are able to cause transcriptional activation of flanking genes [22, 117, 118]. For example, Choi et al. [44] generated activation tagged *P. ginseng* hairy root lines in order to dissect, isolate genes responsible for ginsenoside and triterpene lactone biosynthesis and evaluate the ginsenoside profile of these hairy root lines [119].

Another recent approach based upon the methyl jasmonate inducible system that was applied to hairy root cultures of *Panax ginseng* made possible the identification of new genes through the analysis of 3,134 expressed sequence tags (ESTs) [120]. From the isolated transcripts several would correspond to the genes encoding enzymes as Cytochrome P450, glycosyltransferases, and oxidosqualene cyclase (OSC) involved in the ginsenoside (i.e., triterpene glycosides) biosynthesis [120]. *Ginseng* (or other plant species) ESTs data set can provide important information on genes involved in the regulation and biosynthesis of secondary metabolites, only, when their accumulation can be induced by MeJA treatment [120]. Besides, Jian et al. [121] emphasized the fundamental role of the *A. rhizogenes*-mediated transformation system coupling, at once, genetic transformation and regeneration for gene function testing in *Lotus corniculatus*.

8 Engineering of Terpene Metabolism Pathways

The fortification of numerous steps by overexpression of multiple biosynthesis genes or transcription factors for controlling the expression of genes is a fascinating strategy to alter the accumulation of secondary metabolites [122]. Metabolic engineering of terpenoids biosynthesized in plants can be directed toward two major objectives: to increase the plant resistance to predators and diseases or still to increase the production of medicinal or industrial compounds and to make possible a better knowledge of genes encoding enzymes involved in the terpenoid metabolism pathways or its regulation [8]. Understanding the function of genes involved in terpene production can lead to discovery of novel compounds or metabolic pathways, which might reveal favorable properties for human applications [4]. Thereby *Nicotiana tabacum* plants displayed an increase of terpene emission from leaves after insertion of monoterpene synthase genes [4]. The red pigment present in *Salvia miltiorrhiza* (*Lamiaceae*) roots is composed of many diterpenoid tanshinones. In this plant, diterpenes are synthesized through the MEP pathway in the plastids (Fig. 95.1) and DXR (DXP (1-deoxy-D-xylulose 5-phosphate) reductoisomerase) is an enzyme catalyzing the first step of this pathway. Wu et al. have cloned and sequenced the gene coding for the DXR protein [41]. Functional expression of the *dxr* gene was controlled by genetic complementation with the lethal *dxr*-deficient *E. coli* mutant strain. Besides, the RNA-blot analysis, from hairy root cultures of *S. miltiorrhiza*, elicited with a hyperosmotic stress and/or yeast extract, showed a strong induction of the *dxr* gene transcription that was correlated with the diterpenoid tanshinone increase in these cultures. The same authors concluded that the *dxr* gene involved in the MEP pathway may be a target for the metabolic engineering of diterpenoid tanshinone biosynthesis [41]. The regulation of terpenoid metabolism and, particularly, the regulation of factors controlling genes encoding enzymes of the secondary metabolism require subsequent investigations [123]. *S. miltiorrhiza*, taken as a model plant owing to its small genome, made possible the identification of 40 terpenoid biosynthesis-related genes, of which 27 are novel. It was reported that 20 of the 40 genes could be involved in the tanshinone biosynthesis [124].

9 Hairy Roots: Fascinating Routes to Produce Terpenoids

Secondary metabolites of low molecular weights are widely biosynthesized throughout the plant Kingdom [122] and a large biological diversity of them is accumulated in the roots [85]. In the past few years, a great interest for the natural products led private companies to consider the progresses of the biotechnologies as an alternative strategy for producing biomolecules of high added value. Numerous plant species, genetically transformed with the natural vector *Agrobacterium rhizogenes*, are good candidates to accumulate terpene/terpenoids in organ culture systems [23, 40]. Important research investigations that use differentiated cultures, instead of cell suspension cultures, have been focused on transformed (hairy) roots [23] which offer a flexible and versatile system for a large-scale production of phytochemicals [125]. Hairy roots, because of their high differentiation, allow a stable and extensive production of secondary compounds [126]. Moreover, owing to their culture mode in a sterile, confined, and controlled environment avoiding the transgene dissemination, hairy roots present serious advantages compared with entire plants cultured in the open field [18, 22]. The transformed root cultures can be propagated indefinitely in liquid medium and conserve, at once, a morphological integrity and biochemical stability [86, 115]. Besides, the *rol* genes are now known to possess a stimulating effect on the secondary metabolism with a durable stability, over long-term culture [127]. After its insertion in the plant genome, the T-DNA appears to be a remarkable element in the gene regulation [128]. As described below, some hairy root cultures have shown their capacity to accumulate terpenes/terpenoids in vitro. Thereby, chemical analysis displayed that the contents of three major triterpenoids in hairy roots of *Codonopsis lanceolata* Trautv were similar to those found in wild roots harvested from soil [104]. Sterile 4-week-old shoots of *Salvia sclarea* transformed with the *A. rhizogenes* strain, LBA 9402, led to hairy root cultures from which four diterpenoids (ferruginol, salvipisone, aethiopinone, and 1-oxoaethiopinone) and two ursine-type triterpenoids were isolated [40]. The light enhanced significantly the abietane diterpenoid levels and affected slightly the triterpenoid spectrum in transgenic cultures. Kuzma et al. [40] concluded that *S. sclarea* hairy roots, accumulating terpenoids at higher levels than the roots of field-grown plants, could significantly provide a commercial source of diterpenoids. Hairy roots of *Artemisia dubia* and *Artemisia indica* arising from sterile plantlet-derived stems were transformed with both *A. rhizogenes* strains, LBA 9402 and 8196. Artemisinin levels and fresh root biomass were significantly enhanced in the Ri-roots compared with the control roots and according to the *A. rhizogenes* strain and *Artemisia* species used. Higher levels (0.603 % and 0.753 %) of this natural sesquiterpene lactone were detected in *A. dubia* hairy roots when the stems have been respectively inoculated either with LBA 9402 or 8196 *A. rhizogenes* strains [129].

Despite numerous positive results, some transformed root systems accumulated less secondary metabolites than plant tissues in vivo. The *A. hippocastanum* zygotic embryos are known to be the major site of aescin, a group of chemically related triterpenic glycosides (saponins) involved in the treatment of peripheral vascular

disorders as venous insufficiency [43, 96]. Androgenic embryos of *Aesculus hippocastanum* (horse-chestnut), induced from anthers and inoculated with the *A. rhizogenes* strain A4GUS, led to hairy root clones [43]. The substantial percentages (3.50 % and 4.1 %) of aescin, detected in two hairy root clones, remained inferior to those found in the cotyledonary embryos placed on the hormone-free medium [43]. These authors concluded that the androgenic embryos of *A. hippocastanum* cultured with 2, 4-D is a better strategy for the aescin production on a large scale. Besides, Zdravkovic et al. [96] reported that the hairy root clones of horse chestnut were subsequently able to regenerate transformed plants which may be useful in horticulture for the bonsai establishment. Another tremendous advantage of hairy roots results from the feasibility of regenerating transgenic plants, whereas the regeneration of tissues transformed with *A. tumefaciens* is proved to be often more problematic. Overall secondary metabolites are biosynthesized and accumulated in the roots/hairy roots; nevertheless, some of them can be produced and/or stored in the aerial parts of the plant. Thereby, the total spectrum of secoiridoids that were detected in the non-transformed plants could be provided in vitro at eightfold superior levels in the roots and aerial parts of *Centaureum erythraea* Ri-plants [100]. Likewise, essential oils were mostly detected in the leaves and stems of *Mentha*, but the highest oil levels of *M. piperita* are found in the leaves of Ri-plants compared with those of the wild-type plants [130]. Moreover, the dwarf regenerants of *Mentha* species displayed the hairy root phenotype abnormalities, which does not present any negative impact on the essential oil production [105]. Most hairy root-derived rose-scented geranium plants (*Pelargonium* species) provided a large panel of monoterpenoids (essential oils) similar to those of the wild-mother plant, at the exception of two Ri-plants that biosynthesized two volatile monoterpenoids possessing better olfactory value offering an improvement from a commercial point of view [98]. Green hairy roots can accumulate some secondary products that are normally synthesized in green aerial parts of the plant [130].

10 Elicitation Treatment: A Booster System of Secondary Products

Plants are capable of producing and releasing active allelochemicals targeted in defense against pathogen attacks and stress conditions [2, 131]. Allelochemicals are accumulated under the impact of stress molecules, called elicitors, and then released by the plants into the environment by root exudation, volatile emissions from leaves and other aerial parts [132]. Several natural elicitors have been observed as efficient inducers for boosting the production and secretion of secondary metabolites in cultures in vitro, notably, when they are added to the hairy root cultures [115]. Thereby, the secondary plant messenger methyl jasmonate (MeJA), and its relatives, from the oxylipin family, is a stress hormone able to induce the improvement or optimization of secondary product contents [10, 18, 97]. In 1962, MeJA was discovered as a sweet-smelling biomolecule present in *Jasminum*

grandiflorum L. flower extracts [133, 134] and described as “hormone” because of the cellular responses of elicitation, at low levels, distant from its site of biosynthesis. The jasmonate elicitors are synthesized via the octadecanoid pathways, beginning at linolenic acid and ending at (+) – 7- epi-JA and its conjugates and isomers [134]. There are some examples of elicitation processes for stimulating the secondary metabolite accumulation. Gossypol (sesquiterpenoid) is a toxic phytoalexin biosynthesized in *Gossypium barbadense* (a cotton plant) that possesses anticancer, antiviral, antimicrobial, and antiparasitic properties and is routinely isolated from cottonseed or from cotton roots. This terpene aldehyde interacts with various anticancer agents for increasing the efficiency of their pro-apoptotic proteins, detoxification enzymes, and signaling kinases. Only MeJA was found to be able to stimulate the contents of gossypol and methylated forms in cotton hairy root cultures. Moreover, the distribution of enantiomers was not found to be similar in elicited or control transformed roots of *G. barbadense* [135].

Taxol or Paclitaxel (generic name), a complex diterpenoid, is one of today's better known antineoplastic drugs [56] and acts as a strong antitumor agent owing to its unique mode of action on the cellular microtubules. Its extraction from its natural source and costly synthetic process make it one of the products with the highest added value [58]. Taxotere or Docetaxel (generic name) is a chemical entity semi-synthetically performed via the conversion of non-cytotoxic molecules: baccatin III and 10-deacetylbaccatin III precursors of taxol extracted from the needles of *Taxus baccata* L., the European yew. Paclitaxel can equally be semi-synthesized from its natural precursor, 10-deacetyl baccatin III. As a promising alternative, cell suspension cultures of various *Taxus* species have been harnessed for the efficient production of taxol and its related taxane compounds [57]. The impact of explant source and growth regulators was tested in order to optimize the induction and selection of fast growing callus lines of *Taxus baccata* (European yew tree) [136]. In *Taxus baccata* L. (European yew tree), Ashrafi et al. [136] tested different combinations of growth regulators and various explants (stems, needles, and base of needle) for callus production. Taxol and baccatin III were respectively increased by 3.1- and 5-fold in the callus-derived cell suspensions cultured in medium added with 2, 4-D + Kin, at different concentrations according to both expected secondary metabolites [136]. *Taxus media* cell lines, transformed with *A. rhizogenes* and termed “TXS” and “Rol C” (*txs* corresponding to the transgene encoding enzyme taxadiene synthase of *T. baccata*), were treated with MeJA for optimizing taxane production. The highest taxane level was observed in the “TXS” cell line and resulted from the activity of enzyme taxadiene synthase and MeJA [58]. To date, the most favorable proceeding for a sustainable production of taxol and related taxoids at industrial level is provided by plant cell cultures [58]. Although production has been scaled-up, a serious drawback of the cell suspension cultures is their instability and unpredictability of the expected secondary products [137]. Due to their great genetic stability and potentially unlimited propagation in vitro, hairy roots as well-differentiated organs, are able to accumulate of important taxanes. For example, three strains of *A. rhizogenes* (R1000, A4 and 15834) were compared for the initiation of *Taxus cuspidata* hairy root lines. As regards its

growth capacities, a selected line was treated with MeJA and produced a total paclitaxel content of 52, 5 mg/l over 2 weeks of incubation and at 20-L culture scale. These taxol levels produced after MeJA addition were similar to those detected in the cell suspension cultures of *Taxus* [57]. As another example, taxol and 10-deacetylbaecatin III (10-DAB III) were increased in hairy root cultures of *Taxus x media* var. *Hicksii* Rehd in presence of the BABA precursor, used alone or in combination with MeJA [137]. To date, only two reports have been published for the production of paclitaxel and its related compounds by hairy root cultures [137].

Other molecules, biotic or abiotic, can act as elicitors for successfully boosting the accumulation and/or secretion of secondary products. A non-protein amino acid, that is, β -aminobutyric acid (BABA), used against pathogen attacks and added separately or in combination with the yeast elicitor for enhancing the tanshinone production in *Salvia miltiorrhiza* hairy roots, showed that BABA could strongly potentiate elicitor-induced secondary metabolism in plant organ cultures [138]. From another experimental work, Ge and Wu [139] studied the separate and combined action of biotic (yeast=YE) and abiotic (Ag+) elicitors on the secondary metabolism activities of *Salvia miltiorrhiza* hairy roots. These authors showed that the YE-induced tanshinone production seemed to be mainly derived from a non-MVA pathway encompassing DXS (1-deoxy-D-xylulose 5-phosphate synthase) since YE did not stimulate HMGR (3-hydroxy-3-methylglutaryl CoA reductase). In contrast, Ag+-induced tanshinone production seemed to be dependent on both pathways (MVA and non-MVA pathways), because it was suppressed by the inhibitors of both pathways [139]. Elicitation, but, also cell permeabilization and trapping processes have been investigated to increase the secretion of secondary products in the culture media [18].

11 Hairy Root Cultures on a Large Scale: A True Challenge

Hairy root offers a flexible and versatile culture system that is a promising biotechnology for the large-scale production of valuable phytochemicals [125]. The transfer in bioreactors is the key step toward a commercial production of terpenes/terpenoids by hairy root cultures. Bioreactors offer several advantages for the mass culture of plant cells or organs, for example, a constant regulation of conditions at various stages of the culture [28]. To scale up hairy root cultures, several operational factors, including inoculum size, medium, growth rate, secondary product recovery (intracellular/extracellular), and reproducibility of results, must be overpowered before the technology is ready for commercialization [23]. Thereby a perfect sterility, maintenance of the homogenization of roots as well as biomass measurement, agitation, oxygenation, and nutrition availability are the major goals to reach for scaling-up hairy root cultures [23, 130, 140]. In fact, growth of transgenic roots in liquid medium results in dense and packed root biomass, leading to the formation of stagnation which slows down the fluid nutrient circulation and limits oxygen availability [86]. Mechanical agitation generates wounding of hairy

roots and callus formation [130]. To overcome these limiting hurdles, hairy roots can predominantly be grown in spray or droplet bioreactors and mist bioreactors in which the roots are immobilized on mesh support and exposed to humidified air or a gas mixture and nutrients [86]. Recent bioreactors with their different advantages/drawbacks for producing hairy roots, on a large scale, have been detailed and classified by Srivastava and Srivastava [23] as liquid-gas, gas-phase, or hybrid reactors, for example, airlift bioreactors, bubble column reactors, and nutrient mist bioreactors. Conductivity measurements between small- and large-scale reactor cultures of hairy roots were compared [141]. Scaling studies showed that at 1 L *Artemisia annua* and *Arachis hypogaea* hairy roots had better growth in the mist reactor than shake flasks. The mass production of *Panax ginseng* hairy root cultures has been established in bioreactors [142] and the ginsenoside accumulation from transformed root cultures through large-scale reactor system was assessed to 1–10 t [28, 143]. Roots cultured in bioreactors have been shown to release therapeutically active compounds into the liquid medium of the bioreactor, for example, the antineoplastic drug isolated from various *Taxus* species [29].

12 Therapeutic Merits of Plant Terpene/Terpenoids

Numerous terpenoids, such as taxol, ginkgolides, digoxin, gossypol, and artemisinin, have been found efficient in the prevention and therapy of severe pathologies encompassing cancer and other severe diseases. Nevertheless, a vast proportion of higher plants have not still been screened for medicinal activities. Drug discovery from plant species remains an essential strategy in the search for future medicines [144]. From pharmaceutical point of view, terpenoids achieve a rich pool for harnessing new drugs and leading products. Cancer, after cardiovascular pathologies, is the second cause of death [145]. In fact, numerous terpenoids exhibit cytotoxicity against a wide variety of tumor cells as well as anticancer efficacy in preclinical animal models [146]. Terpenoids, a large class of mevalonate-derived phytochemicals, suppress the multiplication of neoplastic cells and the growth of implanted tumors; for example, geraniol, a volatile isoprenoid, is an acyclic dietary monoterpene which exerts an antineoplastic activity against different cancer cells both in vitro and in vivo. This antineoplastic property has successfully been assessed on TC-118 human tumors transplanted into Swiss nu/nu mice [145]. Likewise, it has been exemplified by diterpenoid drug taxol relating to the cancer [123]. Indeed, taxane biomolecules coming from genus *Taxus* represent one of the most powerful types of compounds among all the chemotherapeutic drugs efficient to treat a variety of cancers (encompassing ovarian, breast, lung, head and neck carcinomas, and the AIDS-related Kaposi's carcinoma) [56, 57]. The antitumor property results of the binding of drug to the human β -tubulin, such a chemical interaction leads to inhibit the depolymerization of the microtubules [147]. Docetaxel (taxotere) and Paclitaxel (taxol) offer some different characteristics in their molecular pharmacology, which are potentially responsible for their clinical activity and toxicity profiles [136, 148]. Taxotere displays the greater

affinity to β -tubulin, targeting centrosome organization and acting on cells in three phases of the cellular cycle (S, G2, M), whereas Taxol causes cell damage by altering the mitotic spindle at the beginning of the mitosis [57, 136]. Artemisinin (a sesquiterpene lactone) isolated from *Artemisia* sp. is an antimalarial drug efficient against multidrug-resistant strains of the malaria parasites: *Plasmodium* sp. [42], notably, *Plasmodium falciparum* that is the responsible agent for one of the world's most severe pathologies causing at least 500 million cases globally each year and resulting in more than one million deaths [149].

Pharmaceutical properties of several plant species can be due to the association of terpene/terpenoids with other active molecules. That will explain the large spectrum of therapeutic activity of these plant species. *Gynostemma pentaphyllum* (Thunb.) or jiaogulan (Chinese name), a member of the family Cucurbitaceae and genus *Gynostemma*, is an oriental medicinal herb characterized by a large spectrum of pharmacological activities, of which antitumor, antioxidant, cholesterol-lowering, immunopotentiating, and hypoglycemic effects could be observed [50]. Genus *Aesculus* (Horse chesnut) encompasses more than 210 compounds from which different classes have been isolated, in particular triterpenoids, triterpenoid glycosides (saponins), flavonoids, and long fatty chain molecules [42]. Standardized therapeutic extracts (HCSE: Horse chesnut seed extract) and aescin (a saponin mixture) of *Aesculus* sp. are considered as the main source of triterpenic glycosides, a group known as aescin and traditionally used against rheumatism, skin, and peripheral vascular disorders [43]. In Europe, the bark, leaves, HCSE, and aescin have been used in the treatment of chronic venous insufficiency, hemorrhoids, and postoperative edema [42]. Several other medicinal properties have been ascribed to saponins, such as immunostimulant, hypocholesterolaemic, and anticarcinogenic effects [150]. But, one of the most interesting properties is the antitumor action of some novel saponins found recently from the *Aesculus pavia* species [42]. The *Aesculus indica* extract, used against the MCF-7 breast cancer cell line, displayed a significant antiproliferative activity, and dose-dependent effect ranging from 34.2 % (at 10 μ -g/ml) to 94 % (at 500 μ -g/ml) could be observed [151]. Inflammation is a natural response of the mammalian body to a variety of pathologies or hostile agents. Rheumatoid arthritis is one of the challenging diseases associated with inflammatory manifestations. In order to treat similar disorders, various molecules have been isolated as, for example, a potent anti-inflammatory analgesic molecule termed "Aspirin" which was harnessed from bark of *Salix alba* L. Recent findings of several species such as *Achillea millefolium* (Asteraceae) or *Aspilia Africana* C.D. Adams (Asteraceae) and *Bacopa monnieri* (Scrophulariaceae) showed an anti-inflammatory activity due to their chemo-profiles revealing the presence of essential oils/sesquiterpenes, terpenoids (α -pinene, limonene), and triterpene, respectively [152]. Aromatic oils, stored in the leaves, are composed of aroma monoterpenes, such as eugenol, thymol, carvacrol, and linalool, and exhibit potent antioxidant properties which may prevent in vivo oxidative damages such as lipid peroxidation associated

with cancer, premature aging, atherosclerosis, and diabetes [36]. Besides, the essential oils, that is, green pesticides, are excellent alternatives to synthetic pesticides in order to reduce negative impacts to human health and the environment [37], notably the essential oils of *Pelargonium graveolens* showed insecticidal effects with a maximal mortality rate of 100 % [153].

Since ancient times, genus *Salvia* (*Lamiaceae*) has been used throughout the world, owing to its large spectrum of beneficial effects, such as its antineoplastic, antiviral, cardioactive, antidiabetic, and antiplasmodial activities [39, 40]. Among these species, *Salvia miltiorrhiza* (Chinese sage) with a rhizome possessing medicinal powers. Different molecules of *S. miltiorrhiza* (or Danshen in Chinese) are widely recommended in modern and traditional medicine for the treatment of various disorders, for example, blood-circulation diseases. The red pigment of its roots is mainly composed of numerous diterpenoid tanshinones, such as tanshinone IIA, major bioactive molecules of the *Salvia* species offering numerous pharmacological activities notably antineoplastic effects [41]. Tanshinone IIA induces apoptosis and inhibits proliferation, migration, and invasion of a number of neoplastic cell types belonging to hepatoma, osteosarcoma, leukemia, prostate cancer, gastric cancer, breast cancer, colon cancer, and glioma. Besides, the prevention and treatment of cerebral infarction by *S. miltiorrhiza* Bunge involves several action modes, including anti-atherosclerosis, anti-hypertensive, anti-platelet aggregation, anti-inflammatory, and anti-oxidative effects. Plant-derived natural products have provided many bioactive terpenoids, some of which have led to important pharmaceuticals available on the market or promoted in clinical trials by the emerging “biotech” companies [144].

13 Conclusion

Plants hold a strong potential for pharmaceutical and industrial applications, resulting from the great richness in secondary metabolites, notably in terpenes/terpenoids which represent the largest group of natural products. Potentially terpenoids/terpenes remain to be identified in numerous plant species, known or still unexplored. Plant transformation by *Agrobacterium rhizogenes* leads to the formation of hairy roots, that is, of well-differentiated roots presenting at once a high genetic stability, an unlimited propagation, and spontaneous regeneration in media devoid of growth regulators. Hairy roots and *A. rhizogenes*-derived plants are potentially promising bioreactors which can be used for producing biomolecules of high added value. Ri-plants make possible the production of secondary metabolites exclusively accumulated in the aerial parts of the plant. In the *Catharanthus roseus* leaves, the final step of terpenoid indole alkaloid (TIA) biosynthesis leads to vinblastine and vincristine (bisindole alkaloids), whereas their two precursors are produced in the roots. Moreover, Ri-regenerants can allow the creation of new varieties in horticulture/floriculture. Bacterial Ri-T-DNA, responsible for the genetic transformation, is a true secretion system based on the genetic horizontal transfer of foreign genes into the host plant genome. Thereby, hairy

roots open large avenues for engineering terpenoid metabolism pathways or revealing novel genes by recent approaches as T-DNA activation tagging or MeJA-inducible system following transcriptome analyses. Nevertheless, due to the great complexity of these pathways, numerous terpenoid biosynthesis-related genes are always to be isolated and identified in plants. Overall, hairy roots produce terpenoids/terpenes at levels comparable to or higher than those found in the wild roots. The cultures of hairy roots assure serious safety advantages compared to those of plants cultured in the open field. Biotic or abiotic elicitors have been used to enhance terpenoid contents. Most often methyl jasmonate was found to be efficient to improve the secondary metabolite synthesis. Nevertheless, the major challenge for commercial exploitation of the hairy roots is scaling up at industrial level. Based on the cell suspension culture biomass concept, specific reactors have been altered to overcome the hurdles limiting the hairy root culture on a large scale. Nowadays medicinal plants provide therapeutically active terpenoids for treating more and less severe pathologies. A vast pool of higher plants biosynthesizing secondary products remain again to be identified. Certain biomolecules of known plants have not been screened for therapeutic properties. In the near future, drug discovery from plant species is a crucial investigation to treat, notably, severe pathologies.

References

1. Poulev A, O'Neal JM, Logendra S, Pouleva RB, Timeva V, Garvey AS, Gleba D, Jenkins IS, Halpern BT, Kneer R, Cragg GM, Raskin I (2003) Elicitation a new window into plant chemodiversity and phytochemical drug discovery. *J Med Chem* 46:2542–2547
2. Chaudhuri KN et al (2009) Transgenic mimicry of pathogen attack stimulates growth and secondary metabolite accumulation. *Transgenic Res* 18:121–128
3. Tholl D, Lee S (2011) Terpene-specialized metabolism in *Arabidopsis thaliana*. *Am Soc Plant Biol*. doi:10.1199/tab.0143
4. Zwenger S, Basu C (2008) Plant terpenoids: applications and future potentials. *Biotechnol Mol Biol Rev* 3(1):1–7
5. Bensaddek L, Villarreal ML, Fliniaux MA (2008) Induction and growth of hairy roots for the production of medicinal compounds. *J Integr Biosci* 3(1):2–9
6. Croteau R, Kutchan TM, Lewis NG (2000) Natural products (secondary metabolites). In: Buchanan B, Gruissem W, Jones R (eds) *Biochemistry and molecular biology of plants*. American Society of Plant Physiologists, Rockville, pp 1250–1318
7. Bohlmann J, Keeling CI (2008) Terpenoid biomaterials. *Plant J* 54:656–669
8. Aharoni A, Jongsma MA, Kim T-Y, Ri M-B, Giri AP, Verstappen FWA, Schwab W, Bouwmeester HJ (2006) Metabolic engineering of terpenoid biosynthesis in plants. *Phytochem Rev*. doi:10.1007/s11101-005-3747-3
9. Dubey VS, Bhalla R, Luthra R (2003) An overview of the non-mevalonate pathway for terpenoid biosynthesis in plants. *J Biosci* 28(5):637–646
10. Ruiz-May E, Galaz-Avalos R, Loyola-Vargas V (2009) Differential secretion and accumulation of terpene indole alkaloids in hairy roots of *Catharanthus roseus* treated with methyl jasmonate. *Mol Biotechnol* 41:278–285
11. Morgan JA, Shanks JV (1999) Inhibitor studies of tabersonine metabolism in *C. roseus* hairy roots. *Phytochemistry* 51:61–68
12. Sevon N, Oksman-Caldentey KM (2002) *Agrobacterium rhizogenes*-mediated transformation: root cultures as a source of alkaloids. *Planta Med* 68:859–868

13. Sudha CG, Reddy BO, Ravishankar GA, Seeni S (2003) Production of ajmalicine and ajmaline in hairy root cultures of *Rauvolfia micrantha* Hook F., a rare and endemic medicinal plant. *Biotechnol Lett* 25:631–636
14. Lorence A, Medina-Bolivar F, Nessler CL (2004) Camptothecin and 10- hydroxycamptothecin from *Camptotheca acuminata* hairy roots. *Plant Cell Rep* 22:437–441
15. Caspeta L, Nieto I, Zamilpa A, Alvarez L, Quintero R, Villarreal ML (2005) *Solanum chrysotrichum* hairy root cultures: characterization, scale-up and production of five antifungal saponins for human use. *Planta Med* 71:1084–1087
16. Ahn JC, Hwang B, Hiromi T, Kanji I, Kazuo S, Koichiro S (1996) Polyacetylenes in hairy roots of *Platycodon grandiflorum*. *Phytochemistry* 42:69–72
17. Medina-Bolivar F, Condori JRA, Hubstenberger J, Shelton K, O'Keefe SF, Mc Dolan BS (2007) Production and secretion of resveratrol in hairy root cultures of peanut. *Phytochemistry* 68:1992–2003
18. Guillon S, Trémouillaux-Guiller J, Pati PK, Rideau M, Gantet P (2006) Hairy root research: recent scenario and exciting prospects. *Curr Opin Plant Biol* 9:341–346
19. Zhou X, Wang Y, Liu B, Xu H (2007) Salidroside production of hairy roots of *Rhodiola sachalinensis* obtained after transformation with *Agrobacterium rhizogenes*. *Biol Pharm Bull* 30:439–442
20. Peebles CAM, Shanks JV, San K-Y (2009) The role of the octadecanoid pathway in the production of terpenoid indole alkaloids in *Catharanthus roseus* hairy roots under normal and UV-B stress conditions. *Biotechnol Bioeng* 103(6):1248–1253
21. Tikhomiroff C, Jolicoeur M (2002) Screening of *Catharanthus roseus* secondary metabolites by high performance liquid chromatography. *J Chromatogr A* 955(1):87–93
22. Guillon S, Trémouillaux-Guiller J, Pati PK, Gantet P (2008) Hairy roots: a powerful tool for plant biotechnological advances. In: Ramawat KG, Mérillon JM (eds) *Bioactive molecules and medicinal plants*. Springer, Berlin, p 19
23. Srivastava S, Srivastava AK (2007) Hairy root culture for mass-production of high-value secondary metabolites. *Crit Rev Biotechnol* 27:29–43
24. Davioud E, Petit A, Tate ME, Ryder MH, Tempé J (1988) Cucumopine – a new T-DNA-Encoded opine in hairy root crown gall. *Phytochemistry* 27:2429–243
25. Trémouillaux-Guiller J, Andreu F, Créche J, Chénieux JC, Rideau M (1987) Variability in tissue cultures of *Choisya ternata*. Alkaloid accumulation in protoclonal and aggregate clones obtained from established strains. *Plant Cell Rep* 6:375–378
26. Trémouillaux-Guiller J, Kodja H, Andreu F, Créche J, Chénieux JC, Rideau M (1988) Variability in tissue cultures of *Choisya ternata*, III Comparing Alkaloid production in cell lines obtained by various strategies. *Plant Cell Rep* 7:456–458
27. Canto-Canche B, Loyola-Vargas VM (1999) Chemicals from roots, hairy roots, and their applications. *Adv Exp Med Biol* 464:235–275
28. Tripathi L, Tripathi JN (2003) Role of biotechnology in medicinal plants. *Trop J Res* 2(2):243–253
29. Khan MY, Saleh A, Kumar V, Rajkumar S (2009) Recent advances in medicinal plant biotechnology. *Indian J Biotechnol* 8:9–22
30. Mc Contey ME, Gershenzon J, Croteau RB (2000) Developmental regulation of mono-terpenes biosynthesis in glandular trichomes of peppermint. *Plant Physiol* 122(1):215–223
31. Martin DM, Toub O, Chiang A, Lo BC, Ohse S, Lund ST, Bohlmann J (2009) The bouquet of grapevine (*Vitis vinifera* L cv Cabernet Sauvignon) flowers arises from the biosynthesis of sesquiterpene volatiles in pollen grains. *Proc Natl Acad Sci USA* 106(17):7245–7250
32. Lückner J, Bowen P, Bohlmann J (2004) *Vitis vinifera* terpenoid cyclases: functional identification of two sesquiterpene synthase cDNAs encoding (+) -valencene synthase and (–) germacrene D synthase and expression of mono- and sesquiterpene synthases in grapevine flowers and berries. *Phytochem* 6:2649–2659
33. Aguiyi JC, Obi CI, Gang SS, Igweh AC (2000) Hypoglycaemic activity of *Ocimum gratissimum* in rats. *Fitoterapia* 71(4):444–446

34. Chaumont JP (2000) Antifungal properties of *Ocimum gratissimum* essential oils (ethyl cinnamate chemotype). *Fitoterapia* 71:567–569
35. Aidam A, Etse KD, Koba K, Raynaud C, Sanda K, Chaumont JP, Trémouillaux-Guiller J (2008) Capacités morphogénétiques in vitro, performance au champ et production d'huiles essentielles chez *Ocimum gratissimum* L. *Acta Bot Gallica* 155:341–354
36. Lee S-J, Umamo K, Shibamoto T, Lee K-G (2005) Identification of volatile components in basil (*Ocimum basilicum* L.) and thyme leaves (*Thymus vulgaris* L.) and their antioxidant properties. *Food Chem* 91:131–137
37. Koul O, Walia S, Dhaliwal GS (2008) Essential oils as green pesticides: potential and constraints. *Biopestic Int* 4(1):63–84
38. Celep F, Kahraman A, Dogan M (2011) A new taxon of the genus *Salvia* (Lamiaceae) from Turkey. *Plant Ecol Evol* 144(1):111–114
39. Fraga BM, Diaz CE, Guadano A, Gonzales-Coloma A (2005) Diterpenes from *Salvia broussonetti* transformed roots and their insecticidal activity. *J Agric Food Chem* 53:5200–5206
40. Kuzma L, Skrzypek Z, Wysokinska H (2006) Diterpenoids and triterpenoids in hairy roots of *Salvia sclarea*. *Plant Cell Tiss Org Cult* 8:171–179
41. Wu SJ, Shi M, Wu JY (2009) Cloning and characterization of the 1-deoxy-D-xylulose 5-phosphate reductoisomerase gene for diterpenoid tanshinone biosynthesis in *Salvia miltiorrhiza* (Chinese sage) hairy roots. *Biotechnol Appl Biochem* 52:89–95
42. Zang Z, Li S, Lian X-Y (2010) An overview of *Genus Aesculus* L.: ethnobotany, phytochemistry, and pharmacological activities. *Pharm Crops* 1:24–51
43. Calic-Dragosavac D, Stevovic S, Zdravkovic-Korac S, Milojevic J, Cingel A, Vinterhalter B (2011) Secondary metabolite of horse chestnut in vitro culture. *Adv Environ Biol* 5(2):267–270
44. Choi DW, Chung H-J, Ko SM, In DS, Song JS, Woo S-S, Liu JR (2009) Mass production and application tagged hairy root lines for functional genomic of secondary metabolism in ginseng. *J Plant Biotechnol* 36:294–300
45. Ernst E (2010) *Panax ginseng*: an overview of the clinical evidence. *J Ginseng Res* 34(4):259–263. doi:10.5142/jgr.2010.34.4.259
46. de Sousa Braz A, Diniz M, de FF M, De Almeida RN (2009) Recent advances in the use of *Panax ginseng* as an analgesic: a systematic review. *Boletín Latinoamericano y del Caribe de Plantas Medicinales y Aromáticas* 8(3):188–194
47. Wang H, Peng D, Xie J (2009) *Ginseng* leaf-stem: bioactive constituents and pharmacological functions. *Chin Med* 4:20–28. doi:10.1186/1749-8546-4-20
48. Pu Z, Zhang P et al (2011) The evaluation of contents of nine ginsenoside monomers in *ginseng* hairy roots by high performance liquid chromatography (HPLC). *J Med Plant Res* 5(23):5513–5516
49. Yue PYK, Mak NK, Cheng YK, Leung KW, Ng TB, Fan DTP, Yeung HW, Wong RNS (2007) Pharmacogenomics and the yin/yang actions of ginseng: anti-tumor, angiomodulating and steroid-like activities of ginsenosides. *Chin Med* 2:6. doi:10.1186/1749-8546-2-6
50. Norberg A, Hoa NK, Liepinsh E, Phan DV, Thuan ND, Jörnvall H, Sillard R, Ostenson C-G (2004) A novel insulin-releasing substance, phanoside, from the *Gynostemma pentaphyllum*. *J Biol Chem* 279:401361–401367
51. Chang CK, Chang KS, Lin YC, Liu SY, Chen CY (2005) Hairy root cultures of *Gynostemma pentaphyllum* Thunb Makino a promising approach for the production of gypenosides as an alternative of ginseng saponins. *Biotechnol Lett* 27:1165–1169
52. Pellicer J, Garnatje T, Vallès J (2011) *Artemisia* (Asteraceae): understanding its evolution using cytogenetic and molecular systematic tools, with emphasis on subgenus *Dracunculus*. In: Munoz-Torrero D (ed) Recent advances in pharmaceutical sciences. Transworld Research Network, Trivandrum, pp 199–222. ISBN: 978-81-7895-528-5

53. Hayat HQ, Ashraf M, Khan MA, Mahmood T, Ahmad M, Jabeen S (2009) Phylogeny of *Artemisia* L: recent developments. *Afr J Biotechnol* 8(11):2423–2428
54. Olofsson L, Engström A, Lundgren A, Brodelius PE (2011) Relative expression of genes of terpene metabolism in different tissues of *Artemisia annua* L. *BMC Plant Biol* 11:45–57. doi:1186/147-2229-11-45
55. Wani MC, Taylor HL, Wall ME, Coggon P, McPhail AT (1971) Plant antitumor agents VI the isolation and structure of taxol a novel antileukemic and antitumor agent from *Taxus brevifolia*. *J Am Chem Soc* 93(9):2325–2327
56. Heinig U, Jennewein S (2009) Taxol: a complex diterpenoid natural product with an evolutionarily obscure origin. *Afr J Biotechnol* 8(8):1370–1385
57. Kim JA, Baek K-H, Son YM, Son SH, Shin H (2009) Hairy root cultures of *Taxus cuspidate* for enhanced production of paclitaxel. *J Korean Soc Appl Biol Chem* 52(2):144–150
58. Exposito O, Syklovska-Baranek K, Moyano E, Onrubia M, Bonfill M, Palazon J, Cusido RM (2010) Metabolic responses of *Taxus media* transformed cell cultures to the addition of methyl jasmonate. *Biotechnol Prog* 26(4):1145–1153
59. Lavelle F, Guéritte-Voegelein F, Guénard D (1993) Le taxotère: des aiguilles d’if à la clinique. *Bull de cancérologie* 80:326–338
60. Riker AJ, Banfield WM, Wright WH, Keitt WH, Sagen HE (1930) *J Agr Res* 41:507–540
61. Willmitzer L, Sanchez-Serrano J, Buschfeld E, Schell J (1982) DNA from *Agrobacterium rhizogenes* is transferred to and expressed in axenic hairy root plant tissues. *Mol Gen Genet* 186:16–22
62. Chilton M-D, Tepfer DA, Petit A, David C, Casse-Delbart F, Tempe J (1982) *Agrobacterium rhizogenes* inserts T-DNA into the genomes of the host plant root cells. *Nature* 295:432–434
63. Bercetche J, Chriqui D, Adam S, David C (1987) Morphogenetic and cellular reorientations induced by *Agrobacterium rhizogenes* (strains 1855, 2659 and 8196) on carrot, pea and tobacco. *Plant Sci* 52:195–210
64. White FF, Nester EW (1980) Hairy root: plasmids encode virulence traits in *Agrobacterium rhizogenes*. *J Bacteriol* 141:1134–1141
65. Tepfer D, Metzger L, Prost R (1989) Use of roots transformed by *Agrobacterium rhizogenes* in rhizosphere research: applications in studies of cadmium assimilation from sewage sludges. *Plant Mol Biol* 13:295–302
66. Veena V, Taylor C (2007) *Agrobacterium rhizogenes* recent developments and promising applications. *In vitro Cell Dev Biol-Plant* 43:383–403
67. Camilleri C, Jouanin L (1991) TR-DNA region carrying the auxin synthesis gene of the *Agrobacterium rhizogenes* agropine type plasmid PRiAA4: nucleotide sequence analysis and introduction into tobacco plants. *Mol Plant Microbe Interact* 4:155–162
68. Slightom JL, Durand-Tardif M, Jouanin L, Tepfer D (1986) Nucleotide sequence analysis of TL-DNA of *Agrobacterium rhizogenes* agropine type plasmid: identification of open-reading frames. *J Biol Chem* 261:108–121
69. Goldmann A, Tempé J, Morel G (1968) Découvertes de nouvelles substances, les opines produites par les cellules des tumeurs. Quelques particularités de diverses souches d’*Agrobacterium tumefaciens*. *CR Sci Biol* 162:630–631
70. Petit A, Delhay S, Tempé J, Morel G (1970) Recherches sur les guanidines des tissus de crown gall. Mise en évidence d’une relation biochimique spécifique entre les souches d’*Agrobacterium tumefaciens* et les tumeurs qu’elles induisent. *Physiol Vég* 8:205
71. Tepfer D, Tempé J (1981) Production d’agropine par de racines formées sous l’action d’*Agrobacterium rhizogenes*, souche A4. *CR Acad Sci Paris* 292:153–156
72. De Paolis A, Mauro HL, Pomponi M, Cardarelli M, Spano L, Constantino P (1985) Localization of agropine-synthesizing functions in the TR regions of the root inducing plasmid of *Agrobacterium rhizogenes* 1865. *Plasmid* 13:1–7
73. Ayadi R, Tremouillaux-Guiller J (2003) Root formation from transgenic calli of *Ginkgo biloba*. *Tree Physiol* 23:437–441

74. Davioud E, Petit A, Tate ME, Ryder MH, Tempé J (1988) Cucumopine – a new T-DNA-encoded opine in hairy root crown gall. *Phytochemistry* 27:2429–2433
75. Tzfira T, Vaidya M, Citovsky V (2004) Involvement of targeted proteolysis in plant genetic transformation by *Agrobacterium*. *Nature* 431:92
76. Tzfira T, Li J, Lacroix B, Citovsky V (2004) *Agrobacterium* T-DNA integration: molecules and models. *Trends Genet* 20:375–383
77. Lacroix B, Vaidya M, Tzfira T, Citovsky V (2005) The VirE3 protein of *Agrobacterium* mimics a host cell function required for plant genetic transformation. *EMBO J* 24: 428–437
78. Gelvin SB (2009) *Agrobacterium* in the genomics age. *Plant Physiol* 150:1665–1668
79. Pitzschke A, Heribert H (2010) New insights into an old story: *Agrobacterium*-induced tumor formation in plants by plant transformation. *EMBO J* 29:1021–1028
80. Pawar PK, Maheshwari VL (2004) *Agrobacterium rhizogenes* mediated hairy root induction in two medicinally important members of family Solanaceae. *India J Biotechnol* 3:414–417
81. Wysokinska H, Chmiel A (1997) Transformed root cultures for biotechnology. *Acta Biotechnol* 17:131–159
82. David C, Chilton MD, Tempé J (1984) Conservation of T-DNA in plants regenerated from hairy root cultures. *Biotechnology* 2:73–76
83. Tepfer D (1984) Transformation of several species of higher plants by *Agrobacterium rhizogenes*: sexual transmission of the transformed genotype and phenotype. *Cell* 37:959
84. Phelep M, Petit A, Martin L, Duhoux E, Tempé J (1991) Transformation and regeneration of a nitrogen-fixing tree, *Allocauarina Verticillata* Lam. *Nat Biotechnol* 9:461–466
85. Flores HE, Vivanco JM, Loyola-Vargas VM (1999) ‘Radicle’ biochemistry: the biology of root-specific metabolism. *Trends Plant Sci* 4:220–226
86. Huang T-K, Mc Donald KA (2011) Bioreactor systems for in vitro production of foreign proteins using plant cell cultures. *Biotechnol Adv* 30(2):398–409
87. Guillon S, Gantet P, Thiersault M, Rideau M, Trémouillaux-Guiller J (2008) Hairy roots of *Catharanthus roseus*: efficient routes to monomeric indole alkaloid production. In: Ramawat KG, Mérillon JM (eds) *Bioactive molecules and medicinal plants*. Springer, Berlin, pp 271–283
88. Rigano MM, Walmsley AM (2005) Expression systems and developments in plant-made vaccines. *Immunol Cell Biol* 83:271–277
89. Subroto MA, Artanti N, Sudrajat D, Djanakum A, Widayat E (2001) *Agrobacterium rhizogenes*-mediated transformation of *Solanum nigrum* L: spontaneous plant regeneration and endogenous IAA contents. *Indon J Agric Sci* 1:53–59
90. Christensen B, Sriskandarajah S, Serek M, Müller R (2008) Transformation of *Kalanchoe blossfeldiana* with rol-genes is useful in molecular breeding towards compact growth. *Plant Cell Rep* 27:1485–1495. doi:1007/s00299-008-0575-0
91. Mishiba KI, Masahiro N, Yoshiko A, Takashi N, Hiromi K, Katsuo K, Toshikazu T, Jun A, Saburo Y (2006) Production of dwarf potted gentian using wild-type *Agrobacterium rhizogenes*. *Plant Biotechnol* 23:33–38
92. Crane C, Wright E, Dixon RA, Wang ZY (2006) Transgenic *Medicago truncatula* plants obtained from *Agrobacterium tumefaciens*-transformed roots and *Agrobacterium rhizogenes*-transformed hairy roots. *Planta* 223:1344–1354
93. Boland W, Garms S (2010) Induced volatiles of *Medicago truncatula*: molecular diversity and mechanistic aspects of a multiproduct sesquiterpene synthase from *M. truncatula*. *Flavour Fragr J* 25:114–116
94. Guerche P, Jouanin L, Tepfer D, Pelletier G (1987) Genetic transformation of oilseed rape (*Brassica napus*) by the Ri T-DNA of *Agrobacterium rhizogenes* and analysis of inheritance of the transformed phenotype. *Mol Gen Genet* 206:382–386
95. Casanova E, Trillas MI, Moysset L, Vainstein A (2005) Influence of rol genes in floriculture. *Biotechnol Adv* 23:3–39

96. Zdravkovic-Korac S, Muhovski Y, Druart P, Calic D, Radojevic L (2004) *Agrobacterium rhizogenes*-mediated DNA transfer to *Aesculus hippocastanum* L and the regeneration of transformed plants. *Plant Cell Rep* 22:698–704
97. Komaraiah P, Reddy GV, Reddy PS, Raghavendra AS, Ramakrishna SV, Reddanna P (2003) Enhanced production of antimicrobial sesquiterpenes and lipoxygenases metabolites in elicitor-treated hairy root cultures of *Solanum tuberosum*. *Biotechnol Lett* 25:593–597
98. Saxena G, Banerjee S, Rahman L, Verma PC, Mallavarapu GR, Kumar S (2007) Rose-scented geranium (*Pelargonium* sp) generated by *Agrobacterium rhizogenes* mediated Ri insertion for improved essential oil quality. *Plant Cell Tiss Org Cult* 90:215–223
99. Jerkovic I, Gaso-Sokac D, Pavlovic H, Marijanovic Z, Gugic M, Petrovic I, Kovac S (2012) Volatile organic compounds from *Centaureum erythraea* Rafn (Croatia) and the antimicrobial potential of its essential oil. *Molecules* 17:2058–2072. doi:10.3390/molecules17022058
100. Piatczak E, Krolicka A, Wysokinska H (2006) Genetic transformation of *Centaureum erythraea* Rafn by *Agrobacterium rhizogenes* and the production of secoiridoids. *Plant Cell Rep* 25:1308–131
101. Yang DC, Choi YE (2000) Production of transgenic plants via *Agrobacterium rhizogenes*-mediated transformation of *Panax ginseng*. *Plant Cell Rep* 19:491–496
102. Gorpenchenko T, Kiselev K, Bulgakov V, Tchernoded G, Bragina E, Khodakovskaya M, Koren OG, Batygina T, Zhuravlev Y (2006) The *Agrobacterium rhizogenes* rolC-gene-induced somatic embryogenesis and shoot organogenesis in *Panax ginseng* transformed calluses. *Planta* 223:457–467
103. Kang HJ, Anbazhagan VR, You XL, Moon HK, Yi JS, Choi YE (2006) Production of transgenic *Aralia elata* regenerated from *Agrobacterium rhizogenes*-mediated transformed roots. *Plant Cell Tiss Org Cult* 85:187–196
104. Kim JA, Kim Y-S, Choi Y-E (2011) Triterpenoid production and phenotypic changes in hairy roots of *Codonopsis lanceolata* and the plants regenerated from them. *Plant Biotechnol Rep* 5:255–263. doi:10.1007/s11816-011-0180-5
105. Inoue F, Sugiura H, Tabuchi A, Kawasawa DE, Minami M (2003) Plant regeneration of peppermint, *Mentha piperita* from the hairy roots generated from microsegment infected with *Agrobacterium rhizogenes*. *Plant Biotechnol* 20:169–172
106. Christey MC (2001) Use of Ri-mediated transformation for production of transgenic plants. In *Vitro Cell Dev Biol Plant* 3:687–700
107. Ackermann C (1977) C pflanzen aus *Agrobacterium rhizogenes*-tumoren aus *Nicotiana tabacum*. *Plant Sci Lett* 8:23–30
108. Vinterhalter B, Ninković S, Cingel A, Vinterhalter D (2006) Shoot and root culture of *Hypericum perforatum* L. transformed with *Agrobacterium rhizogenes* A4M70GUS. *Biol Plant* 50:767–770
109. David C, Petit A, Tempé J (1988) T-DNA length variability in mannopine hairy root: more than 50 kilobasepairs of pRi T-DNA can integrate in plant cells. *Plant Cell Rep* 7:92–95
110. Fründt C, Meyer AD, Ichikawa T, Meins F (1998) A tobacco homologue of the Ri-plasmid *orf 13* gene causes cell proliferation in carrot discs. *Mol Gen Genet* 259:559–568
111. Franche C, Duhoux E (2001) Du transfert d'ADN à l'obtention d'une plante transgénique. In: Elsevier (ed) *La transgénèse végétale Bio campus*, Amsterdam
112. Collier R, Fuchs B, Walter N, Kevin Lutke W, Taylor CG (2005) Ex vitro composite plants: an inexpensive, rapid method for root biology. *Plant J* 43:449–457
113. Bottino PJ, Raineri D, Nester FW, Gordon MP (1989) *Agrobacterium*-mediated DNA transfer. *J Tiss Cult Methods* 12(4):135–138
114. Sheki H, Nishizawa T, Tanaka N, Niwa Y, Yoshida S, Muranaka T (2005) Hairy root-activation tagging: a high-throughput system for activation tagging in transformed hairy roots. *Plant Mol Biol* 59:793–807
115. Voinnet O, Baulcombe DC (1997) Systemic signalling in gene silencing. *Nature* 389:553
116. Busov V, Yordanov Y, Gou J, Meilan R, Ma C, Regan S, Strauss S (2011) Activation tagging is an effective gene tagging system in *Populus*. *Tree Genet Genomes* 7:91–101

117. Guillon S, Trémouillaux-Guiller J, Pati PK, Rideau M, Gantet P (2006) Harnessing the potential of hairy roots: dawn of a new era. *Trends Biotechnol* 24:403–409
118. Ayliffe MA, Pallotta M, Langridge P, Pryor AJ (2007) A barley activation tagging system. *Plant Mol Biol* 64(3):329–347
119. Lee C-Y, Agrawal DC, Chu T-M, Wang C-S, Yu S-M, Chen JJW, Tsay H-S (2008) T-DNA activation tagging as a tool to isolate *Salvia miltiorrhiza* Bunge transgenic lines for higher yields of pharmacological tanshinones. *Planta Med* 74(7):780–786
120. Choi D-W, Jung JD, Ha YI, Park H-W, In DS, Chung HJ, Liu JR (2005) Analysis of transcripts in methyl jasmonate-treated ginseng hairy roots to identify genes involved in the biosynthesis of ginsenosides and other secondary metabolites. *Plant Cell Rep* 23:557–566
121. Jian B, Hou W, Wu C, Liu B, Liu W, Song S, Bi Y, Han T (2009) *Agrobacterium rhizogenes*-mediated transformation of Superroot-derived *Lotus corniculatus* plants: a valuable tool for functional genomics. *BMC Plant Biol* 9:78–92
122. Zang L, Ding R, Chai Y, Bonfil M, Moyano E, Oksman-Caldentey K-M, Xu T, Pi Y, Wang Z, Zhang H, Kai G, Liao Z, Sun X, Tang K (2004) Engineering tropae biosynthetic pathway in *Hyoscyamus niger* hairy root cultures. *Proc Natl Acad Sci USA* 101(17):6786–6791
123. Cheng A-X, Lou Y-G, Mao Y-B, Lu S, Wang L-J, Chen X-Y (2007) Plant terpenoids: biosynthesis and ecological functions. *J Integr Plant Biol* 49(2):179–186
124. Ma Y, Yuan L, Wu B, X'en L, Chen S, Lu S (2012) Genome-wide identification and characterization of novel genes involved in terpenoid biosynthesis in *Salvia miltiorrhiza*. *J Exp Bot* 1–15. doi:10.1093/jxb/err466
125. Faria JMS, Nunes IS, Figueiredo AC, Pedro LG, Trindade H, Barroso JG (2009) Biotransformation of menthol and geraniol by hairy root cultures of *Anethum graveolens*: effect on growth and volatile components. *Biotechnol Lett* 31:897–903. doi:10.1007/s10529-009-9934-3
126. Hu Z-B, Du M (2006) Hairy root and its application in plant genetic engineering. *J Integr Plant Biol* 4(2):121–127
127. Bulgakov VP (2008) Functions of rol genes in plant secondary metabolism. *Biotechnol Adv* 26:318–324
128. Taneja J, Jaggi M, Wankhede DP, Sinha AK (2010) Effect of loss of T-DNA genes on MIA biosynthetic pathway gene regulation and alkaloid accumulation in *Catharanthus roseus* hairy roots. *Plant Cell Rep* 29:1119–1129
129. Mannan A, Shaheen N, Arshad W, Qureshi RA, Zia M, Mirza B (2008) Hairy roots induction and artemisinin *Artemisia dubia* and *Artemisia indica*. *Afr J Biotechnol* 7(18):3288–3292
130. Giri A, Narasu ML (2000) Transgenic hairy roots – recent trends and applications. *Biotechnol Adv* 18:1–22
131. Bais HP, Walker TS, Schweizer HP, Vivanco JM (2002) Root specific elicitation and microbial activity of rosmarinic acid in hairy root cultures of *Ocimum basilicum*. *Plant Physiol Biochem* 40:983–995
132. Uddin MR, Li X, Won OJ, Park SU, Pyon JY (2011) Herbicidal activity of phenolic compounds from hairy root cultures of *Fagopyrum tataricum*. *Weed Res.* doi:10.1111/j.01365-3180.2011.00894.x
133. Demole E, Lederer E, Mercier DE (1962) Isolement et détermination de la structure du jasmonate de méthyle constituant odorant caractéristique de l'essence de jasmin. *Helvetica Chimica Acta* 45:675–685
134. Rohwer CL, Erwin JE (2008) Horticultural applications of jasmonates: a review. *J Horticultural Sci Biotechnol* 83(3):283–304
135. Frankfater CR, Dowd MK, Triplett BA (2009) Effect of elicitors on the production of gossypol and methylated gossypol in cotton hairy roots. *Plant Cell Tiss Org Cult* 98:341–349
136. Ashrafi S, Mofid MR, Otroshi M, Ebrahimi M, Khosroshahli M (2010) Effects of plant growth regulators on the callogenesis and taxol production in cell suspension of *Taxus baccata* L. *Trakia J Sci* 8(2):36–43

137. Syklowska-Baranek K, Pietrosiuk A, Kokoszka A, Furmanowa M (2009) Enhancement of taxane production in hairy root culture of *Taxus x media* var. Hicksii. *J Plant Physiol* 166:1950–1954
138. Ge X, Wu J (2005) Induction and potentiation of diterpenoid tanshinone accumulation in *Salvia miltiorrhiza* hairy roots by β -aminobutyric acid. *Appl Microbiol Biotechnol* 68:183–188
139. Ge X, Wu J (2005) Tanshinone production and isoprenoid pathways in *Salvia miltiorrhiza* hairy roots induced by Ag^+ and yeast elicitor. *Plant Sci* 168:487–491
140. Verpoorte R, van der Heijden R, ten Hoopen HJG, Memelink J (1999) Metabolic engineering of plant secondary metabolite pathways for the production of fine chemicals. *Biotechnol Lett* 21:467–479
141. Sivakumar G, Liu C, Towler MJ, Weathers PJ (2010) Biomass production of hairy roots of *Artemisia annua* and *Arachis hypogaea* in a scaled-up Mist bioreactor. *Biotechnol Bioeng* 107(5):802–813
142. Jeong GT, Park DH, Hwang B, Park K, Kim SW, Woo JC (2002) Studies on mass production of transformed *Panax ginseng* hairy roots in bioreactor. *Appl Biochem Biotechnol* 98:1115–1127
143. Hahn EJ, Kim YS, Yu KW, Jeong CS, Paek KY (2003) Adventitious root cultures of *Panax ginseng* c.v. Meyer and ginsenoside production through large scale bioreactor system. *J Plant Biotechnol* 5:1–6
144. Salim AA, Chin Y-W, Kinghorn AD (2008) Drug discovery from plants. In: Ramawat KG, Mérillon JM (eds) *Bioactive molecules and medicinal plants*. Springer, Berlin, pp 1–24
145. Amin A, Mousa M (2007) Merits of anti-cancer plants from the Arabian Gulf region. *Cancer Ther* 5:55–66
146. Thoppil RJ, Bishayee A (2011) Terpenoids as potential chemopreventive and therapeutic agents in liver cancer. *World J Hepatol* 3(9):228–249
147. Zao K, Sun L, Ma X, Li X, Wang X, Ping W, Zhou D (2011) Improved taxol production in *Nodulisporium sylviforme* derived from inactivated protoplast fusion. *Afr J Biotechnol* 10(20):4175–4182
148. Gligorov J, Lotz J-P (2004) Preclinical pharmacology of the taxanes: implications of the differences. *Oncologist* 9(Suppl 2):3–8
149. Qian Z, Gong K, Zhang L, Lv J, Jing F, Wang Y, Guan S, Wang G, Tang K (2007) A simple and efficient procedure to enhance artemisinin content in *Artemisia annua* L by seeding to salinity stress. *Afr J Biotechnol* 6(12):1410–1413
150. Francis G, Kerem Z, Makkar HPS, Becker K (2002) The biological action of saponins in animal systems: a review. *Br J Nutr* 88:587–605
151. Bibi Y, Nisa S, Zia M, Waheed A, Ahmed S, Chaudhary F (2012) In vitro cytotoxic activity of *Aesculus indica* against breast adenocarcinoma cell line (MCF-7) and phytochemical analysis. *Pak J Pharm Sci* 25(1):183–187
152. Agnihotri S, Wakode S, Agnithori A (2010) An overview on anti-inflammatory properties and chemo-profiles of plants used in traditional medicine. *Indian J Nat Prod Resour* 1(2):150–167
153. Kabera J, Gasogo A, Uwamariya A, Ugirinshuti V, Nyetera P (2011) Insecticidal effects of essential oils of *Pelargonium graveolens* and *Cymbopogon citrates* on *Sitophilus zeamais* (Motsch.). *Afr J Food Sci* 5(6):366–375
154. Sallaud C, Rontein D, Onillon S, Jabès F, Duffé P, Giacalone C, Thoraval S, Escoffier C, Herbet G, Leonhardt N, Causse M, Tissier A (2009) A novel pathway for sesquiterpene biosynthesis from Z, Z-farnesyl pyrophosphate in the wild tomato *Solanum habrochaites*. *Plant Cell* 21:301–317
155. Laurain D, Trémouillaux-Guiller J, Chénieux J-C, van Beek TA (1997) Production of ginkgolide and bilobalide in transformed and gametophyte derived cell cultures of *Ginkgo biloba*. *Phytochemistry* 46:127–130

156. Choi PS, Kim YD, Choi KM, Chung HJ, Choi DW, Liu JR (2004) Plant regeneration from hairy-root cultures transformed by infection with *Agrobacterium rhizogenes* in *Catharanthus roseus*. *Plant Cell Rep* 22:828–831
157. Mehrotra S, Kukreja AK, Khanuja SPS, Mishra BN (2008) Genetic transformation studies and scale up of hairy root culture of *Glycyrrhiza glabra* in bioreactor. *Electronic J Biotechnol* 5:199–222
158. Giri A, Ravindra ST, Dhingra V, Narasu ML (2001) Influence of different strains of *Agrobacterium rhizogenes* on induction of hairy roots and artemisinin production in *Artemisia annua*. *Curr Sci* 81:378–382
159. Sandhu NS, Kaur S, Chopra D (2010) *Equisetum arvense*: pharmacology and phytochemistry – a review. *Asian J Pharm Clin Res* 3(3):146–150
160. Makoto I, Ohta S, Komoto N, Ushijima M, Kodera Y, Hayama M, Shirota O, Sekita S, Kuroyanagi M (2008) Rapid identification of terpenoid saponins in the root of *Codonopsis lanceolata* by liquid chromatography-mass spectrometry. *J Nat Med* 62(4):423–429. doi:10.1007/s11418-008-0270-z
161. Makky EA, Mashitah MY, Ibrahim MM (2012) Impact of medicinal plant phytocomponents against antibiotic resistant bacteria. *J Chem Pharm Res* 4(1):881–893