

Using electrospray laser desorption ionization mass spectrometry to rapidly examine the integrity of proteins stored in various solutions

Yi-Tzu Cho · Min-Zong Huang · Sih-You Wu ·
Ming-Feng Hou · Jianjun Li · Jentaie Shiea

Received: 14 September 2013 / Revised: 5 November 2013 / Accepted: 6 November 2013 / Published online: 18 December 2013
© Springer-Verlag Berlin Heidelberg 2013

Abstract Electrospray laser desorption ionization mass spectrometry (ELDI/MS) allows the rapid desorption and ionization of proteins from solutions under ambient conditions. In this study, we have demonstrated the use of ELDI/MS to efficiently examine the integrity of the proteins stored in various solutions before they were further used for other biochemical tests. The protein standards were prepared in the solutions containing buffers, organic salts, inorganic salts, strong acid, strong base, and organic solvents, respectively, to simulate those collected from solvent extraction, filtration, dialysis, or chromatographic separation. Other than the deposit of a drop of the sample solution on the metallic sample plate in an ELDI source, no additional sample pretreatment is needed. The sample drop was then irradiated with a pulsed laser; this led to desorption of the analyte molecules, which subsequently entered the ESI plume to undergo post-

ionization. Because adjustment of the composition of the sample solution is unnecessary, this technique appears to be useful for rapidly evaluating the integrity of proteins after storage or prior to further biochemical treatment. In addition, when using acid-free and low-organic-solvent ESI solutions for ELDI/MS analysis, the native conformations of the proteins in solution could be detected.

Keywords Electrospray laser desorption ionization mass spectrometry · ELDI · Protein integrity · Native conformation

Introduction

In many circumstances, proteins are stored in different solutions prior to their further use. For example, (a) protein fractions from extraction, filtration, dialysis, or chromatographic separation are stored in various buffers, salts, or organic solvents; (b) integral membrane proteins are extracted and purified in detergent-containing solutions [1]; (c) water-soluble peripheral membrane proteins are stored in high-salt or high-pH solutions for better extraction from membrane pellets [2]; (d) the recoveries of certain types of proteins from inclusion bodies are improved by solubilizing the proteins in high-pH solutions or in detergents (e.g., urea, guanidine hydrochloride, Triton-X 100) [3]; (e) edible proteins from animal muscle are solubilized in alkaline aqueous solutions [4]; and (f) after being captured by immunomagnetic nanoparticles, proteins are sometimes processed and stored in alkaline solutions [5]. Evaluating the structures and integrity of proteins in the purification procedures or after storage is always important. Mass spectrometry is indeed a useful tool for detecting protein molecules; however, various compositions of solutions make the detection of proteins time consuming.

Since its development in 1984, electrospray ionization mass spectrometry (ESI/MS) has played an essential role in

Y.-T. Cho

Department of Cosmetic Applications and Management,
Yuh-Ing Junior College of Health Care & Management,
No.15 Lane 420 Dachang 2nd Rd, Kaohsiung 807, Taiwan
e-mail: ytcho@ms.yuhing.edu.tw

M.-Z. Huang · S.-Y. Wu · J. Shiea (✉)

Department of Chemistry, National Sun Yat-Sen University,
70 Lienhai Rd, 80424 Kaohsiung, Taiwan
e-mail: jetea@fac.nsysu.edu.tw

M.-F. Hou · J. Shiea

Cancer Research Center, Kaohsiung Medical University Hospital,
100, Shih-Chuan 1st Road, 80708 Kaohsiung, Taiwan

J. Li

Institute for Biological Sciences, National Research Council Canada,
6000 Montreal Road, Ottawa ON K1A 0R6, Ontario, Canada

J. Shiea

Department of Medicinal and Applied Chemistry,
Kaohsiung Medical University, 100, Shi-Chuan 1st Road,
Kaohsiung 80708, Taiwan

characterizing biomolecules, including proteins, peptides, and metabolites [6–8]. For the practical analysis of protein or peptide mixtures, ESI/MS is usually coupled with liquid chromatography [9–11], but it can also be used in a standalone manner to characterize proteins or peptides in the solutions [12]. For this purpose, ESI/MS has been used to examine the integrity of proteins—that is, to assess their degrees of denaturation or decomposition, as caused by proteases, acids, bases, organic solvents, or chemical contaminants [13]. Typically, however, sample pretreatment must be performed prior to ESI/MS analysis to prevent ion suppression effects, to remove interference, or to increase the signal intensity of the protein ions. For example, alkaline protein solutions must be adjusted or diluted using electrospray solutions that typically contain organic solvents and organic acids. Although it is possible to generate protein ions having polarity opposite to that of the proteins present in solution, the number of protein ions tend to be lower than that obtained when the ESI polarity is the same as the net charge of the proteins in solution [12]. Thus, to detect proteins from alkaline solutions, either the ESI/MS system should be switched to the negative ion mode or pH adjustment and buffer dilution must be performed. In addition to the effects of alkaline solutions, interference from buffers and detergents are constant problems in ESI/MS analyses of protein samples; typically, these chemical components must be removed prior to analysis [14–18]. Moreover, the presence of moderate amounts of inorganic salts in a sample solution will decrease the stability of the electrospray and significantly affect the intensity of the analyte ion signals, due to ion suppression effects and adduct ion formation [19, 20]. Desalting processes are, therefore, required to obtain good-quality ESI mass spectra; such sample pretreatment procedures are, however, tedious and will make these analyses very time-consuming [14–18]. The development of analytical techniques for characterizing proteins directly from various sample solutions would make it possible to screen large numbers of sample solutions in an efficient manner.

Electrospray laser desorption ionization (ELDI), a two-step desorption ionization process, allows desorption and ionization of chemical and biological compounds directly from sample solutions under ambient conditions [21–30]. It has been applied to characterize non-volatile analyte molecules directly from the surfaces of samples. Irradiating the sample surface with a pulsed laser leads to desorption of the analyte molecules, which subsequently enter the ESI plume to undergo post-ionization [21–30]. In contrast to conventional ESI, the sample introduction and ionization processes in ELDI are two separate events; this feature allows independent control of each set of condition, thereby enhancing the potential use in various practical chemical or biochemical applications. For example, the status of ongoing chemical and biochemical reactions occurring in organic solutions can be monitored continuously using ELDI/MS [27]. In addition, reactive-ELDI provides additional analytical

capabilities by incorporating gas phase reactions prior to ELDI/MS measurement [28].

Herein, we report the use of ELDI/MS under ambient conditions to rapidly produce high-quality mass spectra of proteins stored in the solutions at different pH, organic solvent contents, buffers, salts, or detergents. Comparatively, positively charged proteins ions can be directly detected through ELDI/MS without the need for any sample pretreatment. In addition, by adjusting the composition of the ESI solution in an ELDI/MS analysis, the conformations of proteins stored in solutions with different pH was rapidly determined.

Material and methods

The chemicals and protein standards including cytochrome c, myoglobin, urea, guanidine·HCl, tetrabutylammonium bromide (TBAB), ammonium sulfate (AMS), glycerol, sodium dodecyl sulfate (SDS), 3-[(3-cholamidopropyl)dimethylammonio]-1-propane sulfonate (CHAPS), octyl phenol ethoxylate (Triton X-100), and various buffers and salts were purchased from Sigma and Aldrich (Milwaukee, WI) and used without further purification. The organic solvents (HPLC grade) were purchased from Merck (Darmstadt, Germany); acetic acid (reagent grade) was purchased from J. T. Baker (Phillipsburg, NJ). Distilled deionized water, obtained using a Milli-Q plus system (Millipore; Molsheim, France), was used to prepare the sample solutions. The stock solutions each contained sodium dihydrogen phosphate, sodium acetate, sodium carbonate, and tris(hydroxymethyl)methylamine (25 mM). The pH of the buffered solutions was adjusted by adding HCl (1 M) or NaOH (1 M). The stock protein solutions were diluted in different buffer solutions to obtain protein concentrations of 10^{-4} M.

A drop of the protein sample solution (ca. 5 μ L) was deposited through a micropipette onto a stainless-steel sample plate [5 (L) \times 2 (W) cm^2], which was then positioned on an acrylic plate placed on a XYZ stage and set in front of the sampling capillary of an ion trap mass spectrometer (Esquire 3000 plus, Bruker Daltons). The surface of the sample droplet was irradiated with a pulsed Q-switched Nd:YAG laser (MINILITE I, Continuum, CA, USA) under ambient conditions. The laser was operated at a wavelength of 266 nm, a frequency of 10 Hz, a pulsed energy of approximately 250 μ J, and a pulse duration of 5 ns; the laser beam was focused through an objective lens to create a spot size of approximately $5 \times 10^{-3} \text{ cm}^2$. The strongest ion signal was obtained at an incident laser angle of approximately 45° .

The laser-ablated molecules were post-ionized in an electrospray plume. The capillary electrosprayer was aligned parallel to and approximately 3.5 mm above the acrylic sample plate. The ESI plume was directed toward the ion sampling orifice (i.e., parallel to the sample plate). The electrospray needle was held at +4 kV, with the sampling cone voltage in

Results and discussion

detected by this ELDI/MS. There are, however, several shortcomings when mixing carbon powders with sample solutions: the carbon powder might end up stuck to the MS inlet or inside the mass analyzer, potentially damaging the instrument; the analyte might adsorb permanently onto the surface of the carbon powder, thereby decreasing the detection sensitivity; and an appropriate amount of carbon powder must be measured and added to the sample solution, increasing the analysis time. Therefore, we wished to develop a technique for detecting proteins in various solutions without the need for added carbon powder. Unfortunately, in the absence of carbon powder, the sample solutions were transparent to the UV-laser, resulting in no desorption during laser irradiation [26, 27].

In this study, we examined the capability of ELDI/MS to directly characterize proteins stored in various solutions. However, when we deposited a small sample droplet onto a stainless-steel plate, spread it to form a thin film, and then irradiated it with a UV-laser, we found that the UV-laser energy was absorbed by the stainless-steel plate to produce numerous fine droplets of the sample solution (Fig. 1b). If the sample solution was not spread into a thin film on the plate, large droplets were produced during laser irradiation, and their ionization efficiency during ELDI process was extremely low.

a

Taylor Cone

ESI droplets

MS Inlet

Protein ions

Laser Beam

Sample droplets

Proteins in various solutions

b

It seems the formation of fine sample droplets is a prerequisite for subsequent post-ionization of the analytes in the ESI plume. In this study, the energy required to form analyte-containing droplets arose from the absorption of laser energy by the stainless-steel sample plate. Similar phenomena have been observed in fused-droplet ESI (FD-ESI), another two-step ESI method [31–34]. In FD-ESI, an ultrasonic nebulizer is used to disperse an aqueous sample solution into a fine mist of droplets; the resulting neutral aerosol is then fused with the charged droplets generated by electrospraying an acidic organic solvent.

Figure 2 presents positive ion mode ELDI mass spectra and respective deconvoluted mass spectra of cytochrome c dissolved in aqueous solutions with pH ranging from 0 to 14. To post-ionize the proteins in the droplets generated through laser desorption, we used an ESI solution containing MeOH and water (50 %, v/v) and acetic acid (0.1 %). We detected signals for cytochrome c ions in all of the solutions having pH values of less than 13. Typically, proteins are multi-amphiphilic species that can form net-positively or -negatively charged gaseous ions under a variety of ESI conditions; in this study, we manipulated the composition of the ESI solution for protein post-ionization in the acidic range to enable the formation of positively charged protein ions for analysis in the positive ion mode. Thus, the cytochrome c ($pI=10.0$) molecules desorbed from the buffer solutions (pH from 0 to 13) were fused with the acidic ESI plume (pH 2.7) to form positively charged ESI droplets; the proteins in the droplets were subsequently ionized through the ESI process to generate positively charged protein ions.

In Fig. 2c–g, the cytochrome c ion is the predominant ion present in the deconvoluted mass spectra. In addition, we

detected a protein/phosphate adduct ion from the phosphate buffer solution at pH 2.2 (Fig. 2b) and a protein/chloride adduct ion from the 1 M HCl solution at pH 0 (Fig. 2a). The presence of phosphate in the protein solution fortunately did not significantly suppress the signal of the cytochrome c ion; this situation differs from that of conventional ESI/MS, where the protein ion is heavily suppressed by phosphate in the sample solution. A possible explanation for this phenomenon is that phosphate and HCl were diluted during the fusion process (the desorbed sample droplets fused with charged ESI droplets) [22, 29]. This “dilution” of matrices in the sample solution during fusion is an important feature of two-step ionization. Many factors (e.g., salts, organic solvent) that have the potential to interfere with ion signals will be diluted when the fine sample droplets are fused with the electrospray droplets. The protein ions were readily detected in the solutions containing sodium acetate and citrate (Fig. 2c, d, respectively). Tris, another organic compound that interferes with the detection of proteins in conventional ESI/MS, also did not affect the detection of the protein when using ELDI/MS (Fig. 2e, f). The results obtained here will agree with those from our previous studies using FD-ESI/MS [31–34].

Figure 2a reveals that cytochrome c dissolved in strong HCl solution (pH 0) could be detected using this ELDI approach. Surprisingly, we even detected signals for cytochrome c ions from concentrated (6 and 12 M) HCl solutions (Fig. 3a, b, respectively). In addition to the predominant signals of the molecular ion, however, we also detected a signal for the chloride adduct of the protein ion in the deconvoluted mass spectrum as a result of the extremely high concentration of HCl in the system. Signals for some other protein-like ion




Fig. 2 Positive-ion ELDI mass spectra of cytochrome c dissolved in buffer solutions at various values of pH. *Insets*: Deconvoluted mass spectra of the cytochrome c ions. ESI solution: 50 % MeOH containing 0.1 % acetic acid




Fig. 3 Positive-ion ELDI mass spectra of cytochrome c dissolved in (a), (b) extremely high concentration of HCl, (c) high concentration of Tris buffer, (d)–(f) different buffer solutions at the same pH (pH 9.0). *Insets*:

Deconvoluted mass spectra of the cytochrome c ions. ESI solution: 50 % MeOH containing 0.1 % acetic acid

signals were also evident in the spectra of the 6 and 12 M HCl samples, presumably because some of the amide linkages of the proteins were cleaved in the strongly acidic solutions. In contrast, we detected no protein ions from the solution in 1 M NaOH (pH 14.0); instead, the mass spectrum featured predominant sodium cluster ions, namely $\text{Na}^+(\text{NaOH})_m(\text{H}_2\text{O})_n$ and $\text{Na}^+(\text{NaOAc})_m(\text{H}_2\text{O})_n$ species (Fig. 2h) [35, 36]. The failure to detect protein ions stored in the strongly basic solution suggested that the structure of the protein was destroyed or that the signal of the protein ion was heavily suppressed by basic species (in this case, NaOH).

To further examine the effect of pH on the protein ion signal, we compared the ELDI mass spectra of cytochrome c solutions prepared in different buffer solutions at the same pH (pH 9.0; Fig. 3d–f) with those obtained at different values of pH in the same buffer. Again we detected the signal for the cytochrome c ion in all of the ELDI mass spectra, regardless of the nature of the buffer (Fig. 3c–f). Since we observed the signal for the protein ion in all of the solutions at different values of pH (see Fig. 2), it seems that the pH has no obvious effect on the detection of cytochrome c ions when using this ELDI approach.

Although it appears that the pH and buffer have little effect on the detection of the protein ions in solutions, the ELDI mass spectra in Fig. 2 present cytochrome c ions in their denatured form. This phenomenon arose because the cytochrome c ions were formed through post-ionization in an ESI plume generated from an acidic MeOH/water solution. Our results suggest that regardless of the type of solution in which the protein was stored, contact with acetic acid and MeOH in the ESI plume during post-ionization changed the protein's conformation. It has been reported that the use of 20 % MeOH/water for ESI results in ELDI/MS signals of the native conformation of cytochrome c ion [28]. Figure 4 displays ELDI mass spectra and

respective deconvoluted mass spectra of myoglobin in buffer solutions at various values of pH when using an acid-free MeOH solution (20 % by volume) as the ESI solution. Native and denatured states, corresponding to the holo- and apo-forms, of Mb cannot only be discerned by their charge state distribution, but also by mass due to the loss of the heme group. Based on the calculated molecular weight and charge distribution, we could deduce the conformation of the myoglobin ion. For the sample solutions having values of pH between 5 and 7, the mass spectra featured predominant signals for native myoglobin ions (Fig. 4d, f, j, l), while denatured myoglobin ions were detected when the pH of the sample solution was less than 4.4. These conformations information are delightfully in accordance with those of previous studies [37]. This strongly suggests that ELDI/MS is a direct, rapid, useful, and efficient tool to study the conformations of proteins stored in solutions at various values of pH, which is not achieved by other ionization methods.

The ability to use ELDI/MS to assess the structures of proteins stored in solutions at various pH arises from several unique features of the techniques: (a) droplets containing protein molecules are generated from aqueous sample solutions under laser irradiation (Fig. 1b); (b) mixing the myoglobin-containing droplets with the ESI solvent droplets containing 20 % MeOH does not induce denaturation of myoglobin much; and (c) ESI proceeds from the droplets formed after fusing the myoglobin droplets with the ESI solvent droplets to generate myoglobin ions. These features make ELDI/MS a useful technique to study the integrity and conformations of proteins stored in various solutions.

Organic solvents are commonly used to extract and precipitate proteins from solutions during sample preparation. Organic solutions are also used as mobile phases for chromatographic separation, often in eluting gradients. To study the effects of




Fig. 4 (a)–(f) Positive-ion ELDI mass spectra of myoglobin in buffer solutions at various values of pH. (g)–(i) Deconvoluted mass spectra of the myoglobin ions. ESI solution: 20 % MeOH/water (v/v). Blue circles denatured protein; yellow triangles native protein

organic solvents on the signals of protein ions during ELDI/MS analyses, we prepared protein solutions by dissolving cytochrome c (10^{-4} M) in IPA, EtOH, MeOH, ACN, DMF, and DMSO, respectively. In these cases, the ESI solution was 50 % MeOH with 0.1 % acetic acid. After applying each sample solution to the stainless-steel plate, we subjected it to a pulsed UV laser and analyzed it using ELDI/MS equipped. Figure 5 displays the ELDI mass spectra of the samples; surprisingly, we detected signals for the protein ions from all of the solvents.

IPA is a water-miscible solvent that is commonly used in lipoprotein precipitation, peptide and protein extraction, and as a crucial reagent in Eastern blot analysis [38, 39]. Figure 5a presents the positive-ion ELDI mass spectrum of cytochrome c dissolved in IPA. Although the polarity index of IPA is only 3.9, signals for the cytochrome c ion were still obtained through ELDI/MS.

In addition to their use as reagents for protein precipitation, EtOH, MeOH, and ACN are commonly employed in mobile phases for solvent-gradient HPLC/MS. Baseline drift of the

total ion chromatogram upon increasing the ratio of organic solvent is, however, a common problem for HPLC/MS. Since we detected signals for the protein ions from all of these solvents (Fig. 5b–d), it appears that ELDI/MS might be an alternative useful technique when combined on-line with HPLC for protein separation and characterization. Comparing to conventional HPLC/ESI source, the ions in ELDI source were produced through post-ionized by consistent electrospray solution; it has potential to dissolve the baseline drift problem.

DMSO and DMF are common solvents for preparing stock solutions because of their high polarity (allowing the dissolution of a wide range of chemical compounds), low chemical reactivity, and low vapor pressure. They are also useful solvents for the ESI analysis of hydrophobic compounds, as well as being LC-compatible and capable of improving chromatographic resolution during the analysis of intact proteins [40–43]. Furthermore, the charge enhancement of protein ions by DMSO and DMF has been demonstrated [40–43]. Figure 5e, f displays ELDI mass spectra of cytochrome c dissolved in 100 % DMF and




Fig. 5 Positive-ion ELDI mass spectra of cytochrome c dissolved in various organic solvents. *Insets*: Deconvoluted mass spectra of the cytochrome c ions

DMSO, respectively. The highest charge state of cytochrome c shifted from 18+ or 19+ (in other organic solvents, Fig. 5a–c) to 20+ in DMSO and DMF (Fig. 5e, f).

Another attractive feature of ELDI for protein analysis is its high tolerance toward salts and detergents. In previous studies, we demonstrated the very high salt tolerance [especially for inorganic salts (e.g., NaCl, NaH₂PO₄)] of ELDI/MS and FD-ESI/MS when using acidic MeOH as the ESI solution for protein analysis [21–34]. Because the solubility of inorganic salts in acidic MeOH (ESI solution) is at least ten times lower than that in water, somehow the salt molecules or ions would be excluded from the charged MeOH droplets during ionization. This is one possible reason why the presence of inorganic salts in sample solutions did not affect the detection of proteins through ELDI/MS and FD-ESI/MS analysis. It is also possible that less-polar analyte molecules were positioned on the

surfaces of the neutral sample droplets, whereas the salts were stabilized through increased solvation within the droplets. The analyte molecules presented on the droplets' surfaces would then be easier to extract into the charged MeOH droplets during ionization. Therefore, in this study, we also examined the effects of the salts and detergents that are commonly used for protein storage on the ELDI/MS detection of proteins.

Here, we prepared solutions containing cytochrome c and various types of salts and detergents (urea, guanidine · HCl, glycerol, SDS, CHAPS, Triton X-100). Urea and guanidine · HCl are used widely as chaotropic agents for protein denaturing, solubilizing inclusion bodies, and eluting highly hydrophobic proteins [44, 45]. Protein solutions containing these detergents often require additional cleanup steps prior to analysis through conventional ESI/MS [14–18]. ELDI/MS analysis, however, exhibited extremely high tolerance. Figure 6




Fig. 6 Positive-ion ELDI mass spectra of cytochrome c in solutions spiked with various detergents and salts. *Insets*: Deconvoluted mass spectra of the cytochrome c ions. *Blue circles* signals for cytochrome c ions

display positive-ion ELDI mass spectra of cytochrome c in solutions spiked with various detergents and salts. Surprisingly, the protein signals remain unchanged even in the solution with 10 M urea (Fig. 6a), higher than other ionization methods, presumably because of the lack of solubility of urea in MeOH. In contrast, the removal of guanidine·HCl, which is soluble in both water and MeOH, through the ELDI process was not as favorable as that of urea. Nevertheless, we could still obtain reasonably good signal intensity for the cytochrome c ions from the solution containing 3 M guanidine·HCl when using ELDI/MS (Fig. 6b).

Figure 6c shows the ELDI mass spectrum of cytochrome c stored in glycerol. As generally known, glycerol is a widely used reagent to stabilize proteins in aqueous solution [46]. Nevertheless, the presence of high concentrations of glycerol in sample solutions can strongly affect the ESI/MS performance and decrease the intensities of the signals of analyte ions [47]. In this study, we obtained strong ELDI/MS signals for cytochrome c ions even when the samples were stored in 100 % glycerol. Again, we suspect that this phenomenon arose because the glycerol molecules were diluted during the fusion process.

The presence of detergent in protein solutions is another troublesome factor for conventional ESI/MS analysis. Integral membrane proteins are typically extracted from their native environments using detergent-containing solutions. In addition, the recoveries of certain types of proteins in aqueous solutions can be improved by solubilizing the protein extracts in detergent solutions. Three types of detergents (ionic, zwitterionic, nonionic) are commonly used for this purpose. SDS, an anionic detergent, is one of the most common additives used in bioanalytical assays. Figure 6d reveals that cytochrome c-SDS adducts were evident in the ELDI mass spectrum when the SDS concentration reached 10^{-3} M. Because SDS has high solubility in both water (sample solution) and MeOH (ESI solution), ELDI/MS is not particularly successful at removing its interference. Indeed, the deconvoluted mass spectrum barely revealed the presence of cytochrome c ions and protein-SDS adducts ions.

CHAPS is a zwitterionic detergent commonly used as a non-denaturing solvent during protein purification; it is particularly useful when purifying membrane proteins. CHAPS is also used in conjunction with nonionic detergents (e.g., Triton X-100) to solubilize membrane proteins in their native state. The mass spectrum in Fig. 6e reveals that the signal for cytochrome c ions interfered in the presence of 10 mM CHAPS, but cytochrome c signal is predominated in the deconvoluted mass spectrum, with the presence of minor protein-CHAPS adducts signal. Triton X-100 is a nonionic detergent that features a hydrophilic polyethylene oxide group and a lipophilic or hydrophobic hydrocarbon unit. Although the signals of the protein ions were disrupted by the presence of Triton X-100 at 0.5 % (Fig. 6f), unambiguous characterization of the cytochrome c ion was possible after deconvolution. Because most detergents are also

soluble in MeOH, the detergent tolerance of ELDI/MS was greater than that of ESI/MS presumably because of the diluting effect during the infusion of the fine sample droplets into charged-MeOH ESI droplets. Further studies will be necessary, however, if we are to better understand the mechanisms behind the removal of salt during ELDI/MS.

Conclusion

Positive ion mode ELDI/MS under ambient conditions can be used to detect proteins dissolved in solutions of various compositions. The three-dimensional conformations of the structures of the proteins in solution can be preserved during analysis if the composition of the electrospraying solution is chosen properly. In contrast to the often time-consuming approaches used for analyses in the field of protein biochemistry, ELDI/MS can be a rapid and practical tool for assessing the structures and stabilities of protein molecules in solution. Positively charged proteins ions can be detected directly through ELDI/MS without the need for any sample pretreatment, even when the proteins are dissolved in alkaline solutions, concentrated HCl solutions, or less-polar, moderately polar, or highly polar organic solvents. Furthermore, the salt and detergent tolerance of ELDI/MS is higher than that of conventional ESI/MS, thereby making direct analysis possible. In conclusion, ELDI/MS has great potential for use in the rapid screening of proteins—for their existence and integrity—in various environments. This feature can be used to follow the progress of biochemical procedures and protein purification steps, and to assess their effectiveness. Semi-quantitative information and detection limits of the ELDI/MS for detecting protein in various solutions would be studied discussed in future studies.

Acknowledgments This study was supported financially through a grant (NSC 99-2923-M-110-007-MY3) from the National Science Council, Taiwan.

References

1. Rosenberg IM (2005) Membrane protein. In Protein Analysis and Purification: Benchtop Techniques. 2nd edn. Springer, New York
2. Pieper R, Huang ST, Clark DJ, Robinson JM, Alami H, Parmar PP, Suh MJ, Kuntumalla S, Bunai CL, Perry RD, Fleischmann RD, Peterson SN (2009) Integral and peripheral association of proteins and protein complexes with *Yersinia pestis* inner and outer membranes. *Proc Natl Acad Sci U S A* 7:5
3. Singh SM, Panda AK (2005) Solubilization and refolding of bacterial inclusion body proteins. *J Biosci Bioeng* 99(4):303–310
4. Hultin et al. 2000. High efficiency alkaline protein extraction. United States Patent. 6,136,959
5. Zhang S, Bian Z, Gu C, Zhang Y, He S, Gu N, Zhang J (2007) Preparation of anti-human cardiac troponin I immunomagnetic nanoparticles and biological activity assays. *Colloids Surf B Biointerfaces* 55(2):143–148

6. Fenn JB, Mann M, Meng CK, Wong SF, Whitehouse CM (1989) Electrospray ionization for mass spectrometry of large biomolecules. *Science* 246(4926):64–71
7. Hutton T, Major HJ (1995) Characterizing biomolecules by electrospray ionization-mass spectrometry coupled to liquid chromatography and capillary electrophoresis. *Biochem Soc Trans* 23(4): 924–927
8. Loo JA, Quinn JP, Ryu SI, Henry KD, Senko MW, McLafferty FW (1992) High-resolution tandem mass spectrometry of large biomolecules. *Proc Natl Acad Sci U S A* 89(1):286–289
9. Banks JF Jr, Quinn JP, Whitehouse CM (1994) LC/ESI-MS determination of proteins using conventional liquid chromatography and ultrasonically assisted electrospray. *Anal Chem* 66(21):3688–3695
10. Backstrom M, Thomsson KA, Karlsson H, Hansson GC (2009) Sensitive liquid chromatography-electrospray mass spectrometry allows for the analysis of the O-glycosylation of immunoprecipitated proteins from cells or tissues: application to MUC1 glycosylation in cancer. *J Proteome Res* 8(2):538–545
11. Stone KL, DeAngelis R, LoPresti M, Jones J, Papov VV, Williams KR (1998) Use of liquid chromatography-electrospray ionization-tandem mass spectrometry (LC-ESI-MS/MS) for routine identification of enzymatically digested proteins separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *Electrophoresis* 19(6):1046–1052
12. Pan P, McLuckey SA (2003) Electrospray ionization of protein mixtures at low pH. *Anal Chem* 75(6):1491–1499
13. Banerjee S, Mazumdar S (2012) Electrospray ionization mass spectrometry: a technique to access the information beyond the molecular weight of the analyte. *Int J Anal Chem* 2012:282574
14. Qi J, Isupov MN, Littlechild JA, Anderson LE (2001) Chloroplast glyceraldehyde-3-phosphate dehydrogenase contains a single disulfide bond located in the C-terminal extension to the B subunit. *J Biol Chem* 276(38):35247–35252
15. Bagshaw RD, Callahan JW, Mahuran DJ (2000) Desalting of in-gel digested protein sample with mini-C18 columns for matrix-assisted laser desorption ionization time of flight peptide mass fingerprinting. *Anal Biochem* 284(2):432–435
16. Ekstrom S, Malmstrom J, Wallman L, Lofgren M, Nilsson J, Laurell T, Marko-Varga G (2002) On-chip microextraction for proteomic sample preparation of in-gel digests. *Proteomics* 2(4):413–421
17. Xu Y, Bruening ML, Watson JT (2003) Non-specific, on-probe cleanup methods for MALDI-MS samples. *Mass Spectrom Rev* 22(6):429–4404
18. Tannu NS, Wu J, Rao VK, Gadgil HS, Pabst MJ, Gerling IC, Raghov R (2004) Paraffin-wax-coated plates as matrix-assisted laser desorption/ionization sample support for high-throughput identification of proteins by peptide mass fingerprinting. *Anal Biochem* 327(2):222–232
19. Annesley TM (2003) Ion suppression in mass spectrometry. *Clin Chem* 49(7):1041–1044
20. Hirabayashi A, Ishimaru M, Manri N, Yokosuka T, Hanzawa H (2007) Detection of potential ion suppression for peptide analysis in nanoflow liquid chromatography/mass spectrometry. *Rapid Commun Mass Spectrom* 21(17):2860–2866
21. Shiea J, Huang MZ, Hsu HJ, Lee CY, Yuan CH, Beech I, Sunner J (2005) Electrospray-assisted laser desorption/ionization mass spectrometry for direct ambient analysis of solids. *Rapid Commun Mass Spectrom* 19(24):3701–3704
22. Huang MZ, Hsu HJ, Lee JY, Jeng J, Shiea J (2006) Direct protein detection from biological media through electrospray-assisted laser desorption ionization/mass spectrometry. *J Proteome Res* 5(5):1107–1116
23. Huang MZ, Hsu HJ, Wu CI, Lin SY, Ma YL, Cheng TL, Shiea J (2007) Characterization of the chemical components on the surface of different solids with electrospray-assisted laser desorption ionization mass spectrometry. *Rapid Commun Mass Spectrom* 21(11):1767–1775
24. Peng IX, Shiea J, Ogorzalek Loo RR, Loo JA (2007) Electrospray-assisted laser desorption/ionization and tandem mass spectrometry of peptides and proteins. *Rapid Commun Mass Spectrom* 21(16):2541–2546
25. Lin SY, Huang MZ, Chang HC, Shiea J (2007) Using electrospray-assisted laser desorption/ionization mass spectrometry to characterize organic compounds separated on thin-layer chromatography plates. *Anal Chem* 79(22):8789–8795
26. Shiea J, Yuan CH, Huang MZ, Cheng SC, Ma YL, Tseng WL, Chang HC, Hung WC (2008) Detection of native protein ions in aqueous solution under ambient conditions by electrospray laser desorption/ionization mass spectrometry. *Anal Chem* 80(13):4845–4852
27. Cheng CY, Yuan CH, Cheng SC, Huang MZ, Chang HC, Cheng TL, Yeh CS, Shiea J (2008) Electrospray-assisted laser desorption/ionization mass spectrometry for continuously monitoring the states of ongoing chemical reactions in organic or aqueous solution under ambient conditions. *Anal Chem* 80(20):7699–7705
28. Peng IX, Ogorzalek Loo RR, Shiea J, Loo JA (2008) Reactive-electrospray-assisted laser desorption/ionization for characterization of peptides and proteins. *Anal Chem* 80(18):6995–7003
29. Huang MZ, Jhang SS, Cheng CN, Cheng SC, Shiea J (2010) Effects of matrix, electrospray solution, and laser light on the desorption and ionization mechanisms in electrospray-assisted laser desorption ionization mass spectrometry. *Analyst* 135(4):759–766
30. Cheng SC, Lin YS, Huang MZ, Shiea J (2010) Applications of electrospray laser desorption ionization mass spectrometry for document examination. *Rapid Commun Mass Spectrom* 24(2):203–208
31. Lee CC, Chang DY, Jeng J, Shiea J (2002) Generating multiply charged protein ions via two-step electrospray ionization mass spectrometry. *J Mass Spectrom* 37(1):115–117
32. Chang DY, Lee CC, Shiea J (2002) Detecting large biomolecules from high-salt solutions by fused-droplet electrospray ionization mass spectrometry. *Anal Chem* 74(11):2465–2469
33. Shieh IF, Lee CY, Shiea J (2005) Eliminating the interferences from TRIS buffer and SDS in protein analysis by fused-droplet electrospray ionization mass spectrometry. *J Proteome Res* 4(2): 606–612
34. Shiea J, Sunner J (2007) In: Gross ML, Caprioli RM (eds) *The Encyclopedia of Mass Spectrometry*. Elsevier, Amsterdam
35. Alimi R, Gerber RB (1990) Solvation effects on chemical reaction dynamics in clusters: Photodissociation of HI in XeNHI. *Phys Rev Lett* 64(12):1453–1456
36. Steinbach C, Buck U (2005) Reaction and solvation of sodium in hydrogen bonded solvent clusters. *Phys Chem Chem Phys* 7(5):986–990
37. Postnikova GB, Komarov YE, Yumakova EM (1991) Fluorescence study of the conformational properties of myoglobin structure. 2. pH- and ligand-induced conformational changes in ferric- and ferrousmyoglobins. *Eur J Biochem* 198(1):233–239
38. Marney LC, Laha TJ, Baird GS, Rainey PM, Hoofnagle AN (2008) Isopropanol protein precipitation for the analysis of plasma free metanephrines by liquid chromatography-tandem mass spectrometry. *Clin Chem* 54(10):1729–1732
39. Sen M, Bhattacharyya DK (2000) Nutritional quality of sunflower seed protein fraction extracted with isopropanol. *Plant Foods Hum Nutr* 55(3):265–278
40. Szabo PT, Kele Z (2001) Electrospray mass spectrometry of hydrophobic compounds using dimethyl sulfoxide and dimethylformamide as solvents. *Rapid Commun Mass Spectrom* 15(24):2415–2419
41. Valeja SG, Tipton JD, Emmett MR, Marshall AG (2010) New reagents for enhanced liquid chromatographic separation and charging of intact protein ions for electrospray ionization mass spectrometry. *Anal Chem* 82(17):7515–7519
42. Tjernberg A, Markova N, Griffiths WJ, Hallen D (2006) DMSO-related effects in protein characterization. *J Biomol Screen* 11(2): 131–137

43. Borges V, Henion J (2005) Determination of pharmaceutical compounds in aqueous dimethyl sulfoxide by electrospray ionization mass spectrometry. *Rapid Commun Mass Spectrom* 19(3):415–423
44. Del Vecchio P, Graziano G, Granata V, Barone G, Mandrich L, Rossi M, Manco G (2002) Denaturing action of urea and guanidine hydrochloride towards two thermophilic esterases. *Biochem J* 367(Pt 3): 857–863
45. Monera OD, Kay CM, Hodges RS (1994) Protein denaturation with guanidine hydrochloride or urea provides a different estimate of stability depending on the contributions of electrostatic interactions. *Protein Sci* 3(11):1984–1991
46. Sedgwick H, Cameron JE, Poon WC, Egelhaaf SU (2007) Protein phase behavior and crystallization: effect of glycerol. *J Chem Phys* 127(12):125102
47. Mendes MA, Chies JM, de Oliveira Dias AC, Filho SA, Palma MS (2003) The shielding effect of glycerol against protein ionization in electrospray mass spectrometry. *Rapid Commun Mass Spectrom* 17(7):672–677