

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/243954314>

AnAB initio study of the geometrical structures and vibrational spectra of the free ion NO_2^- and the LiNO_2 molecule

ARTICLE in JOURNAL OF STRUCTURAL CHEMISTRY · NOVEMBER 1985

Impact Factor: 0.51 · DOI: 10.1007/BF00747822

READS

4

2 AUTHORS, INCLUDING:



Victor Solomonik

Ivanovo State University of Chemistry and ...

77 PUBLICATIONS 217 CITATIONS

SEE PROFILE

AN *AB INITIO* STUDY OF THE GEOMETRICAL STRUCTURES AND VIBRATIONAL SPECTRA OF THE FREE ION NO_2^- AND THE LiNO_2 MOLECULE


S. P. Konovalov and V. G. Solomonik

UDC 539.194

The Hartree-Fock-Roothaan method has been used in a two-exponential basis with and without the inclusion of polarization defunctions on the nitrogen and oxygen atoms to calculate the geometry and the vibrational characteristics of the free NO_2^- ion and the LiNO_2 molecule. The results are compared with published data as regards the nitrite ion and the alkali-metal nitrites. The parameters of the LiO_2N rings in LiNO_3 and LiNO_2 are compared. A study has been made of the effects of the polarization defunctions on the calculated parameters of NO_2^- and LiNO_2 . The rigid rotor-harmonic oscillator approximation has been used to calculate the thermodynamic parameters of LiNO_2 in the ideal-gas state.


So far there have been no experimental studies on the geometry and vibrational spectrum of the LiNO_2 molecule. The spectroscopic data [1, 2] and electron diffraction [3, 4] indicate cyclic structures for the molecules MNO_2 ($\text{M} = \text{Na}, \text{K}, \text{Rb}, \text{Cs}$). Nevertheless, no final conclusion has been drawn [1-4] on the symmetry of the equilibrium configurations in these molecules.

In accordance with the data of [1-4], nonempirical calculations on MXO_2 ($\text{M} = \text{Li}, \text{Na}; \text{X} = \text{N}, \text{P}$) [5] have shown that the least energy occurs in a cyclic structure of the form



Also, according to [5], the potential surface for a molecule of this type has

three further local minima corresponding to the structures $\text{Li}-\text{O}-\text{N}=\text{O}$, $\text{Li}=\text{O}-\text{N}=\text{O}$, and



There have been no theoretical studies on the force fields and vibrational

spectra of the alkali metal nitrites.

In previous *ab initio* studies on the free NO_2^- ion (see for example [6] and references in [7]), only the electronic and geometrical structures were considered.

Here we examine the geometrical structure and vibrational spectra of the LiNO_2 molecule and the free NO_2^- ion by the SCF MO LCAO method by the use of fairly broad and flexible bases.

CALCULATION DETAILS

All the calculations were performed with the MOLECULE program [8] at the computing center at Ivanovo Chemical Technology Institute. We used the following sets of grouped gaussian functions: $\text{DZ} - (9s5p/4s2p)_{\text{N,O}}$, $(9s4p/4s2p)_{\text{Li}}$ [9, 10] and $\text{DZ} + \text{P}$ being the same basis supplemented with polarization functions of d type on the nitrogen and oxygen atoms with the exponents 0.80 (N) and 0.85 (O) [11].


The geometrical parameters of the cyclic structure  (C_{2v}) and of the NO_2^- ion

TABLE 1. Characteristics of the Free NO₂ Ion*

Parameter	<i>Ab initio</i> calc.		Published data
	DZ	DZ+P	
$-E$	204,02512	204,12419	
$R_e(\text{N—O})$	1,273	1,235	(1,23±0,05) [7]
$\alpha_e(\text{O—N—O})$	116,7	116,6	(118±5) [7]
$Q(\text{N})$	+0,420	+0,116	
$Q(\text{O})$	-0,710	-0,580	
$q(\text{N—O})$	0,338	0,589	
f_d	8,547	11,587	7,50 [17]
f_{dd}	2,033	3,013	1,66 [17]
f_α	1,614	1,952	1,67 [17]
$f_{d\alpha}$	0,359	0,467	0,394 [17]
$\nu_1(A_1)$	1394	1623	(1330±20) [7]
$\nu_2(A_1)$	800	888	(810±20) [7]
$\nu_3(B_2)$	1354	1554	(1245±40) [7]
$(\partial\mu/\partial Q_1)^2$	1,00	0,90	
$(\partial\mu/\partial Q_2)^2$	0,45	0,29	
$(\partial\mu/\partial Q_3)^2$	17,13	20,65	
μ^\dagger	0,803	0,626	
IP	4,10	3,58	2,36±0,10 ‡ [7]

*Here and in Tables 2 and 5, we use the following symbols and dimensions: E total energy (au), R , internuclear distance (Å), α bond angle (deg), Q charge on an atom (au), q populations in the bonds (au), f force constant (mdyn/Å), ν frequency (cm⁻¹),

$(\partial\mu/\partial Q_i)^2$, square of the derivative of the dipole

moment coordinate Q_i (m² Å⁻² a.m.u.⁻¹), μ dipole moment (D); IP ionization potential (eV), and l mean amplitude (Å).

† The center of mass coincides with the center of the coordinates.

‡ Electron affinity of the NO₂ Molecule.


TABLE 2. Characteristics of the LiNO₂ Molecule Calculated *Ab initio*

Parameter	DZ	DZ+P	Parameter	DZ	DZ+P
$-E$	211,52362	211,63045	$Q(\text{N})$	+0,350	+0,347
$R_e(\text{N—O})$	1,277	1,238	$Q(\text{O})$	-0,569	-0,552
$R_e(\text{Li—O})$	1,934	1,886	$q(\text{N—O})$	0,315	0,544
$R_e(\text{Li ... N})$	2,341	2,255	$q(\text{Li—O})$	0,184	0,168
$\alpha_e(\text{O—N—O})$	111,4	113,6	μ	6,42	6,13
$Q(\text{Li})$	+0,788	+0,757	IP	11,5	11,0

were optimized not only in the DZ basis but also DZ+P. Also, in DZ we optimized the

$\text{Li—O}_b\text{—N} \begin{smallmatrix} \nearrow \text{O}_t \\ \searrow \end{smallmatrix} (C_s)$ structure with the linear Li—O_b—N, fragment, which according to [5] is the isomer having the energy closest to that of the cyclic structure. We obtained the following internuclear distances for the C_s configuration: $R(\text{N—O}_t)$ 1,232, $R(\text{N—O}_b)$ 1,306, $R(\text{Li—O}_b)$ 1,636 Å and the bond angle $\alpha(\text{O}_t\text{—N—O}_b)$ 115,4°. Table 1 and 2 give the calculated characteristics for NO₂⁻ and LiNO₂.

TABLE 3. Force Constants of the LiNO_2 Molecule (mdyn/Å)



Force constant	DZ	DZ+P	Force constant	DZ	DZ+P
f_r	0,701	0,833	f_{dr}	0,260	0,339
f_{rr}	0,194	0,072	f_α	2,113	2,350
f_d	8,411	11,027	$f_{r\alpha}$	-0,214	-0,165
f_{dd}	2,539	3,033	$f_{d\alpha}$	0,691	0,783
$f_{dr'}$	-0,234	-0,309	f_ω	0,214	0,186

TABLE 4. Vibration Frequencies ν_i (cm^{-1}), Values of $(\partial\mu/\partial Q_i)^2$ ($D^2\text{Å}^{-2} \text{ a.m.u.}^{-1}$), Potential-Energy Distribution PED Over the Natural Coordinates, and Isotopic Shifts $\Delta\nu_i$ ($^6\text{Li}-^7\text{Li}$) (cm^{-1}) for the LiNO_2 Molecule

Type	DZ		DZ+P		PED * (DZ)	$\Delta\nu_i(^6\text{Li}-^7\text{Li})$	
	ν_i	$(\partial\mu/\partial Q_i)^2$	ν_i	$(\partial\mu/\partial Q_i)^2$		DZ	DZ+P
A_1	ν_1 1423	0,86	1590	0,74	$0,80d + 0,20\alpha$	0,2	0,4
	ν_2 882	0,51	959	0,36	$0,85\alpha$	0,2	0,5
	ν_3 589	2,79	589	2,90	$0,98r$	40,9	40,8
B_2	ν_4 1266	17,49	1490	21,20	$1,00d$	0,5	0,5
	ν_5 340	0,32	422	0,38	$0,91r$	15,8	19,8
B_1	ν_6 301	1,77	279	1,44	$1,00\omega$	4,3	3,7

*Contributions to PED less than 0.15 have been omitted

[14]* for LiNO_2 . Then a similarity transform was applied to matrix F to find the force constants in the natural coordinates (Table 3).

DISCUSSION

NO_2^- Ion. We found changes in the theoretical values of the geometrical parameters and characteristics for the nitrite ion on including the polarization d functions in the DZ basis analogous to those obtained previously for the BO_2^- , NO_3^- and CO_3^{2-} [16]. The shortening of the $R_e(\text{N}-\text{O})$ internuclear distance is accompanied by reductions in the magnitudes of the charges on the atoms and by increases in the bond populations. The force constants f_d , f_{dd} , f_α , and $f_{d\alpha}$, increase by 36, 48, 20, and 30%, while the frequencies ν_1 , ν_2 , and ν_3 increase by 16, 11, and 15% correspondingly. The calculations with the DZ basis on the whole reproduce the experimental data better than do those with DZ + P. While the force constants obtained with the DZ basis differ from the experimental values [17] by on average 12%, the differences for DZ + P is 42%. The deviations in the frequencies of the normal vibrations are 4.5% (DZ) and 19% (DZ + P).

The isotopic shifts calculated with the DZ basis for the vibrational frequencies of NO_2^- under the substitutions $^{14}\text{N} \rightarrow ^{15}\text{N}$, $^{16}\text{O} \rightarrow ^{18}\text{O}$ also agree well with the experimental values [17] (mean deviation 7%, maximum 11.7%).

LiNO_2 Molecule. The calculations showed that the C_{2v} configuration for the lithium nitrite molecule lies lower in energy than does C_s . The difference $E(C_s) - E(C_{2v})$ is 27.6 kJ in the DZ basis or 62.3 kJ[†] in the DZ + P basis. Therefore, a difference from LiBO_2 [18] is that adding the polarization functions to the basic sets of atoms in the acid residue increases the molecular rigidity as regards the motion of the lithium around the NO_2 fragment in the plane.

*In the case of LiNO_2 , instead of the $S(B_1)$ coordinate adopted in [4] we used $S(B_1) = \sqrt{RD/2}(\omega_1 - \omega_2)$ [15].

[†]Difference in energy between structures whose geometrical parameters were found in the DZ basis.

A check with the DZ basis showed that the C_s structure is not an isomer and does not correspond to any singular point on the potential surface. Movement of the lithium atom that

causes the $\text{Li}-\text{O}_b-\text{N}-\text{O}_t$ to form a ring reduces the energy, while motion in the opposite direction increases it. This result casts doubt on the conclusions of [5] on the existence of two other isomers for the MXO_2 molecule.

We now consider the changes in the LiNO_2 molecule on extending the basis by including polarization d functions. In the case of the NO_2 section, the changes in the geometrical parameters and electron-density distribution are the same as in the NO_2^- ion. The $\text{Li}-\text{O}$ bond resembles $\text{N}-\text{O}$ in shortening when there is the corresponding increase in the force constant f_r . The changes in the nondiagonal force constants means that there is a decrease in the difference between the frequencies of the symmetrical and antisymmetric stretching vibrations of the NO_2 and LiO_2 fragments. The diagonal force constants and vibrational frequencies increase when d functions are included in the basis: ν_1 by 12%, ν_2 by 8%, and ν_3 by 24%. An exception is represented by f_ω and the corresponding frequency ν_6 (B_1), which corresponds to nonplanar ring bending (ν_6 decreases by 8%), while ν_3 hardly alters.

The calculations indicate that the $R_e(\text{N}-\text{O})$ internuclear distances in LiNO_2 and NO_2^- are similar, although there is an altered electron-density distribution in LiNO_2 and a reduction in f_d . The other force constants $f_\alpha, f_{dd}, f_{d\alpha}$ increase. There are certain differences in the changes in $\nu_{\text{sym}}(\text{N}-\text{O})$ on going from NO_2^- to LiNO_2 in the DZ and DZ + P bases: While the value increases in DZ (mainly due to increase in f_{dd}), it decreases in DZ + P.

Table 4 gives the frequencies of the normal vibrations ν_i and the values of $(\partial\mu/\partial Q_i)^2$, which are proportional to the intensities of the IR bands. No experimental value have been published. For other MNO_2 ($\text{M} = \text{Na}, \text{K}, \text{Rb}, \text{Cs}$), measurements have been made only of the wave numbers of the strongest band ν_4 [1, 2], which is close to $\nu_3 = 1245 \pm 40 \text{ cm}^{-1}$ [7] for the nitrite ion, and also $\nu_2 = 802$ and 808 cm^{-1} for KNO_2 [1] (doubling due to band splitting in the matrix), which are close to $\nu_2 = 810 \pm 20 \text{ cm}^{-1}$ [7] in NO_2^- . The constancy of ν_4 in the sequence MNO_2 ($\text{M} = \text{Na}, \text{K}, \text{Rb}, \text{Cs}$) and the similarity of ν_4 and ν_2 to the corresponding frequencies for NO_2^- would indicate that the vibrations of the NO_2 section in the MNO_2 molecules are highly characteristic, including vibrations of totally symmetric type $\nu_{\text{sym}}(\text{N}-\text{O})$. Therefore, a comparison with the measured frequencies for the free NO_2^- ion [7] enables one to estimate the errors in calculating ν_1, ν_2 , and ν_4 for LiNO_2 ; the errors evidently do not exceed 10% in the DZ basis or 20% in DZ + P. The error in calculating $\nu_3(\text{Li}-\text{O})$ in the DZ basis is probably of the same order as for $\nu(\text{Li}-\text{O})$ in LiNO_3 ($\delta = 7.4\%$) [19]. On the whole, as for NO_2^- , the addition of the d functions to the DZ basis leads to an increase in the relative error in determining the frequencies for LiNO_2 . A similar

conclusion can probably be drawn for the frequencies of the fragment $\text{Li}-\text{O}-\text{N}$ in LiNO_3 .

The calculated isotopic shifts for NO_2 due to the substitutions $^{14}\text{N} \rightarrow ^{15}\text{N}$ and $^{16}\text{O} \rightarrow ^{18}\text{O}$ are similar to those for the nitrite ion [17].

We now compare the structure and parameters of MNO_2 with those of the fragment MO_2N in the MNO_3 molecule. Vapor electron diffraction [3, 20, 21] has shown that the effective internuclear distances and the mean vibration amplitudes of the RbNO_2 and CsNO_2 molecules

and of the $\text{M}-\text{O}-\text{N}$ ring in RbNO_3 or CsNO_3 are the same within the limits of the ex-

perimental error. In the interpretation of the internuclear distances for MNO_3 , it was assumed [20, 21] that the NO_3 fragment has D_{3h} symmetry. The internuclear distances $R(\text{N}-\text{O})$ and $R(\text{O} \dots \text{O})$ derived in [20, 21] correspond therefore to certain effective values $\bar{R}(\text{N}-\text{O}) = [R(\text{N}-\text{O}_t) + 2R(\text{N}-\text{O}_b)]/3$ and $\bar{R}(\text{O} \dots \text{O}) = [R(\text{O}_b \dots \text{O}_b) + 2R(\text{O}_b \dots \text{O}_t)]/3$. The quantities $\bar{l}(\text{N}-\text{O})$ and $\bar{l}(\text{O} \dots \text{O})$ were determined analogously.

It is evident from Table 5 that $R_e(\text{N}-\text{O}), l(\text{N}-\text{O})$ in LiNO_2 and $\bar{R}(\text{N}-\text{O}), \bar{l}(\text{N}-\text{O})$ in LiNO_3 are virtually the same. The characteristicity of the $R(\text{M}-\text{O})$ bond lengths for the heavy atoms and the mean amplitudes $\bar{l}(\text{M}-\text{O})$ in the ring [3, 20, 21] apply also for $\text{Li}-\text{O}$. The force

TABLE 5. Comparison of the Parameters of the $\text{Li} \begin{array}{c} \diagup \text{O} \\ \diagdown \text{O} \end{array} \text{N}$ Ring in LiNO_2 and LiNO_3 Calculated in the DZ Basis

Parameter	LiNO_2	LiNO_3 [19]	Parameter	LiNO_2	LiNO_3 [19]
$R(\text{N}-\text{O})$	1,277	1,313	$l(\text{Li} \cdots \text{N})$	0,102	0,102
$\bar{R}(\text{N}-\text{O})$		1,278	$l(\text{O} \cdots \text{O})$	0,056	0,065
$R(\text{Li}-\text{O})$	1,934	1,926	$\bar{l}(\text{O} \cdots \text{O})$		0,063
$R(\text{Li} \cdots \text{N})$	2,341	2,305	f_r	0,701	0,744
$\alpha(\text{O}-\text{N}-\text{O})$	111,4	113,3	f_{rr}	0,194	0,205
$\nu_{\text{sym}}(\text{Li}-\text{O})$	589	567	f_d	8,411	6,084
$\nu_{\text{sym}}(\text{N}-\text{O})$	1423	1013	f_{dd}	2,539	1,230
$\nu_{\text{asym}}(\text{Li}-\text{O})$	340	351	$f_{dr'}$	-0,234	-0,074
$\nu_{\text{asym}}(\text{N}-\text{O})$	1266	1240	f_{dr}	0,260	0,099
ν_{op}	301	185	f_α	2,113	1,801
$l(\text{Li}-\text{O})^*$	0,139	0,133	$f_{r\alpha}$	-0,214	-0,212
$l(\text{N}-\text{O})$	0,047	0,051	$f_{d\alpha}$	0,691	0,298
$\bar{l}(\text{N}-\text{O})$		0,047	f_ω	0,214	0,245

*Values given for $T = 800^\circ\text{K}$

TABLE 6. Thermodynamic Functions for LiNO_2 Ideal Gas in J/mole $\cdot^\circ\text{K}$

T, K	100	298,15	500	1000	1500	2000	2500	3000
c_p	37,01	54,64	65,77	76,99	80,19	81,43	82,03	82,37
Φ^*	177,00	217,80	241,53	280,00	306,04	325,86	341,85	355,28
S^0	211,05	260,17	291,29	341,20	373,13	396,40	414,64	429,63

constants show that NO_2 in LiNO_2 is more rigid than in the $\text{Li} \begin{array}{c} \diagup \text{O} \\ \diagdown \text{O} \end{array} \text{N}$ ring in LiNO_3 : the

values of $f_d, f_{dd}, f_\alpha, f_{d\alpha}$ in the nitrite are larger than those in the nitrate. On the other hand, f_r and f_{rr} are larger in LiNO_3 . Nevertheless, in spite of certain differences, there are analogous force fields and similarity between certain frequencies for lithium nitrite and nitrate. For example, $\nu_{\text{sym}}(\text{Li}-\text{O})$ and $\nu_{\text{sym}}(\text{Li}-\text{O})$ for LiNO_3 and LiNO_2 , which are related to motion of the lithium relative to the acid residue in the plane of the molecule.

We used the molecular parameters of LiNO_2 obtained in the DZ basis (Tables 2 and 4) for the rigid rotor-harmonic oscillator approximation to calculate the thermodynamic functions of LiNO_2 in the ideal-gas state (Table 6), which are required to calculate high-temperature processes involving lithium nitrite vapor.

The *ab initio* calculations give the energy for the dissociation of LiNO_2 into Li^+ and NO_2^- as 690 kJ/mole* in the DZ basis or 709 kJ/mole in the DZ + P. The energies for dissociation into Li plus NO_2 were calculated from data on the ionization potential of the lithium atom ($\text{IP}(\text{Li}) = 5.39178 \pm 0.00001 \text{ eV}$ [22]) and the electron affinity of NO_2 ($\text{EA}(\text{NO}_2) = 2.36 \pm 0.10 \text{ eV}$ [7]) as 397 and 417 kJ/mole in the DZ and DZ + P bases correspondingly.

Therefore, we have derived the complete force field and vibrational spectrum for the lithium nitrite molecule. The calculations show close analogies in structure and properties

between LiNO_2 and the $\text{Li} \begin{array}{c} \diagup \text{O} \\ \diagdown \text{O} \end{array} \text{N}$ ring in LiNO_3 as previously observed for compounds of the

heavy alkali metals [3, 20, 21]. The data will be useful in future theoretical and experimental studies on LiNO_2 and similar molecules.

* $E(\text{Li}^+) = -7.23599 \text{ au}$.

LITERATURE CITED

1. A. A. Belyaeva, V. N. Bukhmarina, Yu. B. Predtechenskii, and L. D. Shcherba, paper deposited at ONIITEKhim No. 132khp-D82 of 13 January 1982.
2. A. M. Shapovalov, PhD Thesis, Moscow University (1980).
3. V. A. Kulikov, V. V. Ugarov, and N. G. Rambidi, Zh. Strukt. Khim., 22, 183 (1981).
4. N. I. Tuseev, E. Z. Zasorin, and V. P. Spiridonov, Zh. Strukt. Khim., 20, 587 (1979).
5. C. J. Marsden, in: Ninth Austin Symposium on Molecular Structure(1982) p. 33.
6. G. V. Pfeiffer and L. C. Allen, J. Chem. Phys., 51 190 (1969).
7. Thermodynamic Parameters of Individual Substances: Handbook [in Russian], Volume 1, Part 1, Nauka, Moscow (1982), p. 295.
8. J. Almlöf, USIP Report 74-29, University of Stockholm (1974).
9. S. Husinaga, J. Chem. Phys., 42, 1292 (1965).
10. T. H. Dunning, J. Chem. Phys., 53, 2823 (1970).
11. T. H. Dunning and P. J. Hay, in: Methods of Electronic Structure Theory, Plenum Press, New York (1977), Chapter I, p. 1.
12. V. G. Solomonik, V. M. Ozerova, and V. V. Sliznev, Zh. Neorg. Khim., 27, 1636 (1982).
13. S. J. Cyvin, Molecular Vibrations and Mean-Square Amplitudes, Elsevier (1968).
14. S. J. Cyvin, B. N. Cyvin, and A. Snelson, J. Chem. Phys., 74, 4338 (1970).
15. A. Snelson, B. N. Cyvin, and S. J. Cyvin, J. Mol. Struct., 24, 165 (1975).
16. S. P. Konovalov and V. G. Solomonik, Zh. Fiz. Khim., 57, 363 (1983).
17. R. Kato and J. Rolfe, J. Chem. Phys., 47, 1901 (1967).
18. S. P. Konovalov and V. G. Solomonik, Zh. Neorg. Khim., 29, No. 7 (1984).
19. V. G. Solomonik and S. P. Konovalov, Zh. Strukt. Khim., 24, No. 5, 10 (1983).
20. V. A. Kulikov, V. V. Ugarov, and N. G. Rambidi, Zh. Strukt. Khim., 22, 196 (1981).
21. V. A. Kulikov, V. V. Ugarov, and N. G. Rambidi, Zh. Strukt. Khim., 22, 168 (1981).
22. V. N. Kondrat'ev (editor), Bond Energies, Ionization Potentials, and Electron Affinities: Handbook [in Russian], Nauka, Moscow (1974).