

See discussions, stats, and author profiles for this publication at:
<https://www.researchgate.net/publication/225336492>

Lichens as a potential natural source of bioactive compounds: a review.

Phytochem Rev

ARTICLE *in* PHYTOCHEMISTRY REVIEWS · JUNE 2010

Impact Factor: 2.41 · DOI: 10.1007/s11101-010-9189-6

CITATIONS

52

READS

278

3 AUTHORS, INCLUDING:



Vertika Shukla

National Botanical Research Institute - ...

32 PUBLICATIONS 218 CITATIONS

SEE PROFILE

Lichens as a potential natural source of bioactive compounds: a review

Vertika Shukla · Geeta Pant Joshi ·
M. S. M. Rawat



Received: 23 March 2010 / Accepted: 5 July 2010 / Published online: 17 July 2010
© Springer Science+Business Media B.V. 2010

Abstract Biological activity of material whether known in folk medicine or observed in planned screening program has been the starting point in the drug research. The general pattern is the isolation of active principles, elucidation their structures, followed by attempts for modulation of its activity potential by chemical modification. Lichens are valuable plant resources and are used as medicine, food, fodder, perfume, spice, dyes and for miscellaneous purposes throughout the world. Lichens are well known for the diversity of secondary metabolites that they produce. Compounds isolated from various lichen species have been reported to display diverse biological activities. Here we review the medicinal efficacy of lichen substances, which intends to explore the pharmaceutical potential of lichen substances.

Keywords Lichen · Lichen substances · Natural product · Bioactivity

Introduction

The use of herbal medicine is a traditional way to cure ailments and has been applied for more than five millennia in several civilizations. Even today, plant materials continue to play a major role in primary health care as therapeutic agents in many developing countries. For example, *Ocimum sanctum* and *Azadirachta indica* are known since ages to cure diseases. Besides using herbal remedies to treat many diseases natural products have contributed enormously to the development of important therapeutic drugs used currently in modern medicine. The major challenge today is the discovery of plants with promising activities and isolation of active principles. Not only plants belonging to the angiosperm but also lower plants like, bryophytes and lichens have potential healing/curing power.

Throughout the ages, lichens have been used for various purposes, in particular as dyes, perfumes and remedies in folk medicines. In Indian spice market lichens are sold by name of 'Chharila', which consists of mixture of two or more species of *Parmelia*, *Usnea longissima*, *Ramalina subcomplanata* and *Heterodermia tremulans*. Chharila has astringent, resolvent, laxative and carminative properties and is also supposed to possess aphrodisiac property. The smoke of chharila is believed to relieve headache. The powdered drug is applied to wounds besides functioning as a good cephalic snuff. Different genera of lichens are used in curing many ailments. Species of

V. Shukla (✉)
Department of Environmental Sciences, Babasaheb
Bhimrao Ambedkar (Central) University, Vidya Vihar,
Raebareilly Road, Lucknow, UP 226 025, India
e-mail: vertika_shukla@rediffmail.com

G. P. Joshi · M. S. M. Rawat
Department of Chemistry, H.N.B. Garhwal University,
Post Box 63, Srinagar Garhwal, Uttarakhand 246 174,
India

Evernia, *Peltigera*, *Parmelia*, *Cladonia*, *Rocella* and *Pertusaria*, were used to control fever, diarrhoea, infections, skin diseases, epilepsy, convulsions and as purgative (Kumar et al. 1996). *Peltigera caninna* is a tonic and used against liver ailments because of its high methionine content (Subramanian and Ramakrishnan 1964).

Medicinal properties of lichens are mentioned in the Ayurvedic system of therapy as they are useful in diseases of blood, heart, bronchitis, scabies, leprosy, enlarged spleen, burning sensation, bleeding pile, thirst, vomiting, asthma while in Unani System lichens find use in curing inflammations, stomach disorder, dyspepsia, vomiting, pain in liver, amenorrhea, vesicular calculus. Their powder is applied to wounds (Kirtikar and Basu 1984). The fame of many lichens derives from the ‘Doctrine of Signature’. These doctrines rely on the concept of ‘like affects like’. *Parmelia sulcata* Taylor was a remedy for cranial maladies; *Peltigera canina* was used for treating rabies. *Peltigera aphosa* (L.) Wild, the thallus of which is dotted with wart like tuberoses was recommended for children who suffered from ‘Thrush’ (Upreti 1994). They have tremendous curative power. The antibiotic aspect of lichens has greatly increased its medicinal importance. This review focuses on the biological activities of the various lichen species along with the individual chemical classes of lichen metabolites responsible for diverse biological activity and their potential pharmaceutical use.

Unique characteristics and chemical composition of lichens

Lichens are stable, self-supporting, mutualistic symbionts involving fungus and microalga and/or a cyanobacterium. They are among the slowest growing plants. The growth rate of 1 cm/year is considered to be high. They have the remarkable ability to tolerate adverse atmospheric condition. Another peculiarity of lichen is the presence of secondary metabolites such as depsides and dibenzofuran (Asahina and Shibata 1954; Culberson 1969; Huneck and Yoshimura 1996), so called lichen substances (Fig. 1). According to their chemical structures, most lichen substances are phenolic compounds (orcinol and β -orcinol derivatives), dibenzofurans (e.g. usnic acid), depsides (e.g. barbatic acid), depsidones (e.g. salazinic acid),

depsones (e.g. picrolichenic acid), lactones (e.g. protolichesterinic acid, nephrosterinic acid), quinones (e.g. parietin), and pulvinic acid derivatives (e.g. vulpinic acid).


Biosynthetic pathways involved in the synthesis of lichen substances

Lichens apparently involved diverse biosynthetic pathways to produce the variety of compounds (Table 1) mainly polymalonate, shikimic acid and mevalonic acid pathways (Boustie and Grube 2005).

- (1) *Polymalonate pathway*: Lichen secondary metabolites synthesized via polymalonyl pathway belong to unique class of chemical compounds mainly, depsides and depsidones. These compounds are synthesized by fungal partner, only when it is in association with alga (lichen symbiosis). These compounds play an important role in successful symbiotic association of lichens.
- (2) *Shikimic acid pathway*: Lichen substances synthesized via shikimate pathway are mainly pulvinic acid and terphenylquinone. These compounds are obtained by fusion of two phenylpyruvate units. These chemicals widely occur in the lichen family Stictaceae.
- (3) *Mevalonic acid pathway*: Mevalonic acid pathway mainly results in the synthesis of terpenes. Sesquiterpenes have not been reported in lichens yet. Diterpenes are also rare (till now 16 α -Hydroxycuarene and Nephrene have been isolated from lichens. Triterpenes are abundant in lichens, 20 different triterpenes have been derived from different species of lichens. Zeorin is the main triene in various species of lichens.

Role of lichen substances

Secondary products from lichens can comprise of up to 20% of the dry weight of thallus although 5–10% is more common. The physiological cost in energy and carbon used to produce these compounds suggests that they have important role in defense mechanism (Dayan and Romagini 2001, 2002). More than 50% lichen substances are synthesized to sustain

Fig. 1 Lichen substances**Table 1** Biosynthetic pathways involved in the synthesis of lichen substances

	Biosynthetic pathways	Lichen substances
1	<i>Acetyl polymalonyl pathway</i>	Depside, Depsidones, Chromone, Xanthone, Dibenzofuran, Emodin, Anthraquinone and Androcrocin
2	<i>Shikimic acid pathway</i>	Pulvinic acid and its derivative Terphenylquinone
3	<i>Mevalonic acid pathway</i>	Terpene

and protect the symbiotic association from various abiotic and biotic factors. Although more than 800 lichen substances are known and their structures have been elucidated (Huneck 2001), many others remain to be characterized. The over-riding interest in these compounds has been in respect to their potential source as prospective pharmacophores (González-Tejero et al. 1995; Huneck 1999).

Biological activities of lichen substances

Secondary metabolites from lichens, especially polymalonyl derived polyketides, have been found to exhibit manifold biological activities in various screenings (Huneck 1999; 2001; Müller 2001). Compounds synthesized via polymalonyl pathway are composed of two or three orcinol and β -orcinol

type phenolic units joined by ester, ether and C–C bond. These phenolic structures could constitute a mechanism of defense against oxidative stress (Hidalgo et al. 1994). The orcinol and β -orcinol derivatives (depsides, depsidones, and dibenzofuran) are especially interesting as they presumably play a role in the establishment of the lichen symbiosis and probably in the interaction between the symbionts and their environments (Armaleo 1995).

Various chemicals have been isolated from different lichen species. Secondary metabolite chemistry of *Xanthoria elegans*, *Lecanora rupicola*, *Graphis scripta* and *G. desquamescens* has been carried out by (Brunauer et al. 2006, 2007; Miyagawa et al. 1994). Lichen acids, mainly atranorin, evernic, physodic and usnic acids, are powerful inhibitors of some metabolic enzymes, interestingly related to polyamine metabolism, such as arginase, arginine decarboxylase, ornithine decarboxylase, etc. (Legaz et al. 2001; Matsubara et al. 1997, 1998; Kinoshita et al. 2002).

Lichen polysaccharides have been receiving much attention for their biological activities, specially antitumor, immunostimulatory, and antiviral properties (Olafsdottir and Ingólfssdottir 2001). Depsidones and fatty acids were present in *Parmelia stygia* (Mischenko et al. 1984). Several species of stictaceae were studied and characterized for their constituents by two-dimensional TLC method (Culberson and Johnson 1976). Aslan et al. (2006) revealed the antioxidant and antimicrobial properties of the lichens *Cladonia foliacea*, *Dermatocarpon minutum*, *Evernia divaricata*, *Evernia prunastri*, and *Neofuscella pulla*. Secondary metabolites of some lichens were separated by HPLC method (Feige et al. 1993). Carotenoids were isolated from the members of the genera *Cladonia*, *Lobaria*, *Nephroma*, *Stereocaulon* and *Sticta* (Czeczuga et al. 1988). The chemical constitution and activities of different lichens have been enlisted in Table 2.

Potentially active depside and depsidone compounds

The structural diversity of depsides and their occurrence in lichens has evoked considerable interest in their biological activities. Many depsides reported in literature have been found to possess important

physiological properties. The antioxidant property has been reported in depsides isolated from various lichen species. It has been found that depsidones are more efficient antioxidants than depsides. The higher efficiency of the depsidones could be related to a larger incorporation into lipidic microdomains (Hidalgo et al. 1994). Bridging at the phenolic group in the *p*-position can increase the antioxidant activity of phenols due to more efficient overlap of the substitute orbital within the aromatic π system (Burton et al. 1985). Depsidones have an ether linkage in addition to the ester linkage of the depsides, resulting in a rigid polycyclic system. Therefore, they are based on an 11*H*-dibenzo[*b,e*][1,4]dioxepin-11-one ring system. This chemical feature was shown to be important for the inhibitory activity of this class of lichen metabolites against HIV-1 integrase (Neamati et al. 1997). Depsidone and depside compounds such as pannarin, 10-chloropannarin and sphaerophorin, tested in cell cultures of lymphocytes, were shown to have a higher cytotoxic effect than colchicine (Correché et al. 2002). The depsidones salazinic acid, stictic acid and psoromic acid were the most apoptotic active derivatives among 15 lichen compounds evaluated on primary cultures of rat hepatocytes (Correché et al. 2004).

Depsides like barbatic and diffractic acids are potential biologically active compounds. Barbatic acid impairs PS II functioning in higher plants while diffractic acid has fungicidal properties (Dayan and Romagani 2001, 2002; Takahagi et al. 2006). Inhibition of photosystem II in tobacco cells by barbatic acid while inhibition of growth of lettuce seedling by diffractic and barbatic acids have been reported by Nishitoba et al. (1987). Lichen acids are known to show antifungal activity (Halama and Van Haluwyn 2004). The inhibition of gastric mucosal lesion, oxidative stress and neutrophil-infiltration in rats by diffractic acid has been studied by Bayir et al. (2006). Diffractic and usnic acids are known to show moderate inhibiting activity towards arachidonate 5-lipoxygenase, antiproliferative activity and antispasmodic properties (Kumar and Müller 1999a, b, c). Atranorin isolated from *Pseudoevernia furfuracea* shows trypsin inhibition mechanism (Proksa et al. 1994). Polysaccharide (GE-3-S) characterized from *Umblicaria esculenta* shows anti-HIV activity (Hirabayashi et al. 1989). Orsellinic acid derivatives from *Evernia prunastri* are nematocidal (Ahad et al. 1991).

Table 2 Chemical constituents and biological activity of lichen species

S. no.	Lichen species	Compounds characterized	Activity	Reference
Cladoniaceae				
1	<i>Cladonia</i>	Usnic acid and other lichen acids	Antibiotic	Vartiä (1973)
2	<i>Cladonia arbuscula</i> (Wallr.) Rabenh	(+)-usnic acid	Antimycobacterial activity	Ingólfssdóttir et al. (1998)
3	<i>Cladonia stellaris</i>	Usnic acid, Perlatolic acid, Atranorin		Falk et al. (2008)
4	<i>Cladonia</i> sp.	44 phenolic lichen substances		Huovinen (1987)
5	<i>Cladonia</i> sp.	(–) usnic acid	Anti tumor	Kupchan and Kopperman (1975)
6	<i>Heterodea muelleri</i> (Hampe) Nyl.	Usnic acid, Diffractic acid and barbatolic acid	UV-C stress and cold temperature stress	Hager et al. (2008)
Stictaceae				
7	<i>Pseudocyphellaria glabra</i> and <i>P. homoeophylla</i>	Usnic acid	Antimicrobial, Antiviral and Cytotoxic	Perry et al. (1999)
Lecanoraceae				
8	<i>Lecanora hybocarpa</i>	A dimer of naphthazarin, the highly substituted pentacyclic hybocarpone	Potent cytotoxic against a murine mastocytoma cell line, with an IC ₅₀ value of 0.27 μM	Ernst-Russell et al. (1999)
Lobariaceae				
9	<i>Lobaria pulmonaria</i>	Depsidones and melanins	Light screening pigments	McEvoy et al. (2007a)
10	<i>Pseudophellaria nudata</i> (Zahlbr.) D.J. Galloway.	Phenolics, depsides, triterpenes		Cuellar et al. (2008)
Nephromataceae				
11	<i>Nephroma arcticum</i>	Usnic acid	UV Screening	McEvoy et al. (2007b)
12	<i>N. laevigatum</i>	Hypericin derivative	Anti-HSV-1	Cohen et al. (1996)
Pannariaceae				
13	<i>Erioderma chilense</i>	Depsidones 1-chloropannarin & pannarin	Antioxidant	Quilhot et al. (1983); Hidalgo et al. (1994)
Parmeliaceae				
14	<i>Alectoria ochroleuca</i>	Vulpinic and (+) (–) usnic acid	Antifungal,	Lauterwein et al. (1995)
15	<i>Cetraria islandica</i>	Funarprotocetraric and protocetraric acid	Inhibit HIV-1 reverse transcriptase	Pengsuparp et al. (1995)
16	<i>C. ericetorum</i>	Orcinol, orsellic acid, orcinol monomethyl ether, everminic acid, evermyl 3,5-dimethoxy toluene, methyl evermate, ethyl evermate, usnic acid, evermic acid and atranorin	Used in perfumery	Heide et al. (1975)
17	<i>Flavoparmelia caperata</i>	Polysaccharide (PC 2)	Long term potentiation enhancement	Smriga et al. (1996)

Table 2 continued

S. no.	Lichen species	Compounds characterized	Activity	Reference
18	<i>Hypotrachyna revoluta</i>	8'-methylmenegazziac acid	Radical scavenging (RSA)	Papadopoulou et al. (2007)
19	<i>Lethariella canariensis</i>	Atranol, chloroatranol, methyl β -orsellinate methyl hematommate ethyl hematommate	Phytotoxicity and Antioxidative	Toledo Marante et al. (2003)
20	<i>Parmelia caperata</i>	Four orsellinic acid derivatives, n-octacosanol, β -sitosterol, 3-D-glucopyranoside		Saraswathy et al. (1990)
21	<i>P. loxodes</i> and <i>P. verruculifera</i>	Orcinol type depside		Culberson and Johnson (1976)
22	<i>P. stenophylla</i> <i>P. scorcia</i> <i>P. sulcata</i>	68 fatty acid		Dembitsky et al. (1992)
23	<i>Parmelia stygia</i>	Fumaroprotocetraric acid depsidones		Mischenko et al. (1984)
24	<i>Peltigera aphosa</i>	Tenuiorin, methyl gyrophorate, methyl evermate, methyl lecanorate, methyl orsellinate, 4-O- methyl gyrophoric and evermic acid	Smooth muscle relaxant	Maass (1975)
25	<i>Protousnea poeppigii</i> (Nees & Flot.) Vain.	Isodivartic acid, 5-propylresorcinol, divartic acid, usnic acid	Antifungal & antiprotozoal	Schmeda-Hirschmann et al. (2007)
26	<i>Pseudevernia furfuracea</i>	Physodic acid, chloroatranorin, atranorin, and olivetonic acid	Antimicrobial	Türk et al. (2006)
27	<i>Protousnea</i> sp.	Resorcinol derivative	Tyrosinase inhibition	Kinoshita et al. (1994)
28	<i>Pseudoevernia furfuracea</i> <i>Evernia prunastri</i> Pertusariaceae	D-Uscic acid, evermic acid and atranorin	Allergic	Goncalo (1987)
29	<i>Thamnomelia subuliformis</i> Ramalinaceae	Rhamnopyranosyl galactofuranan	Immunologically active	Olafsdottir et al. (1999)
30	<i>Ramalina almqvisii</i>	D-Protolichesterinic acid and nephrostermic acid	Anti-tumor	Hirayama et al. (1980)
31	<i>R. farinacea</i> (L.) Ach.	(+)-Uscic acid, evermic and obtusatic acid		Rastogi and Mehrotra (1993)
32	<i>R. farinacea</i>	(+)-Uscic Acid, Norsitic Acid, and Protocetraric Acid	Antimicrobial	Tay et al. (2004)
33	<i>R. subcomplanata</i> .	Secondary metabolites	Antibacterial	Kumar et al. (1995)
34	<i>Roccella belangeriana</i> (Awasthi) Stereocaulaceae	Chloroform extract	Antibacterial	Karthikaidevi et al. (2009)
35	<i>Stereocaulon alpinum</i> Laur	Lobaric acid	Inhibit 5-lipoxygenase from porcine leukocytes, with an IC ₅₀ value of 7 μ M	Ingólfssdóttir et al. (1996)

Table 2 continued

S. no.	Lichen species	Compounds characterized	Activity	Reference
36	<i>Stereocaulon ramulosum</i>	Methyl haematommate	Antifungal	Hickey et al. (1990)
37	Stictaceae <i>Pseudocyphellaria glabra</i> and <i>P. homoeophylla</i>	Usnic acid	Antimicrobial, Antiviral and Cytotoxic	Perry et al. (1999)
38	Teloschistaceae <i>Caloplaca</i> species	Anthraquinone	Broad-spectrum antifungal and antibacterial properties	Manojlovic et al. (2005)
39	<i>Xanthoria parietina</i>	Parietin	UV-protecting agent	Solhaug et al. (2003)
40	Thelotremaaceae <i>Diploschistes muscorum</i>	Secondary metabolites	Oxidative stress	Cuny et al. (2004)
41	Umbilicariaceae <i>Gyrophora exculenta</i>	Secondary metabolites	Inhibitory effect on cyclooxygenase activity of prostaglandin H ₂ synthase	Min et al. (1996)
42	<i>Umbilicaria mammulata</i>	Secondary metabolites	Low temperature stress	Stocker-Wörgötter (2001a)
43	<i>U. Antarctica</i>	Gyrophoric acid, lecanoric acid, and methyl orsellinate	Inhibition of PTP1B	Seo et al. (2009)
44	Usneaceae <i>Usnea campestriis</i>	Usnic acid	Anti-fungal & anti-bacterial	Gutkind et al. (1981)
45	<i>U. diffracta</i>	Diffractaic acid usnic acid	Analgesic & antipyretic	Okuyama et al. (1995)
46	<i>U. diffracta</i>	Diffractaic acid usnic acid	Anti-inflammatory	Otsuka et al. (1972)
47	<i>U. diffracta</i>	Decarboxy stenosporic acid	<i>Staphyrococcus aureus</i>	Yamamoto et al. (1998)
48	<i>U. longissima</i>	Usnic acid, depside	Photosystem II inhibition	Endo et al. (1998)
49	<i>U. longissima</i>	Usnic acid	Plant growth inhibition	Nishitoba et al. (1987)
50	<i>U. longissima</i>	Lichesterinic, (+)- usnic and evermic acid	Epstein-Barr virus activation inhibition	Yamamoto et al. (1995)
51	<i>U. misaminensis</i>	Usnic acid	Smooth muscle relaxant	Mustafa et al. (1995)
52	<i>U. venosa</i>	D-Usnic acid, barbatric acid and norstictic acid		Rastogi and Mehrotra (1991)

Broad spectrum antibiotic compounds

Dibenzofuran, usnic acid is a molecule with wide spectrum of activities (Ingólfðóttir 2002; Cardarelli et al. 1997; Lauterwein et al. 1995; Huovinen and Lampero, 1989; Kupchan and Kopperman 1975; Duman et al. 2008). Usnic acid is known to be present in three forms, i.e. (–)-usnic acid, (+)-usnic acid and isousnic acid (Shibata and Taguchi 1967). Both (–) and (+)-usnic acid are biologically important. Photo system II inhibition is shown by usnic acid (Inoue et al. 1987). Usnic acid is also a non-genotoxic compound with anticancer properties (Mayer et al. 2005). Derivatives of usnic acid are also reported to be antineoplastic (Takai et al. 1979). (–)-usnic acid isolated from *Alectoria ochroleuca* has shown antifungal activity (Proksa et al. 1996).

UV screening compounds

Lichen are also known to synthesize, UV-B-absorbing cortical pigments, like brown melanic compounds in *Lobaria pulmonaria* and orange parietin in *Xanthoria aureola*, which act as uv filter and protects the inner algal layer from harmful effects of uv radiation. These compounds are produced by the mycobiont and induced by UV-B radiation. UV-B has been reported to induce production of usnic acid (Solhaug et al. 2003; McEvoy et al. 2006; Nybakken and Julkunen-Tiitto 2006). Low temperature also can set off an increased production of secondary metabolites, as reported by Stocker-Wörgötter (2001a). According to Swanson and Fahselt (1997) and Begora and Fahselt (2001), UV-A irradiation alone causes the production of higher amount of phenolic compounds in lichens than exposed to both UV-A and UV-B radiation.

Pulvinic acid derivatives

Apart from compounds synthesized via polymalonyl pathway, other compounds synthesized via mevalonate and shikimate pathways are potentially active. Vulpinic acid is also known to possess antimicrobial activity against aerobic and anaerobic microorganisms. It also induces uncoupling of oxidative phosphorylation (Abo-Khatwa et al. 1996).

Hydroxybenzoic acid derivatives

Esters of 4-hydroxybenzoic acid such as methylparaben are used as preservatives in the pharmaceutical preparations. As these agents are rapidly hydrolyzed in vivo to the corresponding acid which is then conjugated and excreted, their toxicity is generally low (Wilson et al. 1998).

Sterol compounds

Due to different types of pharmacological activities like antiinflammatory, antiulcerogenic, antibacterial, antifungal, antirheumatic activities, the vast majority of steroids play an important role in the field of medicines. It has been reported that sterols may also play a pivotal role in the membrane permeability and perhaps influence the transport of materials from algae to fungi in lichens. In general C₂₈ sterols predominate in fungi while C₂₉ sterols are found in algae (Lenton et al. 1973). NMR spectral analysis of sterols is an aid to characterize the structure (Thompson et al. 1972). Sterols have been isolated and characterized from *Xanthoria parietina* (Safe et al. 1975), *Pseudoevernia furfuracea* (Wojciechowski et al. 1973) and *Lobaria pulmonaria*, *L. scrobiculata* and *Usnea longissima* (Safe et al. 1975). Brassicasterol, lichensterol, poriferasterol, β -sitosterol have been reported from *Ramalina africana* (Shukla et al. 2004).

The above-mentioned activities clearly indicate the potential of lichen compounds for pharmaceutical purposes, but some properties of lichens still need consideration. Their slow growth, both in nature and in axenic cultures, may be regarded as a major problem for the production of metabolites. However, culture conditions of lichen mycobionts can be optimized in order to support the synthesis of interesting secondary compounds. Much progress has been achieved recently by specifically addressing these issues (Stocker-Wörgötter 1998, 2001b, 2002a, b, Stocker-Wörgötter and Elix 2004).

Further manifold biological activities of lichen metabolites have now been recognized, but their therapeutic potential has not yet been fully explored. In particular, the depsides and depsidones isolated from lichens are of special interest because of their antiinflammatory, antiproliferative and anti-HIV activities.

Conclusion

Most of the preliminary information on biological activity and potential use of lichens and/or their metabolites are derived from their ethno-botanical lore and ‘Doctrine of Signatures’. But the recent work on the biochemical and physiological activities associated with the individual components of lichen extract reveals tremendous potential of these compounds for mankind. With the aid of modern equipments and methodologies, isolation and characterization of active compounds and elucidation of mechanism of action has become easier.

Chemical structure of most of the lichen substances is simple which facilitates synthesis of these compounds in the laboratory. This practice would provide large amounts of material without affecting ecosystem. In addition, many of these compounds can be used as precursors based on their particular mechanism of action and can then be optimized in the laboratory to fit specific applications. Appropriate chemical modification may increase the stability and reduce the toxicity of the natural products and could lead to an increase in their potency.

Acknowledgments Author (V.S.) wish to thanks Scientific and Engineering Research Council, D.S.T., New Delhi for the financial support vide project no. SR/FT/LS-028/2008.

References

- Abo-Khatwa AN, Al-Robai AA, Al-Jawhari DA (1996) Lichen acids as uncouplers of oxidative phosphorylation of mouse liver mitochondria. *Nat Toxins* 4:96–102
- Ahad AM, Goto Y, Kiuchi F, Tsuda Y, Kondo K, Sato T (1991) Nematocidal principles in “oakmoss absolute” and nematocidal activity of 2, 4-dihydroxybenzoates. *Chem Pharm Bull* 39:1043–1046
- Armaleo D (1995) Factors affecting depside and depsidone biosynthesis in a cultured lichen fungus. *Crypt Bot* 5:14–21
- Asahina Y, Shibata S (1954) Chemistry of lichen substances. Japan Society for the Promotion of Science, Tokyo
- Aslan A, Güllüce M, Sökmen M, Adıgüzel A, Sahin F, Özkan H (2006) Antioxidant and Antimicrobial Properties of the Lichens *Cladonia foliacea*., *Dermatocarpon miniatum*., *Evernia divaricata*., *Evernia prunastri*., and *Neofuscella pulla*. *Pharmaceutical Biol* 44(4):247–252
- Bayir Y, Odabasoglu F, Cakir A, Aslan A, Suleyman H, Halici M, Kazaz C (2006) The inhibition of gastric mucosal lesion, oxidative stress and neutrophil- infiltration in rats by the lichen constituent diffractaic acid. *Phytomedicine* 13:584–590
- Begora MD, Fahselt D (2001) Usnic acid and atranorin concentrations in lichens in relation to bands of UV irradiance. *Bryologist* 104:134–140
- Boustie J, Grube M (2005) Lichens—a promising source of bioactive secondary metabolites. *Plant Genetic Resources* 3(2):273–287
- Brunauer G, Hager A, Krautgartner WD, Turk R, Stocker-Wörgötter E (2006) Experimental studies on *Lecanora rupicola* (L.) Zahlbr.: chemical and microscopical investigations of the mycobiont and re-synthesis stages. *Lichenologist* 38:577–585
- Brunauer G, Hager A, Grube M, Turk R, Stocker-Wörgötter E (2007) Alterations in secondary metabolism of aposymbiotically grown mycobionts of *Xanthoria elegans* and cultured resynthesis stages. *Plant Physiol Biochem* 45:146–151
- Burton GW, Doba T, Gabe EJ, Hughes L, Lee FL, Prasad L, Ingold KU (1985) Autoxidation of biological molecules. 4. Maximizing the antioxidant activity of phenols. *J Am Chem Soc* 107:7053
- Cardarelli M, Serino G, Campanella L, Ercole P, De Cicco Nardone F, Alesiani O, Osiello F (1997) Antimitotic effects of usnic acid on different biological systems. *Cell Mol Life Sci* 53:667–672
- Cohen PA, Hudson JB, Towers GHN (1996) Antiviral activities of anthraquinones, bianthrone and hypericin derivatives from lichens. *Experientia* 52:180–183
- Correché ER, Carrasco M, Giannini F, Piovano M, Garbarino J, Enriz D (2002) Cytotoxic screening activity of secondary lichen metabolites. *Acta Farm Bon* 21:273–278
- Correché ER, Enriz RD, Piovano M, Garbarino J, Gomez-Lechon MJ (2004) Cytotoxic and apoptotic effects on hepatocytes of secondary metabolites obtained from lichens. *Alter Lab Animals* 32:605–615
- Cuellar M, Quilhot W, Rubio C, Soto C, Espinoza L, Carrasco H (2008) Phenolics, depsides, triterpenes from Chilean lichen *Pseudophellaria nudata* (Zahlbr.) D.J Galloway. *J Chilean Chem Soc* 53(3):1624–1625
- Culberson CF (1969) Chemical and botanical guide to lichen products. University of North Carolina Press, Chapel Hill
- Culberson CF, Johnson A (1976) A standardized two dimensional thin-layer chromatographic method for lichen products. *J Chromatogr* 128:253–259
- Cuny D, Van Haluwyn C, Shirali P, Zerimech F, Jerome L, Haguenoer JM (2004) Cellular impact of metal trace elements in terricolous lichen *Diploschistis muscorum* Scop.) R. Sant.—Identification of oxidative stress biomarkers. *Water Air Soil Pollut* 152:55–69
- Czeczuga B, Cifuentes B, Reynaud PA (1988) Carotenoids in lichens from the Canary Islands. *Biochemical systematics and ecology. Biochem Sysemat Eco* 16(2):117
- Dayan FE, Romagini JG (2001) Lichens as a potential source of pesticides. *Pestic Outlook* 12:229–232
- Dayan FE, Romagini JG (2002) Structural diversity of lichen metabolites and their potential for use. In: Upadhyaya R (ed) *Advances in microbial toxin research and its biotechnological Exploration*. Kluwer Academic/Plenum Publisher, New York, p 151
- Dembitsky VM, Bychek IA, Kashin AG (1992) Chemical constituents of some lichen species. *J Hatt Bot Lab* 255–262

- Duman DC, Aras S, Atakol O (2008) Determination of Usnic Acid Content in Some Lichen Species Found in Anatolia. *J Appl Biol Sci* 2(3):41–44
- Endo T, Takahagi T, Kinoshita Y, Yamamoto Y, Sato F (1998) Inhibition of photosystem II of spinach by lichen-derived depsides. *Biosci Biotech Biochem* 62(10):2023
- Ernst-Russell MA, Elix JA, Chai CLL, Willis AC, Hamada N, Nash TH (1999) Hybocarpace, a novel cytotoxic naphthazarin derivative from mycobiont cultures of the lichen *Lecanora hybocarpa*. *Tetrahedron Lett* 40:6321–6324
- Falk A, Green TK, Barboza P (2008) Quantitative determination of secondary metabolites in *Cladina stellaris* and other lichens by micellar electrokinetic chromatography. *J Chromatogr A* 1182:141–144
- Feige GB, Lumbsch HT, Huneck S, Elix JA (1993) The identification of lichen products by a standardized high-performance liquid chromatographic method. *J Chromatogr* 646:417–427
- Goncalo S (1987) Contact sensitivity to lichens and Compositae in *Frullania dermatitis*. *Cont Derm* 16(2):84
- González-Tejero MR, Molero-Mesa J, Casares-Porcel M, Martínez Lirrola MJ (1995) New contributions to the ethnopharmacology of Spain. *J Ethnopharmacol* 45:157–165
- Gutkind GO, Martino V, Grana N, Coussio JD, Torres RA (1981) Screening of South American plants for biological activities. Antibacterial and antifungal activity. *Fitoterapia* 52(5):213
- Hager A, Brunauer G, Türk R, Stocker-Wörgötter E (2008) Production and Bioactivity of Common Lichen Metabolites as Exemplified by *Heterodea muelleri* (Hampe) Nyl. *J Chem Ecol* 34:113–120
- Halama P, Van Haluwyn C (2004) Antifungal activity of lichen extracts and lichenic acids. *Bio Control* 49:95–107
- Heide R, Provatoroff N, Taas PC, Valois J, Plasse N, Wobben HJ, Timmer R (1975) Qualitative analysis of the odoriferous fraction of oakmoss (*Evernia prunastri* (L.) Ach.). *J Agri Food Chem* 23(5):950–957
- Hickey BJ, Lumsden AJ, Cole ALJ, Walker JRL (1990) Antibiotic compounds from New Zealand plants: methyl haematamate, an anti-fungal agent from *Stereocaulon ramulosum*. *New Zealand Nat Sci* 17:49–53
- Hidalgo ME, Fernández E, Quilhot W, Lissi EA (1994) Antioxidant capacity of depsides and depsidones. *Phytochemistry* 37:1585–1587
- Hirabayashi K, Iwata S, Ito M, Shigeta S, Narui T, Mori T, Shibata S (1989) Inhibitory effect of a lichen polysaccharide sulfate, GE-3-S, on the replication of human immunodeficiency virus (HIV) in vitro. *Chem Pharm Bull* 37:2410–2412
- Hirayama T, Fujikawa F, Kasahara T, Otsuka M, Nishida N, Mizuno D (1980) Anti-tumor activities of some lichen products and their degradation products. *Yaku Zasshi* 100:755–759
- Huneck S (1999) The significance of lichens and their metabolites. *Naturwissenschaften* 86:559–570
- Huneck S (2001) New Results on the Chemistry of Lichen Substances. In: Herz W, Falk H, Kirby GW, Moore RE (eds) *Progress in the chemistry of organic natural products*. Springer, Wien, New York, p 81
- Huneck S, Yoshimura I (1996) Identification of lichen substances. Springer, Berlin, pp 1–493
- Huovinen K (1987) A standard HPLC method for the analysis of aromatic lichen in Progress and Problems in Lichenology in the Eighties. *Bibliotheca Lichenolog* 25:457–466
- Huovinen K, Lampero M (1989) Usnic acid as a mitotic inhibitor in the Allium Test. *Planta Med* 55:98
- Ingólfssdóttir K (2002) Usnic acid. *Phytochemistry* 61:729–736
- Ingólfssdóttir K, Gissurarson SR, Müller-Jakic B, Breu W, Wagner H (1996) Inhibitory effects of the lichen metabolite lobaric acid on arachidonate metabolism in vitro. *Phytomedicine* 2:243–246
- Ingólfssdóttir K, Chung GAC, Skúlason VG, Gissurarson SR, Vilhelmsdóttir M (1998) Antimycobacterial activity of lichen metabolites in vitro. *Eur J Pharm Sci* 6:141–144
- Inoue H, Noguchi M, Kubo K (1987) Site of inhibition of usnic acid at oxidizing side of photosystem 2 of spinach chloroplast. *Photosynthetica* 21:88
- Karthikaidevi G, Thirumaran G, Manivannan K, Anantharaman P, Kathiresan K, Balasubramanian T (2009) Screening of the Antibacterial Properties of Lichen *Roccella belangeriana* (Awasthi) from Pichavaram Mangrove (Rhizophora Sp.). *Adv Biol Res* 3(3–4):127–131
- Kinoshita K, Matsubara H, Koyama K, Takahashi K, Youhimura I, Yamamoto Y, Higuchi M, Miura Y, Kinoshita K, Kawai K (1994) New phenolics from *Protousnea* species. *J Hatt Bot Lab* 75:359–364
- Kinoshita K, Saito D, Koyama K, Takahashi K, Sato Y, Okuyama E, Fujimoto H, Yamazaki M (2002) Monoamine oxidase inhibitory effects of some lichen compounds and their synthetic analogues. *J Hatt Bot Lab* 92:277–284
- Kirtikar KR, Basu BD (1984) *Indian Medicinal Plants*. Lalit Mohan Basu Publication 49, Leader Road, Allahabad, India, 4, 2760
- Kumar KCS, Müller K (1999a) Lichen metabolites. 1. Inhibitory action against leukotriene B4 biosynthesis by a nonredox mechanism. *J Nat Pro* 62:817–820
- Kumar KCS, Müller K (1999b) Lichen metabolites. 2. Antiproliferative and cytotoxic activity of gyrophoric, usnic, and diffractaic acid on human keratinocyte growth. *J Nat Pro* 62:821–823
- Kumar KCS, Müller K (1999c) Depsides as non-redox inhibitors of leukotriene B4 biosynthesis and HaCaT cell growth. 1. Novel analogs of barbatic and diffractaic acid. *Eur J Med Chem* 34:1035–1042
- Kumar KCS, Banskota AH, Manandhar MD (1995) Chemical constituents and antibacterial activity of *Ramalina subcomplanata*. *J Nepal Chem Soc* 14:28–35
- Kumar KCS, Banskota AH, Manandhar MD (1996) Isolation and identification of some chemical constituents of *Parmelia nepalensis*. *Planta Med* 62:93–94
- Kupchan SM, Kopperman HL (1975) L-Usnic acid: tumor inhibitor isolated from lichens. *Experientia* 31:625
- Lauterwein M, Oethinger M, Belsner K, Peters T, Marre R (1995) In vitro activities of the lichen secondary metabolites vulpinic acid, (+)-usnic acid, and (–)-usnic acid against aerobic and anaerobic microorganisms. *Antimicrob Agents Chemother* 39:2541–2543
- Legaz ME, Vicente C, Pedrosa MM (2001) Binding of lichen phenolics to purified secreted arginase from the lichen *Evernia prunastri*. *J Biochem Mol Bio* 34:194–200
- Lenton JR, Goad LJ, Goodwin TW (1973) Sterols of Xanthoria parietina evidence for 2 sterol pools and the identification

- of a novel 28 carbon triene erosta-5–8, 22–trien-3- β -ol. *Phytochemistry* 12:1135–1140
- Maass WSG (1975) The phenolic constituents of *Peltigera aphthosa*. *Phytochemistry* 14:2487
- Manojlovic NT, Novakovic M, Stevovic V, Solujic S (2005) Antimicrobial metabolites from three serbian *Caloplaca*. *Pharm Bio* 43(8):718–722
- Matsubara H, Kinoshita K, Koyama K, Ye Y, Takahashi K, Yoshimura I, Yamamoto Y, Miura Y, Kinoshita Y (1997) Antityrosinase activity of lichen metabolites and their synthetic analogues. *J Hatt Bot Lab* 83:179–185
- Matsubara H, Miharuru K, Kinoshita K, Koyama K, Ye Y, Takahashi K, Yoshimura I, Yamamoto Y, Miura Y, Kinoshita Y (1998) Inhibitory effect of lichen metabolites and their synthetic analogues on melanin biosynthesis in cultured B-16 mouse melanoma cells. *Nat Pro Sci* 4:161–169
- Mayer M, O'Neill MA, Murray KE, Santos-Magalhaes NS, Carneiro-Leao AM, Thompson AM, Appleyard VCL (2005) Usnic acid: a non-genotoxic compound with anticancer properties. *Anti Canc Drugs* 16:805–809
- McEvoy M, Nybakken L, Solhaug KA, Gauslaa Y (2006) UV triggers the synthesis of the widely distributed secondary compound usnic acid. *Mycol Prog* 5:221–229
- McEvoy M, Gauslaa Y, Solhaug KA (2007a) Changes in pools of depsidones and melanins, and their function, during growth and acclimation under contrasting natural light in the lichen *Lobaria pulmonaria*. *New Phytol* 175:271–282
- McEvoy M, Solhaug KA, Gauslaa Y (2007b) Solar radiation screening in usnic acid containing cortices of the lichen *Nephroma arcticum*. *Symbiosis* 43:143–150
- Min KR, Kim Y, Kang SH, Mar W, Lee KS, Ro JS, Lee SH, Kim Y (1996) Inhibitory Effects of Herbal Extracts on Cyclooxygenase Activity of Prostaglandin H_2 Synthase from Sheep Seminal Vesicle. *Nat Pro Sci* 2(1):56
- Mischenko NP, Maximo OB, Krivoschekova OE, Stepanenko LS (1984) Depsidones and fatty acids of *Parmelia stygia*. *Phytochemistry* 23(1):180
- Miyagawa H, Hamada N, Sato M, Ueno T (1994) Pigments from the cultured lichen mycobionts of *Graphis scripta* and *G. desquamescens*. *Phytochemistry* 36:1319–1332
- Müller K (2001) Pharmaceutically relevant metabolites from lichens. *Appl Microbiol Biotechnol* 56:9–16
- Mustafa MR, Mohammad R, Din L, Wahid S (1995) Smooth muscle relaxant activities of compounds isolated from Malaysian medicinal plants on rat aorta and guinea-pig ileum. *Phytotherapy Res* 9(8):555
- Neamati N, Hong H, Mazumder A, Wang S, Sunder S, Nicklaus MC, Milne GWA, Proksa B, Pommier Y (1997) Depsides and depsidones as inhibitors of HIV-1 integrase: discovery of novel inhibitors through 3D database searching. *J Med Chem* 40:942–951
- Nishitoba Y, Nishimura H, Nishiyama T, Mizutani J (1987) Lichen acids, plant growth inhibitors from *Usnea longissima*. *Phytochemistry* 26:3181
- Nybakken L, Julkunen-Tiitto R (2006) UV-B induces usnic acid in reindeer lichens. *Lichenologist* 38:477–485
- Okuyama E, Umeyama K, Yamazaki M, Kinoshita Y, Yamamoto Y (1995) Usnic acid and diffractaic acid as analgesic and antipyretic components of *Usnea diffracta*. *Planta Med* 61:113–115
- Olafsdottir ES, Ingólfssdottir K (2001) Polysaccharides from lichens: structural characteristics and biological activity. *Planta Med* 67:99–208
- Olafsdottir ES, Omarsdottir S, Paulsen B, Turcic K, Wagner H (1999) Rhamnopyranosylgalactofuranan, a new immunologically active polysaccharide from *Thamnia subuliformis*. *Phytomed* 6(4):273
- Otsuka H, Komiya T, Tsukumi M, Toyosato T, Fujimura H (1972) Studies on anti-inflammatory drugs. Anti-inflammatory activity of crude drugs and plants. (II). *J Takeda Res Lab* 31:247–254
- Papadopoulou P, Tzakou O, Vagias C, Kefalas P, Roussis V (2007) β -Orcinol Metabolites from the Lichen *Hypotrachyna revolute*. *Molecules* 12:997–1005
- Pengsuparp T, Cai L, Constant H, Fong HHS, Lin L-Z, Kinghorn AD, Pezzuto JM, Cordell GA, Ingólfssdottir K, Wagner H, Hughes SH (1995) Mechanistic evaluation of new plant derived compounds that inhibit HIV-1 reverse transcriptase. *J Nat Prot* 58:1024–1031
- Perry NB, Benn MH, Brennan NJ, Burgess NJ, Ellis D, Galloway DJ, Lorimer SD, Tangney RS (1999) Antimicrobial, Antiviral and Cytotoxic Activity of New Zealand Lichens. *Lichenologist* 31:627–636
- Proksa B, Adamcova J, Sturdikova M, Fуска J. (1994) Metabolites of *Pseudevernia furfuracea* (L.) Zopf. and their inhibition potential of proteolytic enzymes. *Pharmazie* 49:282
- Proksa B, Sturdikova M, Na Pronayova, Liptaj T (1996) (–)-Usnic acid and its derivatives: Their inhibition of fungal growth and enzyme activity. *Pharmazie* 51:195–196
- Quilhot W, Didyk B, Gambaro V, Garbarino JA (1983) Studies on Chilean lichens VI. Depsidones from *Erioderma chilense*. *J Nat Pro* 46(6):942–943
- Rastogi RP, Mehrotra BN. (1991) Compendium of Indian medicinal plants. Central Drug Research Institute, Lucknow and Publication and Information Directorate, New Delhi, 1(1970–1979), 425
- Rastogi RP, Mehrotra BN. (1993). Central Drug Research Institute, Lucknow and Publication and Information Directorate, New Delhi, 2(1970–1979), 577
- Safe S, Safe LM, Maass WSG (1975) Sterols of three lichen species: *Lobaria pulmonaria*, *Lobaria scrobiculata* and *Usnea longissima*. *Phytochemistry* 14:1821–1823
- Saraswathy A, Rajendiran A, Sarada A, Purushothamam KK (1990) Lichen substances of *Parmelia caperata*. *Indian Drugs* 27(9):4602
- Schmeda-Hirschmann G, Tapia A, Lima B, Pertino M, Sortino M, Zacchino S, Rojas de Arias A, Feresin GF (2007) A new antifungal and antiprotazoal depside from the andean lichen *Protosnea poeppigii*. *Phytotherapy Res* 22(3):349–355
- Seo C, Choi Y, Ahn JS, Yim JH, Lee HK, Oh H (2009) PTP1B inhibitory effects of tridepside and related metabolites isolated from the Antarctic lichen *Umbilicaria antarctica*. *J Enz Inhibit Med Chem* 24(5):1133–1137
- Shibata S, Taguchi H (1967) Occurrence of isousnic acid in lichens with reference the “isodihydrousnic acid” derived from dihydrousnic acid. *Tetrahedron Lett* 48:867–871
- Shukla V, Negi S, Rawat MSM, Pant G, Nagatsu A (2004) Chemical Study of *Ramalina africana* (Ramalinaceae) from Garhwal Himalayas. *Biochem Systemat Ecol* 32: 449–453

- Smriga M, Saito H, Shibata S, Narui T, Okuyama T, Nishiyama N (1996) PC-2, Linear Homoglucan with-ALPHA-linkages, Peripherally Enhances the Hippocampal Long-Term Potentiation. *Pharm Res* 13(9):1322–1326
- Solhaug KA, Gauslaa Y, Nybakken L, Bilger W (2003) UV-induction of sun-creening pigments in lichens. *New Phytol* 158:91–100
- Stocker-Wörgötter E (1998) Culture methods and culture of selected mycobionts and photobionts as exemplified by South American lichens. In: Marcelli MP, Seaward MRD (eds) *Lichenology in Latin America: history, current knowledge and applications*. CETESB, São Paulo, p 143
- Stocker-Wörgötter E (2001a) Experimental studies of the lichen symbiosis: DNA-analyses, differentiation and secondary chemistry of selected mycobionts, artificial resynthesis of two- and tripartite symbioses. *Symbiosis* 30:207–227
- Stocker-Wörgötter E (2001b) Experimental lichenology and microbiology of lichens: culture experiments, secondary chemistry of cultured mycobionts, resynthesis and thallus morphogenesis. *The Bryologist* 104:576–581
- Stocker-Wörgötter E. (2002a) Laboratory cultures of selected lichen fungi from Brazil and Chile. *Mitteilungen des Instituts für Allgemeine Botanik, Hamburg* 30:253–270
- Stocker-Wörgötter E (2002b) Analysis of secondary compounds in cultured mycobionts. In: Kranner I, Beckett RP, Varma AK (eds) *Protocols in lichenology, culturing, biochemistry, ecophysiology and use in biomonitoring*. Springer lab manual. Springer, Berlin, pp 296–306
- Stocker-Wörgötter E, Elix JA (2004) Experimental studies of lichenized fungi: formation of rare depsides and dibenzofuranes by the culture mycobiont of *Bunodophoron patagonicum* (Sphaerophoraceae, lichenized Ascomycota). *Bibliotheca Lichenologica* 88:659–669
- Subramanian SS, Ramakrishnan S (1964) Amino acids of *Peltigera canina*. *Curr Sci* 33:522
- Swanson A, Fahselt D (1997) Effects of ultraviolet on polyphenolics of *Umbilicaria americana*. *Can J Bot* 75: 284–289
- Takahagi T, Ikezawa N, Endo T, Ifuku K, Yamamoto Y, Kinoshita Y, Takeshita S, Sato F (2006) Inhibition of PSII in atrazine-tolerant tobacco cells by barbatic acid, a lichen-derived depside. *Biosci Biotech Biochem* 70:266–268
- Takai M, Uehara Y, Beisler JA (1979) Usnic acid derivatives as potential antineoplastic agents. *J Med Chem* 22: 1380–1384
- Tay T, Türk AO, Yılmaz M, Türk H, Kıvanc M (2004) Evaluation of the antimicrobial activity of the acetone extract of the lichen *Ramalina farinacea* and its (+)-Usnic Acid, norstictic acid and protocetraric. *Acid Constituents Z Naturforsch* 59c:384–388
- Thompson MJ, Dutky SR, Patterson G-W, Gooden EL (1972) NMR spectra of C-24 isomeric sterols. *Phytochemistry* 2:1781
- Toledo Marante FJ, Castellano AG, Rosas FE, Aguiar JQ, Barrera JB (2003) Identification and quantitation of allelochemicals from lichen *Lethariella canariensis*: Phytotoxicity and antioxidative activity. *J Chem Ecol* 29(9): 2049–2071
- Türk H, Yılmaz M, Tay T, Türk AO, Kıvanc M (2006) Antimicrobial activity of extracts of chemical races of the lichen *Pseudevernia furfuracea* and their physodic acid, chloroatranorin, atranorin, and olivetoric acid constituents. *Z Naturforsch* 61(7–8):499–507
- Upreti DK (1994) Lichens: The great Benefactors. *Applied Botany Abstract* 14(3):64–75
- Vartia KO (1973) Antibiotics in lichens. In: Ahmadjiian V, Hale ME (eds) *The lichens*, 3rd edn. Academic, New York, pp 547–561
- Wilson CO, Gisvold O, Delgado JN, Remers WA (1998) *Textbook of organic medicinal and pharmaceutical chemistry*, 10th edn. Lippincott-Raven, Philadelphia
- Wojciechowski ZA, Goad LJ, Goodwin TW (1973) Sterols of lichen *Pseudevernia furfuracea*. *Phytochemistry* 12(6): 1433–1436
- Yamamoto Y, Miura Y, Kinoshita Y, Higuchi M, Murakami A, Ohigashi H, Koshimizu K (1995) Screening of tissue cultures and thalli of lichens and some of their active constituents for inhibition of tumor promoter-induced Epstein-Barr virus activation. *Chem Pharm Bull* 43: 1388–1390
- Yamamoto Y, Kinoshita Y, Matsubara H, Kinoshita K, Koyama K, Takahashi K, Kurokawa T, Yoshimura I (1998) Screening of biological activities and isolation of biological active compounds from lichens. *Recent Res Dev Phytochem* 2:23–24