

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/40689596>

# Novel functions of photoreceptor guanylate cyclases revealed by targeted deletion

ARTICLE *in* MOLECULAR AND CELLULAR BIOCHEMISTRY · DECEMBER 2009

Impact Factor: 2.39 · DOI: 10.1007/s11010-009-0322-z · Source: PubMed

---

CITATIONS

33

---

READS

16

3 AUTHORS, INCLUDING:




Wolfgang Baehr

University of Utah

213 PUBLICATIONS 10,157 CITATIONS

SEE PROFILE

**ISSN 0300-8177, Volume 334, Combined 1-2**



**This article was published in the above mentioned Springer issue.  
The material, including all portions thereof, is protected by copyright;  
all rights are held exclusively by Springer Science + Business Media.**

**The material is for personal use only;  
commercial use is not permitted.**

**Unauthorized reproduction, transfer and/or use  
may be a violation of criminal as well as civil law.**

# Novel functions of photoreceptor guanylate cyclases revealed by targeted deletion

Sukanya Karan · Jeanne M. Frederick ·  
Wolfgang Baehr

Received: 10 January 2009 / Accepted: 4 November 2009 / Published online: 9 December 2009  
© Springer Science+Business Media, LLC. 2009

**Abstract** Targeted deletion of membrane guanylate cyclases (GCs) has yielded new information concerning their function. Here, we summarize briefly recent results of laboratory generated non-photoreceptor GC knockouts characterized by complex phenotypes affecting the vasculature, heart, brain, kidney, and other tissues. The main emphasis of the review, however, addresses the two GCs expressed in retinal photoreceptors, termed GC-E and GC-F. Naturally occurring GC-E (*GUCY2D*) null alleles in human and chicken are associated with an early onset blinding disorder, termed “Leber congenital amaurosis type 1” (LCA-1), characterized by extinguished scotopic and photopic ERGs, and retina degeneration. In mouse, a GC-E null genotype produces a recessive cone dystrophy, while rods remain functional. Rod function is supported by the presence of GC-F (*Gucy2f*), a close relative of GC-E. Deletion of *Gucy2f* has very little effect on rod and cone physiology and survival. However, a GC-E/GC-F double knockout (GCdko) phenotypically resembles human LCA-1 with extinguished ERGs and rod/cone degeneration. In GCdko rods, PDE6 and GCAPs are absent in outer segments. In contrast, GC-E<sup>-/-</sup> cones lack proteins of the entire phototransduction cascade. These results suggest that

GC-E may participate in transport of peripheral membrane proteins from the endoplasmic reticulum (ER) to the outer segments.

**Keywords** Membrane guanylate cyclase · Targeted deletions · Rod and cone photoreceptors · Photoreceptor membrane protein transport

## Abbreviations

ER	Endoplasmic reticulum
GC	Guanylate cyclase
GCdko	GC-E/GC-F double knockout
GCAP	Guanylate cyclase-activating protein
LCA	Leber congenital amaurosis
PDE6	cGMP-specific photoreceptor phosphodiesterase
TGN	<i>trans</i> -Golgi network
WT	Wild-type

## Soluble and membrane guanylate cyclases

Guanylate cyclases (GCs) synthesize cyclic GMP (cGMP), a secondary messenger in many pathways, in response to diverse signals, such as nitric oxide (NO), peptide ligands (hormones), and fluxes in intracellular Ca<sup>2+</sup> mediated by Ca<sup>2+</sup>-binding proteins ([Ca<sup>2+</sup>]<sub>i</sub>) [1, 2]. These signals use specific guanylate cyclase receptors and cofactors to initiate the conversion of cytosolic GTP to cGMP. Intracellular cGMP regulates cellular physiology by activating protein kinases, directly gating specific ion channels, or altering intracellular cyclic nucleotide concentrations through regulation of phosphodiesterases (PDEs). Guanylate cyclases are classified as either soluble or membrane (particulate),

S. Karan · J. M. Frederick · W. Baehr (✉)  
Department of Ophthalmology, John A. Moran Eye Center,  
University of Utah Health Science Center,  
65 Mario Capecchi Dr., Salt Lake City, UT 84132, USA  
e-mail: wbaehr@hsc.utah.edu

S. Karan · W. Baehr  
Department of Biology, University of Utah, Salt Lake City,  
UT 84112, USA

W. Baehr  
Department of Neurobiology and Anatomy, University of Utah  
Health Science Center, Salt Lake City, UT 84132, USA

based on both their cellular distribution and structural domains [2, 3]. Soluble guanylate cyclases are heterodimeric proteins consisting of  $\alpha$ - and  $\beta$ -subunits, and are activated by NO, another secondary messenger. Soluble guanylate cyclases are present in various cells in vertebrate retina, and maybe involved in signal transmission/modulation between cells [4]. A role in photoreceptor physiology was envisioned earlier for soluble GCs [5, 6], but no biochemical or genetic evidences are available for a role in modulation of cGMP in phototransduction. Based on phenotypes of photoreceptor GC double knockouts, a specific role for soluble GCs in phototransduction can safely be excluded.

Membrane GC isozymes (GC-A to GC-G, Table 1) exhibit highly conserved domain structures, an extracellular domain (ECD) which comprises a large part of the N-terminal part of the molecule, a single transmembrane (TM) region, an intracellular protein kinase-like homology domain (KHD), a dimerization (hinge) domain (DD), and a C-terminal catalytic domain (CAT). Based on their ligand specificities, membrane GCs have been subdivided into natriuretic peptide receptors (GC-A, GC-B), intestinal peptide-binding receptors (GC-C), olfactory uroguanylin- and guanylin-sensitive receptors (GC-D) and the so-called “orphan” receptors (GC-E/GC-F present in photoreceptors, and GC-G in testis). GC-E and GC-F have no known extracellular ligand (hence, the term “orphan”), but are stimulated by intracellular ligands, the GC-activating proteins (GCAPs). Thus, currently the only real “orphan” receptor is GC-G, a receptor of largely unknown distribution and function.

Non-photoreceptor GCs and cGMP-signaling have been reviewed extensively [1–3, 7]. However, within the last 10 years, the generation of membrane GC knockouts have led to important insights concerning their precise function

(recent review: [8]). The following paragraphs attempt to briefly summarize the phenotypes of non-photoreceptor membrane GC knockouts and of naturally occurring null alleles in human.

### Consequences of non-photoreceptor GC deletions


GC-A (natriuretic peptide receptor A, gene symbol *Npr1*, see Table 1) is expressed in the vasculature, heart, brain, testis, and other tissues, and is stimulated by atrial natriuretic peptides secreted by heart muscles. Mice lacking the *Npr1* gene, produced by placing a neo-cassette in exon 4 [9] (Fig. 1), mimic many of the features of hypertensive heart disease in human patients. *Npr1*<sup>−/−</sup> mice showed multiple phenotypes, including elevated blood pressure, salt-resistant hypertension, progressive cardiac hypertrophy, and sudden death, thereby demonstrating that GC-A is essential for the maintenance of normal blood pressure [9, 10].

GC-B (natriuretic peptide receptor B, gene *Npr2*) is expressed in many different tissues, and its function had been unclear until a knockout was generated [11]. In order to generate the knockouts, exons 3–7 encoding a portion of the ECD and the TM domain, were replaced by a neo-cassette [11] (Fig. 1). *Npr2*<sup>−/−</sup> mice showed a dramatic impairment of endochondral ossification and an attenuation of longitudinal vertebra or limb-bone growth [11]. Female *Npr2*<sup>−/−</sup> mice were infertile, but male mice were not, due to the failure of the female reproductive tract to develop. Null mutations in the *NPR2* gene in humans are associated with autosomal recessive skeletal dysplasia known as “acromesomelic dysplasia, type Maroteaux” (AMDM), characterized by reduced body height. In addition, heterozygous *NPR2* mutations were found to be associated with

**Table 1** Nomenclature of GC enzymes and genes

Mouse ( <i>Gene</i> )	Human ( <i>Gene</i> )	Aliases	Ligand	Tissue/cells	Comments
GC-A ( <i>Npr1</i> )	GCA, NPR-A ( <i>NPR1</i> )	Natriuretic Peptide Receptor A	ANP BNP	Vasculature	Regulates hypertension
GC-B ( <i>Npr2</i> )	GCB, NPR-B ( <i>NPR2</i> )	Natriuretic Peptide Receptor B	CNP	Brain, bone	Null alleles associated with dwarfism
GC-C ( <i>Gucy2c</i> )	GCC ( <i>GUCY2C</i> )		Guanylin uroguanylin enterotoxin	Intestines	Null allele insensitive to enterotoxins
GC-D ( <i>Gucy2d</i> )	GCD ( <i>GUCY2E</i> )	Olfactory neuroepithelia (ONE)-GC	Guanylin uroguanylin	Subset of Olfactory neurons	Pseudogene in primates
GC-E ( <i>Gucy2e</i> )	RetGC-1 ( <i>GUCY2D</i> )	ROS-GC1, GC1	GCAPs	Rods and cones, pineal	Null alleles associated with LCA-1
GC-F ( <i>Gucy2f</i> )	RetGC-2 ( <i>GUCY2F</i> )	ROS-GC2, GC2	GCAPs	Rods	Mutant alleles in cancer cells
GC-G ( <i>Gucy2g</i> )	GCG ( <i>GUCY2G</i> )		Unknown	Mouse testis, kidney	Pseudogene in human

**Fig. 1** Membrane guanylate cyclase knockout representations. *White boxes*—noncoding exons; *black boxes* or *vertical lines*—coding exons. *TM* transmembrane domain. Neo cassettes indicate strategies for knockout constructs. References for knockout mice are: *Npr1* [9]; *Npr2* [11]; *Gucy2c* [15]; *Gucy2d* [18]; *Gucy2e* [78]; *Gucy2f* [31]; *Gucy2g* [21]



short stature in humans [12]. Further, mutations in the *Npr2* gene are responsible for dwarfism, short limbs and tail in the *cn/cn* and *slw* (short-limbed dwarfism) mouse [13, 14].

Heat-stable enterotoxins activate GC-C (gene symbol *Gucy2c*) in intestines and increase levels of cGMP that activate downstream targets causing acute diarrhea. The *Gucy2c* deletion was generated by inserting a neo-cassette into exon 1; the knockout mice developed normally and were resistant to enterotoxin-induced diarrhea [15]. GC-C is the major guanylate cyclase in the mammalian intestines, which has been the focus of a link to colon cancer. However, *Gucy2c* null mice show reduced polyp formation and increased apoptosis, the opposite of proliferation [16].

A small subpopulation of olfactory neurons in the main olfactory epithelium express GC-D, together with CNGA3 (cGMP-gated channel  $\alpha$ -subunit first identified in cones) and cGMP-specific PDE2 [17, 18]. A *Gucy2d* knockout was generated by replacing exons 2 and 3 with a reporter cassette fused to a floxed neo gene [18] (Fig. 1). The knockout permitted the identification of uroguanylin and guanylin as extracellular ligands suggesting that GC-D expressing olfactory neurons may detect cues related to hunger and thirst [18, 19]. The “orphan receptor” GC-G is expressed in mouse testes [20] and kidneys [21], but its distribution in other tissues and possible function are largely unknown. A *Gucy2g* knockout suggested that in

kidneys, GC-G may promote apoptotic and inflammatory responses in ischemia-reperfusion-induced renal injury [21]. Interestingly, the primate *GUCY2E* gene (encoding GC-D) and the human *GUCY2G* gene (encoding GC-G) contain multiple inactivating sequence changes and are non-functional pseudogenes [22]. These data suggest that GC-D has been lost in primates, prior to the divergence of Old World monkeys from New World monkeys (more than 40 million years ago) [22].

These gene targeting experiments clearly demonstrate the power of gene knockouts in pinpointing the exact functions of membrane GCs in vivo. Here, we will discuss the photoreceptor GCs (GC-E and GC-F) expressed in rods and cones, with particular focus on gene knockouts and their consequences for photoreceptor physiology, membrane protein transport, and photoreceptor survival.

### Photoreceptor guanylate cyclases (GC-E and GC-F)

Membrane GCs are key components responsible for the production of cGMP, an essential secondary messenger of phototransduction. During phototransduction, cGMP is rapidly hydrolyzed by a membrane-associated PDE6 located on rod outer segment disk membranes, i.e., coin-like stacks of membranes unconnected to the plasma

membrane. Disappearance of the “messenger” closes cGMP-gated cation channels residing in the cell membrane, effectively hyperpolarizing the cell. In the dark-adapted photoreceptor, the basal GC activity (inhibited by  $\text{Ca}^{2+}$ -bound GC-activating proteins, termed GCAPs) is balanced by the low basal activity of PDE6, adjusting cytoplasmic cGMP to about 1–10  $\mu\text{M}$ . When the intracellular calcium ions decrease from about 500 nM (dark) to <50 nM in light, the GCs are activated by the  $\text{Ca}^{2+}$ -free GCAPs, increasing the catalytic GC activity by about 10-fold. If all the components of phototransduction ( $\text{R}^*$ ,  $\text{G}^*$ ,  $\text{PDE}^*$ ) are “silenced” by returning to the inactive dark state, then GC stimulation leads to restoration of dark cGMP levels [23–26].

GC-E or GC-F each possesses the structural features characteristic of membrane GCs—a signal sequence preceding a large amino-terminal ECD, a single membrane-spanning region, a kinase-like homology domain, and a carboxy-terminal catalytic domain. While the nomenclature of the GCs in various species is somewhat confusing (Table 1), generally accepted abbreviations are retGC-1 and retGC-2 in human [27, 28], ROS-GC1 and ROS-GC2 in bovine [29, 30], and GC1 and GC2 in mouse [31]. Quantitative determination of GC-E and GC-F in outer segment preparations by mass spectrometry showed a ratio of GC-E/GC-F of 3.5/1 [32], but biochemical quantization showed a much higher ratio of GC-E/GC-F of 25/1 [33].


Human GC-E (retGC-1) [34] and human GC-F (retGC-2) [35] show regional localization in the photoreceptor cell bodies and inner segments by in situ hybridization with antisense probes [27, 36]. Human and monkey GC-E were associated with outer segments of rods and cones, but also to a lesser extent with IS and other cells [34, 37]. The mouse GC-E and GC-F polypeptides were shown to be present in rod outer segments, likely as homomers [38].

Mouse GC-E is prominently immunolocalized in outer segments of rods and cones [39], whereas GC-F is detectable only in outer segments of rods (Fig. 2, [31]). Neither GC-E nor GC-F is detected elsewhere in the mouse retina, at least under the conditions of Fig. 2 [31]. However, GC-E was also detected in the synaptic terminals of bovine photoreceptors [40, 41]. Outside the retina, GC-E was detected in the pineal gland [38, 42], an organ developmentally related to the retina, in the olfactory bulb [43], spermatogenic cells of bovine testes [44], as well as the cochlear nerve and the organ of Corti [45]. Interestingly, GC-F was identified in several human cancer cell lines [46] where its function is unknown.

None of the membrane GCs has an affinity for  $\text{Ca}^{2+}$  ions, or conventional  $\text{Ca}^{2+}$ -binding motifs. The activities of photoreceptor cyclases, however, are  $\text{Ca}^{2+}$ -sensitive, a sensitivity that is mediated by guanylate cyclase-activating proteins (GCAPs) at low free  $\text{Ca}^{2+}$ , as well as by S100B and neurocalcin at high free  $\text{Ca}^{2+}$ . GCAPs are  $\text{Ca}^{2+}$ -binding proteins with three high-affinity  $\text{Ca}^{2+}$ -binding sites (EF hands) stimulating GC-E and GC-F by binding to cytoplasmic sites [47–50]. S100B and neurocalcin stimulate GC-E at 17–20  $\mu\text{M}$   $\text{Ca}^{2+}$  [51, 52], a concentration which is too high for a physiological role in outer segments. The peptide ligands of non-photoreceptor GC receptors failed to activate either GC-E or GC-F [53]. GC enzymology, biochemistry, and history of cloning, are reviewed in detail in [29].

### ***GUCY2D* null alleles in human: Leber congenital amaurosis type 1**

Leber congenital amaurosis type 1 (LCA-1), an autosomal recessive, early onset dystrophy defined as a “congenital



**Fig. 2** Distribution of GC1 and GC2 in WT (a, c) and GC knockout (b, d) mouse retina. Cryosections a, b were probed with anti-GC1, and c, d with anti-GC2 antibodies. GC1 is present in rod (intense green staining in the OS area) and cone (arrows) outer segments (a).

GC2 is only detectable in WT rod outer segments (c). GC1 and GC2 are undetectable in the ONL or OPL (synaptic terminals). Faint immunofluorescence in the INL is nonspecific (Color figure online)



**Table 2** GC null alleles in human LCA-1 patients

Exon	Amino acid change	Domain	Enzyme activity	References
2	M1I	ECD	Reduced	[93]
2	W21R	ECD	Yes	[93]
2	L41F	ECD	Yes	[57]
2	C105Y	ECD		[61]
2	H109P	ECD		[93]
2	N129K	ECD	Yes	[57]
3	R313C	ECD	Yes	[57]
3	L325P	ECD		[61]
4	Y351C	ECD		[94]
4	<i>S448ter</i>	ECD	No	[93]
7	R540C	KHD		[93]
8	F565S	KHD	Reduced	[95]
10	R660Q	KHD		[60]
12	R768W	DIM		[94]
13	<i>Q855ter</i>	DIM	No	[96]
13	P858S	DIM		[61]
15	A934P	CAT		[72]
15	L954P	CAT		[61]
15	R976L	CAT		[93]
16	R995W	CAT	No	[93]
16	M1009L	CAT	No	[72]
17	H1019P	CAT	No	[93]
17	<i>Q1036ter</i>	CAT	No	[54]
18	<i>R1040ter</i>	CAT	No	[72]

Italicized mutations produce truncations

*ECD* Extracellular domain, *KHD* kinase homology domain, *DIM* dimerization domain, *CAT* catalytic domain

stationary cone–rod dystrophy with high hypermetropia, panretinal degeneration and highly reduced visual acuity,” is a rare disease (prevalence approximately 1:100,000) [54]. Null mutations in the human GC-E gene (*GUCY2D*, on chromosome 17) are relatively frequent and distributed throughout the ECD, KHD, DIM, and CAT domains (references in Table 2). Mutations in the GC-F gene (*GUCY2F*, on Xq22) [55] have not been linked to a retina disease phenotype. Since *GUCY2F* gene is located on X, only males would be affected by disease-causing mutations. Interestingly, several missense mutations in the peptide leader sequence, the KHD and CAT domains of GC-F have been identified in panels of colon, lung and breast cancer [56]. LCA is one of the most severe forms of inherited retinal dystrophies, accounting for approximately 5% of all inherited retinal dystrophies [57, 58]. Children with LCA have greatly impaired vision and extinguished electroretinograms (ERG), despite seemingly normal fundi at birth or during the first months of life [59].

Immunofluorescent images of human *GUCY2D*-LCA retinas showed that all cones and rods lacked outer segments, as evidenced by the absence of labeling with anti-opsins and anti-rds/peripherin [60].


A number of nonsense and frameshift mutations in the *GUCY2D* gene have been identified in LCA-1 patients (Table 2). Missense mutations in the catalytic domain (e.g., R976L, R995W, M1009L, H1019P and Q1036Z) result in the total abolition of cyclase activity. Conversely, some missense mutations in the extracellular domain (e.g., W21R, L41F, N129, R313C) do not affect cyclase activity in vitro. Truncation mutations (e.g., S448ter) remove the CAT domain, and prevent synthesis of cGMP. Some missense mutations in the extracellular domain (C105Y, L325P) have resulted in 50% decreased cyclase activity. L325 is highly conserved and C105 has been suggested to participate in intramolecular disulfide bond formation and oligomerization of the protein. Thus, GC mutations may alter the enzyme structure and stability [61], and may affect post-biosynthetic folding or retrograde transport of peripheral membrane-associated proteins along microtubules to the cilium (Fig. 8).

### Cone–rod dystrophy (CORD6)

Dominant cone–rod dystrophy (CORD6) also maps to the GC1 locus [62, 63]. Mutations causing dominant cone–rod dystrophy are restricted to the dimerization domain. Cone–rod dystrophy is a progressive disorder, characterized by the initial degeneration of cone photoreceptor cells, causing early loss of visual acuity and color vision, followed by the degeneration of rods leading to progressive night blindness and peripheral visual field loss [62]. Some of the important missense mutations of the dimerization domain are E837D, R838A, R838H, R838C, T839M [64–66]. The three disease mutations at residue 838 are non-equivalent. They exhibit GC activity equal or superior to WT GC at low free  $[Ca]_{free}$  in the order  $R838C < R838H < R838A$  and showed a higher affinity for GCAP1 than WT GC [66]. Interestingly, when expressed in HEK 293 cells, (R838C)GC1 showed a decreased response to GCAP2 suggesting distinct binding sites for the GCAPs [67]. Structural studies have shown that the residue at position at 838 is a key residue that determines the extent of coiled-coil structure responsible for maintaining the retGC1 dimer. Molecular dynamics-based modeling of RetGC1 coiled-coil suggests that Arg838 forms salt bridges which maintain the configuration of the helices producing repulsion between the polypeptide chains. This causes the coils to spread apart as they extend towards the catalytic domain which then determines the dimerization [68].

### A naturally occurring GC-E knockout: the *retinal degeneration (rd)* chicken

The *rd* (*GUCY1\*B*) chicken is a naturally occurring blind mutant discovered in a Rhode Island Red flock about 30 years ago (reviewed in [69]). Photoreceptors in predegenerate *rd/rd* retina appeared normal at the ultrastructural level. Degeneration of rods and cones begins approximately 7 days post-hatch, and progresses relatively slowly over many months. The *rd* chicken carries a null mutation in the gene encoding photoreceptor GC-E exhibiting a disease phenotype resembling LCA [70, 71]. The gene defect is described best as a deletion/insertion event in which a gene fragment containing exons 4–7 was replaced by an inverted portion of exon 9 without interruption of the open reading frame (Fig. 3). Deletion of exon 5, encoding the transmembrane region, is predicted to result in a soluble polypeptide that, if stably expressed, would fail to traffic to the outer segments. Consistent with the deletion of the GC-E gene, levels of cGMP in the mutant retina were very low, unable to sustain phototransduction, presumably leading to permanent closure of the cGMP-gated cation channels and elimination of the dark current (constitutive hyperpolarization). In chicken, as in human LCA type 1 (*Gucy2e* null alleles) [72], only GC-E appears to contribute to the pool of cGMP essential to support phototransduction. It is unclear whether a GC-F ortholog is expressed in chicken retina. A lentivirus-based gene transfer vector carrying a bovine *GUCY2D* cDNA, injected into early-stage *GUCY1\*B* embryos, rescued retinal degeneration to a large part demonstrating convincingly that the *rd* chicken phenotype is caused by deletion of GC-E [73].



**Fig. 3** The *rd* chicken gene defect. *Top*, diagram of the mutant gene in which exons 4–7 are replaced by an inverted fragment of exon 9 (shaded). *Bottom*, predicted WT gene representing exons 3–8. Exon 7a is an additional exon not present in mammalian GC-E genes. Adapted from [70]

Transcription of several genes involved in the phototransduction cascade, including iodopsin and cone PDE $\alpha$ , was shown to be normal [74, 75]. However, protein levels of GCAP1 were reduced by more than 90% in predegenerate *rd/rd* retinas compared to age-matched control retinas, while GCAP2 levels were less affected [76], similarly as seen in GC-E/GC-F double knockouts (see below, and [31]). A regulatory mechanism at the gene level was excluded because transcript levels of both GCAP genes were near normal in the predegenerate *rd* retina [77]. The observed downregulation of GCAP1 was likely caused by mistrafficking of membrane-associated proteins which relies in part on the presence of GC-E (see below).


### Photoreceptor GC knockouts

The genes encoding GC-E (*Gucy2e*, on mouse chromosome 11) and the GC-F (*Gucy2f*, on the X chromosome) genes [55] are closely related in structure, each consisting of 19 exons and featuring an untranslated exon 1. The GC-E knockout mouse was generated by the group of David Garbers at UT Southwestern [78]. His group deleted the *Gucy2e* gene by targeted recombination replacing exon 5, encoding the transmembrane domain, with a neo cassette. The neo cassette effectively disrupted translation in all reading frames, preserving the extracellular domain, but deleting cytoplasmic regions including the catalytic domain so no functional cyclase can be formed. It is unknown whether an extracellular polypeptide (encoded by exons 1–4) is formed in the *Gucy2e*<sup>−/−</sup> photoreceptor. At 1 month of age, mouse cone responses were undetectable while rods continued to function albeit with reduced a- and b-wave amplitudes suggesting a substantial loss of rod function (Fig. 4, [78]). The phenotype of the *Gucy2e*<sup>−/−</sup> mice therefore resembled a recessive cone dystrophy [78, 79] while in human patients, *GUCY2E* null alleles are associated with rod/cone dystrophy (LCA-1).

In *Gucy2e*<sup>−/−</sup> mouse rods, a second cyclase, GC-F, is obviously able to substitute to some extent for the loss of GC-E. To identify the precise function of GC-F in rods, we generated *Gucy2f*<sup>−/−</sup> by replacing exon 2 of the *Gucy2f* gene containing the translation start codon and the leader peptide with a neo cassette. Therefore, a functional GC-F cannot be expressed. The phenotype of *Gucy2f*<sup>−/−</sup> mice was very similar to WT mice (Table 3), but showed slower recovery from intense illumination (Fig. 3 in [31]). In scotopic ERG analyses, the *Gucy2f*<sup>−/−</sup> a-wave amplitude reflecting rod photoreceptor function was identical to WT, while the b-wave amplitude was reduced (Fig. 4). The photopic ERG responses in the *Gucy2f*<sup>−/−</sup> mice were very similar to those of WT mice. We proceeded to generate GC double knockout mice (GCdko) by breeding *Gucy2f*<sup>−/−</sup>



**Fig. 4** Scotopic and photopic ERGs of WT, *Gucy2e*<sup>-/-</sup>, *Gucy2f*<sup>-/-</sup> and GCdko mice at 2.8 log cds m<sup>-2</sup>. Note lack of response in GCdko scotopic ERGs and in *Gucy2e*<sup>-/-</sup> and GCdko photopic ERGs



**Table 3** Effect of GC-E deletion, GC-F deletion, and deletion of both GCs on photoreceptor physiology and outer segment constituents

Rod physiology	Cone physiology	ROS proteins	COS proteins
<i>Gucy2e</i> <sup>-/-</sup>			
Functional; ERG responses attenuated	No response; cones degenerate slowly (autosomal recessive cone dystrophy)	GCAP1, GCAP2 downregulated; GC-F levels not affected	Cone T $\alpha$ , cone T $\gamma$ , GRK1, cone PDE $\alpha'$ undetectable; COS unstable; very little cone pigment present
No degeneration			
<i>Gucy2f</i> <sup>-/-</sup>			
Slower recovery from intense illumination	Normal	Normal	Normal GC-E levels not affected
GCdko			
No response	No response Cones degenerate (recessive LCA)	PDE $\alpha$ , PDE $\beta$ , PDE $\gamma$ , GCAP1, GCAP2 undetectable; T $\alpha$ , T $\gamma$ , arrestin, rhodopsin not affected	Cone T $\alpha$ , cone T $\gamma$ , GRK1, cone PDE $\alpha'$ undetectable; very little cone pigment present
Rods degenerate (recessive LCA)			

mice with *Gucy2e*<sup>-/-</sup> mice. Both full-field ERG (Fig. 4) and rod single-cell recordings (Fig. 4 in [31]) showed that GCdko photoreceptors were insensitive to light.

### Consequence of GC-E /GC-F double knockout (GCdko) in rods


At 1 month, the GCdko photoreceptors are relatively stable but outer segments are slightly reduced in length. Immunolabeling patterns of rhodopsin (Fig. 5b), arrestin (Fig. 5d), T $\alpha$  (Fig. 5f) and T $\gamma$  (Fig. 5h) were apparently normal and similar to WT, but the labeling pattern of rod outer segments was “patchy”, consistent with alternating bands of tubules and dense membranes observed by electron microscopy [31]. However, expression levels of several phototransduction polypeptides were reduced severely, including those of PDE6 subunits (Fig. 5j), GCAP1 (Fig. 5l) and GCAP2 (Fig. 5n). GCAPs activate GCs in low free Ca<sup>2+</sup>, are closely associated with GCs, and may be unstable in the absence of GCs. A similar downregulation of GCAP1 (but not GCAP2) was observed in the *rd* chicken carrying a *GUCY2D* null allele [77]. The diminished expression levels of GCAP1 and GCAP2 were

confined mostly to the inner segments where biosynthesis occurs (Fig. 5l, n).

We were most surprised by absence of PDE6 $\alpha\beta\gamma$  subunits in GCdko ROS (Fig. 5j). PDE6 is a peripheral membrane protein stably attached to the disk membrane surface by farnesyl and geranylgeranyl anchors. PDE $\alpha$  is known to be farnesylated and PDE $\beta$  geranylgeranylated at the C-terminal cysteines [80, 81]. PDE $\alpha\beta$  subunits are prenylated by cytosolic prenyltransferases and then dock to the ER. Once docked, they are insoluble and most likely require vesicular transport to reach their outer segment destination. In contrast to PDE6, transmembrane GCs are synthesized by ER-associated ribosomes, integrate in the ER membrane, and follow the secretory pathway from the ER to the plasma membrane, mediated by IFT through the connecting cilium to the outer segments.

The posttranslational downregulation of PDE and GCAPs, observed by immunoblots [31], and the absence of photoreceptor immunoreactivity for GC-E and GC-F in GCdko retinas suggested a role in protein trafficking, comparable to that of rhodopsin-bearing vesicles [82, 83]. But in this case involving association of PDE6 and GCAPs with GC-bearing vesicles. Based on GC knockout results, enforced by PrBP/ $\delta$  knockouts [84, 85], we developed a

**Fig. 5** Distribution of phototransduction polypeptides in WT (**a,c,e,g,i,k,m,o**) and GCdko (**b,d,f,h,j,l,n,p**) mutant outer and inner segments. Antigens targeted by antibodies (green) are indicated *top right* in the GCdko panels. Nuclei of the ONL are counterstained with propidium iodide (red). The “patchy” appearance of ROS indicates disorganization of the membrane structure [31] (Color figure online)




model describing transport of GCs in vesicles to which PDE subunits are attached peripherally (Fig. 6a).

Like other integral membrane proteins, GC-bearing vesicles are thought to traffic from the ER to the Golgi and emerge from the trans-Golgi network (TGN; Fig. 6a). We previously observed that when the prenyl-binding protein PrBP/ $\delta$  was deleted in rods, traffic of GRK1 and PDE6 to the outer segments was impeded while GC-E trafficked normally. We therefore propose that the GC-bearing vesicles are loaded with PDE6 subunits, aided by PrBP/ $\delta$  (and possibly additional prenyl binding proteins) that transfer PDE6 from the ER to vesicles. For simplicity, in Fig. 7 GC vesicles are charged with PDE post-TGN, but a pre-Golgi

loading of the vesicles cannot be excluded. Results with either GC-E or GC-F knockouts indicate that one GC is sufficient for PDE6 (and GCAP) transport. GC-bearing vesicles loaded with peripheral membrane proteins continue retrograde transport along microtubules toward the minus end, powered by molecular motors (e.g., cytoplasmic dynein). Vesicles likely fuse with the plasma membrane at the base of the cilium where cargo is assembled for intraflagellar transport through the cilium. Heterotrimeric kinesin-2 was suggested to be the molecular motor for IFT in rods based on the observation that rhodopsin accumulated in the distal inner segment when KIF3A, the motor subunit of kinesin-2, was deleted. Intriguingly, GC-E was

**Fig. 6** Putative model of GC and rhodopsin transport following synthesis at the ER.  
**a** WT rod photoreceptor.  
**b** GCdko photoreceptor.  
**a** Integral membrane proteins traffic from the ER to the Golgi/TGN by retrograde transport. Vesicles emerge from the TGN and are charged with the peripheral membrane proteins PDE and GCAPs. Retrograde transport continues to the base of the cilium. **b** Since GC-E and GC-F are not produced, GCAPs and PDE cannot traffic, and are degraded. Rhodopsin, transducin, and GRK1 transport is not affected. For details, see text




suggested to be mislocalized in mice expressing a hypomorphic mutation in IFT88, a protein associated with the axoneme and IFT complexes, but the fate of PDE6 was not investigated [86, 87].

Knockout of both GCs (Fig. 6b) prevents formation of GC-bearing vesicles at the ER and coordinately prevents transport of PDE6 and GCAPs. Since GC-deletion does not

affect rhodopsin and transducin transport to the rod outer segment. GC transport and rhodopsin transport may occur independently. PDE docked and retained in the ER due to the absence of GC carriers, likely presents prolonged cellular stress leading to PDE6 degradation by eliciting the unfolded protein response (UPR) with removal by the proteasome/ubiquitination pathway.



**Fig. 7** Distribution of cone phototransduction polypeptides in WT (a,c,e,g,i,k,m,o) and GC1-/- (b,d,f,h,j,l,n,p) mutant outer and inner segments. Antigens targeted by antibodies (green) are indicated top right in the GCdko panels. Arrows (b, d) indicate disconnected membrane packets containing pigments. Visual pigments, cone transducin, cone PDE, GCAP1 and GRK1 are undetectable in mutant COS (Color figure online)




### Effect of GC-E deletion in cones

It was shown that *Gucy2e*<sup>-/-</sup> cone photoreceptors degenerate gradually [78, 88]. Figure 7i shows that at 1 month of age, *Gucy2e*<sup>-/-</sup> cone outer segments are still present and of near-normal length, but S-opsin and M/L-opsin were present at reduced levels and mislocalized throughout the mutant cone cell. Further, acylated cone Tα (Fig. 7f), farnesylated cone Tγ (Fig. 7h), geranylgeranylated cone PDEα' (Fig. 7j), and farnesylated GRK1 (Fig. 7p) were undetectable in *Gucy2e*<sup>-/-</sup> COSs. These results show that essential cone phototransduction proteins are either absent or below the threshold of detection in *Gucy2e*<sup>-/-</sup> COSs. At 6 months of age, cone cell remnants, particularly synaptic

pedicles and somata, can be identified, but outer segments are absent. This finding is consistent with a postmortem study of an 11-year old LCA patient with a null mutation in the *GUCY2D* gene. The macula and peripheral retina of this patient revealed a substantial number of cones and rods, but both photoreceptor types lacked outer segments [60].

In contrast to GCdko rods where only GCAPs and PDE mistraffic, proteins of the entire phototransduction cascade are absent in *Gucy2e*<sup>-/-</sup> outer segments. Therefore the GC-E polypeptide, or its enzyme activity, must play a key role in retrograde trafficking of peripheral membrane proteins to the base of the cilium. An identical phenotype was observed in *Lrat*<sup>-/-</sup> and *Rpe65*<sup>-/-</sup> cones in which the retinoid cycle is blocked [89, 90]. In these mutants, the

**Fig. 8** Putative model for transport of peripheral membrane proteins in cones. **a** WT cone photoreceptor. **b** GC-E knockout cone photoreceptor. **a** Visual pigments and GC-E traffic from the ER to the Golgi/TGN by retrograde transport along microtubules. Vesicles emerge from the TGN and are charged with the peripheral membrane proteins PDE $\alpha'$ , cone T, GRK1, and GCAPs. Retrograde transport continues to the base of the cilium, followed by IFT to the outer segment. **b** GC-E is not produced. Peripheral proteins remain in the ER owing the absence of GC-E and are degraded



11-*cis*-retinal chromophore cannot be recycled and opsins are synthesized as apoproteins. Central *Lrat*<sup>-/-</sup> cones degenerate rapidly between 2 and 4 weeks postnatally, and immunocytochemistry showed that proteins of the entire phototransduction cascade, including GC-E, GCAPs, and R9AP, are absent in mutant cones. Thus, cone pigments require 11-*cis*-retinal as a cofactor to be competent for post-synthesis trafficking to the outer segments. The interdependence of GC-E, visual pigments, and peripheral membrane proteins supports the idea that a large cargo is assembled at the post-ER or post-TGN levels for co-transport and trafficking to cone outer segments. Further, we recently observed that cone-specific deletion of KIF3A, the obligatory motor subunit of heterotrimeric kinesin-2, prevents intraflagellar transport of visual pigments, GC1, R9AP, and peripheral membrane proteins through the cone cilium [91]. Thus, a common phenotype, i.e. absence of peripheral membrane protein transport, was replicated in four independent deletion models (*Gucy2e*<sup>-/-</sup>, cone *Kif3a*<sup>-/-</sup>, *Lrat*<sup>-/-</sup> and *Rpe65*<sup>-/-</sup> mice); this result argues for the coordinated insertion/assembly of several phototransduction-associated proteins into vesicular membrane for transport from the cone inner segment.

Our working model for cone transport is predicated on the interpretation that cone pigments and GC-E apparently traffic to the Golgi together (Fig. 8). That cone opsins and GC-E co-transport is supported by mislocalization of pigments in the GC-E knockout, and by mislocalization of GC-E in the LRAT/RPE65 knockouts. Emerging from the TGN, vesicles may be formed and loaded with peripheral membrane phototransduction polypeptides, which have been extracted from the ER by prenyl binding proteins. This mechanism is supported by a PrBP/ $\delta$  knockout in which cone PDE and GRK1 are unable to traffic (they are permanently docked to the ER), while cone transducin traffics unimpeded. In WT cones, GC-E and pigment-bearing vesicles are then transported along microtubules to the cilium, where IFT-cargo is assembled. Intraflagellar transport through the cone cilium is powered by the heterotrimeric kinesin-2, an anterograde motor moving towards the distal end of the outer segment (the plus end of microtubules) [91]. Knockout of the motor subunit KIF3A specifically in cones leads to a phenotype mimicking the GC-E and LRAT/RPE65 knockout, thereby preventing integral membrane proteins and membrane-associated proteins of the phototransduction cascade from trafficking to the cone outer segment.

## Conclusion

Deletion of one or both retinal GCs has very different effects on the physiology of rod and cone photoreceptors.

Particularly, the cone phenotype argues that GC-E is essential for trafficking several transmembrane and peripheral membrane proteins to the outer segment. We postulate that absence of GC-E in GUCY2D-LCA accounts for weakening of the outer segment membrane structure and secondarily, that accumulation of peripherally-associated proteins at the ER (mistrafficked) constitutes stress that initiates apoptotic pathways mediated by UPR [92].

**Acknowledgments** This study was supported by RO1 EY08123, RO1 EY019298, P30 EY014800 and a Center Grant from the Foundation Fighting Blindness.

## References

- Kuhn M (2003) Structure, regulation, and function of mammalian membrane guanylyl cyclase receptors, with a focus on guanylyl cyclase-A. *Circ Res* 93:700–709
- Lucas KA, Pitari GM, Kazeronian S, Ruiz-Stewart I, Park J, Schulz S, Chepenik KP, Waldman SA (2000) Guanylyl cyclases and signaling by cyclic GMP. *Pharmacol Rev* 52:375–414
- Potter LR (2005) Domain analysis of human transmembrane guanylyl cyclase receptors: implications for regulation. *Front Biosci* 10:1205–1220
- Sitaramayya A (2002) Soluble guanylate cyclases in the retina. *Mol Cell Biochem* 230:177–186
- Noll GN, Billek M, Pietruck C, Schmidt KF (1994) Inhibition of nitric oxide synthase alters light responses and dark voltage of amphibian photoreceptors. *Neuropharmacology* 33:1407–1412
- Cao L, Blute TA, Eldred WD (2000) Localization of heme oxygenase-2 and modulation of cGMP levels by carbon monoxide and/or nitric oxide in the retina. *Vis Neurosci* 17:319–329
- Garbers DL, Chrisman TD, Wiegand P, Katafuchi T, Albanesi JP, Bielinski V, Barylko B, Redfield MM, Burnett JC Jr (2006) Membrane guanylyl cyclase receptors: an update. *Trends Endocrinol Metab* 17:251–258
- Kuhn M (2005) Cardiac and intestinal natriuretic peptides: insights from genetically modified mice. *Peptides* 26:1078–1085
- Lopez MJ, Wong SK, Kishimoto I, Dubois S, Mach V, Friesen J, Garbers DL, Beuve A (1995) Salt-resistant hypertension in mice lacking the guanylyl cyclase-A receptor for atrial natriuretic peptide. *Nature* 378:65–68
- Oliver PM, Fox JE, Kim R, Rockman HA, Kim HS, Reddick RL, Pandey KN, Milgram SL, Smithies O, Maeda N (1997) Hypertension, cardiac hypertrophy, and sudden death in mice lacking natriuretic peptide receptor A. *Proc Natl Acad Sci USA* 94:14730–14735
- Tamura N, Doolittle LK, Hammer RE, Shelton JM, Richardson JA, Garbers DL (2004) Critical roles of the guanylyl cyclase B receptor in endochondral ossification and development of female reproductive organs. *Proc Natl Acad Sci USA* 101:17300–17305
- Olney RC, Bukulmez H, Bartels CF, Prickett TC, Espiner EA, Potter LR, Warman ML (2006) Heterozygous mutations in natriuretic peptide receptor-B (NPR2) are associated with short stature. *J Clin Endocrinol Metab* 91:1229–1232
- Sogawa C, Tsuji T, Shinkai Y, Katayama K, Kunieda T (2007) Short-limbed dwarfism: slw is a new allele of Npr2 causing chondrodysplasia. *J Hered* 98:575–580
- Tsuji T, Kunieda T (2005) A loss-of-function mutation in natriuretic peptide receptor 2 (Npr2) gene is responsible for disproportionate dwarfism in cn/cn mouse. *J Biol Chem* 280:14288–14292



15. Schulz S, Lopez MJ, Kuhn M, Garbers DL (1997) Disruption of the guanylyl cyclase-C gene leads to a paradoxical phenotype of viable but heat-stable enterotoxin-resistant mice. *J Clin Invest* 100:1590–1595
16. Mann EA, Steinbrecher KA, Stroup C, Witte DP, Cohen MB, Giannella RA (2005) Lack of guanylyl cyclase C, the receptor for *Escherichia coli* heat-stable enterotoxin, results in reduced polyp formation and increased apoptosis in the multiple intestinal neoplasia (Min) mouse model. *Int J Cancer* 116:500–505
17. Juilfs DM, Fulle HJ, Zhao AZ, Houslay MD, Garbers DL, Beavo JA (1997) A subset of olfactory neurons that selectively express cGMP-stimulated phosphodiesterase (PDE2) and guanylyl cyclase-D define a unique olfactory signal transduction pathway. *Proc Natl Acad Sci USA* 94:3388–3395
18. Leinders-Zufall T, Cockerham RE, Michalakakis S, Biel M, Garbers DL, Reed RR, Zufall F, Munger SD (2007) Contribution of the receptor guanylyl cyclase GC-D to chemosensory function in the olfactory epithelium. *Proc Natl Acad Sci USA* 104:14507–14512
19. Duda T, Sharma RK (2008) ONE-GC membrane guanylate cyclase, a trimodal odorant signal transducer. *Biochem Biophys Res Commun* 367:440–445
20. Kuhn M, Ng CK, Su YH, Kilic A, Mitko D, Bien-Ly N, Komuves LG, Yang RB (2004) Identification of an orphan guanylate cyclase receptor selectively expressed in mouse testis. *Biochem J* 379:385–393
21. Lin H, Cheng CF, Hou HH, Lian WS, Chao YC, Ciou YY, Djoko B, Tsai MT, Cheng CJ, Yang RB (2008) Disruption of guanylyl cyclase-G protects against acute renal injury. *J Am Soc Nephrol* 19:339–348
22. Young JM, Waters H, Dong C, Fulle HJ, Liman ER (2007) Degeneration of the olfactory guanylyl cyclase D gene during primate evolution. *PLoS ONE* 2:e884
23. Polans A, Baehr W, Palczewski K (1996) Turned on by  $\text{Ca}^{2+}$ ! The physiology and pathology of  $\text{Ca}^{2+}$  binding proteins in the retina. *Trends Neurosci* 19:547–554
24. Wensel TG (2008) Signal transducing membrane complexes of photoreceptor outer segments. *Vis Res* 48:2052–2061
25. Lamb TD, Pugh EN Jr (2006) Phototransduction, dark adaptation, and rhodopsin regeneration the proctor lecture. *Invest Ophthalmol Vis Sci* 47:5138–5152
26. Arshavsky VY, Lamb TD, Pugh EN Jr (2002) G proteins and phototransduction. *Annu Rev Physiol* 64:153–187
27. Shyjan AW, de Sauvage FJ, Gillett NA, Goeddel DV, Lowe DG (1992) Molecular cloning of a retina-specific membrane guanylyl cyclase. *Neuron* 9:727–737
28. Garbers DL, Lowe DG (1994) Guanylyl cyclase receptors. *J Biol Chem* 269:30741–30744
29. Pugh EN Jr, Duda T, Sitaramayya A, Sharma RK (1997) Photoreceptor guanylate cyclases: a review. *Biosci Rep* 17:429–473
30. Koch KW, Duda T, Sharma RK (2002) Photoreceptor specific guanylate cyclases in vertebrate phototransduction. *Mol Cell Biochem* 230:97–106
31. Baehr W, Karan S, Maeda T, Luo DG, Li S, Bronson JD, Watt CB, Yau KW, Frederick JM, Palczewski K (2007) The function of guanylate cyclase 1 and guanylate cyclase 2 in rod and cone photoreceptors. *J Biol Chem* 282:8837–8847
32. Liu Q, Tan G, Levenkova N, Li T, Pugh EN Jr, Rux JJ, Speicher DW, Pierce EA (2007) The proteome of the mouse photoreceptor sensory cilium complex. *Mol Cell Proteomics* 6:1299–1317
33. Helten A, Saftel W, Koch KW (2007) Expression level and activity profile of membrane bound guanylate cyclase type 2 in rod outer segments. *J Neurochem* 103:1439–1446
34. Dizhoor AM, Lowe DG, Olshevskaya EV, Laura RP, Hurley JB (1994) The human photoreceptor membrane guanylyl cyclase, RetGC, is present in outer segments and is regulated by Calcium and a soluble activator. *Neuron* 12:1345–1352
35. Lowe DG, Dizhoor AM, Liu K, Gu Q, Spencer M, Laura R, Lu L, Hurley JB (1995) Cloning and expression of a second photoreceptor-specific membrane retina guanylyl cyclase (RetGC), RetGC-2. *Proc Natl Acad Sci USA* 92:5535–5539
36. Imanishi Y, Li N, Sowa ME, Lichtarge O, Wensel TG, Saperstein DA, Baehr W, Palczewski K (2002) Characterization of retinal guanylate cyclase-activating protein 3 (GCAP3) from zebrafish to man. *Eur J Neurosci* 15:63–78
37. Liu X, Seno K, Nishizawa Y, Hayashi F, Yamazaki A, Matsumoto H, Wakabayashi T, Usukura J (1994) Ultrastructural localization of retinal guanylate cyclase in human and monkey retinas. *Exp Eye Res* 59:761–768
38. Yang RB, Garbers DL (1997) Two eye guanylyl cyclases are expressed in the same photoreceptor cells and form homomers in preference to heteromers. *J Biol Chem* 272:13738–13742
39. Haire SE, Pang J, Boye SL, Sokal I, Craft CM, Palczewski K, Hauswirth WW, Semple-Rowland SL (2006) Light-driven cone arrestin translocation in cones of postnatal guanylate cyclase-1 knockout mouse retina treated with AAV-GC1. *Invest Ophthalmol Vis Sci* 47:3745–3753
40. Duda T, Koch KW, Venkataraman V, Lange C, Beyermann M, Sharma RK (2002)  $\text{Ca}^{2+}$  sensor S100beta-modulated sites of membrane guanylate cyclase in the photoreceptor-bipolar synapse. *EMBO J* 21:2547–2556
41. Venkataraman V, Duda T, Vardi N, Koch KW, Sharma RK (2003) Calcium-modulated guanylate cyclase transduction machinery in the photoreceptor-bipolar synaptic region. *Biochemistry* 42:5640–5648
42. Venkataraman V, Nagele R, Duda T, Sharma RK (2000) Rod outer segment membrane guanylate cyclase type 1-linked stimulatory and inhibitory calcium signaling systems in the pineal gland: biochemical, molecular, and immunohistochemical evidence. *Biochemistry* 39:6042–6052
43. Duda T, Venkataraman V, Krishnan A, Nagele RG, Sharma RK (2001) Negatively calcium-modulated membrane guanylate cyclase signaling system in the rat olfactory bulb. *Biochemistry* 40:4654–4662
44. Jankowska A, Burczynska B, Duda T, Warchol JB, Sharma RK (2007) Calcium-modulated rod outer segment membrane guanylate cyclase type 1 transduction machinery in the testes. *J Androl* 28:50–58
45. Seebacher T, Beitz E, Kumagami H, Wild K, Ruppersberg JP, Schultz JE (1999) Expression of membrane-bound and cytosolic guanylyl cyclases in the rat inner ear. *Hear Res* 127:95–102
46. Shao RX, Otsuka M, Kato N, Chang JH, Muroyama R, Taniguchi H, Moriyama M, Wang Y, Kawabe T, Omata M (2005) No mutations in the tyrosine kinases of human hepatic, pancreatic, and gastric cancer cell lines. *J Gastroenterol* 40:918
47. Palczewski K, Polans AS, Baehr W, Ames JB (2000) Calcium binding proteins in the retina: structure, function, and the etiology of human visual diseases. *Bioessays* 22:337–350
48. Sokal I, Li N, Verlinde CL, Haeseleer F, Baehr W, Palczewski K (2000)  $\text{Ca}^{2+}$ -binding proteins in the retina: from discovery to etiology of human disease. *Biochim Biophys Acta* 1498:233–251
49. Dizhoor AM, Hurley JB (1999) Regulation of photoreceptor membrane guanylyl cyclases by guanylyl cyclase activator proteins. *Methods* 19:521–531
50. Koch KW (2002) Target recognition of guanylate cyclase by guanylate cyclase-activating proteins. *Adv Exp Med Biol* 514:349–360
51. Duda T, Goraczniak RM, Sharma RK (1996) Molecular characterization of S100A1–S100B protein in retina and its activation mechanism of bovine photoreceptor guanylate cyclase. *Biochemistry* 35:6263–6266
52. Kumar VD, Vijay-Kumar S, Krishnan A, Duda T, Sharma RK (1999) A second calcium regulator of rod outer segment

- membrane guanylate cyclase, ROS-GC1: neurocalcin. *Biochemistry* 38:12614–12620
53. Goracznik RM, Duda T, Sitaramayya A, Sharma RK (1994) Structural and functional characterization of the rod outer segment membrane guanylate cyclase. *Biochem J* 302:455–461
  54. Perrault I, Hanein S, Gerber S, Lebail B, Vlajnik P, Barbet F, Ducroq D, Dufier JL, Munnich A, Kaplan J, Rozet JM (2005) A novel mutation in the GUCY2D gene responsible for an early onset severe RP different from the usual GUCY2D-LCA phenotype. *Hum Mutat* 25:222
  55. Yang RB, Fulle HJ, Garbers DL (1996) Chromosomal localization and genomic organization of genes encoding guanylyl cyclase receptors expressed in olfactory sensory neurons and retina. *Genomics* 31:367–372
  56. Wood LD, Calhoun ES, Silliman N, Ptak J, Szabo S, Powell SM, Riggins GJ, Wang TL, Yan H, Gazdar A, Kern SE, Pennacchio L, Kinzler KW, Vogelstein B, Velculescu VE (2006) Somatic mutations of GUCY2F, EPHA3, and NTRK3 in human cancers. *Hum Mutat* 27:1060–1061
  57. Perrault I, Rozet JM, Gerber S, Ghazi I, Ducroq D, Souied E, Leowski C, Bonnermaison M, Dufier JL, Munnich A, Kaplan J (2000) Spectrum of retGC1 mutations in Leber's congenital amaurosis. *Eur J Hum Genet* 8:578–582
  58. Perrault I, Rozet J-M, Calvas P, Gerber S, Camuzat A, Dollfus H, Chatelin S, Souied E, Ghazi I, Leowski C, Bonnermaison M, Le Paslier D, Frezal J, Dufier J-L, Pittler SJ, Munnich A, Kaplan J (1996) Retinal-specific guanylate cyclase gene mutations in Leber's congenital amaurosis. *Nat Genet* 14:461–464
  59. Yzer S, Leroy BP, De BE, de Ravel TJ, Zonneveld MN, Voeselek K, Kellner U, Ciriano JP, de Faber JT, Rohrschneider K, Roepman R, den Hollander AI, Cruysberg JR, Meire F, Casteels I, van Moll-Ramirez NG, Allikmets R, Van den Born LI, Cremers FP (2006) Microarray-based mutation detection and phenotypic characterization of patients with Leber congenital amaurosis. *Invest Ophthalmol Vis Sci* 47:1167–1176
  60. Milam AH, Barakat MR, Gupta N, Rose L, Aleman TS, Pianta MJ, Cideciyan AV, Sheffield VC, Stone EM, Jacobson SG (2003) Clinicopathologic effects of mutant GUCY2D in Leber congenital amaurosis. *Ophthalmology* 110:549–558
  61. Tucker CL, Ramamurthy V, Pina AL, Loyer M, Dharmaraj S, Li Y, Maumenee IH, Hurley JB, Koenekoop RK (2004) Functional analyses of mutant recessive GUCY2D alleles identified in Leber congenital amaurosis patients: protein domain comparisons and dominant negative effects. *Mol Vis* 10:297–303
  62. Kelsell RE, Gregory-Evans K, Payne AM, Perrault I, Kaplan J, Yang R-B, Garbers DL, Bird AC, Moore AT, Hunt DF (1998) Mutations in the retinal guanylate cyclase (RETGC-1) gene in dominant con-rod dystrophy. *Hum Mol Genet* 7:1179–1184
  63. Perrault I, Rozet JM, Gerber S, Kelsell RE, Souied E, Cabot A, Hunt DM, Munnich A, Kaplan J (1998) A retGC-1 mutation in autosomal dominant cone-rod dystrophy. *Am J Hum Genet* 63:651–654
  64. Payne AM, Morris AG, Downes SM, Johnson S, Bird AC, Moore AT, Bhattacharya SS, Hunt DM (2001) Clustering and frequency of mutations in the retinal guanylate cyclase (GUCY2D) gene in patients with dominant cone-rod dystrophies. *J Med Genet* 38:611–614
  65. Downes SM, Payne AM, Kelsell RE, Fitzke FW, Holder GE, Hunt DM, Moore AT, Bird AC (2001) Autosomal dominant cone-rod dystrophy with mutations in the guanylate cyclase 2D gene encoding retinal guanylate cyclase-1. *Arch Ophthalmol* 119:1667–1673
  66. Wilkie SE, Newbold RJ, Deery E, Walker CE, Stinton I, Ramamurthy V, Hurley JB, Bhattacharya SS, Warren MJ, Hunt DM (2000) Functional characterization of missense mutations at codon 838 in retinal guanylate cyclase correlates with disease severity in patients with autosomal dominant cone-rod dystrophy. *Hum Mol Genet* 9:3065–3073
  67. Tucker CL, Woodcock SC, Kelsell RE, Ramamurthy V, Hunt DM, Hurley JB (1999) Biochemical analysis of a dimerization domain mutation in RetGC-1 associated with dominant cone-rod dystrophy. *Proc Natl Acad Sci USA* 96:9039–9044
  68. Ramamurthy V, Tucker C, Wilkie SE, Daggett V, Hunt DM, Hurley JB (2001) Interactions within the coiled-coil domain of RetGC-1 guanylyl cyclase are optimized for regulation rather than for high affinity. *J Biol Chem* 276:26218–26229
  69. Semple-Rowland SL, Cheng KM (1999) rd and rc chickens carry the same GC1 null allele (GUCY1\*). *Exp Eye Res* 69:579–581
  70. Semple-Rowland SL, Lee NR, Van-Hooser JP, Palczewski K, Baehr W (1998) A null mutation in the photoreceptor guanylate cyclase gene causes the retinal degeneration chicken phenotype. *Proc Natl Acad Sci USA* 95:1271–1276
  71. Semple-Rowland SL, Lee NR (2000) Avian models of inherited retinal disease. *Methods Enzymol* 316:526–536
  72. Hanein S, Perrault I, Gerber S, Tanguy G, Barbet F, Ducroq D, Calvas P, Dollfus H, Hamel C, Loppönen T, Munier F, Santos L, Shalev S, Zafeiriou D, Dufier JL, Munnich A, Rozet JM, Kaplan J (2004) Leber congenital amaurosis: comprehensive survey of the genetic heterogeneity, refinement of the clinical definition, and genotype-phenotype correlations as a strategy for molecular diagnosis. *Hum Mutat* 23:306–317
  73. Williams ML, Coleman JE, Haire SE, Aleman TS, Cideciyan AV, Sokal I, Palczewski K, Jacobson SG, Semple-Rowland SL (2006) Lentiviral expression of retinal guanylate cyclase-1 (RetGC1) restores vision in an avian model of childhood blindness. *PLoS Med* 3:e201
  74. Semple-Rowland SL, Green DA (1994) Molecular characterization of the  $\alpha'$ -subunit of cone photoreceptor cGMP phosphodiesterase in normal and rd chicken. *Exp Eye Res* 59:365–372
  75. Semple-Rowland SL, Green DA (1994) Molecular and biochemical analyses of Iodopsin in rd chick retina. *Invest Ophthalmol Vis Sci* 35:2550–2557
  76. Semple-Rowland S, Gorczyca WA, Buczylo J, Ruiz CC, Subbaraya I, Helekar BS, Palczewski K, Baehr W (1996) Expression of GCAP1 and GCAP2 in the retinal degeneration (rd) chicken retina. *FEBS Lett* 385:47–52
  77. Semple-Rowland SL, Larkin P, Bronson JD, Nykamp K, Streit WJ, Baehr W (1999) Characterization of the chicken GCAP gene array and analyses of GCAP1, GCAP2, and GC1 gene expression in normal and rd chicken pineal. *Mol Vis* 5:14
  78. Yang RB, Robinson SW, Xiong WH, Yau KW, Birch DG, Garbers DL (1999) Disruption of a retinal guanylyl cyclase gene leads to cone-specific dystrophy and paradoxical rod behavior. *J Neurosci* 19:5889–5897
  79. Coleman JE, Zhang Y, Brown GA, Semple-Rowland SL (2004) Cone cell survival and downregulation of GCAP1 protein in the retinas of GC1 knockout mice. *Invest Ophthalmol Vis Sci* 45:3397–3403
  80. Anant JS, Ong OC, Xie H, Clarke S, O'Brien PJ, Fung BKK (1992) In vivo differential prenylation of retinal cyclic GMP phosphodiesterase catalytic subunits. *J Biol Chem* 267:687–690
  81. Qin N, Pittler SJ, Baehr W (1992) In vitro isoprenylation and membrane association of mouse rod photoreceptor cGMP phosphodiesterase  $\alpha$  and  $\beta$  subunits expressed in bacteria. *J Biol Chem* 267:8458–8463
  82. Tai AW, Chuang JZ, Bode C, Wolfrum U, Sung CH (1999) Rhodopsin's carboxy-terminal cytoplasmic tail acts as a membrane receptor for cytoplasmic dynein by binding to the dynein light chain Tctex-1. *Cell* 97:877–887
  83. Deretic D (2006) A role for rhodopsin in a signal transduction cascade that regulates membrane trafficking and photoreceptor polarity. *Vis Res* 46:4427–4433

84. Zhang H, Li S, Doan T, Rieke F, Detwiler PB, Frederick JM, Baehr W (2007) Deletion of PrBP/{delta} impedes transport of GRK1 and PDE6 catalytic subunits to photoreceptor outer segments. *Proc Natl Acad Sci USA* 104:8857–8862
85. Karan S, Zhang H, Li S, Frederick JM, Baehr W (2008) A model for transport of membrane-associated phototransduction polypeptides in rod and cone photoreceptor inner segments. *Vis Res* 48:442–452
86. Insinna C, Besharse JC (2008) Intraflagellar transport and the sensory outer segment of vertebrate photoreceptors. *Dev Dyn* 237:1982–1992
87. Pazour GJ, Baker SA, Deane JA, Cole DG, Dickert BL, Rosenbaum JL, Witman GB, Besharse JC (2002) The intraflagellar transport protein, IFT88, is essential for vertebrate photoreceptor assembly and maintenance. *J Cell Biol* 157:103–113
88. Coleman JE, Semple-Rowland SL (2005) GC1 deletion prevents light-dependent arrestin translocation in mouse cone photoreceptor cells. *Invest Ophthalmol Vis Sci* 46:12–16
89. Fan J, Rohrer B, Frederick JM, Baehr W, Crouch RK (2008) Rpe65<sup>-/-</sup> and Lrat<sup>-/-</sup> mice: comparable models of Leber congenital amaurosis. *Invest Ophthalmol Vis Sci* 49:2384–2389
90. Zhang H, Fan J, Li S, Karan S, Rohrer B, Palczewski K, Frederick JM, Crouch RK, Baehr W (2008) Trafficking of membrane-associated proteins to cone photoreceptor outer segments requires the chromophore 11-cis-retinal. *J Neurosci* 28:4008–4014
91. Avasthi P, Watt CB, Williams DS, Le YZ, Li S, Chen C-K, Marc RE, Frederick JM, Baehr W (2009) Trafficking of membrane proteins to cone but not rod outer segments is dependent on heterotrimeric kinesin-II. *J Neurosci* 29:14287–14298
92. Lin JH, Li H, Yasumura D, Cohen HR, Zhang C, Panning B, Shokat KM, LaVail MM, Walter P (2007) IRE1 signaling affects cell fate during the unfolded protein response. *Science* 318:944–949
93. Rozet JM, Perrault I, Gerber S, Hanein S, Barbet F, Ducroq D, Souied E, Munnich A, Kaplan J (2001) Complete abolition of the retinal-specific guanylyl cyclase (retGC-1) catalytic ability consistently leads to Leber congenital amaurosis (LCA). *Invest Ophthalmol Vis Sci* 42:1190–1192
94. Lotery AJ, Namperumalsamy P, Jacobson SG, Weleber RG, Fishman GA, Musarella MA, Hoyt CS, Heon E, Levin A, Jan J, Lam B, Carr RE, Franklin A, Radha S, Andorf JL, Sheffield VC, Stone EM (2000) Mutation analysis of 3 genes in patients with Leber congenital amaurosis. *Arch Ophthalmol* 118:538–543
95. Duda T, Venkataraman V, Goraczniak R, Lange C, Koch KW, Sharma RK (1999) Functional consequences of a rod outer segment membrane guanylate cyclase (ROS-GC1) gene mutation linked with Leber's congenital amaurosis. *Biochemistry* 38:509–515
96. El-Shanti H, Al-Salem M, El-Najjar M, Ajlouni K, Beck J, Sheffield VC, Stone EM (1999) A nonsense mutation in the retinal specific guanylate cyclase gene is the cause of Leber congenital amaurosis in a large inbred kindred from Jordan. *J Med Genet* 36:862–865