

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/26699835>

^1H , ^{13}C and ^{15}N resonance assignment of phage T4 endoribonuclease RegB

ARTICLE in BIOMOLECULAR NMR ASSIGNMENTS · AUGUST 2007

Impact Factor: 0.76 · DOI: 10.1007/s12104-007-9021-4 · Source: PubMed

READS

13

3 AUTHORS:



Benoit Odaert

French National Centre for Scientific Resea...

25 PUBLICATIONS 355 CITATIONS

SEE PROFILE



Fakhri Saida

General Atomics, United States

15 PUBLICATIONS 118 CITATIONS

SEE PROFILE



François Bontems

French National Centre for Scientific Resea...

52 PUBLICATIONS 1,475 CITATIONS

SEE PROFILE

^1H , ^{13}C and ^{15}N resonance assignment of phage T4 endoribonuclease RegB

Benoît Odaert · Fakhri Saïda · François Bontems

Received: 11 March 2007 / Accepted: 30 April 2007 / Published online: 4 July 2007
© Springer Science+Business Media B.V. 2007

Abstract RegB is involved in the control of the phage T4 life cycle. It inactivates the phage early mRNAs when their translation is no more required. We determined its structure and identified residues involved in substrate binding. For this, all backbone and 90% of side-chain resonance frequencies were assigned.

Keywords RegB · mRNA inactivation · Translation initiation · Toxin-antitoxin modules · Ribosomal S1 protein

Biological context

Inactivation and degradation of messenger RNAs constitute a key process in the regulation of gene expression. They involve two different classes of enzymes: the exoribonucleases and the endoribonucleases. While exoribonucleases are general regulators that processively degrade RNAs, most endoribonucleases act as specific regulators by selecting their targets and cleaving them at specific internal sites. The molecular mechanism of such enzymes remains poorly understood.

The ribonuclease RegB (18 kDa) is encoded by the bacteriophage T4. It is involved in the timing of the phage life cycle through the inactivation of early messenger RNAs produced after viral infection (Sanson et al. 2000). RegB, indeed, preferentially cleaves in the middle of the GGAG sequence found in bacterial translation initiation sites (Jayasena et al. 1996; Uzan 2001). RNA inactivation by RegB may involve the ribosomal complex as suggested by the 10- to 100-fold enhancement observed *in vitro* in the presence of the ribosomal protein S1, whose role is precisely to allow the recognition of the translation initiation region by the ribosome (Ruckman et al. 1994). Stimulation of RegB activity by S1 depends strongly on the sequence located 3' to the GGAG motif, suggesting a crucial role in the selection of RegB substrates (Durand et al. 2006). An *in vitro* study has also indicated that the presence of a stem-loop structure around the GGAG motif could favor RNA cleavage (Lebars et al. 2001).

The absence of sequence similarities to any known protein and the particular properties of RegB have motivated us to decipher its three-dimensional structure and to identify residues involved in the substrate recognition and/or processing. Our results have shown that, despite the absence of any sequence homology, RegB shares structural and functional similarities with the bacterial sequence-specific endoribonucleases YoeB and RelE (Odaert et al. 2007), involved in the translated mRNA inactivation when a rapid change of the bacterial metabolism is required (Codon 2006). To achieve this goal, the ^1H , ^{13}C and ^{15}N resonances of the H48A point mutant of RegB have been assigned using multidimensional heteronuclear NMR experiments. The RegB H48A mutation was necessary to over-express the protein under an inactive, non-cytotoxic but correctly folded form (Saïda et al. 2004).

B. Odaert · F. Saïda · F. Bontems (✉)
ICSN-RMN, Ecole-Polytechnique (équipe CNRS), Palaiseau
91128, France
e-mail: Francois.Bontems@polytechnique.fr

Present Address:

B. Odaert
Institut Européen de Chimie et Biologie, Pessac 33607,
France

Present Address:

F. Saïda
Center for Molecular Genetics, University of California, San
Diego, La Jolla, CA 92093-0634, USA

Methods and experiments

Expression and purification were previously described (Saïda et al. 2004). ^{15}N and ^{15}N , ^{13}C -labeled RegB H48A proteins were expressed in the M9 minimal medium supplemented with ^{15}N ammonium chloride (1 g.l $^{-1}$) and ^{13}C glucose (2 g.l $^{-1}$). NMR samples contained 0.7–1 mM of labeled protein in 50 mM sodium citrate (pH 6.0), 300 mM NaCl, 0.1 mM AESBF protease inhibitor, 2 mM dithiothreitol and 5:95% D $_2$ O/H $_2$ O or 100% D $_2$ O.

NMR data were recorded at 32 °C on a Bruker DRX600 spectrometer equipped with a triple resonance Z-gradient cryoprobe. Data were processed on the Bruker station with XWINNMR 3.0 and analyzed with the Sparky software (University of California San Francisco, Thomas L. Goddard). Backbone assignment was achieved with the 3D triple resonance experiments: HNCA/HN(CO)CA, HNCO/HN(CA)CO, HNHA. Side-chain resonance assignment was achieved with the following experiments: HNCACB, CBCA(CO)NH, HBHA(CBCACO)NH, aliphatic HCCH-TOCSY, ^{15}N -edited TOCSY-HSQC, aromatic ^{13}C -edited HSQC, aromatic ^{13}C -edited NOESY-HSQC and 2D TOCSY/NOESY (60 and 120 ms mixing time) in D $_2$ O. The assignment was confirmed and completed with ^{15}N -edited NOESY-HSQC (60 and 120 ms mixing time) and aliphatic ^{13}C -edited HSQC-NOESY (60 ms mixing time) in H $_2$ O and D $_2$ O.

Extent of assignments and data deposition

The structure determination was achieved by using the INCA software (Savarin et al. 2001) that uses an iterative procedure to simultaneously build the structure and assign the nOe cross-peaks. A critical point of the method is to dispose of precise frequency assignments perfectly coherent with the NOESY experiments used for the reconstruction. We mainly used three NOESY spectra, a ^{15}N -NOESY-HSQC, a ^{13}C -NOESY-HSQC and a two-dimensional NOESY spectrum recorded in D $_2$ O. Accordingly, we determined (and deposited; BMRB accession number 15127) three chemical shift lists corresponding to the frequencies observed in these three spectra. Several assignments derived from scalar experiments (such as HCCH-TOCSY) that could not be confirmed by the presence of unambiguous intra-residual cross-peaks in the NOESY spectra were not retained.

Nearly all N, H $^{\text{N}}$, C $^{\alpha}$, H $^{\alpha}$, C $^{\beta}$ and H $^{\beta}$ resonance frequencies were assigned in the ^{13}C - and ^{15}N -NOESY-HSQC experiments. All Ala, Cys, Asp, Glu, Gly, Leu, Asn, Gln, Ser, Thr, Val hydrogen frequencies were assigned in at least one of the two experiments. The missing resonances mainly concern the Lys (73 hydrogen assigned among 110), Arg (88 among 112), Pro (39 among 49) residues and the aromatic part of His17, Phe67, Phe93, Tyr99 and Phe124 that could not be observed.

Acknowledgments This work was supported by a national grant from the French Department of Scientific Research (Ministère de la Recherche) for Saïda F. and an ICSN postdoctoral fellowship for Odaert B.

References

- Codon C (2006) Shutdown decay of mRNA Mol. Microbiol 61:573–583
- Durand S, Richard G, Bisaglia M, Laalami S, Bontems F, Uzan M (2006) Activation of RegB endoribonuclease by S1 ribosomal protein requires an 11nt conserved sequence. Nucleic Acids Res 34:6549–6560
- Jayasena VK, Brown D, Shtatland T, Gold L (1996) In vitro selection of RNA specifically cleaved by bacteriophage T4 RegB endoribonuclease. Biochemistry 35:2349–2356
- Lebars I, Hu RM, Lallemand JY, Uzan M, Bontems F (2001) Role of the substrate conformation and of the S1 protein in the cleavage efficiency of the T4 endoribonuclease RegB. J Biol Chem 276:13264–13272
- Odaert B, Saïda F, Aliprandi P, Durand S, Crechet JB, Guerois R, Laalami S, Uzan M, Bontems F (2007) Structural and functional studies of RegB, a new member of a family of sequence specific ribonucleases involved in mRNA inactivation on the ribosome. J Biol Chem 282:2019–2028
- Ruckman J, Ringquist S, Brody E, Gold L (1994) The bacteriophage T4 RegB endoribonuclease. Stimulation of the purified enzyme by ribosomal protein S1. J Biol Chem 269:26655–26662
- Saïda F, Odaert B, Uzan M, Bontems F (2004) First structural investigation of the restriction ribonuclease RegB: NMR spectroscopic conditions, $^{13}\text{C}/^{15}\text{N}$ double-isotopic labeling and two-dimensional heteronuclear spectra. Protein Expr Purif 34:158–165
- Sanson B, Hu RM, Troitskayadagger E, Mathy N, Uzan M (2000) Endoribonuclease regB from bacteriophage T4 is necessary for the degradation of early but not middle or late mRNAs. J Mol Biol 297:1063–1074
- Savarin P, Zinn-Justin S, Gilquin B (2001) Variability in automated assignment of NOESY spectra and three-dimensional structure determination: a test case on three small disulfide-bonded proteins. J Biomol NMR 19:49–62
- Uzan M (2001) Bacteriophage T4 RegB endoribonuclease. Methods Enzymol 342:467–480