

## Contents to volume 133 (1990)

### Number 1 – 4 October 1990

#### *Announcement from the publisher*

v

#### *Research reports*

- A novel covalent enzyme-linked immunoassay (CELIA) for simultaneously measuring free and immune complex bound antibodies of defined specificity. I. Application to naturally occurring antipolyamine antibodies in human sera  
*A.M. Roch, J.G. Delcros, J.P. Ripoll, V. Thomas, J. Richard and G. Quash (France)* 1
- A novel covalent enzyme-linked immunoassay (CELIA) for simultaneously measuring free and immune complex bound antibodies with a defined specificity. II. Application to immune complexes containing viral antigens in human sera  
*V. Thomas, S. El Alaoui, A.M. Roch and G. Quash (France)* 13
- A turbidimetric assay in an ELISA reader for the determination of mononuclear phagocyte procoagulant activity  
*T.W. Jungi (Switzerland)* 21
- Isolation of rat peritoneal mononuclear and polymorphonuclear leucocytes on discontinuous gradients of Nycodenz  
*S. Fisker, K. Kudahl and O. Sonne (Denmark)* 31
- Fluorescein-specific hybridomas derived from primary mice exhibit more stringent growth requirements than do hybrids from pre-immune animals  
*J.B. Skeath, M.R. Dalesandro, M.A. Ferguson, G.C. Kinnel and J.A. Owen (U.S.A.)* 39
- A new method for measuring eosinophil activating factors, based on the increased expression of CR3  $\alpha$  chain (CD11b) on the surface of activated eosinophils  
*K.J.I. Thorne, B.A. Richardson, G. Mazza and A.E. Butterworth (U.K.)* 47
- Use of the fluorescence activated cell sorter to enrich dendritic cells from mouse spleen  
*M.T. Crowley, K. Inaba, M.D. Witmer-Pack, S. Gezelter and R.M. Steinman (U.S.A.)* 55
- Immobilization of C3b with retention of functional activity  
*R.E. Edens, R.J. Linhardt and J.M. Weiler (U.S.A.)* 67
- Induction of anti-idiotypic antibodies to a myeloma protein linked covalently to muramyl dipeptide  
*A.E. Eggers and J. Michl (U.S.A.)* 71
- Transfection and gene expression in normal and malignant primary B lymphocytes  
*M. Buschle, M.K. Brenner, I.S.Y. Chen, H.G. Drexler, S.M. Gignac and C.M. Rooney (U.K., U.S.A.)* 77
- New fluorescent dyes for lymphocyte migration studies. Analysis by flow cytometry and fluorescence microscopy  
*S.A. Weston and C.R. Parish (Australia)* 87
- Fetal bovine serum contains an inhibitor of interleukin-1  
*R.C. McKenzie, C.B. Harley, S. Matic and D.N. Sauder (Canada)* 99
- Characterization of polyclonal anti-peptide antibodies specific for transforming growth factor  $\beta_2$   
*A.J.M. Van den Eijnden-Van Raaij, I. Koornneef, H.G. Slager, C.L. Mummery and E.J.J. Van Zoelen (The Netherlands)* 107
- Time resolved fluoroimmunoassay of 7-methyl-2'-deoxyguanosine imidazole (ring open)  
*P. Degan, R. Montesano, A. Abbondandolo and G. Montagnoli (Italy, France)* 119
- Bioassay of interleukin-1 in serum and plasma following removal of inhibitory activity with polyethylene glycol  
*S.J. Hopkins and M. Humphreys (U.K.)* 127
- A single-step method for the purification of anti-FITC antibodies by use of a coumarin immunosorbent  
*D. Samuel and R.A. Abuknesha (U.K.)* 133

#### *Short communication*

- Dissociation rate constant of the biotin-streptavidin complex  
*U. Piran and W.J. Riordan (U.S.A.)* 141

#### *Letter to the editors*

- Non-specific reactivity of monoclonal antibodies in ELISA due to the presence of an antigen-like structure on the alkaline phosphatase molecule  
*M. St. Laurent, F. Martel, S. Verrette and R. Lemieux (Canada)* 145

#### *Book reviews*

147

#### *Notes to authors*

149

*Research reports*

- The sensitivity of dot immunoassay for the peptides helodermin, histidine-isoleucinamide (PHI) and histidine-methioninamide (PHM) increases after peptide cross-linking to proteins prefixed on nitrocellulose  
*P. Gourlet, M. Svoboda, A. Cauvin and J. Christophe (Belgium)* 151
- Antigen protection of monoclonal antibodies undergoing labelling  
*M. Ramjeesingh, M. Zywulko, A. Rothstein, R. Whyte and E.Y. Shami (Canada)* 159
- An immunoassay for the quantitation of cell membrane MHC class II antigen  
*C.V. Whiting and P.W. Bland (U.K.)* 169
- The use of diphenylene iodonium, an inhibitor of NADPH oxidase, to investigate the antimicrobial action of human monocyte derived macrophages  
*A.K. Robertson, A.R. Cross, O.T.G. Jones and P.W. Andrew (U.K.)* 175
- Quantitation of components of the alternative pathway of complement (APC) by enzyme-linked immunosorbent assays  
*M. Oppermann, H. Baumgarten, E. Brandt, W. Gottsleben, C. Kurts and O. Götze (F.R.G.)* 181
- A chemiluminescent, microparticle-membrane capture immunoassay for the detection of antibody to hepatitis B core antigen  
*J. Wolf-Rogers, J.A. Weare, K. Rice, E.F. Robertson, P. Guidinger, O.S. Khalil and G. Madsen (U.S.A.)* 191
- Functional assay of C5-activating and nonactivating cobra venom factor preparations in the mouse system  
*C.W. Van den Berg, P.C. Aerts and H. Van Dijk (The Netherlands)* 199
- Effect of hapten heterology on thyroid hormone immunoassays  
*U. Piran, W.J. Riordan and D.R. Silbert (U.S.A.)* 207
- The detection of anti-Ro/SS-A and anti-La/SS-B antibodies. A comparison of counterimmunoelectrophoresis with immunoblot, ELISA, and RNA-precipitation assays  
*J.F. Meilof, I. Bantjes, J. De Jong, A.P. Van Dam and R.J.T. Smeenk (The Netherlands)* 215
- A comparison of immunoblotting, flow cytometry and ELISA to monitor the binding of anti-lipopolysaccharide monoclonal antibodies  
*D. Nelson, W. Neill and I.R. Poxton (U.K.)* 227
- Gating of the so-called 'lymphocytic' cell population for the quantification of natural killer cells (CD16<sup>+</sup>) by flow cytometry causes loss of CD16 positive cells  
*A.M. Lagaay, C.B. Bosman, M. Van der Keur, G.J. Ligthart, H.R.E. Schuit, H.J. Tanke and W. Hijmans (The Netherlands)* 235
- A method for blocking antigen-independent binding of human IgM to frozen tissue sections when screening human hybridoma antibodies  
*H. Ditzel, K. Erb, B. Nielsen, P. Borup-Christensen and J.C. Jensenius (Denmark)* 245
- Direct blotting with viable cells of protein mixtures separated by two-dimensional gel electrophoresis  
*H. Gulle, B. Schoel and S.H.E. Kaufmann (F.R.G.)* 253
- Qualitative analysis of antibody binding. An in vitro assay for the evaluation and development of vaccines  
*U. Bruderer, E. Furer, S.J. Cryz, Jr. and A.B. Lang (Switzerland)* 263
- Purification of human basophils using mouse monoclonal IgE  
*J.T. Schroeder and L.R. Hanrahan, Jr. (U.S.A.)* 269
- An enzyme-linked immunosorbent assay for the measurement of human interleukin-6  
*N. Ida, S. Sakurai, T. Hosaka, K. Hosoi, T. Kunitomo, Y. Matsuura and M. Kohase (Japan)* 279
- A rapid [<sup>3</sup>H]glycerol radioassay for determination of monocyte-mediated growth inhibition of *Mycobacterium avium*  
*D.K. Blanchard, M.B. Michelini-Norris and J.Y. Djeu (U.S.A.)* 285

*Editorial*

291

*Book review*

293

*Announcements*

295

*Contents to volume 133 (1990)*

297

*Notes to authors*

299