

Available online at www.sciencedirect.com SCIENCE DIRECT*

DOMESTIC ANIMAL ENDOCRINOLOGY

Domestic Animal Endocrinology 28 (2005) 349

www.journals.elsevierhealth.com/periodicals/dae

Erratum

Erratum to "Quantification of insulin-like growth factor binding protein mRNA using real-time PCR in bovine granulosa and theca cells: effect of estradiol, insulin, and gonadotropins" [Domest. Anim. Endocrinol. 26 (2004) 241–258]

J.L. Voge^a, C.A.T. Santiago^a, P.Y. Aad^a, D.W. Goad^b, J.R. Malayer^b, L.J. Spicer^a,*

Received 20 September 2004

The publisher regrets that an error occurred in the above article. The error is corrected as follows. The publisher would like to apologise for any inconvenience caused.

Page 246, Section 2.3.3., paragraph 2, lines 17–20 should read as follows:

An IGFBP-5(335 bp; Accession S52657) forward primer was constructed from bp 193 to 215 with nucleotide sequence of GTTTGCCTGAACGAAAAGAGCTA while the reverse primer was found between bp 275 and 297 with a sequence of CGAGTAGGTCTCCTCTGCCATCT. The IGFBP-5 probe was located from bp 225 to 252 with the sequence of AGCCAAGATCGAAAGAGACTCCCGTGAG.

^a Department of Animal Science, 114 Animal Science Bldg., Oklahoma State University, Stillwater, OK 74078, USA

b Department of Physiological Sciences, 221 McElroy Hall, Oklahoma State University, Stillwater, OK 74078, USA

DOI of original article:10.1016/j.domaniend.2003.11.002

Approved for publication by the Director, Oklahoma Agricultural Experimental Station. This research was supported in part under project H-2329 (to L.J. Spicer).

^{*} Corresponding author. Tel.: +1 405 744 6077; fax: +1 405 744 7390. E-mail address: igf1Leo@okstate.edu (L.J. Spicer).