

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/225564164>

Lignan glycosides and flavonoids from *Saraca asoca* with antioxidant activity

ARTICLE *in* JOURNAL OF NATURAL MEDICINES · SEPTEMBER 2007

Impact Factor: 1.59 · DOI: 10.1007/s11418-007-0182-3

CITATIONS

44

READS

190

5 AUTHORS, INCLUDING:



[Samir Kumar Sadhu](#)

Khulna University

79 PUBLICATIONS 666 CITATIONS

[SEE PROFILE](#)



[Amina Khatun](#)

Manarat International University

26 PUBLICATIONS 138 CITATIONS

[SEE PROFILE](#)

Lignan glycosides and flavonoids from *Saraca asoca* with antioxidant activity

Samir Kumar Sadhu · Amina Khatun ·
Panadda Phattanawasin · Takashi Ohtsuki ·
Masami Ishibashi

Received: 12 April 2007 / Accepted: 13 June 2007
© The Japanese Society of Pharmacognosy and Springer 2007

Abstract Five lignan glycosides, lyoniside, nudiposide, 5-methoxy-9- β -xylopyranosyl-(–)-isolariciresinol, icariside E₃, and schizandriside, and three flavonoids, (–)-epicatechin, epiafzelechin-(4 β →8)-epicatechin and procyanidin B₂, together with β -sitosterol glucoside, were isolated from a methyl alcohol (MeOH) extract of *Saraca asoca* dried bark. Their structures were determined by 1D and 2D nuclear magnetic resonance (NMR) and mass spectroscopic analysis. Antioxidant activities were evaluated by 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical-scavenging assay.

Keywords *Saraca asoca* · Caesalpiniaceae · Lignan glycoside · Flavonoid · Antioxidant activity

Saraca asoca (Roxb.) De Wilde or *S. indica* Linn. (Family: Caesalpiniaceae; local names: Ashok, Anganapriya, etc.) is a medicinal plant of Bangladesh whose bark is astringent and used in menorrhagia, bleeding haemorrhoids and haemorrhagic dysentery [1]. The isolation of tannins [2],

flavonoids [3], proanthocyanidins [4] and leucoanthocyanidins [5] were previously reported from the bark. In our assay method, a methyl alcohol (MeOH) extract of the dried bark showed potent antioxidant activity determined by 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical-scavenging assay [6]. Following this activity, we isolated eight compounds (1–8) from this plant (Fig. 1). This paper describes isolation and identification of isolated compounds together with their antioxidative potential. The bark of *S. asoca* was collected from Satkhira, Bangladesh, in August 2005. A voucher specimen has been deposited in the Laboratory of Natural Products Chemistry, Graduate School of Pharmaceutical Sciences, Chiba University, Japan.

The dried bark of *S. asoca* was ground into a coarse powder (195 g), which was extracted with 1.3 L MeOH to get the extract (14.7 g). Based on the medicinal uses, the extract was tested for antioxidant activity using DPPH radical-scavenging assay that showed a potent effect with an IC₅₀ value of 25 μ g/ml. The extract was then suspended in 440 ml 10% aqueous MeOH and partitioned successively with *n*-hexane, ethyl acetate (EtOAc) and *n*-butanol (*n*-BuOH) that afforded four extracts (*n*-hexane extract: 217 mg; EtOAc extract: 3.11 g; *n*-BuOH extract: 8.74 g; water extract: 2.87 g). Among them, EtOAc and *n*-BuOH extracts showed clear DPPH positive spots on thin-layer chromatography (TLC). These two extracts were then subjected to further separations by repeated-column chromatography. From the *n*-BuOH extract (4.0 g), compounds 1 (20 mg), 2 (16 mg), 3 (4 mg) and 4 (8 mg) were isolated. Compounds 1 (5 mg), 3 (2 mg), 5 (5 mg), 6 (20 mg), 7 (36 mg) and 8 (17 mg), together with β -sitosterol glucoside (12 mg) [7], were isolated from the EtOAc extract (3.0 g). Compounds 1–5 were lignan glycosides, which were identified as lyoniside [α]_D²¹ + 22 (c 1.0, MeOH) [8],

S. K. Sadhu · T. Ohtsuki · M. Ishibashi (✉)
Graduate School of Pharmaceutical Sciences,
Chiba University, 1-33 Yayoi-cho, Inage-ku,
Chiba 263-8522, Japan
e-mail: mish@p.chiba-u.ac.jp

S. K. Sadhu · A. Khatun
Pharmacy Discipline, Life Science School,
Khulna University, Khulna 9208, Bangladesh

P. Phattanawasin
Faculty of Pharmacy, Silpakorn University,
Nakorn Pathom 73000, Thailand




Fig. 1 Structures of compounds **1–8** from *Saraca asoca*

nudiposide [α]_D²² – 52 (c 1.0, MeOH) [9], 5-methoxy-9- β -D-xylopyranosyl-(–)-isolariciresinol [α]_D¹⁹ – 40 (c 1.0, MeOH) [10], icariside E₃ [α]_D²² – 57 (c 2.0, MeOH) [11] and schizandriside [α]_D²³ + 18 (c 1.0, MeOH) [12], respectively (Table 1). The flavonoids **6–8** were identified as (–)-epicatechin [α]_D²⁴ – 42 (c 2.6, acetone) [13], epiafzelechin-(4 β →8)-epicatechin [α]_D²³ + 43 (c 1.7, acetone) [14] and procyanidin B₂ [α]_D²⁴ + 35 (c 1.8, acetone) [15, 16], respectively, by comparison of their spectroscopic data with those in the literature. The lignan glycosides (**1–5**) were obtained from *S. asoca* for the first time. The IC₅₀ values of DPPH radical-scavenging assay for compounds **1–8**, and the positive control quercetin were 104, 85, 44, 75, 55, 50, 55, 40 and 30 μ M, respectively. In conclusion, all the isolated compounds exhibited moderate antioxidant activity that might be responsible together for the therapeutic efficacy of this herb.

Table 1 ¹H and ¹³C nuclear magnetic resonance (NMR) spectroscopic data of compounds **1–5** (δ in ppm, *J* in Hz)

Position	1		2		3		4		5	
	C	H	C	H	C	H	C	H	C	H
1	139.4		139.6		137.9		140.3		138.6	
2	106.9	6.42 (1H, s)	106.8	6.40 (1H, s)	107.8	6.42 (1H, s)	111.7	6.71 (1H, brs)	114.2	6.77 (1H, d, 1.8)
3	149.0		148.9		149.1		153.1		148.9	
4	134.5		134.4		134.9		143.6		145.8	
5	149.0		148.9		149.1		138.6		116.1	6.73 (1H, d, 8.2)
6	106.9	6.42 (1H, s)	106.8	6.40 (1H, s)	107.8	6.42 (1H, s)	120.3	6.71 (1H, brs)	123.1	6.62 (1H, dd, 8.2, 1.8)
7	43.0	4.37 (1H, d, 6.7)	43.4	4.21 (1H, d, 7.0)	49.5	3.80 (1H, overlapped)	33.1	2.63 (2H, dd, 8.2, 7.4)	47.9	4.05 (1H, brd, 10.3)
8	46.7	2.00–2.06 (1H, m)	46.9	1.98–2.03 (1H, m)	45.5	1.90 (1H, tdd, 10.3, 4.3, 2.4)	35.6	1.81 (2H, m)	45.9	1.84 (1H, tt, 10.3, 2.8)
9	71.0	3.41 (1H, dd, 9.8, 3.7) 3.83 (1H, dd, 9.8, 4.9)	70.9	3.57 (1H, dd, 10.0, 4.9) 3.81 (1H, dd, 10.0, 4.6)	70.0	3.61 (1H, dd, 10.1, 2.4) 3.74 (1H, m)	62.2	3.55 (2H, td, 6.4, 0.6)	69.3	3.21 (1H, dd, 10.1, 3.1) 3.97 (1H, dd, 10.1, 2.4)
1'	130.1		130.1		129.2		133.3		129.1	
2'	107.8	6.56 (1H, s)	107.6	6.56 (1H, s)	112.3	6.64 (1H, s)	113.6	6.55 (1H, d, 2.0)	112.3	6.64 (1H, s)
3'	148.6		148.6		147.3		148.4		147.1	
4'	138.9		138.9		145.3		145.3		145.1	
5'	147.6		147.5		117.3	6.19 (1H, s)	115.6	6.54 (1H, d, 8.0)	117.4	6.16 (1H, s)
6'	126.4		126.3		133.6		122.6	6.46 (1H, dd, 8.0, 2.0)	134.3	
7'	33.9	2.63 (1H, dd, 15.0, 11.3) 2.71 (1H, dd, 15.0, 4.6)	34.1	2.68 (2H, dd, 9.8, 6.5)	33.8	2.75 (1H, dd, 15.5, 4.6) 2.87 (1H, dd, 15.5, 11.6)	39.2	2.69 (1H, dd, 13.7, 9.5) 2.97 (1H, dd, 13.7, 5.2)	33.9	2.77–2.86 (2H, m)
8'	40.5	1.66–1.73 (1H, m)	40.6	1.67–1.74 (1H, m)	40.4	1.97–2.04 (1H, m)	42.8	3.92–3.98 (1H, m)	39.6	2.03–2.10 (1H, m)
9'	66.0	3.55 (1H, dd, 11.0, 6.7) 3.65 (1H, dd, 11.0, 4.3)	66.0	3.62 (2H, tlike, 4.3)	65.3	3.72 (1H, m) 3.80 (1H, overlapped)	67.1	3.65 (1H, dd, 10.7, 7.2) 3.75 (1H, dd, 10.7, 6.1)	65.1	3.76 (1H, dd, 11.0, 3.7) 3.70 (1H, dd, 11.0, 6.1)
3-OMe	56.8	3.74 (3H, s)	56.7	3.74 (3H, s)	56.7	3.78 (3H, s)	56.3	3.79 (3H, s)	56.4	3.788 (3H, s)
5-OMe	56.8	3.74 (3H, s)	56.7	3.74 (3H, s)	56.7	3.78 (3H, s)				
3'-OMe	56.6	3.84 (3H, s)	56.5	3.84 (3H, s)	56.3	3.80 (3H, s)	56.2	3.68 (3H, s)	56.3	3.794 (3H, s)

Table 1 continued

Position	1		2		3		4		5	
	C	H	C	H	C	H	C	H	C	H
5'-OMe	60.0	3.31 (3H, s)	59.9	3.28 (3H, s)						
1''	105.5	4.21 (1H, d, 7.6)	105.0	4.09 (1H, d, 7.6)	104.8	4.02 (1H, d, 7.6)	105.6	4.61 (1H, d, 7.4)	105.8	4.04 (1H, d, 7.6)
2''	75.0	3.21 (1H, dd, 8.9, 7.6)	74.9	3.18 (1H, dd, 9.2, 7.6)	74.9	3.16 (1H, dd, 9.1, 7.6)	77.9	3.41 (1H, m)	75.0	3.18 (1H, dd, 9.2, 7.6)
3''	78.0	3.27–3.29 (1H, m)	78.0	3.26 (1H, overlapped)	78.0	3.24 (1H, dd, 9.8, 9.1)	75.9	3.43 (1H, m)	77.9	3.28 (1H, t, 8.9)
4''	71.3	3.47 (1H, ddd, 10.4, 9.0, 5.8)	71.3	3.47 (1H, ddd, 10.4, 8.9, 5.8)	71.3	3.46 (1H, ddd, 10.4, 9.8, 5.5)	71.2	3.37 (1H, m)	71.3	3.42–3.47 (1H, m)
5''	67.0	3.16 (1H, dd, 11.6, 10.4)	67.1	3.12 (1H, dd, 11.3, 10.4)	67.0	3.04 (1H, dd, 11.3, 10.4)	78.1	3.09–3.12 (1H, m)	66.9	3.11 (1H, dd, 11.3, 10.4)
		3.82 (1H, dd, 11.6, 5.8)		3.85 (1H, dd, 11.3, 5.8)		3.81 (1H, overlapped)				3.80 (1H, dd, 11.3, 5.8)
6''							62.5	3.67 (1H, overlapped)		
								3.78 (1H, overlapped)		

NMR was recorded in CD₃OD; ¹H NMR at 500 MHz and ¹³C NMR at 125 MHz

Acknowledgments This work was partly supported by a Grant-in-Aid from Tokyo Biochemical Research Foundation (TBRF), Japan.

References

- Ghani A (2003) Medicinal plants of Bangladesh with chemical constituents and uses. 2nd edn. Asiatic Society of Bangladesh, Dhaka
- Indrani N, Balasubramanian K (1985) Isolation of condensed tannins from *Saraca ashoka*—part II. *Leather Sci* 32:12–13
- Indrani N, Balasubramanian K (1985) Isolation of condensed tannins from *Saraca ashoka*—part I. *Leather Sci* 31:349–350
- Middelkoop TB, Labadie RP (1985) Proanthocyanidins in the bark of *Saraca asoca* Roxb de Wilde. *Z Naturforsch [B]* 40B:855–857
- Duggal JK, Misra K (1980) Leucoanthocyanidins from *Saraca asoca* stem bark. *J Indian Chem Soc* 57:1243
- Sadhu SK, Okuyama E, Fujimoto H, Ishibashi M (2003) Separation of *Leucas aspera*, a medicinal plant of Bangladesh, guided by prostaglandin inhibitory and antioxidant activities. *Chem Pharm Bull* 51:595–598
- Faizi S, Ali M, Saleem R, Irfanullah BS (2001) Complete ¹H and ¹³C NMR assignments of stigma-5-en-3-*O*- β -glucoside and its acetyl derivative. *Magn Reson Chem* 39:399–405
- Dada G, Corbani A, Manitto P, Speranza G, Lunazzi L (1989) Lignan glycosides from the heartwood of European oak *Quercus petraea*. *J Nat Prod* 52:1327–1330
- Inoshiri S, Sasaki M, Kohda H, Otsuka H, Yamasaki K (1987) Aromatic glycosides from *Berchemia racemosa*. *Phytochemistry* 26:2811–2814
- Vecchietti V, Ferrari G, Orsin F, Pelizzoni F (1979) Alkaloid and lignan constituents of *Cinnamosa madagascariensis*. *Phytochemistry* 18:1847–1849
- Miyase T, Ueno A, Takizawa N, Kobayashi H, Oguchi H (1988) Studies on the glycosides of *Epimedium grandiflorum* Morr var *thunbergianum* (Miq.) Nakai III. *Chem Pharm Bull* 36:2475–2484
- Takani M, Ohya K, Takahashi K (1979) Studies on constituents of medicinal plants. XXII. Constituents of *Schizandra nigra* Max (4). *Chem Pharm Bull* 27:1422–1425
- Shen CC, Chang YS, Ho LK (1993) Nuclear magnetic resonance studies of 5,7-dihydroxyflavonoids. *Phytochemistry* 34:843–845
- Kashiwada Y, Iizuka H, Yoshioka K, Chen RF, Nonaka G, Nishioka I (1990) Tannins and related compounds. XCIII. Occurrence of enantiomeric proanthocyanidins in the Leguminosae plants, *Cassia fistula* L. and *C javanica* L. *Chem Pharm Bull* 38:888–893
- Fletcher AC, Porter LJ, Haslam E, Gupta RK (1977) Plant proanthocyanidins. Part 3. Conformational and configurational studies of natural procyanidins. *J Chem Soc Perkin Trans I*:1628–1637
- Nonaka GI, Nishioka I, Nagasawa T, Oura H (1981) Tannins and related compounds I Rhubarb. *Chem Pharm Bull* 29:2862–2870