

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/232390986>

# Calculation of rate coefficients of electron energy transfer processes for molecular nitrogen and molecular oxygen

ARTICLE *in* ADVANCES IN SPACE RESEARCH · JANUARY 2004

Impact Factor: 1.36 · DOI: 10.1016/j.asr.2003.06.014

---

CITATIONS

26

---

READS

24

## 1 AUTHOR:



[Andrey S. Kirillov](#)

Polar Geophysical Institute

**59** PUBLICATIONS **176** CITATIONS

SEE PROFILE

# Calculation of rate coefficients of electron energy transfer processes for molecular nitrogen and molecular oxygen

A.S. Kirillov

*Polar Geophysical Institute, Apatity, Murmansk region 184200, Russia*

Received 10 September 2002; received in revised form 12 March 2003; accepted 15 June 2003

## Abstract

Cross sections and rate coefficients of electronic quenching for triplet and singlet states of molecular nitrogen and molecular oxygen in molecular inelastic collisions  $N_2-N_2$  and  $O_2-O_2$  were calculated from analytical expressions. Electronic quenching is considered as a sum of quasi-resonant intramolecular and intermolecular electron energy transfer processes in inelastic interactions. In most conditions good agreement with experimental data is obtained.

© 2003 COSPAR. Published by Elsevier Ltd. All rights reserved.

**Keywords:** Ionosphere; Molecular nitrogen; Molecular oxygen; Electron energy transfer processes

## 1. Introduction



The total rate of the excitation of triplet and singlet states of molecular nitrogen and molecular oxygen in low temperature plasma (Konovalov and Son, 1987) and in auroral ionosphere (Sergienko and Ivanov, 1993) is comparable with the rate of ion–electron pair production. The spontaneous transitions between the states of  $N_2$  cause emission of Vegard–Kaplan (transition  $A^3\Sigma_u^+ \rightarrow X^1\Sigma_g^+$ ), first positive ( $B^3\Pi_g \rightarrow A^3\Sigma_u^+$ ), second positive ( $C^3\Pi_u \rightarrow B^3\Pi_g$ ), Wu–Benesch ( $W^3\Delta_u \rightarrow B^3\Pi_g$ ), Lyman–Birge–Hopfield ( $a^1\Pi_g \rightarrow X^1\Sigma_g^+$ ), etc. bands. The spontaneous transitions between the states of  $O_2$  cause the radiation of infrared atmospheric, atmospheric, Herzberg I, II, III, etc. bands (transitions from  $a^1\Delta_g$ ,  $b^1\Sigma_g^+$ ,  $A^3\Sigma_u^+$ ,  $c^1\Sigma_u^-$ ,  $A'^3\Delta_u$  states to  $X^3\Sigma_g^-$ , respectively).

Electronic and vibrational excitation of  $N_2$  and  $O_2$  could increase the rate coefficients of chemical reactions with ions and neutrals influencing atmospheric chemical kinetics. The collisional quenching rates of some triplet and singlet states are comparable with radiational ones in the region of the lower thermosphere–mesosphere.

Main purpose of the paper is the calculation of cross sections and rate coefficients of electron energy transfers and quenching processes for triplet and singlet states of molecular nitrogen and molecular oxygen in molecular inelastic collisions  $N_2-N_2$  and  $O_2-O_2$ . To this end we apply the set of formulas derived by Kirillov (2003).



## 2. Electron energy transfers between triplet states of $N_2$

Collision-induced coupling within the triplet manifold of  $N_2$  has been studied in Bachmann et al. (1992, 1993), Ottinger et al. (1995). The investigations have shown that the processes





have large cross sections due to energy resonance between the reactant and product vibrational levels. Moreover, it was obtained that in the experiments the off-resonance processes

E-mail address: [kirillov@pgi.kolasc.net.ru](mailto:kirillov@pgi.kolasc.net.ru) (A.S. Kirillov).



were found rather important. Bachmann et al. (1993) attributed this to absorption of excess energy by target vibration as an additional degree of freedom. We, however, feel the processes (R3) and (R4) are intermolecular electron energy transfers to target molecule, namely:



Moreover, an isotopic study of electron energy transfer processes (Ottinger et al., 1995) by separating of intermolecular from competing intramolecular collisional coupling has shown that intermolecular coupling of triplet states in  $\text{N}_2\text{--N}_2$  collisions occurs with cross sections of similar magnitudes as the intramolecular processes.

We suggest to calculate the quenching coefficients of intramolecular (Eqs. (R1) and (R2)) and intermolecular (Eqs. (R5) and (R6)) processes according to analytical formulas obtained by Kirillov (2003) applying Landau–Zener and Rosen–Zener approximations. In this approach the processes (R3) and (R4) are unimportant because the Franck–Condon factor for the molecule disappears in Eq. (27) of Kirillov (2003).

By the way, here we use conclusions of Bachmann et al. (1993) that a large cross section of the exothermic process equation (R1) compared to the one of endothermic process equation (R2) points to a crossing of exchange potentials in the case of  $\text{A}^3\Sigma_u^+, v' \rightarrow \text{B}^3\Pi_g, v$  exothermic transfer. Suggested rate coefficients for  $\text{A}^3\Sigma_u^+, v' \rightarrow \text{B}^3\Pi_g, v$ ,  $\text{W}^3\Delta_u, v' \rightarrow \text{B}^3\Pi_g, v$ ,  $\text{B}^3\Sigma_u, v' \rightarrow \text{B}^3\Pi_g, v$  transitions are

$$k = k_{LZ} q_{v'v}, \quad (1)$$

$$k = k_{RZ} q_{v'v} \quad (2)$$

for intramolecular exothermic and endothermic energy transfer processes, respectively, and

$$k = k_{LZ} q_{v'v''} q_{0v}, \quad (3)$$

$$k = k_{RZ} q_{v'v''} q_{0v} \quad (4)$$

for intermolecular exothermic and endothermic energy transfer processes, respectively.  $k_{RZ}$  and  $k_{LZ}$  have to be calculated according to Eqs. (20) and (22) of Kirillov

(2003). Franck–Condon factors  $q$  were taken according to Gilmore et al. (1992).

We took also account of the fact that  $\text{A}^3\Sigma_u^+$ ,  $\text{W}^3\Delta_u$ ,  $\text{B}^3\Sigma_u^-$  states have similar configuration of electron orbitals  $1\sigma_g^2 1\sigma_u^2 2\sigma_g^2 2\sigma_u^2 1\pi_u^3 3\sigma_g^2 1\pi_g$  and there is the  $\sigma_g \rightarrow \pi_u$  change in the transitions to  $\text{B}^3\Pi_g$  state. Moreover, it is suggested that all three states have similar repulsive exchange potentials and the parameters  $\gamma$  and  $\beta$  in Eqs. (20) and (22) of Kirillov (2003) have the same magnitudes for  $\text{A}^3\Sigma_u^+, v' \rightarrow \text{B}^3\Pi_g, v$ ,  $\text{W}^3\Delta_u, v' \rightarrow \text{B}^3\Pi_g, v$  and  $\text{B}^3\Sigma_u^-, v' \rightarrow \text{B}^3\Pi_g, v$  electron energy transfers. Only the parameter  $k_0$  is two times greater for transitions from the  $\text{W}^3\Delta_u$  state because of  $\Lambda$ -doubling of this initial state.

Experimental data of Bachmann et al. (1992, 1993) for 14  $\text{A}^3\Sigma_u^+, v' \rightarrow \text{B}^3\Pi_g, v$  and seven  $\text{W}^3\Delta_u, v' \rightarrow \text{B}^3\Pi_g, v$  electron transitions were normalized on the expressions (1)–(4) to obtain the mentioned parameters. The method of least squares was used in the normalizing. Corre-




Fig. 1. Calculated cross sections (crosses) for  $\text{A}^3\Sigma_u^+, v' \rightarrow \text{B}^3\Pi_g, v$  transitions are compared with experimental data of Bachmann et al. (1993, squares and triangles for exothermic and endothermic processes, respectively).





Fig. 2. Calculated cross sections (crosses) for  $\text{W}^3\Delta_u, v' \rightarrow \text{B}^3\Pi_g, v$  transitions are compared with experimental data of Bachmann et al. (1992, 1993, squares and triangles for exothermic and endothermic processes, respectively).

sponding to experimental conditions of Bachmann et al. (1992, 1993) temperature  $T$  was taken equal to 657 K. Every energy transfer process was considered as a sum of possible intramolecular and intermolecular processes. The normalization has given  $\gamma = 106 \text{ cm}^{-1}$  and  $\beta = 0.446 \text{ cm}^{-1}/\text{K}$ .  $k_0$  is equal to  $5.1 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$  for the transitions from the  $A^3\Sigma_u^+$  state (also from the  $B'^3\Sigma_u^-$  state) and  $10.2 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$  for the transitions from the  $W^3\Delta_u$  state. Results of our calculations for cross sections are compared with experimental data of Bachmann et al. (1992, 1993) for 14  $A^3\Sigma_u^+, v' \rightarrow B^3\Pi_g, v$  transitions in Fig. 1 and for seven  $W^3\Delta_u, v' \rightarrow B^3\Pi_g, v$  transitions in Fig. 2.



### 3. Quenching rate coefficients of triplet and singlet states of molecular nitrogen in inelastic collisions $N_2$ – $N_2$

Applying Kirillov (2003) here we consider every inelastic quenching of electronically excited triplet and singlet states of molecular nitrogen as a sum of all possible effective intramolecular and intermolecular energy transfer processes.

For the  $B^3\Pi_g$  state we took account of the intramolecular



and intermolecular



processes. The summation includes all possible  $v'$  in the case of intramolecular transfers and all possible  $v', v''$  in the case of intermolecular transfers. Contributions of energetic non-resonant processes in the summation are negligible due to exponential dependence on  $|\Delta E|$ . The rates of exothermic and endothermic processes were calculated according to Eqs. (2), (4) and Eqs. (1), (3), respectively. The results of our calculation at temperature  $T = 300 \text{ K}$  are compared in Fig. 3 with experimental data of Piper (1988) and Shemansky (1976). Good agreement was found with experimental data. Note that the calculation is made with parameters  $k_0 = 10.2 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$ ,  $\beta$  and  $\gamma$  obtained by normalizing the analytical expressions with experimental data (Bachmann et al., 1992, 1993).

For the  $a^1\Pi_g$  state we took account of the intramolecular





Fig. 3. Calculated quenching rate coefficients for the  $B^3\Pi_g$  state of  $N_2$  at temperature  $T = 300 \text{ K}$  are compared with experimental data of Piper (1988, squares) and Shemansky (1976, triangles).






Fig. 4. Calculated quenching rate coefficients for the  $a^1\Pi_g$  state of  $N_2$  at temperature  $T = 300 \text{ K}$  are compared with experimental data of van Veen et al. (1982, squares).

and intermolecular



processes. The value of  $k_0 = 3.4 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$  was taken three times lower than for the  $B^3\Pi_g$  state because of the multiplicity  $2S + 1 = 1$ . Values of  $\beta$  and  $\gamma$  were taken as for the  $B^3\Pi_g$  state. The rates of exothermic and endothermic processes were calculated according to Eqs. (2), (4) and Eqs. (1), (3), respectively. The similarity of the calculations for  $a^1\Pi_g$  and  $B^3\Pi_g$  states is explained by configuration similarity of their electron orbitals. Also all five  $A^3\Sigma_u^+$ ,  $W^3\Delta_u$ ,  $B'^3\Sigma_u^-$ ,  $a'^1\Sigma_u^-$ ,  $w^1\Delta_u$ , states are identical. The results of our calculation at temperature

$T = 300$  K are compared in Fig. 4 with experimental data of van Veen et al. (1982).

#### 4. Franck–Condon factors for $O_2$

Our calculations differ from similar ones for molecular nitrogen in applying Franck–Condon factors obtained by Morse approximation. The Morse potential  $U$  is expressed as

$$U(r) = D_e \{1 - \exp[-a(r - r_e)]\}^2, \quad (5)$$

where  $r_e$  is the equilibrium value of internuclear distance  $r$ ,  $D_e$  and  $a$  are parameters. Eq. (5) is a very good approximation for real potentials in most cases (Kuzmenko et al., 1984). Moreover, the Schrodinger equation with the potential has an exact solution for nuclear wavefunctions:

$$\psi_v(r) = N_v e^{-z/2} z^{\beta/2} L_v^\beta(z), \quad (6)$$


where  $z = \exp[-a(r - r_e)]/x_e$ ,  $\beta = 1/x_e - 2v - 1$ ,  $x_e$  is anharmonic constant,  $L_v^\beta$  are Laguerre polynomials.

Spectroscopic constants for  $X^3\Sigma_g^-$ ,  $a^1\Delta_g$ ,  $b^1\Sigma_g^+$  states were taken according to Huber and Herzberg (1979) and for  $c^1\Sigma_u^-$ ,  $A'^3\Delta_u$ ,  $A^3\Sigma_u^+$  states according to Slinger and Cosby (1988). Franck–Condon factors were calculated for all considered electron transitions.



#### 5. Quenching rate coefficients of $a^1\Delta_g$ and $b^1\Sigma_g^+$ states of molecular oxygen in inelastic collisions $O_2$ – $O_2$

A lot of vibrational levels of  $X^3\Sigma_g^-$ ,  $a^1\Delta_g$ ,  $b^1\Sigma_g^+$  states are in good energetic resonance. Moreover, since all three  $X^3\Sigma_g^-$ ,  $a^1\Delta_g$ ,  $b^1\Sigma_g^+$  states have same configuration of electron orbitals  $1\sigma_g^2 1\sigma_u^2 2\sigma_g^2 2\sigma_u^2 3\sigma_g^2 1\pi_u^4 1\pi_g^2$ , we believe that all exchange potentials are parallel and apply only Eqs. (2) and (4) for the calculation of quenching rate coefficients of the states. Distinct from the calculations for states of  $N_2$  we have no any information about parameter  $\gamma$ .  $\gamma$  was taken equal to  $106 \text{ cm}^{-1}$  in the calculations for triplet and singlet states on  $N_2$ . So similar values were adopted for  $O_2$  and we have made our calculations for states of  $O_2$  at  $\gamma = 90$  and  $120 \text{ cm}^{-1}$ . Parameter  $k_0$  was taken 3 and 6 times smaller for  $a^1\Delta_g$  and  $b^1\Sigma_g^+$  states, respectively, than for the  $B^3\Pi_g$  state of  $N_2$  because of degeneration of these states.


For the  $a^1\Delta_g$  state we took account of intramolecular






and intermolecular



processes. The summation includes all possible  $v'$  in the case of intramolecular transfers and all possible  $v', v''$  in the case of intermolecular transfers. Results of the calculation have shown that processes (R15) with  $v' = 0$  and  $v'' = v$  or  $v'' = v - 1$  dominate. Calculated quenching rate coefficients with  $k_0 = 3.4 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$  are presented in Fig. 5. As it is seen from this figure there appears a local dip in the calculated rate coefficients at  $v = 7$ – $8$ . The presence of the dip could be explained by greatest mismatch  $|\Delta E| \sim 700$ – $800 \text{ cm}^{-1}$  in processes (R15) for the vibrational levels.



and intermolecular



processes. Results of the calculation have shown that processes (R18) with  $v' = 0$  and  $v'' = v$  or  $v'' = v - 1$  dominate for low values of  $v$ . But the contribution of processes (R17) with  $v' = 0, 1$  and  $v'' = v + 2$  or  $v'' = v + 3$  increases for higher values of  $v$ . Calculated quenching rate coefficients with  $k_0 = 1.7 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$  are compared in Fig. 6 with experimental data of Bloemink et al. (1998), Kalogerakis et al. (2002), Yan-




Fig. 5. Calculated quenching rate coefficients for the  $a^1\Delta_g$  state of  $O_2$  (solid and dashed lines are for  $\gamma = 90$  and  $120 \text{ cm}^{-1}$ , respectively).






Fig. 6. Calculated quenching rate coefficients for the  $b^1\Sigma_g^+$  state of  $O_2$  are compared with experimental data of Bloemink et al. (1998, squares), Kalogerakis et al. (2002, squares), Yankovskii (1991, crosses), Kebabian and Freedman (1997, triangle), Lee and Slanger (1978, star). Solid and dashed lines are for  $\gamma = 90$  and  $120 \text{ cm}^{-1}$ , respectively.




kovskii (1991), Kebabian and Freedman (1997) and Lee and Slanger (1978). As in the case of the  $a^1\Delta_g$  state the calculated rate coefficients (Fig. 6) are decreasing from  $v = 1$  to  $v = 4$  with a local dip at  $v = 4$  and an increase thereafter up to  $v = 8$ . The decrease is in agreement with experimental data and also could be explained by greatest mismatch  $|\Delta E|$  in processes (R19) for  $v = 4$ .

## 6. Quenching rate coefficients of $A^3\Sigma_u^+$ and $c^1\Sigma_u^-$ states of molecular oxygen in inelastic collisions $O_2-O_2$


For the  $A^3\Sigma_u^+$  state we took account of intramolecular






and intermolecular



processes. For the  $c^1\Sigma_u^-$  state we took account of intramolecular



and intermolecular



processes.

Since  $c^1\Sigma_u^-$ ,  $A'^3\Delta_u$ ,  $A^3\Sigma_u^+$  states have the same configuration of electron orbitals  $1\sigma_g^2 1\sigma_u^2 2\sigma_g^2 2\sigma_u^2 3\sigma_g^2 1\pi_u^3 1\pi_g^3$ , so transitions to lower  $X^3\Sigma_g^-$ ,  $a^1\Delta_g$ ,  $b^1\Sigma_g^+$  states cause the “gerade”  $\pi$ -orbital to change into “ungerade” one. Therefore we suggest parallel exchange potentials and apply Eqs. (2) and (4) in all calculations for the states.




Fig. 7. Calculated quenching rate coefficients for the  $A^3\Sigma_u^+$  state of  $O_2$  in processes (R20) and (R23) (a and b, respectively) are compared with experimental data of Knutsen et al. (1994, squares) and Kenner and Ogryzlo (1983, triangle). Solid and dashed lines are for  $\gamma = 90$  and  $120 \text{ cm}^{-1}$ , respectively.




Fig. 8. Calculated quenching rate coefficients for the  $c^1\Sigma_u^-$  state of  $O_2$  in processes (R24) and (R27) (a and b, respectively) are compared with experimental data of Copeland et al. (1996, square) and Kenner and Ogryzlo (1983, triangle). Solid and dashed lines are for  $\gamma = 90$  and  $120\text{ cm}^{-1}$ , respectively.

The calculations of quenching rate coefficients for the  $A^3\Sigma_u^+$  state have shown that intermolecular processes (R21) and (R22) can be neglected. Calculated values for processes (R20) and (R23) with  $k_0 = 5.1 \times 10^{-10}\text{ cm}^3\text{ s}^{-1}$  are presented in Fig. 7(a) and (b), respectively. Experimental data of Knutsen et al. (1994) and Kenner and Ogryzlo (1983) have been compared with the results of the calculation. Similar results are obtained for the  $c^1\Sigma_u^-$  state. Calculated values for Eqs. (R24) and (R27) with  $k_0 = 1.7 \times 10^{-10}\text{ cm}^3\text{ s}^{-1}$  are presented in Fig. 8(a) and (b), respectively. Experimental data of Copeland et al. (1996) and Kenner and Ogryzlo (1983) have been compared with our calculation. It is seen from Figs. 7 and 8 that our calculations of quenching rate coefficients correlate with experimental data only for intramolecular processes.

## 7. Conclusions

1. Analytical expressions obtained by Kirillov (2003) were applied to calculate rate coefficients of intramolecular and intermolecular energy transfer processes between triplet states of molecular nitrogen in molecular inelastic collisions  $N_2-N_2$ . Experimental data of Bachmann et al. (1992, 1993) were normalized on the expressions to obtain all necessary parameters for the calculation. Every energy transfer process between triplet states were considered as a sum of possible intramolecular and intermolecular processes. Results of our calculations for cross sections were compared with experimental data of Bachmann et al. (1992, 1993) for 14  $A^3\Sigma_u^+, v' \rightarrow B^3\Pi_g, v$  and for seven  $W^3\Delta_u, v' \rightarrow B^3\Pi_g, v$  transitions.
2. Rate coefficients of electronic quenching in molecular collisions  $N_2-N_2$  were calculated for  $B^3\Pi_g$  and  $a^1\Pi_g$

states of  $N_2$  and compared with experimental data. Good agreement of calculated rate coefficients with experimental data was obtained for the  $B^3\Pi_g$  state.

3. Rate coefficients of electronic quenching in molecular collisions  $O_2-O_2$  were calculated for  $a^1\Delta_g$  and  $b^1\Sigma_g^+$  states of  $O_2$  and were compared with experimental data for the  $b^1\Sigma_g^+$  state. Dips found in the results of the calculation and in experimental data are explained by energy mismatch for some vibrational levels of the states. It is pointed that intermolecular energy transfer processes are giving the main contribution to the electronic quenching.
4. Rate coefficients of electronic quenching in molecular collisions  $O_2-O_2$  were calculated for  $A^3\Sigma_u^+$  and  $c^1\Sigma_u^-$  states of  $O_2$  and compared with experimental data. It was shown that a correlation of the results of the calculation with experimental data could be obtained only when intramolecular energy transfer processes were dominant.

## Acknowledgements

The author thanks an unknown referee for his evaluation of the paper. Special thanks go to Prof. K.M. Rawer for his help in the correct writing of the paper. The work is supported by the grant 02-05-64114 of Russian Foundation of Fundamental Investigations.

## References

- Bachmann, R., Li, X., Ottinger, Ch., Vilesov, A.F. Molecular-beam study of the collisional intramolecular coupling of  $N_2(B^3\Pi_g)$  with  $N_2(A^3\Sigma_u^+)$  and  $N_2(W^3\Delta_u)$  states. *J. Chem. Phys.* 96, 5151–5164, 1992.
- Bachmann, R., Li, X., Ottinger, Ch., Vilesov, A.F., Wulfmeyer, V. Vibrational-state-to-state intramolecular energy transfer  $N_2(A^3\Sigma_u^+, v'' \rightarrow B^3\Pi_g, v')$ . *J. Chem. Phys.* 98, 8606–8625, 1993.

- Bloemink, H.I., Copeland, R.A., Slanger, T.G. Collisional removal of  $O_2(b^1\Sigma_g^+, v=1, 2)$  by  $O_2$ ,  $N_2$ , and  $CO_2$ . *J. Chem. Phys.* 109, 4237–4245, 1998.
- Copeland, R.A., Knutsen, K., Onishi, M.E., Yalcin, T. Collisional removal of  $O_2(c^1\Sigma_u^-, v=9)$  by  $O_2$ ,  $N_2$ , and He. *J. Chem. Phys.* 105, 10349–10355, 1996.
- Gilmore, F.R., Laher, R.R., Espy, P.J. Franck–Condon factors, r-centroids, electronic transition moments, and Einstein coefficients for many nitrogen and oxygen band systems. *J. Phys. Chem. Ref. Data* 21, 1005–1107, 1992.
- Huber, K.P., Herzberg, G. *Molecular spectra and molecular structure*, in: *Constants of Diatomic Molecules*, vol. 4. Van Nostrand Reinhold, New York, 1979.
- Kalogerakis, K.S., Copeland, R.A., Slanger, T.G. Collisional removal of  $O_2(b^1\Sigma_g^+, v=2, 3)$ . *J. Chem. Phys.* 116, 4877–4885, 2002.
- Kebabian, P.L., Freedman, A. Rare gas quenching of metastable  $O_2(b^1\Sigma_g^+)$  at 295 K. *J. Phys. Chem.* 101, 7765–7767, 1997.
- Kenner, R.D., Ogryzlo, E.A. Quenching of  $O_2(c^1\Sigma_u^-)$   $v=0$  by  $O(^3P)$ ,  $O_2(a^1\Delta_g)$ , and other gases. *Can. J. Chem.* 61, 921–926, 1983.
- Kirillov, A.S. Application of Landau–Zener and Rosen–Zener approximations in the calculation of rate coefficients of intramolecular and intermolecular electron energy transfer processes for atmospheric gases, *Adv. Space Res.*, this issue.
- Knutsen, K., Dyer, M.J., Copeland, R.A. Laser double-resonance study of the collisional removal of  $O_2(A^3\Sigma_u^+, v=6, 7, \text{ and } 9)$  with  $O_2$ ,  $N_2$ ,  $CO_2$ , Ar, and He. *J. Chem. Phys.* 101, 7415–7422, 1994.
- Konovalov, V.P., Son, E.E. Degradation spectra of electrons in gases. *Chem. Plasma* 14, 194–227, 1987 (in Russian).
- Kuzmenko, N.E., Kuznetsova, L.A., Kuzjakov, Yu.A. Franck–Condon factors of diatomic molecules. Moscow university, Moscow, 1984 (in Russian).
- Lee, L.C., Slanger, T.G. Observations on  $O(^1D \rightarrow ^3P)$  and  $O_2(b^1\Sigma_g^+ \rightarrow X^3\Sigma_g^-)$  following  $O_2$  photodissociation. *J. Chem. Phys.* 69, 4053–4060, 1978.
- Ottinger, Ch., Vilesov, A.F., Xu, D.D. Isotopic study of the intermolecular versus intramolecular energy transfer in the  $N_2(W,A)+N_2(X)$  reactions. *J. Chem. Phys.* 102, 1673–1680, 1995.
- Piper, L.G. State-to-state  $N_2(A^3\Sigma_u^+)$  energy pooling reactions. II. The formation and quenching of  $N_2(B^3\Pi_g, v'=1-12)$ . *J. Chem. Phys.* 88, 6911–6921, 1988.
- Shemansky, D.E.  $A^3\Sigma_u^+$  molecules in the  $N_2$  afterglow. *J. Chem. Phys.* 64, 565–580, 1976.
- Sergienko, T.I., Ivanov, V.E. A new approach to calculate the excitation of atmospheric gases by auroral electron impact. *Ann. Geophys.* 11, 717–727, 1993.
- Slanger, T.G., Cosby, P.C.  $O_2$  spectroscopy below 5.1 eV. *J. Phys. Chem.* 92, 267–282, 1988.
- van Veen, N., Brewer, P., Das, P., Bersohn, R. Detection of the  $a^1\Pi_g(v'=0-1) \leftarrow X^1\Sigma_g^+(v''=0)$  transition in  $N_2$  by laser-induced fluorescence. *J. Chem. Phys.* 77, 4326–4329, 1982.
- Yankovskii, V.A. Electronic-vibrational relaxation of  $O_2(b^1\Sigma_g^+, v=1, 2)$  molecules in collisions with ozone atoms and molecules of oxygen. *Chem. Phys.* 10, 291–306, 1991 (in Russian).