

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/274092890>

Odorant-binding proteins display high affinities for behavioral attractants and repellents in the natural predator *Chrysopa pallens*

ARTICLE *in* COMPARATIVE BIOCHEMISTRY AND PHYSIOLOGY. PART A, MOLECULAR & INTEGRATIVE PHYSIOLOGY · MARCH 2015

Impact Factor: 1.97 · DOI: 10.1016/j.cbpa.2015.03.011 · Source: PubMed

READS

81

6 AUTHORS, INCLUDING:



Zhao-Qun Li

Nanjing Agricultural University

16 PUBLICATIONS 62 CITATIONS

[SEE PROFILE](#)



Shuai Zhang

Institute of Cotton Research (ICR), Chinese ...

19 PUBLICATIONS 91 CITATIONS

[SEE PROFILE](#)



Si-bao Wang

Nanjing Agricultural University

3 PUBLICATIONS 0 CITATIONS

[SEE PROFILE](#)



Shuang-Lin Dong

Nanjing Agricultural University

54 PUBLICATIONS 442 CITATIONS

[SEE PROFILE](#)



Odorant-binding proteins display high affinities for behavioral attractants and repellents in the natural predator *Chrysopa pallens*

Zhao-Qun Li^{a,b}, Shuai Zhang^a, Jun-Yu Luo^a, Si-Bao Wang^b, Shuang-Lin Dong^{b,*}, Jin-Jie Cui^{a,**}

^a State Key Laboratory of Cotton Biology, Institute of Cotton Research, Chinese Academy of Agricultural Sciences, Henan 455000, China

^b College of Plant Protection, Nanjing Agricultural University, Key Laboratory of Integrated Management of Crop Diseases and Pests, Ministry of Education, Nanjing 210095, China

ARTICLE INFO

Article history:

Received 22 January 2015

Received in revised form 12 March 2015

Accepted 16 March 2015

Available online 22 March 2015

Keywords:

Behavioral trial

Chrysopa pallens

Fluorescence binding assay

Odorant-binding protein

ABSTRACT

Chrysopa pallens is an important natural predator of various pests in many different cropping systems. Understanding the sophisticated olfactory system of insect antennae is crucial for studying the physiological bases of olfaction and could also help enhance the effectiveness of *C. pallens* in biological control. However, functional studies of the olfactory genes in *C. pallens* are still lacking. In this study, we cloned five odorant-binding protein (OBP) genes from *C. pallens* (CpalOBPs). Quantitative RT-PCR results indicated that the five CpalOBPs had different tissue expression profiles. Ligand-binding assays showed that farnesol, farnesene, cis-3-hexenyl hexanoate, geranylacetone, beta-ionone, octyl aldehyde, decanal, nerolidol ($K_i < 20 \mu\text{M}$), and especially 2-pentadecanone ($K_i = 1.19 \mu\text{M}$) and 2-hexyl-1-decanol ($K_i = 0.37 \mu\text{M}$) strongly bound to CpalOBP2. CpalOBP15 exhibited high binding affinities for beta-ionone, 2-tridecanone, trans-nerolidol, and dodecyl aldehyde. Behavioral trials using the 14 compounds exhibiting high binding affinities for the CpalOBPs revealed that nine were able to elicit significant behavioral responses from *C. pallens*. Among them, farnesene and its corresponding alcohol, farnesol, elicited remarkable repellent behavioral responses from *C. pallens*. Our study provides several compounds that could be selected to develop slow-release agents that attract/repel *C. pallens* and to improve the search for strategies to eliminate insect pests.

© 2015 Elsevier Inc. All rights reserved.

1. Introduction

A sophisticated olfactory system is a key physiological trait in insect behaviors, including host seeking, mating, oviposition, toxin avoidance, and negative taxis (Zhou, 2010). In insects, olfactory genes are expressed in and around olfactory receptor (OR) neurons, which are located at the sensilla in antennae (Schultze et al., 2013, 2014), mouthparts (Syed and Leal, 2007; Wasserman and Itagaki, 2003), and ovipositors (Anderson and Hallberg, 1990). Odorant-binding proteins (OBPs) comprise a family of small, water-soluble, extracellular proteins that are found in abundance within the sensillum lumen, and are thought to affect insect chemoreception by enhancing the solubility of exogenous odorant molecules and delivering them to the chemosensory receptors (Leal, 2012; Vogt, 2003, 2004; Zhou, 2010). Two models have been proposed for the mechanisms of OR activation (Leal, 2012). The first model suggests that a ligand per se activates its OR in moths (Leal et al., 2005) and mosquitoes (Wogulis et al., 2006). In contrast,

the second model hypothesizes that a pheromone-binding protein, such as Lush in *Drosophila melanogaster*, forms an OBP/odorant complex that directly activates the OR (Laughlin et al., 2008).

The first OBP was identified as a pheromone-binding protein in the giant moth *Antheraea polyphemus*, and it was specifically bound to a radioactive sex pheromone (Vogt and Riddiford, 1981). Thereafter, a large number of proteins and associated genes were identified in a wide range of insects in the orders of the Neoptera (Pelosi et al., 2005, 2006; Pelosi and Maida, 1995; Zhou, 2010). In addition, many papers have been published on functional and structural studies in Lepidoptera (Liu et al., 2013, 2015), Diptera (Lagarde et al., 2011a; Laughlin et al., 2008), Hymenoptera (Spinelli et al., 2012), and Hemiptera (He and He, 2014; He et al., 2011; Zhou et al., 2014). The typical characteristics of an insect OBP are six highly conserved cysteines, with a specific spacing among them, that form three disulfide bridges crucial for stabilizing the three-dimensional structure (Lagarde et al., 2011b; Tsitsanou et al., 2012). In addition to the classical OBPs (six conserved cysteines), some non-classical OBPs have been identified, including Plus-C OBPs (eight conserved cysteines), Minus-C OBPs (only four conserved cysteines), Dimer OBPs (two six-cysteine signatures), and Atypical OBPs (9–10 cysteines and a long C-terminus) (Hekmat-Scafe et al., 2002; Li et al., 2013a; Zhou et al., 2004).

The lacewing *Chrysopa pallens* (Neuroptera: Chrysopidae) is an important natural predator of various pests, such as aphids, coccids,

* Correspondence to: S.-L. Dong, College of Plant Protection, Nanjing Agricultural University, No. 1, Weigang, Nanjing 210095, China.

** Correspondence to: J.-J. Cui, Plant Protection Department, Institute of Cotton Research, Chinese Academy of Agricultural Sciences, No. 38, Huanghe Road, Anyang, Henan 455000, China.

E-mail addresses: sldong@njau.edu.cn (S.-L. Dong), cuijinjie@126.com (J.-J. Cui).

thrips, mites, planthoppers, and the eggs and young larvae of lepidopteran insects (Winterton and de Freitas, 2006), in many different cropping systems including maize, cotton, and rice (Bai et al., 2005; Brooks, 1994). In agreement with the feeding habits of larval and adult lacewings, aphid sex pheromone compounds, enantiomers of nepetalactol and nepetalactone [(1R,4aS,7S,7aR)-nepetalactol and (4aS,7S,7aR)-nepetalactone], were shown to attract lacewings and parasitoids (Boo et al., 1998; Hooper et al., 2002; Koczor et al., 2010).

To enhance the effectiveness of *C. pallens* in biological control using the “push–pull” strategy in which combinations of repellent and attractive stimuli are used to regulate the populations of insect pests and natural enemies (Cook et al., 2007), studies on olfactory gene function and olfactory behavior are essential. In our previous study, 14 OBPs were identified from the *C. pallens* transcriptome (Li et al., 2013b). Compared with the many available reports on functional and structural studies of olfactory genes in pests, such data from *C. pallens* are rare. Therefore, we cloned five OBP genes from *C. pallens*. Quantitative PCR (qPCR) results indicated that these five OBPs had different tissue expression profiles, suggesting different functions in chemoreception. The binding properties of the five OBPs with plant volatiles revealed that 14 compounds exhibited high binding affinities for the *C. pallens* OBPs (CpalOBPs), of which nine were able to elicit significant behavioral responses from *C. pallens*. Among them, farnesene and its corresponding alcohol, farnesol, elicited remarkable repellent behavioral responses from *C. pallens*. Our study provides several compounds that could be selected to develop slow-release agents that attract/repel *C. pallens* and to improve the search for strategies to eliminate insect pests.

2. Material and methods

2.1. Insect rearing and collection

The *C. pallens* individuals used in this study were collected from a cotton field at the Institute of Cotton Research (CAAS, Henan, China). The experimental insects were the offspring of a single female and were reared in a laboratory on *Acyrtosiphon pisum*. The rearing conditions were 25 ± 1 °C, 14-h/10-h light/dark cycle, and $65 \pm 5\%$ relative humidity. The pupae were kept in separate cages for eclosion, after which the adults were supplied with *A. pisum*. For cDNA cloning and qPCR, antennae, heads, thoraxes, abdomens, legs, and wings were collected from 3-day-old virgin female and male *C. pallens* individuals. The tissues were immediately frozen and stored at -80 °C until RNA isolation.

2.2. cDNA cloning, sequence alignment, and phylogenetic analysis

Total RNA was extracted from the antennae of *C. pallens* using TRIzol reagent (Invitrogen, Carlsbad, CA, USA), and cDNA templates were synthesized using a Reverse Transcription System (Promega, Madison, WI, USA) according to the manufacturer's instructions. The full-length sequences were confirmed by end-to-end PCR with specific primers designed using Primer Premier 5.0. The primer sequences are listed in Table S1. An amino acid sequence alignment of the corresponding OBPs was created using ClustalX 2.0 (Larkin et al., 2007), and visualized using Jalview 2.4.0 b2 (Waterhouse et al., 2009). Five CpalOBPs were chosen for phylogenetic tree construction along with the OBPs from other insects, which were downloaded from NCBI (<http://www.ncbi.nlm.nih.gov/>). A phylogenetic tree was constructed based on the amino acid sequences using the neighbor-joining algorithm implemented in MEGA6 (Tamura et al., 2011), with the default settings and 1000 bootstrap replicates. The sequences used in the phylogenetic tree are listed in Table S2.

2.3. Real-time qPCR and data analysis

Total RNA was isolated using an SV Total RNA Isolation System (Promega) according to the manufacturer's instructions. Single-stranded cDNA templates were synthesized using 1 µg of total RNA from various samples using a Reverse Transcription System (Promega) according to the manufacturer's instructions. qPCR was performed in a Mastercycler ep realplex (Eppendorf, Hamburg, Germany) with primers designed based on *C. pallens* nucleotide sequences using Beacon Designer 7.7 (Table S1). The *C. pallens* GTP-binding protein and ribosomal protein genes were used as reference genes. The expression levels of the tested mRNAs were determined using GoTaq qPCR Master Mix (Promega) according to the manufacturer's instructions and calculated using the comparative $2^{-\Delta\Delta CT}$ method (Pfaffl, 2001). Each reaction involved three technical replicates performed on three independent biological replicates.

2.4. Escherichia coli expression and purification of recombinant proteins

An *E. coli* expression system was used to express the CpalOBPs. Recombinant pET-32a(+)/CpalOBPs were constructed using an In-Fusion HD Cloning Kit (Clontech, Temecula, CA, USA) according to the manufacturer's instructions, and the recombinant plasmids were transformed into competent *E. coli* BL21 cells. Positive clones were incubated at 37 °C following a previously described protocol (Li et al., 2013a). Recombinant proteins in the supernatants were purified using HisTrap FF affinity columns (GE Healthcare, Piscataway, NJ, USA). The His-tags were cleaved from the proteins using 40 µl of enterokinase (Dingguo, Beijing, China) while loaded on the HisTrap FF columns at 37 °C for 10 h. After completion of the cleavage reaction, the recombinant proteins were eluted with phosphate-buffered saline. Finally, the eluted proteins were desalted using a HiTrap Desalting Kit (GE Healthcare), lyophilized, and stored at -80 °C until use.

2.5. Competitive fluorescence binding assay

Emission fluorescence spectra were measured on an F-4500 fluorescence spectrophotometer (Hitachi, Tokyo, Japan) in a right-angle configuration with a 1-cm light path quartz cuvette. The proteins were dissolved in 50 mM Tris–HCl buffer (pH 7.4), whereas all ligands used in the binding experiments were added as 1 mM methanol solutions.

Aliquots of 2 µM OBP solutions were titrated with 1 mM 1-N-phenyl naphthylamine (1-NPN) to a final concentration of 20 µM to measure the OBP affinities for the fluorescent probe. The 1-NPN fluorescent probe was excited at 337 nm, and emission spectra were recorded from 360 to 500 nm. OBP dissociation constants for 1-NPN were calculated from Scatchard plots of the binding data. The comparative binding of other ligands was measured using 2 µM 1-NPN, and the corresponding fluorescence intensities were recorded. Dissociation constants (K_i) of the OBPs for each ligand were calculated from the corresponding concentrations of ligands halving the initial fluorescence value of 1-NPN (IC_{50} values) based on the binding curves, using the following equation: $K_i = IC_{50} / (1 + C_{1-NPN} / K_{1-NPN})$. In this equation, C_{1-NPN} is the free concentration of 1-NPN, and K_{1-NPN} is the dissociation constant of the complex protein/1-NPN (Wei et al., 2008). Each reaction was repeated in triplicate.

2.6. Behavioral trials

The behavioral responses of *C. pallens* adults (5 days after emergence) to the 14 putative ligands for recombinant CpalOBP5, CpalOBP7, and/or CpalOBP10 were examined using a glass Y-tube olfactometer. The central tube was 25 cm (length) \times 30 mm (diameter), with arms of 20 cm (length) \times 30 mm (diameter), and an angle of 75° between the arms. Two filter paper strips were placed in the arms of the olfactometer; one held a control and the other held the test chemical.

A constant airflow (0.2 l/min), which was filtered through activated-charcoal and deionized water, was introduced into each arm of the olfactometer.

C. pallens individuals received 5 min to choose between the two arms of the Y-tube olfactometer, and the insect was then discarded. A response was identified when *C. pallens* walked into one arm of the olfactometer beyond 5 cm. After testing one insect, the arms were exchanged to avoid an asymmetric bias effect and the test compound was replaced with a new one (10 µl on filter paper; 10 mg/ml). To minimize the visual distraction for *C. pallens*, the Y-tube olfactometer was placed in darkness. At least 40 insects were used. Before behavioral measurements, the insects were starved for ~12 h. All tests were conducted at 24 ± 2 °C and $60 \pm 5\%$ relative humidity. All olfactometers used were washed with ethanol and dried at 80 °C for 5 min. The chi-squared test was used to analyze the behavioral assay data.

3. Results

3.1. Gene cloning and sequence analysis of CpalOBPs

Five OBP genes were cloned and designated *CpalOBP1*, *CpalOBP2*, *CpalOBP5*, *CpalOBP13*, and *CpalOBP15* (GenBank accession numbers KP082905, KP082903, KP403738, KP403734, and KP403736, respectively). The predicted open reading frames of *CpalOBP1*, *CpalOBP2*, *CpalOBP5*, *CpalOBP13*, and *CpalOBP15* had lengths of 450, 417, 408, 633, and 771 bp, respectively, which encoded 149, 138, 135, 210, and 256 amino acids, respectively. The initial 19, 19, 21, 20, and 16 amino acids of the respective sequences were predicted signal peptides (Fig. 1). The predicted protein molecular weights and isoelectric points of *CpalOBP1*, *CpalOBP2*, *CpalOBP5*, *CpalOBP13*, and *CpalOBP15* were 16.47 kDa and 5.36, 15.72 kDa and 4.6, 15.12 kDa and 4.89, 23.76 kDa and 5.29, and 29.63 kDa and 5.45, respectively. The phylogenetic tree of the five *CpalOBPs* and 35 OBPs from other insects indicated four possible protein subfamilies (Fig. 2). The results showed that *CpalOBP1* and *CpalOBP2* belonged to the Classic OBPs, *CpalOBP5* to the Minus-C OBPs, *CpalOBP13* to the Plus-C OBPs, and *CpalOBP15* to the Atypical OBPs.

3.2. Tissue expression profiles of the CpalOBPs

Tissue expression patterns can often provide functional information. The expression levels of the *CpalOBPs* in different adult tissues were detected by qPCR (Fig. 3). The *CpalOBPs* in our study displayed various tissue expression profiles. *CpalOBP1*, *CpalOBP2*, and *CpalOBP13* were highly expressed in the antennae of both sexes. Among these genes, *CpalOBP2* expression was restricted to the antennae, while *CpalOBP1* and *CpalOBP13* had low levels of expression in the other tissues examined. *CpalOBP5* was highly expressed in the wings. *CpalOBP15* was ubiquitously expressed in most of the tissues examined at relatively high levels.

3.3. In vitro expression and purification of the five CpalOBPs

The five *CpalOBPs* were successfully expressed as soluble proteins using a bacterial system. A His-tag affinity column was used to purify




Fig. 2. Phylogenetic tree of candidate OBPs from *C. pallens* and other species. Tcas: *Tribolium castaneum*; Dpon: *Dendroctonus ponderosae*; Bhor: *Batocera horsfieldi*; Aaeg: *Aedes aegypti*; Aalb: *Aedes albopictus*; Ham: *Helicoverpa armigera*; Sexi: *Spodoptera exigua*; Csup: *Chilo suppressalis*; Aips: *Agrotis ipsilon*; Dhou: *Dendrolimus houi*; Msex: *Manduca sexta*; Cqui: *Culex quinquefasciatus*; Asin: *Anopheles sinensis*; Llin: *Lygus lineolaris*; Dmel: *Drosophila melanogaster*; Asut: *Adelphocoris suturalis*; Alin: *Adelphocoris lineolatus*; Eher: *Euschistus heros*; Bmor: *Bombyx mori*; and Dple: *Danaus plexippus*. Values at nodes indicate the bootstrap percentages based on 1000 replicates, and branches with bootstrap values above 50% are marked. OBPs from *C. pallens* are highlighted in red. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

the proteins, followed by treatment with an enterokinase to remove the His-tag. More than 10 mg of each purified *CpalOBP* was obtained from each 1.0 L culture. The purified proteins were identified by SDS-PAGE (Fig. 4).

3.4. Ligand-binding assays of the five CpalOBPs

To investigate and compare the binding specificities of the five *CpalOBPs* with 87 ligands, fluorescence levels were measured as reported previously (Li et al., 2013a). The dissociation constants for *CpalOBP1*, *CpalOBP2*, *CpalOBP5*, *CpalOBP13*, and *CpalOBP15* combined with the fluorescent probe 1-NPN were 15.26, 4.08, 32.34, 12.72, and 6.70, respectively, based on the binding curves and Scatchard plots (Fig. 5A).




Fig. 1. Alignment of *CpalOBP* amino acid sequences. Predicted signal peptides are boxed, and conserved cysteines are labeled with red pentagrams. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)




Fig. 3. Relative mRNA expression levels of the five *CpalOBPs* in different adult tissues. The relative expression levels were normalized using GTP-binding protein and ribosomal protein. Error bars represent standard errors. H, heads; T, thoraxes; Ab, abdomens; L, legs; W, wings; A, antennae.

In a subsequent experiment, 1-NPN was used as the fluorescent reporter to determine the binding activities of the five *CpalOBPs* to different odorants. The IC_{50} and K_i values for all 87 compounds tested were calculated based on the binding curves (Fig. 5B and C). In general, *CpalOBP1*, *CpalOBP5*, and *CpalOBP13* displayed a weak binding ($K_i > 20 \mu M$) to most of the 87 ligands tested. However, there were 10 ligands that showed relatively higher binding affinities ($K_i < 20 \mu M$) for *CpalOBP2* and four ligands for *CpalOBP15* (Table 1). *CpalOBP15* exhibited high binding affinities for beta-ionone, 2-tridecanone, trans-nerolidol, and dodecyl aldehyde. Farnesol, farnesene, cis-3-hexenyl hexanoate, geranylacetone, beta-ionone, octyl aldehyde, decanal, nerolidol ($K_i < 20 \mu M$), and especially 2-pentadecanone ($K_i = 1.19 \mu M$) and 2-hexyl-1-decanol ($K_i = 0.37 \mu M$) bound strongly to *CpalOBP2*. Meanwhile, *CpalOBP1* only showed a binding affinity for phlorizin dehydrate ($K_i = 15.93 \mu M$).

3.5. Behavioral trials

The behavioral responses to the 14 compounds that exhibited high binding affinities ($K_i < 20 \mu M$) for the five *CpalOBPs* were examined in a Y-tube olfactometer. Nine of the 14 compounds were able to elicit behavioral responses in *C. pallens* (Fig. 6). *C. pallens* showed a significant attraction to cis-3-hexenyl hexanoate ($\chi^2 = 6.26$, $P < 0.05$), decanal ($\chi^2 = 8.26$, $P < 0.01$), and 2-hexyl-1-decanol ($\chi^2 = 13.02$, $P < 0.01$).

In contrast, *C. pallens* showed a significant aversion to 2-pentadecanone ($\chi^2 = 6.04$, $P < 0.05$), 2-tridecanone ($\chi^2 = 12.12$, $P < 0.01$), geranylacetone ($\chi^2 = 11.28$, $P < 0.01$), octyl aldehyde ($\chi^2 = 6.04$, $P < 0.05$), farnesol ($\chi^2 = 4.32$, $P < 0.05$), and farnesene ($\chi^2 = 8.26$, $P < 0.01$).

4. Discussion

C. pallens is an important natural enemy of pests in different agro-ecosystems (Bai et al., 2005; Brooks, 1994; Winterton and de Freitas, 2006). Understanding the olfactory behavior of *C. pallens*, including hunting, mating, and oviposition, can improve biological control in sustainable agriculture. However, most studies have focused on pest olfaction (Gu et al., 2013; Sparks et al., 2014; Vogt et al., 2009) and studies on natural enemy insects are scarce. Consequently, five *CpalOBPs* were cloned and functionally characterized in this study. The present findings could increase the understanding of the olfactory processes of *C. pallens* during hunting.

To date, four OBP subfamilies have been found in the insect, namely the Classic, Minus-C, Plus-C, and Atypical OBP subfamilies (Zhou, 2010). A phylogenetic tree of the five *CpalOBPs*, together with 27 OBPs from other insects, showed that the *CpalOBPs* were well clustered into the four distinct subfamilies, with *CpalOBP1* and *CpalOBP2* belonging to the Classic OBPs, *CpalOBP5* to the Minus-C OBPs, *CpalOBP13* to




Fig. 4. SDS-PAGE analyses showing the expression and purification of the recombinant *CpalOBPs*. M: molecular markers; CK and I: bacterial cells before and after induction by IPTG, respectively; P: purified protein after cleavage by thrombin.




Fig. 5. Ligand-binding assays of the CpalOBPs. (A) Binding curve for 1-NPN and Scatchard plot of CpalOBPs. (B) Binding curves of selected ligands. (C) Comparisons of binding affinities (indicated by $1/K_i$) of CpalOBPs with 14 components.

the Plus-C OBPs, and CpalOBP15 to the Atypical OBPs. These findings suggested that the five CpalOBPs play different roles in odorant recognition, and that rapid evolution has occurred through gene duplication.

To understand the physiological functions of the five CpalOBPs, we examined their tissue expression profiles in *C. pallens* adults. The qPCR data revealed that CpalOBP1, CpalOBP2, and CpalOBP13 were highly expressed in the antennae of both sexes. Antennae are the major chemosensory organs of insects. Therefore, the antennae-enriched genes may be involved in insect olfaction. Consistent with previous studies, CpalOBP5 was highly expressed in the wings (Gu et al., 2011; Hua et al., 2012, 2013), whereas CpalOBP15 was ubiquitously expressed in most of the tissues examined. Olfactory sensilla have been found on the wings and legs of some insects (Leal, 2012), but no such sensilla or chemosensory functions of the wings and legs have been described in *C. pallens*. The high expression of CpalOBP5 in the wing and ubiquitous expression of CpalOBP15 deserve further study.

A fluorescence competitive binding assay was used to determine the binding activities of the five CpalOBPs for 87 volatile compounds, including 37 from cotton (Yu et al., 2007), 10 from maize (Carroll et al., 2006; De Moraes et al., 1998; Itoh et al., 2002), seven from tobacco (De Moraes et al., 1998; Yan et al., 2005), 14 from faba bean (Webster et al., 2008), and the remainder from other plants (Obata et al., 1983). The results showed that CpalOBP2 had high binding capacities for 10 plant volatiles, suggesting that it plays a role in the transportation of plant volatiles. In particular, seven ligands, 2-hexyl-1-decanol, 2-pentadecanone, octyl aldehyde, nerolidol, volatiles from cotton and rice, farnesene and its corresponding alcohol, farnesol, showed a good affinity ($K_i < 10 \mu$ M) for CpalOBP2 only. These data indicate that CpalOBP2 is involved in detecting and recognizing the alarm pheromone in aphid species. CpalOBP15 exhibited high binding affinities for 2-Tridecanone, released by cauliflower leaf (Seenivasagan and Paul, 2011), and beta-ionone, a fragrant component of flowers and other tissues of many plants (Gora and Brud, 1983; Werkhoff et al.,

Table 1
Binding data for the recombinant CpalOBPs with different plant volatiles.

Ligands	OBP1		OBP2		OBP15	
	IC ₅₀	Ki	IC ₅₀	Ki	IC ₅₀	Ki
Cis-3-hexenyl hexanoate	>20	>20	16.63	11.90	>20	>20
2-Tridecanone	>20	>20	>20	>20	9.82	7.76
Farnesol	>20	>20	5.80	4.15	>20	>20
2-Pentadecanone	>20	>20	1.67	1.19	>20	>20
2-Hexyl-1-decanol	>20	>20	0.52	0.37	>20	>20
Geranylacetone	>20	>20	15.24	10.90	>20	>20
Farnesene	>20	>20	12.41	8.87	>20	>20
Beta-ionone	>20	>20	15.03	10.74	9.92	7.85
Octyl aldehyde	>20	>20	13.59	9.72	>20	>20
Decanal	>20	>20	19.80	14.16	>20	>20
Nerolidol	>20	>20	11.66	8.34	>20	>20
Trans-nerolidol	>20	>20	>20	>20	12.74	10.08
Fodecyl aldehyde	>20	>20	>20	>20	15.07	11.92
Phlorizin dehydrate	17.90	15.93	>20	>20	>20	>20

Notes: IC₅₀ values of <20 μM were obtained based on the binding curves. Other tested compounds (not listed) that have high Ki values (>20 μM) for all of the OBPs are beta-caryophyllene, alpha-phellandrene, (+/–)-alpha-pinene, (+)-beta-pinene, ocimene, (–)trans-caryophyllene, ethylbenzene, indole, naphthalene, cumene, tridecane, tetradecane, undecane, dodecane, tetradecane, octadecane, heptadecane, tridecane, benzyl alcohol, 1-hexanol, cis-3-hexen-1-ol, cis-2-hexen-1-ol, geraniol, (+/–)-linalool, eucalyptol, alpha-ionol, 2-heptanol, (+)-cedrol, linalool, trans-3-hexen-1, methyl anthranilate-ol, butyl formate, caproyl acetate, pentyl acetate, ethyl propionate, ethyl benzoate, octyl acetate, cis-3-hexenyl acetate, trans-2-hexenyl acetate, phenylacetaldehyde, beta-cyclocitral, (+)-carvone, damascenone, 6-methyl-5-hepten-2-one, 2-heptanone, (±)-camphor, acetophenone, hexyl butyrate, (+)-limonene oxide, (E3)-hexen-1-ol, camphene, (R)-(+)-limonene, phenethyl alcohol, 3-hexanol, cis-3-hexenol, benzyl acetate, phenethyl acetate, geranyl acetate, cis-3-hexenyl acetate, ethyl butyrate, trans-2-hexenyl butyrate, heptyl acetate, methyl salicylate, butyl acetate, nonyl acetate, isoamyl acetate, cis-3-hexenyl butyrate, benzaldehyde, hexanol, trans-2-hexenal, cis-3-hexenal, 3-hexanone, 2-hexanone, and undecanal.

1992). Meanwhile, CpalOBP1, CpalOBP5, and CpalOBP13 did not show strong binding affinities for the ligands tested, based on the Ki values (Ki > 20 μM). These differences in the binding properties are probably caused by different roles in chemoreception among the CpalOBPs.

Of the 14 compounds that exhibited high binding affinities (Ki < 20 μM) for the CpalOBPs, six elicited significant attractant behavioral responses and three elicited remarkable repellent behavioral responses from *C. pallens* in the behavioral trials. Among these nine volatiles, *C. pallens* showed a significant aversion behavior to farnesene




Fig. 6. Behavioral responses of *C. pallens* in a Y-tube olfactometer bioassay when given a choice between different odorant chemicals and hexane (control). The chi-squared test was used to determine significant differences in the numbers of insects choosing a particular odor, with the following levels of significance: **P* < 0.05 and ***P* < 0.01.

and its corresponding alcohol, farnesol. Since beta farnesene is the aphid alarm pheromone (Bowers et al., 1972), the aphid will leave an area containing farnesene, making it unnecessary for *C. pallens* to search the area. Thus, *C. pallens* might recognize and use the aphid alarm pheromone to improve its hunting efficiency for aphids. The compounds that elicited significant behavioral responses could be selected to develop slow-release agents to attract/repel *C. pallens* and to improve the search for strategies, such as the “push–pull” strategy, in which combinations of repellent and attractive stimuli are used to alter the populations of insect pests and natural enemies to eliminate insect pests (Cook et al., 2007). For future studies, RNAi or knockdowns of CpalOBPs are needed to confirm the roles of these compounds and their influence on the behavior of *C. pallens*, as are efficacy evaluations of the screened potential attractants/repellents in the field.

In this study, we cloned five CpalOBP genes from *C. pallens*. qPCR showed that the five genes had diverse expression profiles. Ligand-binding assays and behavioral trials showed that 14 compounds exhibited high binding affinities for the CpalOBPs, of which nine elicited significant behavioral responses from *C. pallens*. The nine compounds that elicited significant behavioral responses could be selected to develop slow-release agents to attract/repel *C. pallens* and to improve the search for strategies to eliminate insect pests.

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.cbpa.2015.03.011>.

Acknowledgment

This study was funded by a research grant from the Ministry of Agriculture of China (2014ZX08011-002).

References

Anderson, P., Hallberg, E., 1990. Structure and distribution of tactile and bimodal taste tactile sensilla on the ovipositor, tarsi and antennae of the flour moth, *Ephestia kuehniella* (Zeller) (Lepidoptera, Pyralidae). *Int. J. Insect Morphol. Embryol.* 19, 13–24.

Bai, Y.Y., Jiang, M.X., A, C.J., 2005. Effects of transgenic cry1Ab rice pollen on the oviposition and adult longevity of *Chrysoperla sinica* Tjeder. *Acta Phys. Sin.* 32 (3), 225–230 (In Chinese with English abstract).

Boo, K.S., Chung, I.B., Han, K.S., Pickett, J.A., J., W.L., 1998. Response of the lacewing *Chrysopa cognata* to pheromones of its aphid prey. *J. Chem. Ecol.* 24, 631–643.

Bowers, W.S., Nault, L.R., Webb, R.E., Dutky, S.R., 1972. Aphid alarm pheromone: isolation, identification, synthesis. *Science* 177, 1121–1122.

Brooks, S.J., 1994. A taxonomic review of the common green lacewing genus *Chrysoperla* (Neuroptera: Chrysopidae). *Bull. Brit. Mus. (Nat. Hist.) Entomol. Sin.* 63, 137–210.

Carroll, M.J., Schmelz, E.A., Meagher, R.L., Teal, P.E., 2006. Attraction of *Spodoptera frugiperda* larvae to volatiles from herbivore-damaged maize seedlings. *J. Chem. Ecol.* 32, 1911–1924.

Cook, S.M., Khan, Z.B., Pickett, J.A., 2007. The use of push–pull strategies in integrated pest management. *Annu. Rev. Entomol.* 52, 375–400.

De Moraes, C.M., Lewis, W.J., Pare, P.W., Alborn, H.T., Tumlinson, J.H., 1998. Herbivore-infested plants selectively attract parasitoids. *Nature* 393, 570–573.

Gora, J., Brud, W., 1983. Progress in synthesis of sensory important trace components of essential oils and natural flavours. *Nahrung* 27, 413–428.

Gu, S.-H., Wang, S.-P., Zhang, X.-Y., Wu, K.-M., Guo, Y.-Y., Zhou, J.-J., Zhang, Y.-J., 2011. Identification and tissue distribution of odorant binding protein genes in the lucerne plant bug *Adelphocoris lineolatus* (Goeze). *Insect Biochem. Mol. Biol.* 41, 254–263.

Gu, S.H., Zhou, J.J., Wang, G.R., Zhang, Y.J., Guo, Y.Y., 2013. Sex pheromone recognition and immunolocalization of three pheromone binding proteins in the black cutworm moth *Agrotis ipsilon*. *Insect Biochem. Mol. Biol.* 43, 237–251.

He, M., He, P., 2014. Molecular characterization, expression profiling, and binding properties of odorant binding protein genes in the whitebacked planthopper, *Sogatella furcifera*. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 174, 1–8.

He, P., Zhang, J., Liu, N.Y., Zhang, Y.N., Yang, K., Dong, S.L., 2011. Distinct expression profiles and different functions of odorant binding proteins in *Nilaparvata lugens* Stal. *PLoS One* 6, e28921.

Hekmat-Scafe, D.S., Scafe, C.R., McKinney, A.J., Tanouye, M.A., 2002. Genome-wide analysis of the odorant-binding protein gene family in *Drosophila melanogaster*. *Genome Res.* 12, 1357–1369.

Hooper, A.M., Donato, B., Woodcock, C.M., Park, J.H., Paul, R.L., Boo, K.S., Hardie, J., Pickett, J.A., 2002. Characterization of (1'R,4S,4aR,7S,7aR)-dihydronepetalactol as a semiochemical for lacewings, including *Chrysopa* spp. and *Peyerimhoffia gracilis*. *J. Chem. Ecol.* 28, 849–864.

Hua, J.F., Zhang, S., Cui, J.J., Wang, D.J., Wang, C.Y., Luo, J.Y., Lv, L.M., 2012. Identification and binding characterization of three odorant binding proteins and one chemosensory protein from *Apolygus lucorum* (Meyer-Dur). *J. Chem. Ecol.* 38, 1163–1170.

- Hua, J.F., Zhang, S., Cui, J.J., Wang, D.J., Wang, C.Y., Luo, J.Y., Lv, L.M., Ma, Y., 2013. Functional characterizations of one odorant binding protein and three chemosensory proteins from *Apolygus lucorum* (Meyer-Dur) (Hemiptera: Miridae) legs. *J. Insect Physiol.* 59, 690–696.
- Itoh, S., Kuwahara, S., Hasegawa, M., Kodama, O., 2002. Synthesis of the (17R)- and (17S)-isomers of volicitin, an elicitor of plant volatiles contained in the oral secretion of the beet armyworm. *Biosci. Biotechnol. Biochem.* 66, 1591–1596.
- Kocor, S., Szentkiralyi, F., Birkett, M.A., Pickett, J.A., Voigt, E., Toth, M., 2010. Attraction of *Chrysoperla carnea* complex and *Chrysopa* spp. lacewings (Neuroptera: Chrysopidae) to aphid sex pheromone components and a synthetic blend of floral compounds in Hungary. *Pest Manag. Sci.* 66, 1374–1379.
- Lagarde, A., Spinelli, S., Qiao, H., Tegoni, M., Pelosi, P., Cambillau, C., 2011a. Crystal structure of a novel type of odorant binding protein from *Anopheles gambiae*, belonging to the C+ class. *Biochem. J.* 437, 423–453.
- Lagarde, A., Spinelli, S., Tegoni, M., He, X., Field, L., Zhou, J.J., Cambillau, C., 2011b. The crystal structure of odorant binding protein 7 from *Anopheles gambiae* exhibits an outstanding adaptability of its binding site. *J. Mol. Biol.* 414, 401–412.
- Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A., McWilliam, H., Valentin, F., Wallace, I.M., Wilm, A., Lopez, R., Thompson, J.D., Gibson, T.J., Higgins, D.G., 2007. Clustal W and Clustal X version 2.0. *Bioinformatics* 23, 2947–2948.
- Laughlin, J.D., Ha, T.S., Jones, D.N., Smith, D.P., 2008. Activation of pheromone-sensitive neurons is mediated by conformational activation of pheromone-binding protein. *Cell* 133, 1255–1265.
- Leal, W.S., 2012. Odorant reception in insects: roles of receptors, binding proteins, and degrading enzymes. *Annu. Rev. Entomol.* 58, 373–463.
- Leal, W.S., Chen, A.M., Ishida, Y., Chiang, V.P., Erickson, M.L., Morgan, T.I., Tsuruda, J.M., 2005. Kinetics and molecular properties of pheromone binding and release. *Proc. Natl. Acad. Sci. U. S. A.* 102, 5386–5391.
- Li, Z.Q., Zhang, S., Luo, J.Y., Cui, J.J., Ma, Y., Dong, S.L., 2013a. Two Minus-C odorant binding proteins from *Helicoverpa armigera* display higher ligand binding affinity at acidic pH than neutral pH. *J. Insect Physiol.* 59, 263–272.
- Li, Z.Q., Zhang, S., Ma, Y., Luo, J.Y., Wang, C.Y., Lv, L.M., Dong, S.L., Cui, J.J., 2013b. First transcriptome and digital gene expression analysis in Neuroptera with an emphasis on chemoreception genes in *Chrysopa pallens* (Rambur). *PLoS One* 8, e67151.
- Liu, N.Y., Liu, C.C., Dong, S.L., 2013. Functional differentiation of pheromone-binding proteins in the common cutworm *Spodoptera litura*. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* 165, 254–262.
- Liu, N.Y., Yang, K., Liu, Y., Xu, W., Anderson, A., Dong, S.L., 2015. Two general-odorant binding proteins in *Spodoptera litura* are differentially tuned to sex pheromones and plant odorants. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* 180, 23–31.
- Obata, T., Koh, H.S., Kim, M., Fukami, H., 1983. Constituents of planthopper attractant in rice plant. *Appl. Entomol. Zool.* 18, 161–169.
- Pelosi, P., Maida, R., 1995. Odorant-binding proteins in insects. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 111, 503–514.
- Pelosi, P., Calvello, M., Ban, L., 2005. Diversity of odorant-binding proteins and chemosensory proteins in insects. *Chem. Senses* 30 (Suppl. 1), i291–i292.
- Pelosi, P., Zhou, J.J., Ban, L.P., Calvello, M., 2006. Soluble proteins in insect chemical communication. *Cell. Mol. Life Sci.* 63, 1658–1676.
- Pfaffl, M.W., 2001. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* 29, e45.
- Schultze, A., Pregitzer, P., Walter, M.F., Woods, D.F., Marinotti, O., Breer, H., Krieger, J., 2013. The co-expression pattern of odorant binding proteins and olfactory receptors identify distinct trichoid sensilla on the antenna of the malaria mosquito *Anopheles gambiae*. *PLoS One* 8, e69412.
- Schultze, A., Breer, H., Krieger, J., 2014. The blunt trichoid sensillum of female mosquitoes, *Anopheles gambiae*: odorant binding protein and receptor types. *Int. J. Biol. Sci.* 10, 426–437.
- Seenivasagan, T., Paul, A.V., 2011. Gas–chromatography and electroantennogram analysis of saturated hydrocarbons of cruciferous host plants and host larval body extracts of *Plutella xylostella* for behavioural manipulation of *Cotesia plutellae*. *Indian J. Exp. Biol.* 49, 375–386.
- Sparks, J.T., Bohbot, J.D., Dickens, J.C., 2014. The genetics of chemoreception in the labella and tarsi of *Aedes aegypti*. *Insect Biochem. Mol. Biol.* 48C, 8–16.
- Spinelli, S., Lagarde, A., Iovinella, I., Legrand, P., Tegoni, M., Pelosi, P., Cambillau, C., 2012. Crystal structure of *Apis mellifera* OBP14, a C-minus odorant-binding protein, and its complexes with odorant molecules. *Insect Biochem. Mol. Biol.* 42, 41–50.
- Syed, Z., Leal, W.S., 2007. Maxillary palps are broad spectrum odorant detectors in *Culex quinquefasciatus*. *Chem. Senses* 32, 727–738.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., Kumar, S., 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28, 2731–2739.
- Tsitsanou, K.E., Thireou, T., Drakou, C.E., Koussis, K., Keramioti, M.V., Leonidas, D.D., Eliopoulos, E., Iatrou, K., Zographos, S.E., 2012. *Anopheles gambiae* odorant binding protein crystal complex with the synthetic repellent DEET: implications for structure-based design of novel mosquito repellents. *Cell. Mol. Life Sci.* 69, 283–297.
- Vogt, R.G., 2003. Biochemical diversity of odor detection: OBPs, ODEs and SNMPs. *Insect Pheromone Biochemistry and Molecular Biology*, pp. 391–445.
- Vogt, R.G., 2004. Molecular Basis of Pheromone Detection in Insects.
- Vogt, R.G., Riddiford, L.M., 1981. Pheromone binding and inactivation by moth antennae. *Nature* 293, 161–163.
- Vogt, R.G., Miller, N.E., Litvack, R., Fandino, R.A., Sparks, J., Staples, J., Friedman, R., Dickens, J.C., 2009. The insect SNMP gene family. *Insect Biochem. Mol. Biol.* 39, 448–456.
- Wasserman, S.L., Itagaki, H., 2003. The olfactory responses of the antenna and maxillary palp of the fleshfly, *Neobellieria bullata* (Diptera: Sarcophagidae), and their sensitivity to blockage of nitric oxide synthase. *J. Insect Physiol.* 49, 271–280.
- Waterhouse, A.M., Procter, J.B., Martin, D.M., Clamp, M., Barton, G.J., 2009. Jalview Version 2—a multiple sequence alignment editor and analysis workbench. *Bioinformatics* 25, 1189–1191.
- Webster, B., Bruce, T., Dufour, S., Birkemeyer, C., Birkett, M., Hardie, J., Pickett, J., 2008. Identification of volatile compounds used in host location by the black bean aphid, *Aphis fabae*. *J. Chem. Ecol.* 34, 1153–1161.
- Wei, Y., Brandazza, A., Pelosi, P., 2008. Binding of polycyclic aromatic hydrocarbons to mutants of odorant-binding protein: a first step towards biosensors for environmental monitoring. *Biochim. Biophys. Acta* 1784, 666–671.
- Werkhoff, Peter, Brennecke, Stefan, Bretschneider, Wilfried, Giinter, Matthias, Hopp, Rudolf, Surburg, H., 1992. Chiro-specific analysis in essential oil, fragrance and flavor research. *Z. Lebensm. Unters. Forsch.* 196, 307–328.
- Winterton, S., de Freitas, S., 2006. Molecular phylogeny of the green lacewings (Neuroptera: Chrysopidae). *Aust. J. Entomol.* 45, 235–243.
- Wogulis, M., Morgan, T., Ishida, Y., Leal, W.S., Wilson, D.K., 2006. The crystal structure of an odorant binding protein from *Anopheles gambiae*: evidence for a common ligand release mechanism. *Biochem. Biophys. Res. Commun.* 339, 157–164.
- Yan, Z.G., Yan, Y.H., Wang, C.Z., 2005. Attractiveness of tobacco volatiles induced by *Helicoverpa armigera* and *Helicoverpa assulta* to *Campoplex chloridae*. *Chin. Sci. Bull.* 50, 1334–1341.
- Yu, H.L., Zhang, Y.J., Pan, W.L., Guo, Y.Y., Gao, X.W., 2007. Identification of volatiles from field cotton plant under different induction treatments. *Ying Yong Sheng Tai Xue Bao* 18, 859–864.
- Zhou, J.J., 2010. Odorant-binding proteins in insects. *Vitam. Horm.* 241–272 (2010/09/14 ed).
- Zhou, J.J., Huang, W., Zhang, G.A., Pickett, J.A., Field, L.M., 2004. “Plus-C” odorant-binding protein genes in two *Drosophila* species and the malaria mosquito *Anopheles gambiae*. *Gene* 327, 117–129.
- Zhou, S.S., Sun, Z., Ma, W., Chen, W., Wang, M.Q., 2014. De novo analysis of the *Nilaparvata lugens* (Stål) antenna transcriptome and expression patterns of olfactory genes. *Comp. Biochem. Physiol. D Genomics Proteomics* 9, 31–39.