

See discussions, stats, and author profiles for this publication at:  
<https://www.researchgate.net/publication/258637582>

# Ion-neutral complexes in gas-phase dissociations of protonated nitriles: n-C<sub>3</sub>H<sub>7</sub>CNH<sup>+</sup> and i-C<sub>3</sub>H<sub>7</sub>CNH<sup>+</sup>

ARTICLE *in* JOURNAL OF THE AMERICAN SOCIETY FOR MASS SPECTROMETRY · APRIL 1991

Impact Factor: 2.95 · DOI: 10.1016/1044-0305(91)80013-W · Source: PubMed

---

CITATIONS

5

---

READS

7

3 AUTHORS, INCLUDING:



Henryk Wincel

Instytut Chemii Fizycznej PAN

23 PUBLICATIONS 240 CITATIONS

SEE PROFILE

# Ion-Neutral Complexes in Gas-Phase Dissociations of Protonated Nitriles: $n\text{-C}_3\text{H}_7\text{CNH}^+$ and $i\text{-C}_3\text{H}_7\text{CNH}^+$

H. Wincel

Polish Academy of Science, Institute of Physical Chemistry, Warszawa, Poland

R. H. Fokkens and N. M. M. Nibbering

Institute of Mass Spectrometry, University of Amsterdam, Amsterdam, The Netherlands

(*J Am Soc Mass Spectrom* 1991, 2, 178–179)

In the last decade ion-neutral complexes have been proposed frequently as intermediates in the unimolecular dissociation of many organic ions in the gas phase [1]. For example, for the unimolecularly fragmenting ions  $n\text{-C}_3\text{H}_7\text{R}^+$  and  $i\text{-C}_3\text{H}_7\text{R}^+$ , where a.o.  $\text{R} = \text{CH}_2\text{O}$ ,  $\text{CH}_3\text{CHO}$  and  $(\text{CH}_3)_2\text{CO}$  [2],  $\text{CH}_2 = \text{NH}$  [3a] and  $\text{CH}_3\text{CH} = \text{NH}$  [3b] and  $\text{CO}$  [4], ion-neutral complexes have been invoked to rationalize the intramolecular isomerization of  $[n\text{-C}_3\text{H}_7^+ \dots \text{R}]$  into  $[i\text{-C}_3\text{H}_7^+ \dots \text{R}]$  before elimination of  $\text{R}$  or, in appropriate systems, the derived alkene( $\text{R-H}$ ) [2–4]. However, for  $\text{R} = \text{CO}$  the presumed involvement of such complexes is very difficult to verify experimentally, as in that case a possible interaction between the  $\text{C}_3\text{H}_7^+$  and  $\text{CO}$  species prior to their separation cannot be probed by  $\text{H/D}$  exchange.

We report here on the occurrence of ion-neutral complexes in the unimolecular fragmentation of the metastable ions  $n\text{-C}_3\text{H}_7\text{CNH}^+$ , **1**, and  $i\text{-C}_3\text{H}_7\text{CNH}^+$ , **2**, which are iso-electronic with  $n\text{-C}_3\text{H}_7\text{CO}^+$  and  $i\text{-C}_3\text{H}_7\text{CO}^+$ , respectively. The proton (deuteron) on nitrogen in these ions serves as a valuable tool to probe the interaction between the  $\text{C}_3\text{H}_7^+$  ion and the  $(\text{H,C,N})$  neutral before they separate from each other.

Experiments were performed by the mass-analyzed ion kinetic energy spectrometry technique [5] using a VG ZAB-2HF mass spectrometer. The ions **1** and **2** were prepared by chemical ionization using various protonating agents in combination with deuterium and  $^{13}\text{C}$ -labeling. Our results (Table 1) show that both **1** and **2** predominantly dissociate by loss of  $(\text{H,C,N})$  to give  $\text{C}_3\text{H}_7^+$  ( $m/z$  43). The data of  $^{13}\text{C}$ -labeling studies on  $n\text{-C}_3\text{H}_7\text{CNH}^+$  establish that the  $^{13}\text{C}$  atom is fully retained in the expelled  $(\text{H,C,N})$  neutral. This

implies that there is no intramolecular rearrangement via cyclic intermediates prior to dissociation as we observed for  $\text{CH}_3\text{CNH}^+$  [6].

The kinetic energy release, measured as  $T_{0.5}$  [5] from the metastable peak width at half-height, for loss of  $(\text{H,C,N})$  from **1** ( $T_{0.5} = 36.9$  meV) is higher than from **2** ( $T_{0.5} = 27.6$  meV). This effect is analogous to that found [4] for the  $n\text{-C}_3\text{H}_7\text{CO}^+$  and  $i\text{-C}_3\text{H}_7\text{CO}^+$  ions, and may be assigned to the isomerization of  $n\text{-C}_3\text{H}_7^+$  to  $i\text{-C}_3\text{H}_7^+$  before or during loss of  $(\text{H,C,N})$  from **1**. This process is expected to occur when the interspecies distance in  $[n\text{-C}_3\text{H}_7^+ \dots \text{CNH}]$  is lengthened to allow the isomerization (Ab initio calculations performed for  $n\text{-C}_4\text{H}_{10}^+$  predict that isomerization of  $n\text{-to-}i\text{-C}_3\text{H}_7^+$  occurs at a  $[n\text{-C}_3\text{H}_7^+ \dots \text{CH}_3]$  separation of ca. 2.1 Å.) [7]. Relevant to the occurrence of ion-neutral complexes during dissociation of ions **1** and **2** is the formation of abundant  $\text{C}_3\text{H}_6\text{D}^+$  ( $m/z$  44) from  $n\text{-C}_3\text{H}_7\text{CND}^+$  and  $i\text{-C}_3\text{H}_7\text{CND}^+$  (Table 1), which indicates that loss of  $(\text{H,C,N})$  is accompanied by a very large deuterium incorporation into the  $\text{C}_3\text{H}_7^+$  product ion in both cases.

All these results can be explained by a mechanism illustrated in Scheme 1 which involves the intermediate formation of an ion-neutral complex **2a**, the  $i\text{-C}_3\text{H}_7^+$  partner of which serves as a protonated reagent [the proton affinity  $\text{PA}(\text{HNC}) > \text{PA}(\text{C}_3\text{H}_6)$ , see [8a]] for  $\text{HNC}$  (**2a**  $\rightarrow$  **2b**). The extensive  $(\text{H,C,N})$ -to- $\text{C}_3\text{H}_7$   $\text{H/D}$  exchange suggests that the interspecies separation in the ion-neutral complex **2b** is large enough to allow a reorientation of the partners into a configuration which permits  $\text{D}^+$ -transfer from  $\text{HCND}^+$  to  $\text{CH}_3\text{CH} = \text{CH}_2$  [ $\text{PA}(\text{HCN}) < \text{PA}(\text{C}_3\text{H}_6)$ , see [8a]]. [ $\Delta H_f(\text{C}_3\text{H}_6\text{CN}) = 45$  kcal/mol is estimated on the basis of  $\Delta H_f(\text{C}_3\text{H}_6\text{CN}) = \Delta H_f(\text{C}_3\text{H}_7\text{CN}) - \Delta H_f(\text{H}) + \text{D}(\text{H-C}_3\text{H}_6\text{CN})$  (assuming  $\text{D}(\text{H-C}_3\text{H}_6\text{CN})$  is that of  $\text{D}(\text{H-CH}(\text{CH}_3)\text{CN}) \approx 90$  kcal/mol see [8b]) thermochemical data from ref. 8a].

This interpretation is in harmony with the low

Address reprint requests to N. M. M. Nibbering, Institute of Mass Spectrometry, University of Amsterdam, Nieuwe Achtergracht 129, 1018 WS Amsterdam, The Netherlands.


**Table 1.** Partial mass-analyzed ion kinetic energy spectra of the  $n\text{-C}_3\text{H}_7\text{CNX}^+$  and  $i\text{-C}_3\text{H}_7\text{CNX}^+$  ( $X = \text{H}$  or  $\text{D}$ ) metastable ions generated by the gas-phase reactions  $\text{MX}^+ + \text{C}_3\text{H}_7\text{CN} \rightarrow \text{C}_3\text{H}_7\text{CNX}^+ + \text{M}$

Initial structure	MX <sup>+</sup>	-H <sub>f</sub> <sup>a</sup> (kcal/mol)	Mass to charge ratio of daughter ions <sup>b</sup>									
			27	28	29	39	40	41	42	43	44	
n-C <sub>3</sub> H <sub>7</sub> CNH <sup>+</sup>	n-C <sub>3</sub> H <sub>7</sub> CN <sup>+</sup>	49 <sup>c</sup>	4	8		3	2	9	2	70		
	SH <sub>3</sub> <sup>+</sup>	23	7	6	2	4	2	11	1	58		
n-C <sub>3</sub> H <sub>7</sub> <sup>13</sup> CNH <sup>+</sup>	H <sub>3</sub> <sup>+</sup>	92	1		8			1	2	90		
	(CH <sub>3</sub> ) <sub>2</sub> OH <sup>+</sup>	2	2		6			2	2	88		
n-C <sub>3</sub> H <sub>7</sub> CND <sup>+</sup>	D <sub>3</sub> <sup>+</sup>	92			7			1	1	11	79	
	CD <sub>3</sub> CND <sup>+</sup>	2	3	2	7	4	1	4	7	15	50	
i-C <sub>3</sub> H <sub>7</sub> CNH <sup>+</sup>	i-C <sub>3</sub> H <sub>7</sub> CN <sup>+</sup>	49 <sup>c</sup>		2					1	97		
	SH <sub>3</sub> <sup>+</sup>	23	2	4		3		2	2	84		
i-C <sub>3</sub> H <sub>7</sub> CND <sup>+</sup>	D <sub>3</sub> <sup>+</sup>	92								5	95	
	CD <sub>3</sub> CND <sup>+</sup>	2			4	1		1	2	5	85	

<sup>a</sup> Nominal heat of reaction calculated by using the heats of formation taken from ref. 8a.

<sup>b</sup> Intensities of peaks are expressed as percentage of the sum of intensities of the peaks due to daughter ions.

<sup>c</sup> Nominal heat of reaction calculated by using the heats of formation taken from ref. 8a and the estimated heat of formation of  $\text{C}_3\text{H}_6\text{CN}$  from page 178.



stabilization energy of  $2b^1$ . The interaction of  $i\text{-C}_3\text{H}_7^+$  with HCN at moderate separations (3.5-5 Å) is expected to be in the range of 17-8 kcal/mol. Since the rearrangements of  $1a$  and  $2a$  into  $2c$  (Scheme 1) are exothermic,<sup>2</sup> it is very plausible that part of the complexes  $2c$  formed from both precursors will exceed the  $i\text{-C}_3\text{H}_7^+ \dots \text{NCH}$  bond strength leading to the product  $i\text{-C}_3\text{H}_7^+ + \text{HCN}$ . A fraction of the exothermicity appears as kinetic energies of these products. These are reflected in the  $T_{0.5}$  values discussed above, where the higher  $T_{0.5}$  value for  $1$  is in line with the higher energy content of complex  $2c$  when arising from  $1$ . Another argument for the occurrence of ion-neutral complexes in the dissociation of ions  $1$  and  $2$  is provided by the observed small yields of  $\text{HCNH}^+$  ( $m/z$  28) from unlabeled  $1$  and  $2$  as well as  $\text{H}^{13}\text{CNH}^+$  and  $(\text{H}_7\text{C}_3\text{N}_2\text{D})^+$  ( $m/z$  29) from the labeled precursors (Ta-

ble 1). They suggest that some of the ion-neutral complexes  $2b$  have sufficient energy to dissociate into the products  $\text{HCNH}^+ + \text{C}_3\text{H}_6$ .

Although ions  $1$  and  $2$  are isoelectronic with  $n\text{-C}_3\text{H}_7\text{CO}^+$  and  $i\text{-C}_3\text{H}_7\text{CO}^+$ , the extension of our findings as an indication for the involvement of ion-neutral complexes in the unimolecular dissociation of the latter ions seems not to be straightforward because of the large differences in the dipole moments of HNC ( $\mu = 2.65$  D), HCN ( $\mu = 2.98$  D) and CO ( $\mu = 0.10$  D) (see [9]).

## Acknowledgments

H. W. thanks The Netherlands Organization of Scientific Research (NWO) for providing a visitor's grant, and the Polish program CPBP-01.19 for financial support. R. H. F. and N. M. M. N. thank The Netherlands Organization of Scientific Research (SON/NWO) for continuous support.

## References

- (a) McAdoo, D. J. *Mass Spectrom. Rev.* **1988**, *7*, 363. (b) Morton, T. H. *Tetrahedron* **1982**, *38*, 3195.
- Bowen, R. D.; Williams, D. H. *Int. J. Mass Spectrom. Ion Phys.* **1979**, *29*, 47.
- (a) Bowen, R. D. *J. Chem. Soc. Perkin Trans. II.* **1980**, 1219. (b) Bowen, R. D. *J. Chem. Soc. Perkin Trans. II.* **1989**, 913.
- Williams, D. H.; Stapleton, B. J.; Bowen, R. D. *Tetrahedron Lett.* **1978**, 2919.
- Cooks, R. G.; Beynon, J. H.; Caprioli, R. M.; Lester, G. R. *Metastable Ions*; Elsevier, Amsterdam, 1973.
- Wintel, H.; Fokkens, R. H.; Nibbering, N. M. M. *Int. J. Mass Spectrom. Ion Processes* **1989**, *88*, 241.
- Takeuchi, T.; Yamamoto, M.; Nishimoto, K.; Tanaka, H.; Hirota, K. *Int. J. Mass Spectrom. Ion Phys.* **1983**, *52*, 139.
- (a) Lias, S. G.; Bartmess, J. E.; Liebman, J. F.; Holmes, J. L.; Levin, R. D.; Mallard, W. G. *J. Phys. Chem. Ref. Data* **1988**, *17*, supp. no. 1. (b) McMillen, D. F.; Golden, D. M. *Ann. Rev. Phys. Chem.* **1982**, *33*, 521.
- (a) *Handbook of Chemistry and Physics*, 64th ed.; West, R. C. Ed.; 1983-84; p. E59. (b) Meot-Ner (Mautner), M.; Karpas, Z.; Deakynne, C. A. *J. Am. Chem. Soc.* **1986**, *108*, 3913.

<sup>1</sup> Stabilization energies relative to the energies of the separated products for complexes  $2b$  and  $2c$  were estimated from the expression for ion-dipole interaction (see, for example, ref 2) using the dipole moments of 2.98 and 0.37 D for HCN and  $\text{CH}_3\text{CH}=\text{CH}_2$ , respectively, taken from ref 9. Complex  $2b$  is stabilized by 1.5 kcal/mol at a distance of 4 Å.

<sup>2</sup> The exothermicities of the rearrangement of  $1a$  and  $2a$  into  $2c$  calculated as the differences  $\Delta H_f(n\text{-C}_3\text{H}_7^+ + \text{HNC}) - \Delta H_f(i\text{-C}_3\text{H}_7^+ + \text{HCN}) = 36$  kcal/mol and  $\Delta H_f(i\text{-C}_3\text{H}_7^+ + \text{HNC}) - \Delta H_f(i\text{-C}_3\text{H}_7^+ + \text{HCN}) = 16$  kcal/mol, respectively. The heats of formation were taken from ref 8a.