

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/229160160>

Bioactive secondary metabolites from *Nigrospora* sp. LLGLM003, an endophytic fungus of the medicinal plant *Moringa oleifera* Lam

ARTICLE *in* WORLD JOURNAL OF MICROBIOLOGY AND BIOTECHNOLOGY (FORMERLY MIRCEN JOURNAL OF APPLIED MICROBIOLOGY AND BIOTECHNOLOGY) · MAY 2012

Impact Factor: 1.78 · DOI: 10.1007/s11274-012-1015-4 · Source: PubMed

CITATIONS

9

READS

339

5 AUTHORS, INCLUDING:



Chulong Zhang

Zhejiang University

35 PUBLICATIONS 453 CITATIONS

SEE PROFILE

Bioactive secondary metabolites from *Nigrospora* sp. LLGLM003, an endophytic fungus of the medicinal plant *Moringa oleifera* Lam.

J. H. Zhao · Y. L. Zhang · L. W. Wang ·
J. Y. Wang · C. L. Zhang

Received: 10 October 2011 / Accepted: 30 January 2012 / Published online: 12 February 2012
© Springer Science+Business Media B.V. 2012

Abstract An endophytic fungus was isolated from the root of the medicinal plant *Moringa oleifera* Lam. Based on analyzing the rDNA sequence, the fungus was identified as *Nigrospora* sp. This is the first report of the isolation of endophytic *Nigrospora* from *M. oleifera*. By bioassay-guided fractionation, four antifungal secondary metabolites were isolated from liquid cultures of the fungus *Nigrospora* sp. LLGLM003, and their chemical structures were determined to be griseofulvin (**1**), dechlorogriseofulvin (**2**), 8-dihydroramulosin (**3**) and mellein (**4**) on the basis of spectroscopic analyses. Compound **2**, **3** and **4** were isolated from *Nigrospora* sp. for the first time. In vitro antifungal assay showed that griseofulvin displayed clear inhibition of the growth of 8 plant pathogenic fungi. Dechlorogriseofulvin and mellein exhibited only weak antifungal activities, whereas 8-dihydroramulosin displayed no antifungal activities.

Keywords Endophytic fungus · *Nigrospora* sp. · Secondary metabolites · Antifungal activity

Introduction

An endophytic fungus is a microorganism that colonized living, internal tissues of its host plants, without causing any apparent symptoms of disease (Strobel 2002). There are approximately 300,000 different plant species on earth. Generally, each individual plant is host to many endophytes, thus providing a rich reservoir of endophytic fungi (Schulz et al. 2002; Strobel 2003; Strobel et al. 2004; Zhang et al. 2006). Many endophytic fungi have proven to be a rich source of bioactive secondary metabolites which are of interest for proper medicinal or agrochemical applications (Strobel and Daisy 2003).

Since 2002, we have started a survey of the diversity of fungal endophytes distributed in China and its secondary metabolites with antifungal and antitumor activities. We have isolated many bioactive metabolites, including cercosporamide along with new α -tetralone derivative (3*S*)-3, 6, 7-trihydroxy- α -tetralone from endophytic *Phoma* sp. of *Arisaema erubescens* (Wang et al. 2011), trichodermin from endophytic *Trichoderma* sp. of *Taxus mairei* (Chen et al. 2008), etc. This paper reports the isolation and bioactive metabolites of an endophytic fungus *Nigrospora* sp. LLGLM003 from the medicinal plant *Moringa oleifera* Lam. *M. oleifera*, a small deciduous tree, is the most widely known species of *Moringaceae* family and widely distributed in Asia Minor, Africa, the Indian subcontinent (Morton 1991). Traditionally, its root, bark, pod, leave are used in medicine for the treatment of a variety of human ailments whereas pods and young leaves are used as vegetables (Siddhuraju and Becker 2003). There are few

J. H. Zhao · Y. L. Zhang
College of Pharmaceutical Sciences, Zhejiang University of
Technology, Hangzhou 310014, People's Republic of China

J. H. Zhao
Institute of Pesticide and Environmental Toxicology, Zhejiang
University, Hangzhou 310058, People's Republic of China

L. W. Wang
Department of Pharmaceutical Science, College of Medicine and
Public Health Management, Hangzhou Normal University,
Hangzhou 310036, People's Republic of China

J. Y. Wang · C. L. Zhang (✉)
Key Laboratory of Molecular Biology of Crop Pathogens and
Insects, Ministry of Agriculture, Institute of Biotechnology,
Zhejiang University, Hangzhou 310058, People's Republic of
China
e-mail: clzhang@zju.edu.cn

studies on secondary metabolites of endophytic fungi isolated from *M. oleifera* (Ke et al. 2006). The crude extract of the endophytic fungus *Nigrospora* sp. LLGLM003 showed interesting antifungal activities. Investigation of the extract led to the isolation of four active compounds. Their structures were identified by spectroscopic data. Their antifungal activities were also evaluated.

Materials and methods

Isolation and purification of the endophytic fungi

The endophytic fungus strain LLGLM003 was isolated from a root of *M. oleifera*, collected in Xiamen municipality, Fujian Province, People's Republic of China. The general isolated procedures adopted followed the method described by preceding researchers (Gu 2005). Specifically, the roots of *M. oleifera* were washed softly with running tap water, then sterilized with 75% ethanol for 30 s and 1.5% sodium hypochlorite for 10 min, then rinsed in sterile water for three times. The sterilized root tissue were then cut into pieces of 0.5 cm length and six of them placed on a plate containing potato dextrose agar (PDA) supplemented with penicillin (100 µg/ml) and streptomycin sulphate (100 µg/ml) to prevent bacterial growth, and incubated in the dark at 25°C. After 1 week, the emerging hyphae from segments were cut and transferred into new PDA Petri dish for purification. The isolated endophytic fungi were stored by covering a culture on PDA slants with sterile liquid paraffin at 25°C.

Identification of the endophytic fungal strain LLGLM003

The fungal strain LLGLM003 was incubated in PDA medium at 25°C in darkness for 6–8 days. Mycelia were scraped from the plate and grounded to a powder under liquid nitrogen. Fungal DNA was extracted using the Multisource Genomic DNA Miniprep Kit (Axygen Bioscience, Inc.) following the manufacturer's instructions. Primers ITS1 (5'-TCCGTAGGTGAACCTGCGG-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-3') (White et al. 1990) were used to amplify a fragment of rDNA internal transcribed spacer (ITS) regions 1 and 2. Each 50-µl reaction contained 1× polymerase chain reaction (PCR) buffer, 2.5 mM Mg²⁺, 100 µM dNTPs, 0.5 µM each primer, 10 ng total DNA, and 2 U Taq polymerase. PCR thermocycling was carried out at 94°C for 3 min, followed by 35 cycles of 94°C for 40 s, 54°C for 50 s, and 72°C for 60 s, followed by a final extension at 72°C for 10 min. The PCR products were purified using a DNA gel extraction kit (Axygen Incorporation, China) and sequenced on an ABI 3730

sequencer (Applied Bio-systems, USA). The sequence was submitted to GenBank (accession no. JN387909) and analyzed by BLAST. Phylogenetic analysis was done by neighbor-joining in MEGA 4, using 1,000 bootstrap replicates.

Fermentation and metabolites isolation

The fungal strain *Nigrospora* sp. LLGLM003 was grown in 1,000 ml Erlenmeyer flasks containing 500 ml of potato dextrose broth (PDB), and then incubated in the dark on a rotary shaker (150 rpm; 25°C; 7 days). After filtration, the culture filtrate (totaling 20 l) was extracted three times with equal volumes of ethyl acetate (EtOAc). The EtOAc layer was evaporated to dryness (3.57 g) and mixed with 7 g silica gel, dried at 50°C, and then loaded on a silica gel column (40 × 1,000 mm) containing 210 g silica gel (200–300 mesh). The column was eluted with petroleum ether: ethyl acetate (4:1, 2:1, 1:1, and 1:2, v/v) firstly, and then the eluent was changed into chloroform: methanol (15:1, v/v). Five fractions were collected and labeled F1, F2, F3, F4, F5, which were detected in antifungal assay with *Botrytis cinerea* as target microorganism. Fractions F1, F4, F5 with antifungal activity were purified thereafter. The F1 fraction (140 mg) was purified by CC over SephadexLH-20 using a mixture of methanol and chloroform in a ratio of 1 to 1 as eluent to give compounds **3** (35 mg) and **4** (18 mg). The F4 (89 mg) fraction was subjected to the same purification method as the F1 fraction to give compound **2** (43 mg). The F5 (500 mg) fraction was subjected to crystallization in petroleum ether: ethyl acetate (2:1, v/v) to give compound **1** (450 mg).

Spectral measurements

To elucidate the chemical structures of four antifungal compounds, they were analyzed using mass and nuclear magnetic resonance (NMR) spectrometry. Mass spectra were recorded on a Bruker Esquire 3000^{plus} mass spectrometer. All NMR data were collected on a Bruker AMX-500(500 MHz) NMR spectrometer using TMS as internal standard, the chemical shifts were expressed in δ (ppm) and coupling constants *J* in Hz. Melting points were determined on a Beijing X4 micro melting point apparatus and were uncorrected. Silica gel (200–300 mesh) for column chromatography was produced by Qingdao Marine Chemical Factory, Qingdao, China. SephadexLH-20 was produced by Pharmacia Biotech, Sweden.

Compound 1

Griseofulvin, white solid, molecular formula C₁₇H₁₇ClO₆; m.p. 221–223°C; ESI-MS (*m/z*): 353 [*M* + *H*]⁺; ¹H NMR

(500 MHz, CDCl_3 , J in Hz): δ 6.14 (1H, s, 5-H), δ 5.55 (1H, s, 2'-H), δ 4.04 (3H, s, 4-OCH₃), δ 4.04 (3H, s, 6-OCH₃), δ 3.63 (3H, s, 1'-OCH₃), δ 3.04 (1H, dd, J = 16.7 Hz, 4.6, 4'-H), δ 2.86 (1H, m, 5'-H), δ 2.44 (1H, dd, J = 16.0 Hz, 3.8, 4'-H), δ 0.97 (3H, d, J = 6.8 Hz, 5'-CH₃); ^{13}C NMR (125 MHz, CDCl_3): δ 197.1 (C-3'), δ 192.5 (C-3), δ 169.5 (C-7a), δ 164.6 (C-6), δ 170.8 (C-1'), δ 157.7 (C-4), δ 105.1 (C-3a), δ 104.8 (C-2'), δ 97.2 (C-7), δ 90.7 (C-2), δ 89.4 (C-5), δ 57.0 (4-OCH₃), δ 56.7 (1'-OCH₃), δ 56.4 (6-OCH₃), δ 40.0 (C-4'), δ 36.4 (C-5'), δ 14.2 (5'-CH₃). The ^1H NMR and ^{13}C NMR data are consistent with the literature reported (Park et al. 2005).

Compound 2

Dechlorogriseofulvin, white solid, molecular formula $\text{C}_{17}\text{H}_{18}\text{O}_6$; m.p. 179–181°C; ESI-MS (m/z): 319 $[\text{M} + \text{H}]^+$ and 336 $[\text{M} + \text{NH}_4]^+$; ^1H NMR (500 MHz, CDCl_3 , J in Hz): δ 6.24 (1H, d, J = 1.8 Hz, 5-H), δ 6.05 (1H, d, J = 1.7 Hz, 7-H), δ 5.55 (1H, s, 2'-H), δ 3.91 (3H, s, 1'-OCH₃), δ 3.90 (3H, s, 6-OCH₃), δ 3.63 (3H, s, 4-OCH₃), δ 3.05 (1H, dd, J = 16.8 Hz, 13.4, 4'-H), δ 2.75 (1H, m, 5'-H), δ 2.41 (1H, dd, J = 16.8 Hz, 4.7, 4'-H), δ 0.97 (3H, d, J = 6.7 Hz, 5'-CH₃); ^{13}C NMR (125 MHz, CDCl_3): δ 197.4 (C-3'), δ 192.5 (C-3), δ 176.0 (C-7a), δ 171.3 (C-6), δ 170.3 (C-1'), δ 159.1 (C-4), δ 104.7 (C-3a), δ 104.3 (C-2'), δ 93.3 (C-7), δ 89.9 (C-2), δ 88.5 (C-5), δ 57.0 (4-OCH₃), δ 56.6 (1'-OCH₃), δ 56.1 (6-OCH₃), δ 40.0 (C-4'), δ 36.5 (C-5'), δ 14.2 (5'-CH₃). The ^1H NMR and ^{13}C NMR data were identical with literature reported (Park et al. 2005).

Compound 3

8-dihydroramulosin, white solid, molecular formula $\text{C}_{10}\text{H}_{16}\text{O}_3$; m.p. 142–144°C; ESI-MS (m/z): 185 $[\text{M} + \text{H}]^+$ and 207 $[\text{M} + \text{Na}]^+$; ^1H NMR (500 MHz, CDCl_3 , J in Hz): δ 4.39 (1H, m, 3-H), δ 2.23 (1H, ddd, J = 14.1 Hz, 6.4, 3.8, 4-H), δ 1.23 (1H, m, 4-H), δ 2.13 (1H, m, 4a-H), δ 1.61 (1H, brd, J = 12.7 Hz, 5-H), δ 0.92 (1H, qd J = 13.0 Hz, 3.2, 5-H), δ 1.70 (1H, m, 6-H), δ 1.23 (1H, m, 6-H), δ 1.87 (1H, dt, J = 13.0 Hz, 4.0, 7-H), δ 1.72 (1H, m, 7-H), δ 3.61 (1H, dt, J = 11.3 Hz, 5.2, 8-H), δ 2.95 (1H, brt, J = 5.3 Hz, 8a-H), δ 1.31 (3H, d, J = 6.2 Hz, 9-H); ^{13}C NMR (125 MHz, CDCl_3): δ 174.9 (C-1), δ 73.4 (C-3), δ 36.5 (C-4), δ 31.9 (C-4a), δ 30.9 (C-5), δ 23.4 (C-6), δ 32.8 (C-7), δ 70.1 (C-8), δ 44.2 (C-8a), δ 20.6 (C-9). The ^1H NMR and ^{13}C NMR data are consistent with the literature reported (Stierle et al. 1998).

Compound 4

Mellein, white solid, molecular formula $\text{C}_{10}\text{H}_{10}\text{O}_3$; m.p. 55–56°C; ESI-MS (m/z): 179 $[\text{M} + \text{H}]^+$; ^1H NMR

(500 MHz, CDCl_3 , J in Hz): δ 11.03 (1H, s, OH), δ 7.41 (1H, dd, J = 8.5 Hz, 0.5, 6-H), δ 6.89 (1H, d, J = 8.5 Hz, 5-H), δ 6.69 (1H, dd, J = 0.5 Hz, 8.5, 7-H), δ 4.73 (1H, m, 3-H), δ 2.93 (2H, d, J = 7.5 Hz, 4-H), δ 1.53 (3H, d, J = 6.0 Hz, Me); ^{13}C NMR (125 MHz, CDCl_3): δ 169.9 (C-1), δ 116.2 (C-7), δ 162.1 (C-8), δ 108.2 (C-8a), δ 139.3 (C-4a), δ 117.9 (C-5), δ 136.1 (C-6), δ 76.1 (C-3), δ 34.6 (C-4), δ 20.8 (3-Me). The ^1H NMR and ^{13}C NMR data are in full agreement with literature reported (Dimitriadis et al. 1997).


Antifungal assay

The four metabolites dissolved in methanol were tested for fungistatic activity in vitro against 8 plant pathogenic fungi: *B. cinerea*, *Colletotrichum orbiculare*, *Fusarium oxysporum* f.sp. *cucumerinum*, *Fusarium oxysporum* f.sp. *melonis*, *Pestalotia diospyri*, *Pythium ultimum*, *Rhizoctonia solani* and *Sclerotinia sclerotiorum*. To this end, a series of different concentrations of the metabolite were diluted with melted PDA medium and poured into a Petri dish. After cooling, agar discs (5 mm in diameter) of test fungi were placed at the center of plates containing the fungistatic compound of 100, 50, 25, 10, 5, 2.5, and 1.25 $\mu\text{g}/\text{ml}$ concentrations. Three replicate plates of each concentration for each fungus were incubated in the dark at 25°C for all test fungi. Plates containing media mixed with isometric methanol were included as control. The diameter of the growing colony was measured at least three times during the following days, and plotted against the incubation time. The part of the curve which showed a linear relationship was used to calculate the growth rate (millimeter per day). This was then plotted against the respective metabolite concentration. The effective concentrations (EC_{50}) are defined as the concentration of the test compound that allows the test organism to grow at 50% of its rate under the same conditions in the absence of the metabolite.

Results and discussion

In order to identify the fungus isolated from the root of miracle trees (*M. oleifera*), its internal transcribed spacer 1 (ITS1), 5.8S ribosomal RNA gene, and internal transcribed spacer 2 (ITS2) were sequenced and submitted to BLASTN at the NCBI nucleotide database. It was showed that similar sequences belong to *Nigrospora*. We chose some sequences located in *Trichosphaeriales* also with one from *Achroceratosphaeria potamia* as the out-group to be aligned with that of LLGLM003 using ClustalX1.8. The alignment was then subjected to neighbor-joining analysis, placing LLGLM003 within the clade including *Nigrospora sphaerica* and *Nigrospora oryzae* (Fig. 1), indicating that

Fig. 1 Neighbor-joining tree based on ITS sequence



this isolate LLGLM003 is a *Nigrospora* sp. closely related to these taxa.

The *Nigrospora* species are common endophytes in plants, previously isolated from *Taxus globosa* (Ruiz-Sanchez et al. 2010), *Rhizophora mucronata* (Xu et al. 2011), *Azadirachta indica* (Wu et al. 2009), *Smilax sonchifolius* (Lopes and Pupo 2011). The genus *Nigrospora* was also a rich source of biologically active secondary metabolites, such as phytotoxic and antibacterial nigrosporins (Tanaka et al. 1997), herbicidal lactones (Fukushima et al. 1998), phomalactone (Kim et al. 2001) and phytoxin epoxyxerohilone (Cutler et al. 1991). This is the first report of the isolation of endophytic *Nigrospora* from *M. oleifera*.

The endophytic fungus *Nigrospora* sp. LLGLM003 was selected for chemical and biological investigations because of the strong antifungal activity of the crude extract against the plant pathogenic fungi *B. cinerea* (EC_{50} = 6.09 μ g/

ml). The EtOAc extract of the fermentation broth was isolated and purified repeatedly by silica gel, SephadexLH-20 column and re-crystallization to yield four compounds (Fig. 2), which were identified as griseofulvin (1), dechlorogriseofulvin (2), 8-dihydramulosin (3), mullein (4), respectively according to mass and NMR spectral data. So far as we know, compounds 2–4 were isolated from *Nigrospora* sp. for the first time.

The antifungal efficacy experiments of compounds 1–4 were tested in vitro against 8 plant pathogenic fungi and the results were listed in Table 1. Among them, compound 1 exhibited strong activity against *B. cinerea* and *C. orbiculare* with the EC_{50} of 0.20 and 0.49 μ g/ml, respectively, and moderate activity against other test fungi. Compound 2 displayed weak activity against *B. cinerea*, *R. solani*, *C. orbiculare*, and *S. sclerotiorum* with the EC_{50} of 40.0, 43.9, 81.7 and 28.5 μ g/ml, respectively, and inactive to other test fungi. But compound 3 showed inactive to all test

Fig. 2 Structure of the compounds 1–4




Table 1 The antifungal activities of compounds **1–4** obtained from *Nigrospora* sp

Test plant pathogenic fungi	EC ₅₀ (μg/ml)			
	1	2	3	4
<i>Botrytis cinerea</i>	0.20	40.0	>100	49.2
<i>Colletotrichum orbiculare</i>	0.49	81.7	>100	>100
<i>Fusarium oxysporum</i> f.sp. <i>cucumerinum</i>	4.08	>100	>100	>100
<i>Fusarium oxysporum</i> f.sp. <i>melonis</i>	8.69	>100	>100	>100
<i>Pestalotia diospyri</i>	42.7	>100	>100	>100
<i>Pythium ultimum</i>	6.08	>100	>100	>100
<i>Rhizoctonia solani</i>	17.6	43.9	>100	>100
<i>Sclerotinia sclerotiorum</i>	5.78	28.5	>100	>100

fungi and compound **4** only exhibited weak activity against *B. cinerea* with the EC₅₀ of 49.2 μg/ml.

Griseofulvin (**1**) has been used as an antifungal antibiotic for the treatment of diseases of humans caused by pathogenic fungus (Gupta et al. 1997; Knasmüller et al. 1997) and inflammatory diseases (Asahina et al. 2001). Many fungus of *Penicillium* sp. (Mantle et al. 1984), *Aspergillus versicolor* (Kingston et al. 1976) and *Xylaria* sp. (Park et al. 2005) have presently been considered as typical griseofulvin producing fungi. Although chemical synthesis of griseofulvin was reported by former researchers (Stork and Tomasz 1962; Danishefsky and Walker 1979), it is economically infeasible. Therefore, griseofulvin is commercially produced by fermentation processes instead of by chemical synthesis (Dasu et al. 2002a; Dasu et al. 2002b). In this aspect, the discovery of more fungal species that can produce griseofulvin is very important. Dechlorogriseofulvin (**2**) showed weak antifungal activity and was previously obtained from the endophytic fungi *Penicillium janthinellum* (Xue et al. 2006) and *Xylaria* sp. (Cafêu et al. 2005; Park et al. 2005).

Both 8-dihydroramulosin (**3**) and mellein (**4**) were isocoumarin derivatives and widely distributed in fungi. 8-dihydroramulosin (**3**) was formerly reported as metabolite of the endophyte *Canoplea elegantula* (Findlay et al. 1995) and *Botrytis* sp. (Stierle et al. 1998) and toxic to spruce budworm larvae (Findlay et al. 1995). Mellein (**4**) was previously isolated from many endophytic fungi, including *Botryosphaeria rhodina* (Rukachaisirikul et al. 2009), *Botryosphaeria mamane* (Pongcharoen et al. 2007), *Prumnopitys andina* (Schmeda-Hirschmann et al. 2005), *Plectrophomella* sp., *Pezicula livida* and *Cryptosporiopsis malicorticis* (Krohn et al. 1997). The biological properties of mellein include: antibacterial activity (Krohn et al. 1997), antifungal activity (Höller et al. 1998), phytotoxic activity (Parisi et al. 1993) and bioactivity against the protease of hepatitis C virus (HCV) (Dai et al. 2001).

In conclusion, this is the first report on the isolation of endophytic *Nigrospora* from *M. oleifera*. The compounds from the strain LLGLM003 displayed strong or weak antifungal activity, which indicated that endophytic *Nigrospora* are promising sources of natural bioactive and novel metabolites and with great potential for further study.

Acknowledgments This work was supported by Special Fund for Agro-scientific Research in the Public Interest (200903052) to Chulong Zhang.

References

- Asahina A, Tada Y, Nakamura K, Tamaki K (2001) Griseofulvin has a potential to modulate the expression of cell adhesion molecules on leukocytes and vascular endothelial cells. *Int Immunopharmacol* 1(1):75–83
- Cafêu MC, Silva GH, Teles HL, Bolzanie VS, Araújo AR, Young MCM, Pfening LH (2005) Antifungal compounds of *Xylaria* sp., an endophytic fungus isolated from *Palicourea marcgravii* (Rubiaceae). *Quim Nova* 28(6):991–995
- Chen SY, Zhang CL, Chen YZ, Lin FC (2008) Trichodermin (4β-acetoxy-12, 13-epoxytrichothec-9-ene). *Acta Cryst E* 64:o702
- Cutler HG, Hoogsteen K, Littrell RH, Arison BH (1991) Epox-yexserohilone, a novel metabolite from *Nigrospora Sphaerica*. *Agric Biol Chem* 55(8):2037–2042
- Dai JR, Carté BK, Sidebottom PJ, SekYew AL, Ng SB, Huang YC, Butler MS (2001) Circumdatin G, a new alkaloid from the fungus *Aspergillus ochraceus*. *J Nat Prod* 64(1):125–126
- Danishefsky S, Walker FJ (1979) A stereospecific synthesis of griseofulvin. *J Am Chem Soc* 101(23):7018–7020
- Dasu V, Pandan VT, Chidambaram M (2002a) Development of medium for griseofulvin production: part I. Screening of medium constituents using the Plackett-Burman experimental design. *J Microbiol Biotechnol* 12(3):355–359
- Dasu V, Panda VT, Chidambaram M (2002b) Development of medium for griseofulvin production: part II. Optimization of medium constituents using central composite design. *J Microbiol Biotechnol* 12(3):360–366
- Dimitriadis C, Gill M, Harte MF (1997) The first stereospecific approach to both enantiomers of mellein. *Tetrahedron Asym* 8(13):2153–2158
- Findlay JA, Buthelezi S, Lavoie R, Peña-Rodríguez L, Miller JD (1995) Bioactive isocoumarins and related metabolites from conifer endophytes. *J Nat Prod* 58(11):1759–1766
- Fukushima T, Tanaka M, Gohbara M, Fujimori T (1998) Phytotoxicity of three lactones from *Nigrospora sacchari*. *Phytochemistry* 48(4):625–630
- Gu W (2005) Bioactive metabolites from *Alternaria brassicicola* ML-P08, an endophytic fungus residing in *Malus halliana*. *World J Microbiol Biotechnol* 25(9):1677–1683
- Gupta AK, Sibbald RG, Lynde CW, Hull PR, Prussick R, Shear NH, De Doncker P, Daniel CR, Elewski BE (1997) Onychomycosis in children: Prevalence and treatment strategies. *J Am Acad Dermatol* 36(3):395–402
- Höller U, König GM, Wright AD (1998) Three New Metabolites from Marine-Derived Fungi of the Genera *Coniothyrium* and *Microsphaeropsis*. *J Nat Prod* 62(1):114–118
- Ke Y, Chen Z, Mao JB, Hang XM, Zeng SR (2006) Studies of isolation of endophytic fungi from *Moringa* sp. and antibiotic active substance produced by the *Aspergillus* sp. ly14. *J Hunan Agric University (Natural Sciences)* 32(5):521–523(in Chinese)

- Kim JC, Choi GJ, Park JH, Kim HT, Cho KY (2001) Activity against plant pathogenic fungi of phomalactone isolated from *Nigrospora sphaerica*. *Pest Manag Sci* 57(6):554–559
- Kingston DGI, Chen PN, Vercellotti JR (1976) Metabolites of *Aspergillus-Versicolor*: 6,8-di-O-methylnidurufin, griseofulvin, dechlorogriseofulvin, and 3,8-dihydroxy-6-methoxy-1-methylxanthone. *Phytochemistry* 15(6):1037–1039
- Knasmüller S, Parzefall W, Helma C, Kassie F, Ecker S, Schulte-Hermann R (1997) Toxic effects of griseofulvin: disease models, mechanisms, and risk assessment. *Crit Rev Toxicol* 27(5):495–537
- Krohn K, Bahramsari R, Flörke U, Ludewig K, Kliche-Spory C, Michel A, Aust HJ, Draeger S, Schulz B, Antus S (1997) Dihydroisocoumarins from fungi: Isolation, structure elucidation, circular dichroism and biological activity. *Phytochemistry* 45(2):313–320
- Lopes AA, Pupo MT (2011) Biosynthesis of Aphidicolin Proceeds via the Mevalonate Pathway in the Endophytic Fungus *Nigrospora sphaerica*. *J Braz Chem Soc* 22(1):80–85
- Mantle PG, Laws I, Tan MJL, Tizard M (1984) A novel process for the production of penitrem mycotoxins by submerged fermentation of *Penicillium-Nigricans*. *J Gen Microbiol* 130:1293–1298
- Morton JF (1991) The horseradish tree, *Moringa pterygosperma* (Moringaceae): a boon to arid lands? *Econ Bot* 45(3):318–333
- Parisi A, Piattelli M, Tringali C, Di San Lio GM (1993) Identification of the phytotoxin mellein in culture fluids of *Phoma Tracheiphila*. *Phytochemistry* 32(4):865–867
- Park JH, Choi GJ, Lee HB, Kim KM, Cho KY, Jung HS, Lee SW, Jang KS, Cho KY, Kim JC (2005) Griseofulvin from *Xylaria* sp strain F0010, an endophytic fungus of *Abies holophylla* and its antifungal activity against plant pathogenic fungi. *J Microbiol Biotechnol* 15(1):112–117
- Pongcharoen W, Rukachaisirikul V, Phongpaichit S, Sakayaroj J (2007) A new dihydrobenzofuran derivative from the endophytic fungus *Botryosphaeria mamane* PSU-M76. *Chem Pharm Bull* 55(9):1404–1405
- Ruiz-Sanchez J, Flores-Bustamante ZR, Dendooven L, Favela-Torres E, Soca-Chafre G, Galinder-Mayer J, Flores-Cotera LB (2010) A comparative study of Taxol production in liquid and solid-state fermentation with *Nigrospora* sp a fungus isolated from *Taxus globosa*. *J Appl Microbiol* 109(6):2144–2150
- Rukachaisirikul V, Arunpanichlert J, Sukpondma Y, Phongpaichit S, Sakayaroj J (2009) Metabolites from the endophytic fungi *Botryosphaeria rhodina* PSU-M35 and PSU-M114. *Tetrahedron* 65(51):10590–10595
- Schmeda-Hirschmann G, Hormazabal E, Astudillo L, Rodriguez J, Theoduioz C (2005) Secondary metabolites from endophytic fungi isolated from the Chilean gymnosperm *Prumnopitys andina* (Lleuque). *World J Microbiol Biotechnol* 21(1):27–32
- Schulz B, Boyle C, Draeger S, Römmert AK, Krohn K (2002) Endophytic fungi: a source of novel biologically active secondary metabolites. *Mycol Res* 106(9):996–1004
- Siddhuraju P, Becker K (2003) Antioxidant properties of various solvent extracts of total phenolic constituents from three different agroclimatic origins of drumstick tree (*Moringa oleifera* Lam.). *J Agric Food Chem* 51(8):2144–2155
- Stierle DB, Stierle AA, Kunz A (1998) Dihydromulose from *Botrytis* sp. *J Nat Prod* 61(10):1277–1278
- Stork G, Tomasz M (1962) A stereospecific total synthesis of griseofulvin. *J Am Chem Soc* 84(2):310–312
- Strobel GA (2002) Rainforest endophytes and bioactive products. *Crit Rev Biotechnol* 22(4):315–333
- Strobel GA (2003) Endophytes as sources of bioactive products. *Microbes Infect* 5(6):535–544
- Strobel G, Daisy B (2003) Bioprospecting for microbial endophytes and their natural products. *Microbiol Mol Biol Rev* 67(4):491–502
- Strobel G, Daisy B, Castillo U, Harper J (2004) Natural products from endophytic microorganisms. *J Nat Prod* 67(2):257–268
- Tanaka M, Fukushima T, Tsujino Y, Fujimori T (1997) Nigrosporins A and B, new phytotoxic and antibacterial metabolites produced by a fungus *Nigrospora oryzae*. *Biosci Biotechnol Biochem* 61(11):1848–1852
- Wang LW, Xu BG, Wang JY, Su ZZ, Lin FC, Zhang CL, Kubicek CP (2011) Bioactive metabolites from *Phoma* species, an endophytic fungus from the Chinese medicinal plant *Arisaema erubescens*. *Appl Microbiol Biotechnol*. doi:10.1007/s00253-011-3472-3
- White TJ, Bruns S, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TL (eds) PCR protocols: a guide to methods and applications. Academic Press, San Diego, pp 315–322
- Wu SH, Chen YW, Shao SC, Wang LD, Yu Y, Li ZY, Yang LY, Li SL, Huang R (2009) Two New Solanapyrone Analogues from the Endophytic Fungus *Nigrospora* sp YB-141 of *Azadirachta indica*. *Chem Biodivers* 6(1):79–85
- Xu J, Aly AH, Wray V, Proksch P (2011) Polyketide derivatives of endophytic fungus *Pestalotiopsis* sp isolated from the Chinese mangrove plant *Rhizophora mucronata*. *Tetrahedron Lett* 52(1):21–25
- Xue CM, Li T, Deng ZW, Fu HZ, Lin WH (2006) Janthinolide A-B, two new 2,5-piperazinedione derivatives from the endophytic *Penicillium janthinellum* isolated from the soft coral *Dendronephthya* sp. *Pharmazie* 61(12):1041–1044
- Zhang HW, Song YC, Tan RX (2006) Biology and chemistry of endophytes. *Nat Prod Rep* 23(5):753–771