

# Pulmonary vein cellular electrophysiology and atrial fibrillation: Does basic research help us understand clinical pulmonary-vein arrhythmogenesis?

Stanley Nattel, MD

*From the Department of Medicine and Research Center, Montreal Heart Institute and University of Montreal, Montreal, Quebec, Canada.*

Arguably the most important discovery in the clinical electrophysiology of atrial fibrillation (AF) has been the recognition, largely through the work of Haissaguerre's group,<sup>1</sup> of the enormous importance of pulmonary vein (PV) myocardial sleeves in AF initiation. Numerous clinical investigators have confirmed the value of PV focal-source ablation and/or PV-left atrial disconnection in the management of AF, particularly of the paroxysmal variety.

Because of the substantial clinical importance of PV activity in AF, basic investigators have made great efforts to understand what specific properties of the PVs make them susceptible to ectopic impulse formation. In this issue of *Heart Rhythm*, Miyauchi et al<sup>2</sup> provide a novel contribution by showing that heterogeneous PV repolarization properties are associated with a propensity to early-after depolarization (EAD) and triggered-activity initiation by isoproterenol and rapid pacing. The mechanism(s) of this phenomenon is (are) not elucidated, but heterogeneous ion-current distribution and differential action-potential restitution are suggested by the investigators as potential contributors. Whether the abnormal activity is due to local reentry, to  $\text{Ca}^{2+}$ -induced triggered activity as has been suggested by previous investigators,<sup>3</sup> or to  $\text{Ca}^{2+}$ -current reactivation as for classical EADs<sup>4</sup> remains unknown.

These observations add an additional candidate mechanism for the basis of PV arrhythmia vulnerability, along with marked tissue anisotropy,<sup>5</sup>  $\text{Ca}^{2+}$ -handling abnormalities,<sup>3</sup> arrhythmogenic responses to local autonomic activity,<sup>6</sup> and peculiarities of PV ion-channel distribution.<sup>7</sup> However, these observations pose challenges to clinical and basic investigators alike. If normal PVs, as assessed in all of these studies, are so good at generating ectopic activity, why don't we all have paroxysmal AF, or at least frequent atrial ectopic activity?

This work was supported by grants from the Canadian Institutes of Health Research, the Quebec Heart and Stroke Foundation, and the Mathematics of Information Technology and Complex Systems (MITACS) Network of Centers of Excellence.

**Address reprint requests and correspondence:** Dr. Stanley Nattel, Montreal Heart Institute Research Center, 5000 Belanger Street East, Montreal, Quebec, Canada H1T 1C8.

E-mail address: Stanley.nattel@icm-mhi.org

What is it about the PVs that make them such good arrhythmia generators in AF patients? Is there an underlying PV predisposition to arrhythmogenesis that requires only a specific additional insult to unmask (the "multiple-hit" theory)? If so, what is the insult that moves PV arrhythmogenesis from latent to manifest? Or is there some specific pathology to which clinically arrhythmogenic PVs are susceptible that simply never occurs in normal PVs? Although studies in normal experimental animals and their tissues give useful insights into basic differences between PV myocardial sleeves and atria, they may not reveal the path to clinical arrhythmogenesis. Until we understand the crucial mechanistic factor(s) that differentiate clinically normal from arrhythmogenic PVs, we will not understand the role of PVs in AF.

## References

1. Haissaguerre M, Jais P, Shah DC, Takahashi A, Hocini M, Quiniou G, Garrigue S, Le Mouroux A, Le Metayer P, Clementy J. Spontaneous initiation of atrial fibrillation by ectopic beats originating in the pulmonary veins. *N Engl J Med* 1998;339:659–666.
2. Miyauchi Y, Hayashi H, Miyauchi M, Okuyama Y, Mandel WJ, Chen PS, Karagueuzian HS. Heterogeneous pulmonary vein myocardial repolarization. Implications for reentry and triggered activity. *Heart Rhythm* 2005;2:1339–1345.
3. Honjo H, Boyett MR, Niwa R, Inada S, Yamamoto M, Mitsui K, Horiuchi T, Shibata N, Kamiya K, Kodama I. Pacing-induced spontaneous activity in myocardial sleeves of pulmonary veins after treatment with ryanodine. *Circulation* 2003;107:1937–1943.
4. Viswanathan PC, Rudy Y. Pause induced early after depolarizations in the long QT syndrome: a simulation study. *Cardiovasc Res* 1999;42:530–542.
5. Hocini M, Ho SY, Kawara T, Linnenbank AC, Potse M, Shah D, Jais P, Janse MJ, Haissaguerre M, De Bakker JM. Electrical conduction in canine pulmonary veins: electrophysiological and anatomic correlation. *Circulation* 2002;105:2442–2448.
6. Patterson E, Po SS, Scherlag BJ, Lazzara R. Triggered firing in pulmonary veins initiated by in vitro autonomic nerve stimulation. *Heart Rhythm* 2005;2:624–631.
7. Melnyk P, Ehrlich JR, Pourrier M, Villeneuve L, Cha TJ, Nattel S. Comparison of ion channel distribution and expression in cardiomyocytes of canine pulmonary veins versus left atrium. *Cardiovasc Res* 2005;65:104–116.