

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/260373642>

Mesenchymal stem cell delivery strategies to promote cardiac regeneration following ischemic injury

ARTICLE *in* BIOMATERIALS · FEBRUARY 2014

Impact Factor: 8.56 · DOI: 10.1016/j.biomaterials.2014.01.075 · Source: PubMed

CITATIONS

11

READS

135

5 AUTHORS, INCLUDING:



[Valerio Russo](#)

University of British Columbia - Vancouver

5 PUBLICATIONS 29 CITATIONS

SEE PROFILE



[Andrew Hamilton](#)

Queen's University

27 PUBLICATIONS 238 CITATIONS

SEE PROFILE



[Brian Amsden](#)

Queen's University

120 PUBLICATIONS 3,472 CITATIONS

SEE PROFILE



[Lauren E Flynn](#)

The University of Western Ontario

30 PUBLICATIONS 777 CITATIONS

SEE PROFILE



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/authorsrights>



Contents lists available at ScienceDirect

Biomaterials

journal homepage: www.elsevier.com/locate/biomaterials

Review

Mesenchymal stem cell delivery strategies to promote cardiac regeneration following ischemic injury



Valerio Russo^{a,b,1}, Stuart Young^{a,b,1}, Andrew Hamilton^c, Brian G. Amsden^{a,b},
Lauren E. Flynn^{a,d,e,*}

^a Department of Chemical Engineering, Queen's University, 19 Division Street, Kingston, Ontario, Canada K7L 3N6

^b Human Mobility Research Centre, Kingston General Hospital, 76 Stuart Street, Kingston, Ontario, Canada K7L 2V7

^c Division of Cardiac Surgery, Department of Surgery, Kingston General Hospital, Kingston, Ontario, Canada K7L 2V7

^d Department of Chemical and Biochemical Engineering, Thompson Engineering Building, The University of Western Ontario, London, Ontario, Canada N6A 5B9

^e Department of Anatomy and Cell Biology, Schulich School of Medicine & Dentistry, The University of Western Ontario, London, Ontario, Canada N6A 5C1

ARTICLE INFO

Article history:

Received 23 January 2014

Accepted 30 January 2014

Available online 21 February 2014

Keywords:

Mesenchymal stem cell

Cell encapsulation

Injectable biomaterials

Cardiac tissue engineering

Angiogenesis

ABSTRACT

Myocardial infarction (MI) is one of the leading causes of mortality worldwide and is associated with irreversible cardiomyocyte death and pathological remodeling of cardiac tissue. In the past 15 years, several animal models have been developed for pre-clinical testing to assess the potential of stem cells for functional tissue regeneration and the attenuation of left ventricular remodeling. The promising results obtained in terms of improved cardiac function, neo-angiogenesis and reduction in infarct size have motivated the initiation of clinical trials in humans. Despite the potential, the results of these studies have highlighted that the effective delivery and retention of viable cells within the heart remain significant challenges that have limited the therapeutic efficacy of cell-based therapies for treating the ischemic myocardium. In this review, we discuss key elements for designing clinically translatable cell-delivery approaches to promote myocardial regeneration. Key topics addressed include cell selection, with a focus on mesenchymal stem cells derived from the bone marrow (bMSCs) and adipose tissue (ASCs), including a discussion of their potential mechanisms of action. Natural and synthetic biomaterials that have been investigated as injectable cell delivery vehicles for cardiac applications are critically reviewed, including an analysis of the role of the biomaterials themselves in the therapeutic scheme.

© 2014 Elsevier Ltd. All rights reserved.

1. Introduction and clinical scope

Cardiovascular disease (CVD) is the leading cause of mortality in Canada, the United States, and many European countries. In fact, CVD is the most lethal human medical condition in the developed world, accounting for 17.4 million deaths in 2008, or 30% of all deaths worldwide [1]. In the USA alone, more than 2150 people die each day from conditions associated with CVD, which is more than the combined total for cancer, chronic lower respiratory disease and accidents [2]. In addition to these dramatic mortality rates, the treatment of CVD places a significant economic burden on the healthcare system. In the USA, the direct and indirect costs

of treating CVD were estimated to be more than \$312.6 billion in 2009 [2].

CVD encompasses a broad range of medical conditions that affect the heart and vasculature. Focusing on coronary heart disease (CHD), the most common adverse event is myocardial infarction (MI). When the vascular flow is interrupted in MI, a portion of the heart experiences ischemic necrosis and is surrounded by a “marginal zone”, which can recover function, but is susceptible to irreversible damage [3]. As a result of subsequent inflammation during the early phase of post-infarction remodeling, the marginal zone is expanded into the surrounding tissue. In the late remodeling stage, collagen scar maturation causes tissue fibrosis as well as ventricular dilation and wall-thinning. The process continues until the tensile strength of the scar balances the distending forces of the heart [3]. The infarcted tissue does not contribute to pressure generation during systole, leading to impaired cardiac function and increasing the work and energy consumption of the remaining viable segments of the ventricular myocardium. The rate of electrical signal

* Corresponding author. Department of Chemical and Biochemical Engineering, Thompson Engineering Building, The University of Western Ontario, London, Ontario, Canada N6A 5B9 Tel.: +1 519 661 2111x87226; fax: +1 519 661 3498.

E-mail address: lauren.flynn@uwo.ca (L.E. Flynn).

¹ These authors contributed equally to this publication.

propagation in the marginal zone may be heterogeneous and may lead to development of arrhythmias, such as ventricular tachycardia, with resultant inefficient cardiac contraction. In more serious cases, multiple chaotic electrical wavefronts can cause ventricular fibrillation where effective cardiac output is impossible [4].

Although current therapies (Fig. 1) have decreased mortality rates from acute MI, many survivors are left with chronic ischemia or complications related to myocardial necrosis such as congestive heart failure. It has been estimated that at least 12% of patients undergoing cardiac catheterization at tertiary medical centers for angina or cardiac ischemia have obstructive coronary disease not amenable to standard revascularization therapies [5]. Attempting further traditional revascularization in this group has not resulted in decreases in morbidity or mortality [6]. The present treatment of congestive heart failure mainly targets its symptoms, failing to address the underlying weakness of the ventricular muscle. When this approach becomes inadequate, ventricular assist devices and cardiac transplantation are considered, although potential demand for these expensive and resource-intensive therapies far exceeds availability [7].

As such, there is an evolving need for new strategies to promote non-traditional coronary revascularization and restoration of functional cardiac tissue following ischemic necrosis. Cell-based strategies involving the delivery of cells with reparative or regenerative potential into the afflicted regions of the heart have shown promise for the treatment of cardiac ischemia. Primarily, these

approaches seek to replace damaged cardiomyocytes, promote revascularization, reduce fibrosis and improve ventricular compliance, pressure, and hemodynamics.

Numerous studies in animal models of MI have demonstrated functional benefits following the injection of bone marrow-derived mesenchymal stem cells (bMSCs) and adipose-derived stem cells (ASCs). The lack of significant *in vivo* cardiomyogenic differentiation and cell engraftment observed in the majority of these studies suggests that paracrine action may play a pivotal role in the improved cardiac function. Cell delivery scaffolds may have the potential to enhance the outcomes of MSC therapy by overcoming the limitations of poor localization, retention and viability associated with injecting cell suspensions [8]. With this view, recent research has focused on injectable hydrogels as delivery systems, which can be designed to provide temporary passive reinforcement to the myocardium and improve MSC retention within the site of delivery.

In this review, we will discuss the key elements that are critical for the development of an effective cell-based therapeutic strategy with MSCs for myocardial regeneration, including cell selection and elucidation of their biological mechanisms of action. Additionally, we will address a major challenge in the clinical translation of this strategy: effectively delivering and retaining viable cells in the heart. Biomaterials that could serve as injectable delivery vehicles for cardiac applications are critically reviewed, including an analysis of the role of biomaterials themselves in the therapeutic scheme.




Fig. 1. Overview of the current clinical approaches for the prevention and treatment of myocardial ischemic injury.

2. MSCs for cardio-regenerative applications

For clinical applications, a regenerative cell source should be (i) abundant, (ii) accessible, (iii) easy to deliver within the scope of a single operation, (iv) tested and optimized in appropriate animal models, (v) non-tumorigenic and non-immunogenic, and (vi) should stimulate regeneration in a predictable manner with a well-characterized dose response. With this view, autologous cell therapies have advantages, particularly if the cells can be isolated and available for immediate use in the procedure suite during the time frame of a planned cardiac intervention. As the number of cells required for cardiac cell therapy is predicted to be high ($\sim 10^8$ – 10^9 per application [9,10]), an ideal source would yield a large number of the desired cell type with a harvesting process that is neither complex nor painful.

To date, many different cell types have been investigated both *in vitro* and *in vivo* for cardiac regeneration, including skeletal myoblasts, embryonic stem cells (ESCs), cardiac stem cells (CSCs), and MSCs [11]. Clinical trials with skeletal myoblasts have demonstrated only modest improvements in cardiac function and remodeling [12,13]. In rat models, transplanted ESCs have improved myocardial function, but the effects were temporary. Further, ESCs demonstrate immunological incompatibility, may form teratomas, and are associated with ethical issues that have limited their broad-scale application [14]. The clinical use of CSCs is impeded by the limited availability and accessibility of cardiac tissue, which contains only a small population of regenerative cells, necessitating large-scale cell expansion. The SCIPIO clinical trial is investigating autologous CSCs extracted from a portion of the atrial appendage resected during cardiac surgery. However, this can be risky in the elderly or in patients who have fragile atria or high right-side venous pressure [15–17]. While studies to date with MSCs have not convincingly demonstrated extensive differentiation into functional cardiomyocytes, their injection into the damaged myocardium can yield improved functional outcomes in the treatment of cardiac ischemia.

MSCs are multipotent adult stem cell populations that have differentiation potential typically associated with their mesodermal lineage (adipogenic, chondrogenic, osteogenic, myogenic). MSCs have been successfully harvested from a variety of adult tissues including bone marrow [18], adipose tissue [19], periosteum

[20], synovium [21], skeletal muscle [22], and deciduous teeth [23]. In this review, we will focus on two promising MSC sources for cardiac tissue engineering: bMSCs and ASCs (Table 1).

bMSCs have been one of the most extensively studied stem cell populations for regenerative approaches since they were first described by Friedenstein et al. in 1976 [48]. The development of protocols for bMSC isolation, purification and culturing have been reviewed in detail previously [18,49,50]. Typically, bMSCs are extracted from the bone marrow of the iliac crest. This procedure can be challenging for patients with end-stage ischemic heart disease. However, the promise of regeneration with bMSCs shown in animal models of MI (Table 2) has motivated further research and several clinical trials.

ASCs represent an attractive alternative due to their relative abundance, accessibility, multipotent differentiation capacity and demonstrated lack of immunogenicity [62]. Most patients would be able to donate a sample of fat for stem cell isolation with minimal adverse effects. In adult bone marrow, the fraction of MSCs to total nucleated cells is estimated to range between 1:50,000 and 1:1 million, and the limited volume of bone marrow that can be safely harvested per donor may necessitate significant *in vitro* expansion [24–26,63,64]. In contrast, ASC frequency in adult human adipose tissue is estimated to range between 1:30–1:100 per total nucleated cells [27,28]. As a potential allogenic cell source, large volumes of adipose tissue are routinely discarded during lipo-reduction surgeries. Due to the promising availability of adipose tissue as a regenerative cell source, numerous groups have been investigating the therapeutic potential of ASCs for the treatment of MI with encouraging results (Table 3).

Recently, our group has isolated and characterized ASCs derived from pericardial adipose tissue, which is a depot that would be readily accessible and expendable for intraoperative ASC isolation during open heart surgery [47]. As outlined in Table 1, while MSCs isolated from adipose tissue and bone marrow express similar cell surface markers, there is variability in the specific levels of expression between the sources that may have implications in terms of their regenerative capacity [30–36,75].

Importantly in the context of cardiac regeneration, injected bMSCs and ASCs migrate to sites of injury or ischemic tissues and contribute to regeneration [66,72,76]. While the mechanisms involved are not yet fully resolved, gradients of chemotactic factors

Table 1
Key characteristics of bMSCs and ASCs in the context of tissue regeneration.

	bMSCs	Both	ASCs
Cell yield (MSCs:total nucleated cells)	1:50,000 – 1×10^6 [24–26]		1:100 [27,28]
Immunophenotype	<ul style="list-style-type: none"> CD34⁺ and CD73⁺ in freshly isolated cells [29] CD106⁺, CD49f⁺, PODXL⁺, CD49d⁺ and CD54⁺ upon culturing [30–34] 	<ul style="list-style-type: none"> CD29⁺, CD44⁺, CD73⁺, CD90⁺, CD105⁺ and CD166⁺ upon culturing [35,36] 	<ul style="list-style-type: none"> Expression of CD34 in freshly isolated cells [29,30] CD49d⁺, CD54⁺, CD106⁺, CD49f⁺ and PODXL⁺ upon culturing [30–34]
Strengths	<ul style="list-style-type: none"> Demonstrated short-term safety in allogenic cardiovascular applications [37] Extensively characterized 	<ul style="list-style-type: none"> Multilineage differentiation potential and CFU-F formation [35] Immuno-modulatory activity 	<ul style="list-style-type: none"> Abundant, easily harvestable and uniquely expendable cell source associated with low donor site morbidity High cell yield Higher proliferation [38,39], angiogenic capacity [40] and immunomodulation [41–43] and lower senescence ratio [35] compared to bMSCs
Limitations	<ul style="list-style-type: none"> Procedure sometimes complex and not suitable for patients with end-stage ischemic disease Conflicting results in clinical trials Risk of ectopic bone formation [44] Limited quantity of aspirates collectable without adverse effects Cell expansion challenging 	<ul style="list-style-type: none"> Low engraftment when delivered as a cell suspension Limited <i>in vivo</i> differentiation Influence of donor age and health status on cell response currently not well characterized 	<ul style="list-style-type: none"> Variability in the ASC population with gender, body mass index, and source depot [28,45–47] Population not as extensively characterized to date; relative potency compared to bMSCs not fully elucidated Safety for allogenic applications not yet demonstrated

Table 2
Summary of key studies investigating cell therapy with bMSCs in animal models of MI.

Animal model/ delivery method	Cell population	Time-points	Quantification of engraftment/retention	Differentiation into cardio-vascular components	Effects on cardiac function and/or ventricular remodeling	Ref.
Rat/i.m.	Autologous, (i) freshly isolated bMSCs, (ii) 5-aza treated bMSCs and (iii) untreated bMSCs	5 wks	<ul style="list-style-type: none"> Engrafted cells at 5 wks (not quantified) 	<ul style="list-style-type: none"> Donor cells TnI⁺ ↑ Capillary density in all MSC-treated groups 	<ul style="list-style-type: none"> ↑ LVSP and LVDP in 5-aza treated group ↓ Scar area and LV chamber size 	[51]
Rat/i.m.	Autologous bMSCs + CMEC	3 – 12 wks	n.d.	<ul style="list-style-type: none"> Donor cells factor VIII⁺ ↑ Capillary density and # of collateral vessels 	<ul style="list-style-type: none"> ↑ Myocardial perfusion and LVEF to 45 ± 2.2% 	[52]
Rat/i.m.	Autologous bMSCs	8 wks	n.d.	<ul style="list-style-type: none"> Donor cells vWF⁺ ↑ Capillary density 	<ul style="list-style-type: none"> ↑ LVEF and perfusion in scar 	[53]
Rat/i.m.	Autologous bMSCs or peripheral blood mononuclear cells (PB-MNC)	1 – 60 days	n.d.	<ul style="list-style-type: none"> Donor cells factor VIII⁺ ↑ Capillary density 	<ul style="list-style-type: none"> bMSCs restored LVD and LVEDV to baseline ↑ FS ↓ Collagen deposition for bMSC-treated group only 	[54]
Rat/i.m.	Allogenic bMSCs	1, 2 and 4 wks	<ul style="list-style-type: none"> Engrafted cells at 1–2 and 4 wks (not quantified) 	<ul style="list-style-type: none"> Donor cells CD31⁺, α-SMA⁺ and TnT⁺ (low number) at 2 wks ↑ Capillary density 	<ul style="list-style-type: none"> ↑ LVSP and wall thickness ↓ LVEDP and infarct size 	[55]
Mouse/i.m.	Allogenic c-kit ⁺ bMSCs	9 ± 2 days	n.d.	<ul style="list-style-type: none"> Donor cells cardiac myosin⁺, factor VIII⁺ and α-SMA⁺ Genetic expression of cx-43, MEF2 and GATA-4 	<ul style="list-style-type: none"> ↑ LVDP ↓ LVEDP and infarct area 	[56]
Mouse/i.m.	Allogenic lin [−] bMSCs	2 wks	<ul style="list-style-type: none"> Engrafted cells at 2 wks (not quantified) 	<ul style="list-style-type: none"> Donor cells α-sarcomeric actin⁺ New cardiomyocytes in infarct region bMSCs incorporated into blood vessels (low number) 	<ul style="list-style-type: none"> ↓ Infarct size and fibrosis 	[57]
Mouse/i.m.	Allogenic c-kit ⁺ bMSCs	5 – 10 days	<ul style="list-style-type: none"> Engrafted cells at 10 days (not quantified) 	<ul style="list-style-type: none"> Donor cells positive for GATA-4, Nkx2.5, MEF2C, α-sarcomeric actin, cardiac MHC, TnI, desmin, cx-43, GATA-6, ETS-1, vWF and α-SMA ↑ Capillary density and # of arterioles 	<ul style="list-style-type: none"> ↓ LVEDP and infarct size 	[58]
Mouse/i.m.	Allogenic c-kit ⁺ bMSCs	30 days	<ul style="list-style-type: none"> Engrafted cells at 2 days (~25% of donor cells detected in border zone) 	<ul style="list-style-type: none"> Donor cells vWF⁺ and α-SMA⁺ Cx-43 and Nkx2.5 expressed at 48 h Gap junctions between donor and host cells 	<ul style="list-style-type: none"> Restored contraction ↑ LVDP and wall thickness ↓ LVEDP, scar formation and chamber volume 	[59]
Pig/i.m.	Autologous bMSCs	4 wks	<ul style="list-style-type: none"> “Significant” engraftment at 6 months (not quantified) 	<ul style="list-style-type: none"> Donor cells TnT⁺, MHC⁺ and tropomyosin⁺ 	<ul style="list-style-type: none"> Preservation of wall thickness 	[60]
Dog/i.m.	Allogenic bMSCs	30 days	n.d.	<ul style="list-style-type: none"> Donor cells factor VIII⁺ and α-SMA⁺ ↑ Vascular density 	<ul style="list-style-type: none"> ↑ LVEF ↓ Collagen fibrosis 	[61]

5-aza, 5-azacytidine; α-SMA, alpha smooth muscle actin; CMEC, coronary microvascular endothelial cells; FS, fractional shortening; i.m., intramyocardial; i.c., intracoronary; n.d., no data; LV, left ventricle; LVEDP, left ventricular end diastolic pressure; LVEDV, left ventricular end diastolic volume; LVEF, left ventricular ejection fraction; LVDP, left ventricular developed pressure; LVSP, left ventricular systolic pressure; LVD, left ventricular dysfunction; MHC, myosin heavy chain; TnI, troponin I; TnT, troponin T; vWF, von Willebrand factor; cx-43, connexin-43.

that are up-regulated in the ischemic myocardium can promote the homing of exogenous MSCs in cell-based therapies and stimulate the migration of endogenous stem cells [77]. MSCs express a variety of membrane receptors involved in the recognition of chemo-attractant cytokines, including stromal derived factor-1 (SDF-1), transforming growth factor-β1 (TGF-β1) and tumor necrosis factor-

α (TNF-α), which are expressed in inflamed and injured tissues [78,79]. In particular, signaling through SDF-1/CXCR4 and HGF/c-MET has been identified as being of importance in stimulating the migration of reparative cells to the ischemic heart [79–83]. MSCs also secrete a range of growth factors and cytokines that play pivotal roles in mediating inflammation, immune responses, cell

Table 3
Summary of key studies investigating cell therapy with ASCs in animal models of MI.

Animal model/ delivery method	Cell population	Time points	Quantification of engraftment/retention	Differentiation into cardiovascular components	Effects on cardiac function and/or ventricular remodeling	Ref.
Rat/implanted cell sheet	Autologous ASCs	4 – 8 wks	<ul style="list-style-type: none"> Thickening of cell sheet at 4 wks <1% TUNEL positive cells at 48 h 	<ul style="list-style-type: none"> Donor cells cardiac troponin⁺, vWF⁺ and α-SMA⁺ 	<ul style="list-style-type: none"> ↑ LV maximum dP/dt ↓ LVEDP Inhibition of LV enlargement 	[65]
Rat/i.m.	Allogenic ASCs, bMSCs, or AD-CMG	1 – 4 wks	<ul style="list-style-type: none"> Low ASC and bMSC engraftment at 1 wk No engraftment for any treatment group at 4 wks 	<ul style="list-style-type: none"> Donor cells MCL2V⁺, α-SMA⁺ and BSL-I lectin⁺ at 1 and 4 wks ↑ Capillary density 	<ul style="list-style-type: none"> ↑ LVEF at 4 wks in ASC group only ↓ Infarct size in all treatment groups 	[66]
Rat/i.m.	Allogenic ASCs	12 wks	<ul style="list-style-type: none"> Engrafted cells in 2 of 11 rats at 12 wks (not quantified) 	<ul style="list-style-type: none"> Donor cells cardiac troponin⁺ (very rare) ↑ Capillaries density and # of arterioles 	<ul style="list-style-type: none"> ↑ LVEF, SV, CO and LV end-diastolic dimension 	[67]
Rat/i.m.	Allogenic ASCs	1 – 4 wks	<ul style="list-style-type: none"> n.d. 	<ul style="list-style-type: none"> Donor cells sarcomeric α-actinin⁺ and α-MHC⁺ (low levels) ↑ Capillary density 	<ul style="list-style-type: none"> ↑ LVEF and % viable myocardium in infarct region Preservation of LV wall thickness 	[68]
Athymic nude rat/i.m.	Human ASCs	4 wks	<ul style="list-style-type: none"> Engrafted cells in 3 of 10 rats at 4 wks (not quantified) 	<ul style="list-style-type: none"> Donor cells α-sarcomeric actin⁺ and α-SMA⁺ at 4 wks ↑ # of arterioles 	<ul style="list-style-type: none"> ↑ LVEF and FS ↓ LVESV and LVEDV and infarct size 	[69]
Nude rat and SCID mouse/i.m.	Human ASCs from epicardial fat	15 – 30 days	n.d.	<ul style="list-style-type: none"> Donor cells TnI⁺ and CD31⁺ 	<ul style="list-style-type: none"> ↑ EF, FS and wall thickness ↓ Infarct size 	[70]
SCID mouse/i.m.	Freshly-isolated or cultured human ASCs	1 – 4 wks	<ul style="list-style-type: none"> Engraftment of both types of cells at 28 days (not quantified) 	<ul style="list-style-type: none"> Donor cells TnI⁺, vWF⁺ and α-SMA⁺ Cx-43 between donor and host cells ↑ Capillary density in both treatment groups 	<ul style="list-style-type: none"> ↑ LVEF ↓ LVESV 	[71]
Pig/i.c.	Autologous ASCs or bMSCs	30 days	<ul style="list-style-type: none"> Engrafted cells at 30 days (not quantified) 	<ul style="list-style-type: none"> Donor cells cardiac TnI⁺, vWF⁺, α-SMA⁺ and desmin⁺ ↑ Capillary density 	<ul style="list-style-type: none"> ↑ LVEF and wall thickness with both populations 	[72]
Pig/i.c.	Autologous freshly-isolated ASCs	4 – 8 wks	n.d.	<ul style="list-style-type: none"> ↑ Capillary density in infarct border 	<ul style="list-style-type: none"> ↑ LVEF, myocardial salvage and wall thickness ↓ Myocardial perfusion defects 	[73]
Rabbit/i.m.	Autologous ASCs or 5-aza treated ASCs	5 wks	<ul style="list-style-type: none"> Engrafted cells at 5 wks (not quantified) 	<ul style="list-style-type: none"> Donor cells α-sarcomeric actin⁺, cardiac-specific TnT⁺ and vWF⁺ ↓ Capillary density 	<ul style="list-style-type: none"> ↑ LVEF ↓ Infarct size (more significant with 5-aza treatment) 	[74]

5-aza, 5-azacytidine; α -SMA, alpha smooth muscle actin; CO, cardiac output; dP/dt, rate of rise of left ventricular pressure; FS, fractional shortening; EF, ejection fraction; i.c., intracoronary; i.m., intramyocardial; n.d., no data; LV, left ventricle; LVEDP, left ventricular end diastolic pressure; LVEDV, left ventricular end diastolic volume; LVEF, left ventricular ejection fraction; LVESV, left ventricular end systolic volume; LVDP, left ventricular developed pressure; LVSP, left ventricular systolic pressure; MHC, myosin heavy chain; SV, stroke volume; TnI, troponin I; TnT, troponin T; vWF, von Willebrand factor; cx-43, connexin-43; MLC2V, myosin light chain 2v; AD-CMG, adipose-derived cardiomyoblasts; BSL-I lectin, *Bandeiraea simplicifolia* lectin I.

migration and importantly, in the context of cardiovascular regeneration, angiogenic and vasculogenic factors. This natural secretome has been exploited by many groups assessing cell-based therapies for the treatment of ischemia using induced MI and ischemic hind limb animal models [52,84–86].

Another important feature of MSCs in the context of cell-based therapies is their immunoprivileged status, which may enable the application of allogenic cells with fewer concerns about the initiation of undesired immune responses. This characteristic, which is believed to be associated with very low expression of the major histocompatibility complex II (MHC II) antigen, has been examined both *in vitro* and *in vivo*. Furthermore, results with bMSCs and ASCs indicate that both cell populations modulate T-cell activation, proliferation, migration and inflammatory cytokine secretion [43,87–90]. Recently,

Melief et al. showed that ASCs elicit a more potent suppression of peripheral blood mononuclear cell (PBMC) proliferation and a more effective induction of a shift from Th1 response to Th2 response *in vitro* when compared to bMSCs [43]. The assessment of the specific mechanisms by which MSCs mediate the immune response is an area that warrants further investigation, to develop a deeper understanding of how MSCs may promote a more regulatory versus inflammatory condition within the transplant environment.

2.1. Mechanisms of MSC-based cardiac regeneration

Several possible mechanisms by which exogenous MSCs can enhance cardiac regeneration have been proposed: (i) the mobilization of endogenous stem cells capable of cardiomyogenesis to the

site of injury through chemotactic signaling, (ii) the reduction of fibrosis and improvement in cardiomyocyte survival mediated in part through neo-angiogenesis and vasculogenesis, (iii) the mechanical reinforcement of the infarct scar, allowing endogenous repair mechanisms to improve left ventricular function, and (iv) the reduction in the initial tissue damage through the modulation of the inflammatory response.

It is important that *in vivo* studies are designed to assess the mechanisms of MSC-mediated regeneration. Key endpoints that should be considered in assessing the efficacy of a cell-based therapy for MI include:

- i. The engraftment, retention, and survival of the transplanted cells
- ii. The re-population of lost cardiomyocytes through *de novo* functional cardiomyogenesis and/or the promotion of endogenous cardiomyocyte proliferation
- iii. The integration of *de novo* cardiomyocytes with the host electromechanical network
- iv. The induction of angiogenesis and vasculogenesis through paracrine-mediated effects or direct differentiation of the transplanted cells into vascular structures
- v. Improvements in cardiac function, as measured by left ventricle stroke volume (LVS), ejection fraction (LVEF), and fractional shortening (LVFS), both at rest and under stress
- vi. The prevention or attenuation of left ventricular remodeling in terms of tissue fibrosis, infarct size and ventricular chamber dilation, measured by end diastolic and end systolic diameters (EDD, ESD)

For bMSCs (Table 2), in 2001, Orlic et al. first demonstrated that intramyocardial injection of undifferentiated lineage-negative (Lin[−]) bMSCs in a mouse model of MI resulted in the formation of new cardiac tissue and also improved cardiac contractility [56]. Two years later, Kudo et al. demonstrated that bMSC transplantation reduced fibrosis and infarct size, with evidence of bMSC differentiation into cardiomyocytes and endothelial cells within the ischemic myocardium [57]. Similar results were reported by Zhang et al. in a study comparing the effects of intramyocardial injection of bMSCs versus PBMcs that demonstrated functional improvements with only bMSCs [54]. In a more recent study, a highly proliferative bMSC subpopulation was isolated from human bone marrow by Yoon et al. Intramyocardial injection of this subpopulation in a nude rat model resulted in a higher degree of engraftment and differentiation as compared to unselected bMSCs, and also induced endogenous cardiomyogenesis [91].

Similarly for ASCs (Table 3), Valina et al. first compared the effect of intracoronary injection of ASCs and bMSCs in a porcine model. Although cardiac differentiation was not detected, some exogenous cells expressed endothelial lineage markers. Furthermore, both cell types induced comparable improvements in cardiac function and increased capillary density within the infarct zone [72]. Zhang et al. observed similar outcomes in a rat model of MI, comparing ASCs induced towards a cardiomyocyte phenotype using 5-azacytidine (5-aza) with untreated ASCs. In their model, LVEF and capillary density were dramatically increased in both treatment groups as compared to the controls. Additionally, cardiomyogenic differentiation of the ASCs was observed, along with a reduction in the infarct size, with indications that 5-aza pre-treatment was beneficial in reducing the infarcted region [74].

2.1.1. *In vitro* differentiation potential

In vitro studies have demonstrated the potential for MSCs to differentiate into the major cellular components of cardiac tissue when the culture medium is supplemented with specific factors, as

outlined in Table 4. It is important to note that this induced *in vitro* differentiation may not be reflective of the typical cell response following *in vivo* transplantation. Further, it has been postulated that the use of some differentiation factors may result in unpredictable reprogramming patterns that could increase the potential for the malignant transformation of the cells [106].

Co-culture systems with MSCs and cardiomyocytes have allowed scientists to develop models to better understand the cell–cell interactions that can occur at the site of transplantation. Both bMSCs [107] and ASCs [108] have been induced to undergo cardiomyogenic differentiation using this approach. This microenvironment-induced differentiation has been hypothesized to be mediated by cell–cell contact [109,110], although similar results have been obtained through indirect co-culturing [108] as well as exposure to cardiomyocyte lysates [111] or ischemic cardiac tissue-conditioned medium [112]. These results suggest that *in vivo*, where a broad array of microenvironmental cues are present, MSC cardiomyogenic differentiation could result from multiple pathways.

2.1.2. *In vivo* regeneration through MSC engraftment and differentiation

The results of several *in vivo* studies support the possibility that transplanted MSCs could contribute to regeneration through direct engraftment and differentiation. Cell fusion, rather than trans-differentiation, has been proposed as a driving mechanism for these events [58,113]. However, in strategies to date, cardiomyogenic differentiation is often not observed [61,66,69,72] or at best occurs at extremely low levels. Even when there is evidence of differentiation, most studies include only limited qualitative assessments. In seeking to quantify the findings, based on two studies, only 0.5–5% of the engrafted cells have been reported to differentiate [65,68]. Further, histological evidence of MSC differentiation has not always been paralleled by a functional benefit [51]. In particular, although the coupling between *de novo* and host cardiomyocytes through gap junction formation has been documented with both bMSCs [59] and ASCs [71], transplanted bMSCs failed to generate spontaneous and electrically-evoked intracellular calcium transients 9–10 days after delivery to the peri-infarct region in a mouse MI model [114].

Neovascularization must proceed in parallel with cardiomyogenesis to compensate for the ischemic conditions. Although at low levels, integration of transplanted MSCs into vascular structures has been reported in many animal models of MI (Tables 2 and 3). The critical importance of vasculogenic differentiation in improving cardiac function has been recently demonstrated by Yoon et al. Through an elegant suicide gene-based system, it was shown that depletion of cells differentiating toward the vascular lineages drastically reduced functional benefits, whereas elimination of cardiomyogenic cells had no significant effect [115]. In a few studies that have compared ASCs and bMSCs, capillarization and ventricular remodeling have been reported to be more prominent following ASC injection [66,72].

2.1.3. Regeneration through paracrine-mediated effects

Despite the modest levels of engraftment and differentiation, the injection of bMSCs or ASCs increases capillarization and neo-angiogenesis, in some cases almost doubling vascular density as compared to controls [51,53,61]. Similarly, echocardiographic analysis has shown that MSC-based therapies can significantly improve left ventricular diastolic and systolic pressure, LVEF, ESD and EDD, as well as vascular perfusion in the scar. These functional improvements are likely associated with a reduction in infarct size, preservation of wall thickness, and reduction in tissue fibrosis, which all contribute to enhanced ventricular compliance. The

Table 4
Factors investigated for the *in vitro* differentiation of MSCs towards cardiovascular lineages.

Lineage	Cell population	Supplements	Differentiation response	Ref.
Cardiomyogenic	bMSCs/mouse	3 μM /L 5-aza + 20% FBS	• Spontaneously beating cardiomyocytes	[92]
Cardiomyogenic	bMSCs/human	10 μM /L 5-aza + 10 μg /L bFGF + 0.25 mg/L amphotericin + 10% FBS	• MEF-2A and MEF-2D gene expression	[93]
Cardiomyogenic	bMSCs/human	1.0 mg/mL insulin + 10^{-9} M dex + 10^{-4} M ascorbate phosphate + 0.47 μg /mL linoleic acid + 0.55 mg/mL transferrin + 0.5 μg /mL sodium selenite + 10% FBS	• Expression of GATA-4 and MEF2, cardiac Tnl, cx-43, cardiac-specific α/β -MHC, and L-type cardiac Ca^{2+} channel	[94]
Cardiomyogenic	ASCs/rabbit	1, 3, 6, and 9 μM /L 5-aza + 15% FCS	• Spontaneous beating	[95]
Cardiomyogenic	ASCs/human	9 μM 5-aza + 0.12 $\mu\text{g}/\text{cm}^2$ laminin coating or 2.5 $\mu\text{g}/\text{cm}^2$ fibronectin coating + 15% FBS	• Expression of MHC, actinin and Troponin-I	[96]
Cardiomyogenic	ASCs/mouse and human	1% methylcellulose medium + 10 ng/mL IL-3 + 10 ng/mL IL-6 + 50 ng/mL SCF + 100 μM 2-mercaptoethanol + 2 mM L-glutamine + 10 μg /mL recombinant human insulin + 200 μg /mL transferrin + 15/30% FBS	• Contractile activity	[97,98]
Endothelial	bMSCs/human	50 ng/mL VEGF + 12% FCS	• Expression of vWF	[99]
Endothelial	bMSCs/human	100 ng/mL VEGF + 50 ng/mL EGF + 1 μg /mL hydrocortisone + 5% FBS	• Ability to form capillaries in semi- solid medium	[100]
Endothelial	ASCs/human and rat	20 – 50 ng/mL VEGF + 10 – 2% FCS	• Expression of vWF, flK-1 and VE-cadherin	[55,101]
Endothelial	ASCs/rat	EGM2 media + 1 μg /mL ascorbic acid + 22.5 μg /mL heparin + 0.2 μg /mL hydrocortisone + 5 ng/mL EGF + 10 ng/mL FGF2 + 20 ng/mL IGF-1 + 0.5 ng/mL VEGF + 2% FBS	• Ability to uptake acLDL	[102]
Smooth muscle	bMSCs/human	1 ng/mL TGF- β 1 + 2 ng/mL ITS + 10% FBS	• Expression of vWF and ability to uptake acLDL	[103]
Smooth muscle	ASCs/human	100 units/mL heparin + 1% FBS	• Expression of CD31	[104]
Sinus-node like cells	bMSCs and ASCs/human	10 μM 5-aza + 10% FBS	• Expression of endothelial markers CD31, vWF, and eNOS	[105]
			• Formation of endothelial-like tube structures	
			• Expression of α -SMA	
			• Expression of α -SMA, smoothelin, calponin, caldesmon	
			• Dose dependent contraction in response to muscarinic agonists	
			• Expression of HCN2 and HCN4	

5-aza, 5-azacytidine; dex, dexamethasone; ITS, insulin, transferrin, selenium; α -SMA, alpha smooth muscle actin; MHC, myosin heavy chain; MLC, myosin light chain; vWF, von Willebrand factor; acLDL, acetylated low density lipoprotein.

positive effects observed in the absence of long-term engraftment and differentiation suggest that the MSCs primarily function through paracrine-mediated effects to establish a more regenerative milieu within the host tissues. This theory is supported by the work of Kinnaird et al. who showed that injecting bMSC-conditioned medium increased the collateral perfusion in a murine model of peripheral ischemia [84].

Table 5 highlights some key factors secreted by MSCs and their potential biological effects within the ischemic heart. Among the many proteins secreted by MSCs, VEGF is potentially one of the most important mediators of angiogenesis in the ischemic heart. Tang et al. assessed the presence of VEGF in the infarct region through immunohistochemistry and detected its up-regulation following bMSC injection [53]. The critical role of VEGF in MSC-induced myocardial revascularization was further demonstrated by Markel et al., who showed that small interference RNA (siRNA) directed against VEGF negatively affected the recovery of myocardial function in a rat model of MI [126].

While genetic engineering approaches with MSCs are beyond the scope of the current review, the importance of chemotactic signaling pathways has been demonstrated by studies where the forced over expression of chemotactic receptors or ligands, including CXCR4, insulin-like growth factor-1 (IGF-1) and chemokine receptor-1 (CCR1), have enhanced the homing, viability and/or engraftment of injected MSCs [127–131].

Finally, MSCs respond to variations in oxygen tension through hypoxia-inducible factors (HIFs), which can up-regulate the expression of a broad array of genes associated with metabolism [132,133], proliferation and viability [84,118,134–136], angiogenesis

and vascular remodeling [137–145], and cellular motility [146,147]. As such, a number of groups are trying to better understand the biological mechanisms of this hypoxia-induced gene up-regulation, in order to harness it as a means to improve the regenerative capacity of the transplanted cells in the context of the ischemic heart [142,144–146,148].

2.2. Clinical trials with MSCs for myocardial regeneration

Based on the positive results observed in animal models, several clinical trials investigating bMSCs in the treatment of MI have been

Table 5
Paracrine factors secreted by MSCs in the context of tissue regeneration in ischemic tissues.

Biological effect	Factors	References
Angiogenesis	VEGF bFGF Ang-1 PDGF	[52,53,65,70,84,86,116–120] [52,70,86,118] [52,119,120] [84,119,121]
Inflammation mediation/ angiogenesis	IL-1, 6, 7, 8, 11, 12, 14 CINC MCP-1 TGF- β TNF- α	[70,84,86,121,122] [86] [86,121] [84,117,119] [70,84]
Cytoprotection and chemotaxis	HGF IGF SDF	[65,117,123,124] [116,118,125] [118,124]

Ang-1, angiopoietin 1; CINC, cytokine-induced neutrophil chemoattractant; MCP-1, monocyte chemotactic protein 1.

initiated. An initial short-term trial demonstrated that intracoronary injection of autologous bMSCs was safe and beneficial in terms of improved cardiac function at 3 months [149]. However, longer-term trials with autologous bMSCs have yielded conflicting data. The TOPCARE-AMI (8.5% treatment vs. 2.5% control), REPAIR-AMI (5.5% vs. 3.0%), and BOOST (5.9% vs. 3.1%) trials have all shown increased LVEF compared to control groups at 4, 12 and 18 months post-treatment, respectively [150–152]. The TOPCARE-AMI trial also compared the effects of bMSC treatment to cardiac progenitor cell (CPC) infusion, with the bMSCs showing a greater improvement in terms of left ventricular contractility [153]. However, a study conducted in Belgium, as well as the ASTAMI and BONAMI trials, demonstrated no significant improvement in cardiac function between control and bMSC-treated patients [154–156]. Further, a 5-year follow up of the patients enrolled in the BOOST trial showed that the beneficial effects observed at 18 months post-treatment in terms of cardiac contractility were not sustained. In contrast, the 5-year follow up of the TOPCARE-AMI study showed a significant improvement in LVEF in the bMSC-treated group, as well as reductions in end-systolic volume and end-diastolic volume. The observed differences between the trials may result from donor variability, the diverse methods of cell isolation and delivery employed, as well as different baselines for assessing left ventricular function [157,158].

Numerous ongoing clinical trials, including the REVITALIZE (NCT00874354), BAMI (NCT01569178), REGEN-AMI (NCT00765453) and REPAIR-ACS (NCT00711542) trials, are seeking to further elucidate the effects of autologous bMSCs in the treatment of MI through multinational, multicenter studies. Given the potential benefits of allogenic transplantation, the results from the first phase I clinical trial to test the safety and efficacy of intravenously-injected allogenic bMSCs were reported in 2009. Promisingly, in 53 patients who suffered MI, there were no rejection issues and heart function was improved, as measured by LVEF that showed a $6.7 \pm 2.2\%$ increase at 6 months. Nevertheless, when patients' physical performances were assessed using the 6 min walk test, no differences between the treated and control groups were observed [37].

Clinical trials for the treatment of MI with autologous human ASCs have also been initiated. More specifically, the APOLLO [159] and the PRECISE [160] trials have investigated the transplantation of freshly-isolated ASCs in 14 patients with MI that have demonstrated acute ST-elevation by electrocardiography, as well as 27 patients with non-revascularizable cardiac ischemia. Although statistically limited by the low number of subjects, these two trials have demonstrated the feasibility, safety and efficacy of autologous ASC administration and motivated the initiation of ATHENA (NCT01556022), a PHASE II clinical trial enrolling 45 patients, in June 2012. Additionally, MyStromalCell is a PHASE II clinical trial initiated in April 2010 that was designed to assess the efficacy of expanded autologous ASCs stimulated with VEGF-A prior to injection [161].

3. Cell delivery to the ischemic heart

The efficacy of cell therapy depends on the ability of the delivery mechanism to provide sufficient localization, survival, and retention of the donor cells in the affected tissues [162]. These challenges are particularly apparent in the field of cardiac tissue regeneration following MI, where the host environment presents numerous unique obstacles to cellular transplantation. Nevertheless, in the vast majority of clinical trials, an isotonic saline solution was routinely employed as the cell delivery vehicle [37,163]. These solutions possess physiological osmolarity but do not have the

capacity to localize or retain the administered cells within the target site while maintaining long-term cell viability.

3.1. Challenges of cell delivery: localization, survival, and retention

Compressive forces produced by continuous, periodic systolic contraction, as well as shear forces caused by hemodynamic flow, can cause the dissemination of cells injected in saline. One hour following delivery of PBMCs via intramyocardial, intracoronary, and interstitial retrograde coronary venous injection, retention rates were $11 \pm 3\%$, $2.6 \pm 0.3\%$, and $3.2 \pm 1\%$, respectively, with the greatest fraction lost to the pulmonary circulation [164]. Further, confocal microscopy of green fluorescent protein (GFP)-labeled MSCs injected intramyocardially has shown delivery in saline results in variable cellular distribution throughout the depth of the ventricle, with a decrease in retention from 31% to 1% over five days post-injection [8]. An uneven distribution could further exacerbate the already disrupted alignment of the surviving cardiomyocytes, interrupting the conduction of electrical and mechanical signals through the parallel bundles of cardiomyocytes, potentially producing a pro-arrhythmic heterogeneous milieu [165].

The short-term survival of the retained cells represents a further challenge [166]. In the post-MI heart, ischemia results in localized hypoxia and nutrient depletion [167], while the acute inflammatory reaction is associated with leukocyte and macrophage infiltration, fibrotic tissue deposition, oxidative load, and the generation of apoptotic cytokines [168]. Postponing delivery may avoid some inflammatory effects of acute-MI, but later stages of MI present additional challenges: advancing tissue fibrosis, ventricular dilation and wall thinning, and reduced vascularization all result in an apoptotic environment deficient in oxygen and nutrients [169]. Continuing cardiomyocyte death contributes to the release of apoptotic factors that further threaten the survival and function of the transplanted cell population [170].

Considering these events, timing of cell delivery emerges as an important factor in regenerative efficacy. A study by Hu et al. concluded that delivery of MSCs one week post-MI provided greater functional benefit as compared to delivery after 1 h or 2 weeks [171]. However, there is motivation to initiate stem cell therapy before the localized inflammatory response has subsided, as inflammatory cytokines are known to be involved in the homing, differentiation, and activation of MSCs [172].

3.2. Designing a cell delivery vehicle

Recent efforts have focused on biomaterial scaffolds as a means of delivering cells to the heart. An ideal scaffold would (i) introduce a critical cell population at the site of infarction, (ii) maintain their localization and viability over the course of the wound healing response, and (iii) promote the sustained production and release of beneficial paracrine factors into the ischemic region. Two general methods have been explored: surgical implantation of cells in pre-formed scaffolds, and the injection of cells in a polymer precursor solution that forms a cell-encapsulating hydrogel scaffold *in situ*. While implantation allows greater control over physical structure, it also necessitates open surgery. Injectable scaffolds represent a less invasive approach that could minimize damage to the already injured tissue, require shorter procedure and recovery times, and utilize existing catheter delivery technology to expedite translation to clinical practice. Emphasizing the role of cells as trophic mediators, this review focuses on injectable biomaterials that have demonstrated potential for delivering cells to the heart, their required material properties in terms of chemical, physical, and biological action, and their lifecycle in the body, from delivery to degradation.

Table 6Representative natural biomaterials investigated *in vivo* as potential injectable cell-delivery scaffolds for cardiac cell therapy.

Material class	Cell type/animal model	Experimental design	Gelation mechanism	Ref.
Alginate	Cell-free/rat	i.m. (100–150 μ L) 7 or 60 days post-MI	Ca ²⁺ dependent physical gelation	[173]
Alginate	Cell-free/swine	i.c. (1,2, or 4 mL) 4 days post-MI	Ca ²⁺ dependent physical gelation	[174]
Alginate (\pm RGD, \pm YIGSR)	Cell-free/rat	i.m. (130 μ L) 7 days post-MI	Ca ²⁺ dependent physical gelation	[175]
Alginate microspheres (+RGD)	Human bMSCs/nude rat	Closed chest, echocardiography guided injections (100 μ L) 7 days post-MI	Ca ²⁺ dependent physical gelation	[176]
Chitosan	Mouse ESCs/rat	i.m. at infarct border (1 \times 10 ⁷ ESCs in 100 μ L) 1 wk post-MI	Temperature responsive, glycerol phosphate-dependent physical gelation	[177]
Chitosan	Allogenic ASCs/rat	i.m. at infarct border (4 \times 10 ⁶ ASCs in 100 μ L)	Temperature responsive, glycerol phosphate-dependent physical gelation	[178]
Collagen	Cell-free/rat	i.m. at infarct centre (100 μ L) 1 wk post-MI	Physical fibrillogenesis of Zyderm [®] type II	[179]
Collagen	Allogenic bMSCs/rat	i.m. injection at infarct centre (2 \times 10 ⁶ bMSCs in 70 μ L) 1 wk post-MI	Physical fibrillogenesis of Zyderm [®] type I	[180]
Fibrin	Allogenic skeletal myoblasts/rat	i.m. (5 \times 10 ⁶ myoblasts in 50 μ L) 1 wk post-MI	Physical assembly of fibrin after co-injection of Tisseel VH sealant with thrombin and CaCl ₂	[181]
Fibrin (PEGylated)	Allogenic BMNCs/mouse	i.m. in infarct and border zone (5 \times 10 ⁵ BMNCs in 50 μ L) immediately post-MI	Physical assembly of fibrin after injection of PEGylated fibrinogen pre-mixed with thrombin	[182]
Fibrin	Allogenic ASCs/rat	i.m. to infarct border (5 \times 10 ⁵ ASCs in 100 μ L) 1 wk post-MI	Physical assembly of fibrin after co-injection of Tisseel VH sealant with thrombin and CaCl ₂	[183]
Hyaluronic acid (methacrylated)	Cell-free/sheep	i.m. to infarct area (20 \times 0.3 mL) 30 min post-MI	Covalent radical polymerization with APS and TEMED initiators	[184]
Hyaluronic acid (HEMA)	Cell-free/sheep	i.m. to infarct area (20 \times 0.3 mL) 30 min post-MI	Covalent radical polymerization with APS and TEMED initiators	[185]
Matrigel	Allogenic ESCs/mouse	i.m. to infarct (1 \times 10 ⁶ in 50 μ L) immediately post-MI	Temperature responsive physical gelation	[186]
Matrigel (+collagen)	Allogenic neonatal cardiomyocytes/rat	i.m. to infarct (2–3 \times 10 ⁶ in 150 μ L) 3 wks post-MI	Temperature responsive physical gelation	[187]
Matrigel (+pro-survival factors)	hESC-derived cardiomyocytes/ athymic nude rat	i.m. to infarct (10 \times 10 ⁶ hESCs in 70 μ L) 4 days post-MI	Temperature responsive physical gelation	[188]

i.m., intramyocardial; i.c. intracoronary; PEG, Poly(ethylene glycol); HEMA, hydroxyl ethyl methacrylate.

Tables 6 and 7 highlight the major classes of injectable biomaterials that have been identified as potential delivery scaffolds.

3.3. Material requirements

3.3.1. Injectability

The injectable scaffold approach introduces several design requirements:

- The material and cell suspension must be a catheter-transferable liquid for either intracoronary, epicardial, or transendocardial injection;
- Once injected, the solution must quickly undergo a gelling process to prevent dissemination of the material and cells without damaging surrounding tissues or reducing stem cell viability and function;
- The gelled scaffold must maintain structural integrity over the course of the cell therapy to negate the risk of peripheral or cerebral embolism;
- To obviate the requirement of surgical removal, the biomaterial should ultimately degrade in a manner that does not harm the local tissue or adversely affect its function, while yielding non-toxic and readily metabolized or eliminated degradation products.

The three routes of injection feature unique risks and benefits. Intracoronary delivery is similar to balloon angioplasty, and therefore utilizes well-established and common tools and techniques. While it is the least invasive approach and does not directly puncture the myocardium, it provides the lowest control over the amount and location of the delivered material and raises the risk of

microembolization [198]. Epicardial injection allows for precise control over delivery and good visualization; however, the necessary open-chest or endoscopic surgery requires the puncture or removal of the pericardium, significantly increasing the risk to the patient. The endocardial route provides similar access directly to the ischemic tissue, while the use of a catheter bypasses the need for surgery. However, visualization requires indirect imaging, and injecting material into the beating myocardial wall is technically challenging, increasing the risk of material dissemination into the ventricle [199]. Particularly in the cardiac setting, material dissemination must be tightly controlled to mitigate the potentially critical risk of systemic embolization.

Various non-cytotoxic mechanisms of gel-formation have been investigated, including two-component chemical cross-linking, UV photo-cross-linking, and thermal- or pH-induced physical gelling. The timescale and conditions of gelation are critical factors. If gelation occurs prematurely or too rapidly, the catheter could become clogged; if gelation is delayed, the biomaterial would provide no benefit over liquid-phase delivery. Ideally, the gelation mechanism would allow for consecutive injections at multiple sites.

3.3.2. Mechanical properties

Appropriate mechanical properties of the gelled scaffold are critical for supporting the damaged ventricular wall. A scaffold providing strength and thickness to the myocardium could temporarily compensate for the loss of contractile tissue, thereby reducing the extent of remodeling and scar formation required to balance hemodynamic forces [200,201]. Many groups have focused on designing gels with stiffness matching healthy myocardium, based on an end-diastolic elastic modulus of approximately 10 kPa

Table 7

Representative synthetic biomaterials investigated *in vivo* as potential injectable cell-delivery scaffolds for cardiac cell therapy.

Material class	Cell type/animal model	Experimental design	Gelation mechanism	Ref.
Self-assembling peptides (with IGF-1)	Allogenic CPCs/rat	i.m. to infarct border (1×10^5 CPCs in $5 \mu\text{L} + 5 \mu\text{L}$ peptides) immediately post-MI	Physical self-assembly	[189]
Self-assembling peptides	Autologous BMNCs/swine	i.m. to infarct (1×10^8 BMNCs in 2 mL) immediately post-MI	Physical self-assembly	[190]
mPEG-PCL-mPEG with α -cyclodextrin	Allogenic bMSCs/rabbit	i.m. to infarct (2×10^7 bMSCs in $4 \times 50 \mu\text{L}$) 7 days post MI	Physical assembly of copolymer within α -cyclodextrin cavities	[191]
8a-PEG-VS + cysteine-flanked peptide	hESCs-derived ELCs and SMLCs/nude rat	i.m. to infarct (0.66×10^6 ELC and 0.33×10^6 SMLC in $60 \mu\text{L}$) 1 h post-MI	Covalent Michael type addition between peptide thiol groups and vinyl sulfone groups	[192]
Oligo (PEG fumarate) + PEGDA	Mouse ESCs/rat	i.m. to infarct border (1×10^6 ESCs in $3 \times 33 \mu\text{L}$) 1 wk post-MI	Covalent radical polymerization with APS and TEMED initiators	[193]
PVL-PEG-PVL	Cell-free/rat	i.m. to infarct border ($4 \times 25 \mu\text{L}$) 7 days post-MI	Physical gelation via temperature-dependent hydrophobic aggregation	[194]
PNIPAAm-co-(HEMA-PCL -grafted dextran)	Cell-free/rabbit	i.m. ($4 \times 50 \mu\text{L}$) 4 days post-MI	Physical gelation via temperature-dependent hydrophobic aggregation	[195]
PNIPAAm-co-AAC-co-HEMApTMC	Cell-free/rat	i.m. into infarct and border ($5 \times 100 \mu\text{L}$) 2 wks post-MI	Physical gelation via temperature-dependent hydrophobic aggregation	[196]
PNIPAAm-co-AAC + peptide cross-linker + RGD-grafted pAAc	Allogenic bMSCs/mouse	i.m. injection into infarct border (1×10^5 bMSCs in $10 \mu\text{L}$) immediately post-MI	Physical gelation via temperature-dependent hydrophobic aggregation	[197]

i.m., intramyocardial; i.c., intracoronary; CPCs, cardiac progenitor cells; PEG, Poly(ethylene glycol); PCL, poly(ϵ -caprolactone); 8a-PEG-VS, 8-arm PEG vinyl sulfone; ELC, endothelial-like cells; SMLC, smooth muscle-like cells; PEGDA, PEG diacrylate; PVL, poly(δ -valerolactone); pNIPAAm, poly(*N*-isopropyl acrylamide); AAC, acrylic acid; HEMA, hydroxyl ethyl methacrylate; pTMC, poly(trimethylene carbonate).

[200]. There is evidence that stiffer gels with elastic moduli over 40 kPa are more effective at preventing maladaptive remodeling [184]. This approach somewhat counter-intuitively mimics the properties of fibrotic scar, as a stiffer material may better stabilize the myocardium and bear more wall stress, leading to reduced infarct expansion and LV dilation [184]. An optimal biomaterial should provide the mechanical stiffness to passively balance peak distending forces during end-systole, since these forces are a cause of the maladaptive remodeling process [184]. Subsequent material degradation should occur in response to cellular infiltration, vascularization, and formation of contractile tissue, such that the load originally borne by the scaffold is transferred to the regenerated myocardium. The scaffold must also be pliable and robust, withstanding fatigue failure at least until the tissue has recovered. Aligning mechanical properties with the surrounding tissues may additionally minimize physical irritation, promote integration at the site of delivery, and recreate a myocardial-like microenvironment for the delivered MSCs.

3.3.3. Cell–material interactions

Complex signaling interactions between the scaffold, the surrounding extracellular matrix (ECM), and the cells (both transplanted and host) must be considered across a range of length scales:

- Direct contact interactions: including mechanical properties such as stiffness of the material and bioactive surface properties, whether inherent or engineered.
- Microenvironmental interactions: including the microstructure and porosity of the scaffold; paracrine signaling molecules released by transplanted cells or the scaffold itself; and the formation of a regenerative milieu by locally concentrating cytokines.
- Systemic interactions: the recruitment of host stem cells from the systemic circulation and the prevention of donor cell migration; promotion of vascularization within and surrounding the scaffold; mediation of the host immune response.

Scaffold properties are influential in directing the cellular response, including migration [202], phenotype [203], and differentiation [204]. The microstructural properties of the scaffold,

including cross-link density and porosity, also play critical roles in cell therapy by influencing the diffusion of nutrients, oxygen, and cytokines [205,206].

4. Injectable biomaterials as scaffolds for MI cell therapy

4.1. Alginate

Alginate, a natural polysaccharide derived from algae, is composed of (1-4)-linked β -D-mannuronic acid (M unit) and α -L-guluronic acid (G unit) arranged in block and atactic regions (Fig. 2a). The addition of divalent cations such as Ca^{2+} causes the binding of G unit blocks on adjacent chains, rapidly forming an ionic cross-linked network. The elevated concentration of calcium ions in the post-MI environment has been exploited as a location-dependent *in situ* cross-linking mechanism [173]. Dissociation of G unit blocks occurs due to the outward flux of ions to the surrounding medium in a process that can potentially result in uncontrolled and typically slow hydrogel degradation kinetics. This mechanism, however, is well suited to myocardial applications: as the post-infarction environment recovers, the local calcium ion concentration decreases, increasing the rate of alginate dissociation and clearance from the region. Dissociated alginate that enters the circulation may be cleared by the kidneys if below the renal excretion limit of $48 \text{ kg} \cdot \text{mol}^{-1}$ weight average molecular weight [207]. Degradation rates may be increased by combining alginates with cross-linking G blocks of varying length [208] and by partial oxidation to facilitate hydrolysis [209]. As the hydrophilic nature of alginate limits protein adsorption, scaffolds are commonly functionalized with cell-adhesive peptides [175,176,210]. The functionalizable and non-thrombogenic [173,175] nature of alginate has motivated its use as an injectable cardiac biomaterial, both cell-free and as a delivery vehicle for cell therapy.

Yu et al. compared the injection of alginate and fibrin glue in a chronic (5 week old) LV aneurysm model [211]. Both materials increased arteriogenesis compared to saline controls, thickened the infarct wall, and improved LV function 2 days after administration. After 5 weeks, alginate, which was still present, improved LVFS, while fibrin glue, which had resorbed, resulted in LVFS similar to controls. Morphometric improvements remained however, indicating a beneficial effect of fibrin even after resorption.




Fig. 2. Chemical structures of common naturally-derived hydrogels explored as cell delivery vehicles for cardiac regeneration. (a) Alginate, composed of (1-4)-linked β -D-mannuronic acid (M) and α -L-guluronic acid (G). (b) Chitosan, composed of β -(1,4)-linked D-glucosamine and N-acetyl-D-glucosamine. (c) Hyaluronic acid, composed of β -(1,4)-D-glucuronic acid and β -(1,3)-N-acetyl-D-glucosamine.

Landa et al. injected an aqueous solution of $30\text{--}50\text{ kg}\cdot\text{mol}^{-1}$ sodium alginate combined with calcium gluconate via catheter into recent (7 day) and aged (2 month) infarcts in a rat MI model [173]. Histological results indicated that the material had been replaced by connective tissue within 6 weeks. In both groups, after 8 weeks alginate resulted in a thicker scar and attenuation of systolic and diastolic dysfunction compared to saline controls. When delivered in recent infarcts, alginate also reduced ventricular dilation. This material was also used to evaluate the efficacy of intracoronary injection in a pig model, 3–4 days after MI. The alginate diffused from the leaky coronary vasculature into the infarcted myocardium, without generating thrombi or further infarction. By 60 days, the material was replaced by myofibroblasts and collagen, reversed LV enlargement, and increased scar thickness.

Alginate has been modified with synthetic peptide sequences in attempts to induce cellular infiltration, adhesion, matrix

deposition, and vascularization at the infarct site. These attempts have met with mixed results. Tsur-Gang et al. functionalized $30\text{--}50\text{ kg}\cdot\text{mol}^{-1}$ alginate with RGD and YIGSR motifs at a degree of substitution of 0.2% of uronic acid monomer, which improved cardiofibroblast adhesion *in vitro*, but increased solution viscosity when cross-linked with calcium gluconate [175]. Injected in a rat MI model, the modified alginate gel reduced the therapeutic efficacy in terms of scar thickness, ventricle diameter, and function, and did not induce any cellular infiltration or angiogenesis. The reduced physical benefit could potentially result from the poor material distribution upon injection due to increased viscosity, while the lack of cellular response to the peptide modification was likely due to the low level of functionalization. Yu et al. mixed alginate with a commercially available high-density RGD-alginate complex at a ratio of 1:4 [210]. Injected with CaCl_2 5 weeks post MI in a rat model, alginate with and without RGD-modification

improved LV function, while RGD-modification induced a greater angiogenic response. While the level of RGD functionalization was not reported, this study demonstrates the importance of cell–material interactions for angiogenesis. The same material was used to deliver human MSCs via microsphere encapsulation in a rat MI model [176]. Both encapsulated MSCs and microspheres alone maintained LV shape, limited remodeling, reduced the infarct area, and increased arteriole density. Compared to MSCs delivered in saline, which were only retained for one day, MSCs delivered in microspheres were detected after up to two weeks, although the overall retention rate was not reported.

4.2. Chitosan

Chitosan is the deacetylated form of the linear polysaccharide chitin, a component of crustacean exoskeletons. Its cationic molecular structure of random β -(1,4)-linked D-glucosamine and N-acetyl-D-glucosamine facilitates electrostatic interactions with anionic glycosaminoglycans (GAGs) and proteoglycans of the ECM (Fig. 2b) [212]. *In vivo*, chitosan undergoes biodegradation primarily by lysozyme at a rate inversely related to the degree of deacetylation. Degradation produces oligosaccharides [213], which have been suggested to promote vascular endothelial cell migration and angiogenesis [214]. Combined with innate biocompatibility, antibacterial, and wound healing properties, the amenability of chitosan to chemical modification has drawn interest to it as an injectable biomaterial.

Chenite et al. developed a temperature-responsive hydrogel system by combining chitosan with a polyol-phosphate salt at physiologic pH [215]. The resulting liquid solution gels when heated to body temperature through hydrogen bonding and electrostatic and hydrophobic interactions. Lu et al. used this mechanism to encapsulate ESCs and deliver them 1 week post-infarction in a rat MI model [177]. Compared to PBS, chitosan reportedly improved ESC retention in terms of cellular graft area from $9.91 \pm 1.80\%$ to $17.48 \pm 3.30\%$ (24 h) and from $6.41 \pm 1.15\%$ to $12.93 \pm 1.90\%$ (4 weeks). At 4 weeks, the scaffold, both with and without cells, significantly improved heart function including ESD, EDD, LVEF, wall thickness, and microvessel density within the infarct. Liu et al. utilized the same material to deliver ASCs in a rat MI model, and investigated the ability of chitosan to protect the ASCs from reactive oxygen species (ROS) generated in the ischemic zone [178]. They found that ROS, specifically H_2O_2 , inhibited the expression of matrix adhesion molecules $\beta 1$ and αV , as well as other cell-adhesion pathways, and contributed to apoptosis. Chitosan attenuated these effects, improving adhesion and significantly increasing ASC viability. *In vivo*, ASCs delivered in chitosan resulted in increased stem cell homing and revascularization. The authors suggest that GAGs electrostatically associated with the chitosan may have enhanced the paracrine effects of ASCs by improving retention and concentration of cytokines and growth factors.

4.3. Collagen

Collagen has been used in numerous tissue engineering applications and is available in several commercial forms. Injectable collagen hydrogels are prepared by acid-solubilization of collagen, primarily type I [216]. Solution stability is determined by the electrostatic and hydrophobic interactions between amino acid groups, which are respectively affected by solution pH/ionic strength and temperature. Upon *in vivo* injection at neutral pH and 37°C , collagen undergoes entropy-driven self-assembly through fibrillogenesis to form a physically cross-linked hydrogel capable of encapsulating cells [217]. While supporting cell attachment,

viability, and proliferation, a drawback of injectable collagen is that the process of solubilization results in a loss of structural properties, including stiffness and elasticity. Collagen hydrogels formed under physiological conditions typically have compressive moduli between 7 and 14 kPa, depending on salt concentration [218].

Using radio-labeled MSCs injected in a rat MI model, Dai demonstrated that the commercially available collagen scaffold Zyderm at 35 mg/mL reduced MSC dissemination compared to delivery in saline [180]. Specifically, when collagen scaffold delivery was compared to saline, fewer of the 13 rats had MSCs present in the lungs (2 vs. 3), liver (0 vs. 4), spleen (4 vs. 11), and non-infarcted myocardium (5 vs. 9) at 4 weeks following injection. However, compared to the LVEF resulting from saline treatment ($54.9 \pm 3.4\%$), MSCs in collagen failed to improve LVEF ($53.2 \pm 1.4\%$), while MSCs in saline, as well as collagen alone increased the LVEF ($58.6 \pm 2.4\%$, $58.1 \pm 2.5\%$ respectively) [180]. The authors postulated that the collagen may have reduced long-term MSC survival by impairing the diffusion of oxygen and nutrients to the encapsulated cells, and by limiting microvascular ingrowth into the interstitial space.

4.4. Fibrin

Fibrin is a natural polymer involved in the coagulation cascade that contains intrinsic sites for cell binding. A fibrin hydrogel is formed when fibrinogen is cleaved by the enzyme thrombin, yielding insoluble fibrin peptides, which rapidly aggregate to form a fibril network mesh capable of encapsulating cells [219]. The physical properties of fibrin depend on the fibrinogen concentration and self-assembly kinetics. Higher thrombin concentrations result in faster gelation times and produce a tighter network of finer, thinner fibers [220]. By varying thrombin concentrations, cell-seeded fibrin scaffolds have been produced with elastic moduli of approximately 10–30 kPa, a range suitable for myocardial tissue applications [221]. Depending on fibrin concentration, fibrinolysis occurs naturally over days to weeks, and can be slowed by inclusion of aprotinin, a protease inhibitor [219]. Degradation produces non-immunogenic amino acid products with known angiogenic properties [222]. Under anoxic and glucose-deprived conditions *in vitro*, fibrin encapsulation of marrow-derived CSCs resulted in significantly improved survival rates over 7 days [223].

Christman et al. compared injecting bovine serum albumin (BSA) as a control to fibrin, with and without skeletal myoblasts, in a rat MI model [224]. With and without cells, fibrin reduced the percentage of infarcted ventricle wall ($17.5 \pm 3.4\%$ and $19.7 \pm 3.8\%$, respectively, compared to $26.5 \pm 2.2\%$ for cell-free BSA), and also increased arteriole density in the infarct and border zone ($13 \pm 3\text{ mm}^{-2}$ and $16 \pm 1\text{ mm}^{-2}$, compared to $6 \pm 2\text{ mm}^{-2}$ for cells in BSA).

Zhang et al. utilized a two-component fibrin/thrombin material with aprotinin to deliver ASCs in a rat MI model. ASCs delivered in fibrin covered significantly more of the infarct compared to ASCs delivered in PBS after 24 h ($19.10 \pm 3.13\%$ vs. $14.16 \pm 2.73\%$), and this significant difference persisted after four weeks ($11.52 \pm 2.34\%$ vs. $5.85 \pm 1.35\%$). After four weeks, ASCs in fibrin compared to ASCs in PBS and fibrin alone resulted in significant reductions in EDD ($5.91 \pm 0.24\text{ mm}$ vs. $6.60 \pm 0.38\text{ mm}$ and $6.70 \pm 0.41\text{ mm}$) and ESD ($4.24 \pm 0.26\text{ mm}$ vs. $5.04 \pm 0.22\text{ mm}$ and $5.24 \pm 0.35\text{ mm}$), as well as modest improvements in LVEF ($61.77 \pm 4.14\%$ vs. $55.75 \pm 3.41\%$ and $53.14 \pm 3.59\%$) and LVFS ($28.16 \pm 2.23\%$ vs. $24.11 \pm 1.97\%$ and $23.24 \pm 2.44\%$) [183].

4.5. Hyaluronic acid

Hyaluronic acid (HA) is an anionic linear glycosaminoglycan disaccharide polymer of β -(1,4)-D-glucuronic acid and β -(1,3)-N-

acetyl-D-glucosamine, ranging in size from 5 to 20,000 kg·mol⁻¹ *in vivo* (Fig. 2c). It is a widely distributed component of the ECM in mammalian tissues, and has been chemically modified to produce numerous biomaterials [225]. HA is involved in many cellular processes including proliferation and differentiation, and its abilities to promote angiogenesis and suppress fibrous tissue formation have drawn attention to its potential as a cardiac biomaterial [226].

In 2007, Kim et al. developed an acrylated 170 kg·mol⁻¹ HA derivative, which when combined with 10 kg·mol⁻¹ tetrathiolated four-armed poly(ethylene glycol), underwent a Michael-type addition reaction to form a hydrogel in 10 min under physiological conditions possessing an elastic modulus of 1.8 kPa [227]. Originally intended for bone regeneration, *in vivo* tests indicated the gel was a suitable scaffold for MSC and growth factor delivery. Despite its low modulus, this material was tested *in vivo* as a cell-free scaffold in a rat MI model, injected 2 weeks post-infarction [228]. Compared to untreated rats at 4 weeks post-implantation, the gel significantly increased wall thickness by 200%, decreased infarct area by 53%, and enhanced generation of arterioles and capillaries in the border region by 152% and 148%, respectively.

In 2010, Ilkovits et al. adapted a process used to methacrylate dextran [229] to produce a methacrylated 74 kg·mol⁻¹ HA (MeHA) which underwent radical polymerization when initiated with ammonium persulfate (APS) and tetramethylethylenediamine (TEMED) [184]. At low and high levels of methacrylation (30% and 60% substitution), the elastic modulus increased from 8 kPa to 43 kPa. By increasing the initiator concentration from 5.0 to 12.5 mM APS and 5.0–6.25 mM TEMED, the gelation time was reduced from 4.5 to 2.5 min. While not used to deliver cells, these investigations illustrate the influence of scaffold mechanical properties. The lower initiator concentrations facilitated deeper tissue penetration during injection, independent of methacrylation levels. This enhanced infiltration allowed for evaluation of the effects of hydrogel mechanical properties on tissue remodeling, independent of degradation rates and other material properties, which were consistent between the materials. MeHA was injected at 4% w/v in an ovine MI model, 30 min post-infarction. While both hydrogels significantly increased apex wall thickness (7.02 and 6.54 mm, respectively, versus 2.13 mm in controls), only the higher-modulus gel significantly increased the elastic modulus of the infarcted myocardium (from 5.8 ± 1.5 kPa to ~40 kPa). The slow gelation of the less-modified MeHA resulted in a wider distribution through the myocardium and no change in myocardial stiffness. Only the higher modulus gel produced significant reductions in infarct area (23.9% versus 28.6% in controls) and improvements in end-diastolic and end-systolic volumes (increasing only 1.7 and 1.9 times over baseline, respectively, versus 2.1 and 2.4 times in controls).

Similarly, Tous et al. produced hydroxyethyl methacrylate HA (HeMA-HA) hydrogels that underwent tuneable hydrolytic degradation [185]. At low and high levels of functionalization, *in vitro* degradation increased from approximately 3–10 weeks, and the elastic modulus increased from 8 to 33 kPa, respectively. Using the same ovine MI model, HeMA-HA gels were compared to hydrolytically stable MeHA gels. 8 weeks after injection, HeMA-HA with low functionalization was no longer present in the tissue, and only small amounts of HeMA-HA with high functionalization remained. While both HeMA-HA increased inflammation, potentially due to degradation product release, all HA gels increased vessel formation. The stable MeHA gels maintained myocardial thickness more effectively. While both high-modulus gels prevented LV enlargement at 2 weeks, only the stable MeHA maintained the effects at 8 weeks, implying that long-term wall stabilization is required for volume maintenance.

4.6. Matrigel™

Matrigel™ is a commercialized material composed of an assortment of ECM proteins, primarily laminin, collagen IV, and entactin, derived from mouse tumor cells. Based on their previous success in directly injecting ESC-derived human cardiomyocytes to uninjured hearts in nude rats [230], Laflamme et al. investigated a similar cell delivery approach in an infarcted rat heart model [188]. Cell retention was much lower in the MI model, where only 18% of hearts contained grafted cells compared to 90% in the uninjured model, and the observed grafts were generally much smaller. This failure was attributed to cell death from mitochondrial damage, ischemic shock, and anoikis. Cardiomyocyte delivery was reattempted with a cocktail of pro-survival factors and a Matrigel™ delivery vehicle to prevent anoikis. Consistent with previous results, no engraftment was observed when the cells were delivered alone. When delivered in Matrigel™, 100% of the infarcts had detectable grafts, which were four-fold larger when combined with the delivery of pro-survival factors. Transplanted cardiomyocytes persisted and were proliferating after 4 weeks, with the vast majority localized in the central regions of the infarct. Despite these promising results, the sarcoma cell line origin of Matrigel™ limits its clinical translatability [231].

4.7. Self-assembling peptides

Self-assembling peptides (SAPs) are oligopeptides composed of alternating hydrophilic and hydrophobic amino acids, 8 to 16 residues in length, capable of forming cell-supportive scaffolds under physiological conditions. At low pH and osmolarity, SAPs remain in solution, but when exposed to physiological conditions upon injection, rapidly self-assemble into fibers forming a stable macroporous membranous matrix [232]. SAPs can be modified to present growth factors and cellular signals to further promote cell attachment, growth, and differentiation, as well as matrix vascularization. Further, SAPs are capable of binding clinically relevant levels of growth factors such as platelet-derived growth factor (PDGF), VEGF, basic fibroblast growth factor (bFGF), and angiopoietin-1, making them potential candidates for growth factor delivery [233].

Davis et al. developed a SAP system that provided sustained, controlled release of IGF-1 tethered to the scaffold via a biotin-tetravalent streptavidin complex [234]. In a rat MI model, CPCs delivered in the IGF-1 tethered peptides significantly attenuated ventricular dilation and LVFS compared to CPCs alone and peptides with tethered IGF-1 alone. Compared to CPCs alone and peptides with IGF-1 alone, the combined CPC and IGF-1 therapy also increased the number of newly formed myocytes by 32% and 230%, respectively, and increased the length and density of newly formed arterioles by 73% and 83%, respectively [189].

4.8. Injectable hydrogel scaffolds based on synthetic polymers

Synthetic polymers offer several advantages over natural materials, including batch-to-batch consistency, precise control over mechanical and degradation properties, and amenability to chemical functionalization [235]. However, synthetic polymers often lack the innate bioactivity of natural materials, affecting cellular interactions, including migration, adhesion, and remodeling.

4.8.1. Poly(ethylene glycol)-based hydrogels

Poly(ethylene glycol) (PEG) is a water soluble polymer exhibiting low protein adsorption and immunogenicity and is widely used as a synthetic matrix in tissue engineering [235]. Wu et al.

developed a supramolecular tri-block methoxy PEG-*b*-poly(ϵ -caprolactone)-*b*-methoxy PEG (MPEG-PCL-MPEG) hydrogel with corresponding block weights of 5000–6110–5000 Da [236]. Upon mixing with α -cyclodextrin, a matrix was formed by linear MPEG-PCL-MPEG threading into cavities of α -cyclodextrin. Injected in a rabbit MI model, this material prevented scar expansion and wall thinning, and also improved ESD and EDD after 4 weeks. However, unlike most natural biomaterials, it did not induce neovascularization [237]. This hydrogel was used to deliver bMSCs in a rabbit MI model. After four weeks, delivery via scaffold compared to culture media significantly improved cell retention (2150 ± 235 cells/mm² vs. 845 ± 156 cells/mm²), LVEF (62% vs. 47%), EDD (145 mm vs. 175 mm), and infarct size (20% vs. 30%), and the hydrogel was completely absent from the infarct, likely by chain dissociation and diffusion [191].

Kraehenbuehl et al. have developed a hydrogel system using a multi-arm PEG vinyl sulfone that undergoes *in situ* cross-linking via Michael addition with a cysteine residue-flanked peptide sequence [238]. The peptide sequences were engineered to include a matrix metalloproteinase (MMP) substrate site, allowing for complete hydrogel degradation within 6 weeks, as well as an RGDSP cell adhesive ligand. This material was used to deliver endothelial- and smooth muscle-like cells derived from human ESCs, along with thymosin β 4 (T β 4), a pro-angiogenic and pro-survival factor, in a rat MI model 1 h post-infarction [192]. *In vivo*, there was approximately 80% release within 3 days of the 2.5 μ g of T β 4, and 95% after 6 weeks. At both time points, combined ESC and growth factor delivery, the hydrogel was found to preserve contractile function performance, attenuate LV dilation, and decrease infarct size. The density of small vessels (2–50 μ m in perimeter) in the infarct area was approximately three-fold greater than in PBS-treated controls, potentially due in part to the elevation in the human ESC-derived cytokines VEGF, EGF, and HGF that were detected after 1 day.

4.8.2. Poly(*N*-isopropylacrylamide)

Poly(*N*-isopropylacrylamide) (PNIPAAm)-based polymers have been investigated as hydrogel scaffolds due to their thermo-gelling nature. With a lower critical solution temperature slightly below body temperature, PNIPAAm hydrogels undergo a phase change into an insoluble gel when injected into the body. However, as PNIPAAm is non-biodegradable, the eventual solubilization *in vivo* may produce high molecular weight fragments that cannot be cleared by the kidneys. For this reason, PNIPAAm is often combined with degradable copolymers to improve biocompatibility [196,239].

Wang et al. developed a thermosensitive composite hydrogel by combining PNIPAAm with the polysaccharide dextran [195]. First, 60–90 kg·mol⁻¹ dextran was grafted with hydrophobic poly(ϵ -caprolactone)-2-hydroxyl ethyl methacrylate to form an amphiphilic Dex-PCL-HEMA, which was subsequently copolymerized with NIPAAm. In solution at 8 wt%, the composite underwent reversible phase transition to gel in 30 s when raised from 20 to 37 °C. The products of *in vivo* degradation were found to be below 10 kg·mol⁻¹, and thus expected to be cleared from renal filtration. The hydrogel was first tested 4 days post-infarction in a rabbit MI model [195]. After 30 days, the material prevented scar expansion and wall thinning, and improved LV morphology and hemodynamic performance; however, no neovascularization was reported. The same material was used to deliver bone-marrow-derived mononuclear cells (BM-MNCs) in a similar animal model [240]. While the polymer alone produced similar benefits, when combined with BM-MNCs it also enhanced neovascularisation and prevented scar expansion, illustrating the positive influence cells can have when combined with an otherwise non-angiogenic material.

Wall et al. addressed biodegradability by designing a semi-interpenetrating network (sIPN) based on PNIPAAm and acrylic acid (AAc) copolymers [197]. NIPAAm and AAc at 95:5 were combined with 0.3 mol % of a diacrylated, MMP-labile peptide cross-linker, and a range of concentrations of 450 kg·mol⁻¹ poly(AAc) functionalized with a cell-binding RGD sequence. These components were initiated with APS to form the thermo-responsive sIPN. The sIPN was used to deliver bMSCs to a rat infarct and compared to saline alone, bMSCs in saline, sIPN alone, and bMSCs in Matrigel™. All injections with bMSCs versus cell-free controls resulted in significant functional improvements after 2 weeks in LVFS and LVEF. However, these benefits decreased after 6 weeks to levels no different than saline-treated rats. Over 6 weeks, only sIPN alone improved LV function to a level not significantly different from sham-operated, non-infarcted rats. This result could be due to increased degradation of the sIPN or Matrigel™ by encapsulated bMSCs, reducing its structural benefit to the myocardium. A higher percentage of hearts (38%) contained bMSCs when delivered in sIPNs as opposed to Matrigel™ (25%) or saline (0%). Separately, these results indicate the sIPN polymer has potential to improve LV function and promote bMSC retention and survival. Together, they reveal the challenge of timing the degradation of a scaffold to match tissue recovery, and indicate that many benefits seen in other studies may be transient during early time points.

5. The future of biomaterials for cell delivery to the ischemic heart

Regardless of the particular application, the first factor to take into consideration when designing a cell-based therapy is the selection of an appropriate regenerative cell population. Studies in animal models have demonstrated significant benefits following the injection of either bMSCs or ASCs, primarily in terms of increased vascularization and modest improvements in cardiac function. When directly comparing the two MSC populations, the abundance, ease of harvest, and high regenerative cell yield of adipose tissue are advantageous, and the presence of fat depots in the intra-thoracic cavity could allow for intraoperative isolation without requiring a secondary harvest site [47,70,241]. If *ex vivo* cell expansion was necessary, the higher proliferation rate of ASCs would minimize the culture times required as compared to bMSCs to obtain a clinically-relevant cell population [39]. Studies in animal models have also challenged the original hypothesis that direct differentiation of the exogenous populations into cardiovascular components represents the primary mechanism in stem cell-mediated tissue regeneration. In fact, while engraftment and differentiation of administered cells have been demonstrated to be possible, these events are rare and occur at best at very low levels. However, studies have consistently demonstrated improvements in cardiac function and ventricular remodeling following MSC therapy, providing evidence that paracrine-mediated effects may be the driving mechanism in cardiac regeneration in this context. Further, there is a growing body of work that suggests that treatment efficacy could be augmented through enhanced cellular retention and/or engraftment.

A fair appraisal of regenerative cell types depends on a delivery strategy that can consistently deliver and retain a clinically relevant number of cells to the heart. In this role, injectable biomaterials have repeatedly demonstrated therapeutic efficacy towards improving cellular localization and retention in the infarcted myocardium in a minimally invasive manner. In small and large animal models, injected hydrogels consistently ameliorated negative effects on cardiac morphology and function caused by MI. However, at present, the field faces several challenges. Most critically, the therapeutic mechanisms of injectable hydrogels are

unclear. Research has variously focused on structurally reinforcing the myocardium, activating the intrinsic healing response through bioactive molecules or degradation products, encouraging host cell infiltration, and delivering therapeutic cells. Until the relative contributions of these factors are explored, it is difficult to optimize any specific material for stem cell delivery. In this context, because of the vast inconsistency in research methods it is difficult to compare studies, and to select the most promising approaches that should be advanced to clinical consideration. Therefore, broader studies should be designed keeping consistency in animal models, methodology and time points of both administration and assessment, and only varying the material and cells employed.

While natural polymers offer innate biological activity, synthetic polymers are emerging as a highly-customizable platform for hydrogel formulation. Though the scaffolds described here have all exhibited some degree of efficacy in delivering cells, appropriate mechanical properties for physically supporting the myocardium during the course of healing has emerged as a critical factor, and should also be a focus of future delivery strategies. Future materials will likely borrow from both fields, combining the pro-angiogenic activity of natural polymers with the consistent and tailorable degradation and mechanical properties of synthetics.

Furthermore, when employing natural polymers, batch-to-batch variability and regulatory approval are among the largest challenges; synthetic polymers, however, face issues in scaling-up intricate reaction processes and sourcing some precursors. Due to specific gelation kinetics and conditions, new injection equipment may be required, as current catheter technology is inappropriate for many of these systems. Special conditions must be met to maintain the liquid polymer before injection, facilitate rapid gelation, and provide repeated deliveries during the period of operation. Entirely new equipment may need to be designed, including catheterized double-barrel injectors for two-component mixing, cooled catheters for thermal gelation, and UV light guides for photo-cross-linking. Ultimately, the cyto-compatibility of these processes will be the determinant factor. Researchers must consider these criteria, and the input from medical professionals, when designing gelation mechanisms.

Finally, clinical translation must be the ultimate consideration throughout development of the biomaterial scaffold, specifically in terms of material selection and delivery method. While functional benefits are observed in terms of LVEF, it is unclear if these metrics will translate into any clinical benefit for a patient, such as overall health, activity, and life expectancy. Also, the clinical significance of improved LVEF has been questioned since it is often not paralleled by variations in other parameters such as left ventricular end systolic volume and ejection fraction [152,242]. The end point of any therapy is to improve the patient's quality of life, ideally restoring the pre-pathology health status. In the particular case of heart conditions, this means allowing a patient to perform routine physical activities with no adverse effects. Therefore, clinical trials should always include physical performance analysis, as the success of a treatment should not only be evaluated by measuring changes in cardiac parameters but also by assessing if there are paralleled improvements in patient exercise tolerability.

References

- [1] Alwan A. Global status report on noncommunicable diseases 2010. World Health Organization; 2011.
- [2] Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Borden WB, et al. Executive summary: heart disease and stroke statistics—2013 update: a report from the American Heart Association. *Circulation* 2013;127:143–52.
- [3] Sutton MG, Sharpe N. Left ventricular remodeling after myocardial infarction: pathophysiology and therapy. *Circulation* 2000;101:2981–8.
- [4] Weiss JN, Qu Z, Chen P-S, Lin S-F, Karagueuzian HS, Hayashi H, et al. The dynamics of cardiac fibrillation. *Circulation* 2005;112:1232–40.
- [5] Mukherjee D, Bhatt DL, Roe MT, Patel V, Ellis SG. Direct myocardial revascularization and angiogenesis—how many patients might be eligible? *Am J Cardiol* 1999;84:598–600.
- [6] Kiernan TJ, Boilson BA, Sandhu GS, Lennon RJ, Roger VL, Barsness GW, et al. Nonrevascularizable coronary artery disease following coronary artery bypass graft surgery: a population-based study in Olmsted County, Minnesota. *Coron Artery Dis* 2009;20:106–11.
- [7] Velagaleti RS, Pencina MJ, Murabito JM, Wang TJ, Parikh NI, D'Agostino RB, et al. Long-term trends in the incidence of heart failure after myocardial infarction. *Circulation* 2008;118:2057–62.
- [8] Collins MC, Moore Jr JL, Burrows BJ, Kypson AP, Muller-Borer BJ. Early cell loss associated with mesenchymal stem cell cardiomyoplasty. *Open Tissue Eng Regen Med J* 2012;5:17–24.
- [9] Traverse JH, McKenna DH, Harvey K, Jorgensen BC, Olson RE, Bostrom N, et al. Results of a phase 1, randomized, double-blind, placebo-controlled trial of bone marrow mononuclear stem cell administration in patients following ST-elevation myocardial infarction. *Am Heart J* 2010;160:428–34.
- [10] Wöhrle J, Merkle N, Mailänder V, Nusser T, Schauwecker P, Scheidt von F, et al. Results of intracoronary stem cell therapy after acute myocardial infarction. *Am J Cardiol* 2010;105:804–12.
- [11] Lee J, Terracciano CM. Cell therapy for cardiac repair. *Br Med Bull* 2010;94:65–80.
- [12] Menasche P, Alfieri O, Janssens S, McKenna W, Reichenspurner H, Trinquart L, et al. The Myoblast Autologous Grafting in Ischemic Cardiomyopathy (MAGIC) trial: first randomized placebo-controlled study of myoblast transplantation. *Circulation* 2008;117:1189–200.
- [13] Veltman CE, Soliman Oll, Geleijnse ML, Vletter WB, Smits PC, Cate ten FJ, et al. Four-year follow-up of treatment with intramyocardial skeletal myoblasts injection in patients with ischaemic cardiomyopathy. *Eur Heart J* 2008;29:1386–96.
- [14] Nussbaum J, Minami E, Laflamme MA, Virag JAI, Ware CB, Masino A, et al. Transplantation of undifferentiated murine embryonic stem cells in the heart: teratoma formation and immune response. *FASEB J* 2007;21:1345–57.
- [15] Bolli R, Chugh AR, D'Amario D, Loughran JH, Stoddard MF, Ikram S, et al. Cardiac stem cells in patients with ischaemic cardiomyopathy (SCPIO): initial results of a randomised phase 1 trial. *Lancet* 2011;378:1847–57.
- [16] Bearzi C, Rota M, Hosoda T, Tillmanns J, Nascimbene A, De Angelis A, et al. Human cardiac stem cells. *Proc Natl Acad Sci U S A* 2007;104:14068–73.
- [17] Hosoda T. C-kit-positive cardiac stem cells and myocardial regeneration. *Am J Cardiovasc Dis* 2012;2:58–67.
- [18] Pittenger MF, Mackay AM, Beck SC, Jaiswal RK, Douglas R, Mosca JD, et al. Multilineage potential of adult human mesenchymal stem cells. *Science* 1999;284:143–7.
- [19] Gimble J, Guilak F. Adipose-derived adult stem cells: isolation, characterization, and differentiation potential. *Cytotherapy* 2003;5:362–9.
- [20] Fukumoto T, Sperling JW, Sanyal A, Fitzsimmons JS, Reinholz GG, Conover CA, et al. Combined effects of insulin-like growth factor-1 and transforming growth factor-beta1 on periosteal mesenchymal cells during chondrogenesis in vitro. *Osteoarthritis Cartil* 2003;11:55–64.
- [21] De Bari C, Dell'Accio F, Tylzanowski P, Luyten FP. Multipotent mesenchymal stem cells from adult human synovial membrane. *Arthritis Rheum* 2001;44:1928–42.
- [22] Jankowski RJ, Deasy BM, Huard J. Muscle-derived stem cells. *Gene Ther* 2002;9:642–7.
- [23] Miura M, Gronthos S, Zhao M, Lu B, Fisher LW, Gehron Robey P. SHED: stem cells from human exfoliated deciduous teeth. *Proc Natl Acad Sci U S A* 2003;100:5807–12.
- [24] Castro-Malaspina H, Ebell W, Wang S. Human bone marrow fibroblast colony-forming units (CFU-F). *Prog Clin Biol Res* 1984;154:209–36.
- [25] D'Ippolito G, Schiller PC, Ricordi C, Roos BA, Howard GA. Age-related osteogenic potential of mesenchymal stromal stem cells from human vertebral bone marrow. *J Bone Miner Res* 1999;14:1115–22.
- [26] Muschler GF, Nitto H, Boehm CA, Easley KA. Age- and gender-related changes in the cellularity of human bone marrow and the prevalence of osteoblastic progenitors. *J Orthop Res* 2001;19:117–25.
- [27] Mitchell JB, McIntosh K, Zvonick S, Garrett S, Floyd ZE, Kloster A, et al. Immunophenotype of human adipose-derived cells: temporal changes in stromal-associated and stem cell-associated markers. *Stem Cells* 2006;24:376–85.
- [28] Fraser JK, Wulur I, Alfonso Z, Zhu M, Wheeler ES. Differences in stem and progenitor cell yield in different subcutaneous adipose tissue depots. *Cytotherapy* 2007;9:459–67.
- [29] Maumus M, Peyrafitte J-A, D'Angelo R, Fournier-Wirth C, Bouloumié A, Casteilla L, et al. Native human adipose stromal cells: localization, morphology and phenotype. *Int J Obes (Lond)* 2011;35:1141–53.
- [30] Noël D, Caton D, Roche S, Bony C, Lehmann S, Casteilla L, et al. Cell specific differences between human adipose-derived and mesenchymal-stromal cells despite similar differentiation potentials. *Exp Cell Res* 2008;314:175–84.
- [31] De Ugarte DA, Morizono K, Elbarbary A, Alfonso Z, Zuk PA, Zhu M, et al. Comparison of multi-lineage cells from human adipose tissue and bone marrow. *Cells Tissues Organs* 2003;174:101–9.
- [32] De Ugarte DA, Alfonso Z, Zuk PA, Elbarbary A, Zhu M, Ashjian P, et al. Differential expression of stem cell mobilization-associated molecules on multi-lineage cells from adipose tissue and bone marrow. *Immunol Lett* 2003;89:267–70.

- [33] Yañez R, Lamana ML, García-Castro J, Colmenero I, Ramírez M, Bueren JA. Adipose tissue-derived mesenchymal stem cells have in vivo immunosuppressive properties applicable for the control of the graft-versus-host disease. *Stem Cells* 2006;24:2582–91.
- [34] Pachón-Peña G, Yu G, Tucker A, Wu X, Vendrell J, Bunnell BA, et al. Stromal stem cells from adipose tissue and bone marrow of age-matched female donors display distinct immunophenotypic profiles. *J Cell Physiol* 2011;226:843–51.
- [35] Kern S, Eichler H, Stoeve J, Klüter H, Bieback K. Comparative analysis of mesenchymal stem cells from bone marrow, umbilical cord blood, or adipose tissue. *Stem Cells* 2006;24:1294–301.
- [36] Rebelatto CK, Aguiar AM, Moretao MP, Senegaglia AC, Hansen P, Barchiki F, et al. Dissimilar differentiation of mesenchymal stem cells from bone marrow, umbilical cord blood, and adipose tissue. *Exp Biol Med (Maywood)* 2008;233:901–13.
- [37] Hare JM, Traverse JH, Henry TD, Dib N, Strumpf RK, Schulman SP, et al. A randomized, double-blind, placebo-controlled, dose-escalation study of intravenous adult human mesenchymal stem cells (prochymal) after acute myocardial infarction. *J Am Coll Cardiol* 2009;54:2277–86.
- [38] Izadpanah R, Trygg C, Patel B, Kriedt C, Dufour J, Gimble JM, et al. Biologic properties of mesenchymal stem cells derived from bone marrow and adipose tissue. *J Cell Biochem* 2006;99:1285–97.
- [39] Paul A, Srivastava S, Chen G, Shum-Tim D, Prakash S. Functional assessment of adipose stem cells for xenotransplantation using myocardial infarction immunocompetent models: comparison with bone marrow stem cells. *Cell Biochem Biophys* 2013;67:263–73.
- [40] Kim Y, Kim H, Cho H, Bae Y, Suh K, Jung J. Direct comparison of human mesenchymal stem cells derived from adipose tissues and bone marrow in mediating neovascularization in response to vascular ischemia. *Cell Physiol Biochem* 2007;20:867–76.
- [41] Bochev I, Elmadjian G, Kyurkchiev D, Tzvetanov L, Altankova I, Tivchev P, et al. Mesenchymal stem cells from human bone marrow or adipose tissue differently modulate mitogen-stimulated B-cell immunoglobulin production in vitro. *Cell Biol Int* 2008;32:384–93.
- [42] Ivanova-Todorova E, Bochev I, Mourdjeva M, Dimitrov R, Bukarev D, Kyurkchiev S, et al. Adipose tissue-derived mesenchymal stem cells are more potent suppressors of dendritic cells differentiation compared to bone marrow-derived mesenchymal stem cells. *Immunol Lett* 2009;126:37–42.
- [43] Melief SM, Zwaginga JJ, Fibbe WE, Roelofs H. Adipose tissue-derived multipotent stromal cells have a higher immunomodulatory capacity than their bone marrow-derived counterparts. *Stem Cells Transl Med* 2013;2:455–63.
- [44] Breitbach M, Bostani T, Roell W, Xia Y, Dewald O, Nygren JM, et al. Potential risks of bone marrow cell transplantation into infarcted hearts. *Blood* 2007;110:1362–9.
- [45] Baglioni S, Cantini G, Poli G, Francalanci M, Squecco R, Di Franco A, et al. Functional differences in visceral and subcutaneous fat pads originate from differences in the adipose stem cell. *PLoS One* 2012;7:e36569.
- [46] Park HT, Lee ES, Cheon Y-P, Lee DR, Yang K-S, Kim YT, et al. The relationship between fat depot-specific preadipocyte differentiation and metabolic syndrome in obese women. *Clin Endocrinol* 2012;76:59–66.
- [47] Russo V, Yu C, Belliveau P, Hamilton A, Flynn LE. Comparison of human adipose-derived stem cells isolated from subcutaneous, omental, and intrathoracic adipose tissue depots for regenerative applications. *Stem Cells Transl Med*. <http://dx.doi.org/10.5966/sctm.2013-0125>. Available from: <http://stemcellstm.alphamedpress.org/content/early/2013/12/19/sctm.2013-0125>; 2013.
- [48] Friedenstien AJ, Gorska JA, Kulagina NN. Fibroblast precursors in normal and irradiated mouse hematopoietic organs. *Exp Hematol* 1976;4:267–74.
- [49] Reyes M, Lund T, Lenvik T, Aguiar D, Koodie L, Verfaillie CM. Purification and ex vivo expansion of postnatal human marrow mesodermal progenitor cells. *Blood* 2001;98:2615–25.
- [50] Mosna F, Sensebé L, Krampera M. Human bone marrow and adipose tissue mesenchymal stem cells: a user's guide. *Stem Cells Dev* 2010;19:1449–70.
- [51] Tomita S, Li RK, Weisel RD, Mickle DAG, Kim EJ, Sakai T, et al. Autologous transplantation of bone marrow cells improves damaged heart function. *Circulation* 1999;100:II-247–II-256.
- [52] Kamihata H, Matsubara H, Nishie T, Fujiyama S, Tsutsumi Y, Ozono R, et al. Implantation of bone marrow mononuclear cells into ischemic myocardium enhances collateral perfusion and regional function via side supply of angioblasts, angiogenic ligands, and cytokines. *Circulation* 2001;104:1046–52.
- [53] Tang YL, Zhao Q, Zhang YC, Cheng L, Liu M, Shi J, et al. Autologous mesenchymal stem cell transplantation induce VEGF and neovascularization in ischemic myocardium. *Regul Pept* 2004;117:3–10.
- [54] Zhang S, Guo J, Zhang P, Liu Y, Jia Z, Ma K, et al. Long-term effects of bone marrow mononuclear cell transplantation on left ventricular function and remodeling in rats. *Life Sci* 2004;74:2853–64.
- [55] Tang J, Xie Q, Pan G, Wang J, Wang M. Mesenchymal stem cells participate in angiogenesis and improve heart function in rat model of myocardial ischemia with reperfusion. *Eur J Cardiothorac Surg* 2006;30:353–61.
- [56] Orlic D, Kajstura J, Chimenti S, Jakoniuk I, Anderson SM, Li B, et al. Bone marrow cells regenerate infarcted myocardium. *Nature* 2001;410:701–5.
- [57] Kudo M, Wang Y, Wani MA, Xu M, Ayub A, Ashraf M. Implantation of bone marrow stem cells reduces the infarction and fibrosis in ischemic mouse heart. *J Mol Cell Cardiol* 2003;35:1113–9.
- [58] Kajstura J, Rota M, Whang B, Cascapera S, Hosoda T, Bearzi C, et al. Bone marrow cells differentiate in cardiac cell lineages after infarction independently of cell fusion. *Circ Res* 2005;96:127–37.
- [59] Rota M, Kajstura J, Hosoda T, Bearzi C, Vitale S, Esposito G, et al. Bone marrow cells adopt the cardiomyogenic fate in vivo. *Proc Natl Acad Sci U S A* 2007;104:17783–8.
- [60] Shake JC, Gruber PJ, Baumgartner WA, Senechal G, Meyers J, Redmond JM, et al. Mesenchymal stem cell implantation in a swine myocardial infarct model: engraftment and functional effects. *Ann Thorac Surg* 2002;73:1919–25. discussion 1926.
- [61] Silva GV, Litovsky S, Assad JAR, Sousa ALS, Martin BJ, Vela D, et al. Mesenchymal stem cells differentiate into an endothelial phenotype, enhance vascular density, and improve heart function in a canine chronic ischemia model. *Circulation* 2005;111:150–6.
- [62] Gimble JM, Katz AJ, Bunnell BA. Adipose-derived stem cells for regenerative medicine. *Circ Res* 2007;100:1249–60.
- [63] Nishida S, Endo N, Yamagiwa H, Tanizawa T, Takahashi HE. Number of osteoprogenitor cells in human bone marrow markedly decreases after skeletal maturation. *J Bone Miner Metab* 1999;17:171–7.
- [64] Stenderup K, Justesen J, Eriksen EF, Rattan SI, Kassem M. Number and proliferative capacity of osteogenic stem cells are maintained during aging and in patients with osteoporosis. *J Bone Miner Res* 2001;16:1120–9.
- [65] Miyahara Y, Nagaya N, Kataoka M, Yanagawa B, Tanaka K, Hao H, et al. Monolayered mesenchymal stem cells repair scarred myocardium after myocardial infarction. *Nat Med* 2006;12:459–65.
- [66] Mazo M, Planat-Bénard V, Abizanda G, Pelacho B, Léobon B, Gavira JJ, et al. Transplantation of adipose derived stromal cells is associated with functional improvement in a rat model of chronic myocardial infarction. *Eur J Heart Fail* 2008;10:454–62.
- [67] Schenke-Layland K, Strem BM, Jordan MC, Deemedio MT, Hedrick MH, Roos KP, et al. Adipose tissue-derived cells improve cardiac function following myocardial infarction. *J Surg Res* 2009;153:217–23.
- [68] Wang L, Deng J, Tian W, Xiang B, Yang T, Li G, et al. Adipose-derived stem cells are an effective cell candidate for treatment of heart failure: an MR imaging study of rat hearts. *Am J Physiol Heart Circ Physiol* 2009;297:H1020–31.
- [69] Cai L, Johnstone BH, Cook TG, Tan J, Fishbein MC, Chen P-S, et al. IFATS collection: human adipose tissue-derived stem cells induce angiogenesis and nerve sprouting following myocardial infarction, in conjunction with potent preservation of cardiac function. *Stem Cells* 2009;27:230–7.
- [70] Bayes-Genis A, Soler-Botija C, Farré J, Sepúlveda P, Raya A, Roura S, et al. Human progenitor cells derived from cardiac adipose tissue ameliorate myocardial infarction in rodents. *J Mol Cell Cardiol* 2010;49:771–80.
- [71] Bai X, Yan Y, Song YH, Seidensticker M, Rabinovich B, Metzler R, et al. Both cultured and freshly isolated adipose tissue-derived stem cells enhance cardiac function after acute myocardial infarction. *Eur Heart J* 2010;31:489–501.
- [72] Valina C, Pinkernell K, Song YH, Bai X, Sadat S, Campeau RJ, et al. Intracoronary administration of autologous adipose tissue-derived stem cells improves left ventricular function, perfusion, and remodeling after acute myocardial infarction. *Eur Heart J* 2007;28:2667–77.
- [73] Alt E, Pinkernell K, Scharlau M, Coleman M, Fotuhi P, Nabzdyk C, et al. Effect of freshly isolated autologous tissue resident stromal cells on cardiac function and perfusion following acute myocardial infarction. *Int J Cardiol* 2010;144:26–35.
- [74] Zhang D-Z, Gai L-Y, Liu H-W, Jin Q-H, Huang J-H, Zhu X-Y. Transplantation of autologous adipose-derived stem cells ameliorates cardiac function in rabbits with myocardial infarction. *Chin Med J (Engl)* 2007;120:300.
- [75] Gronthos S, Franklin DM, Leddy HA, Robey PG, Storms RW, Gimble JM. Surface protein characterization of human adipose tissue-derived stromal cells. *J Cell Physiol* 2001;189:54–63.
- [76] Pittenger MF, Martin BJ. Mesenchymal stem cells and their potential as cardiac therapeutics. *Circ Res* 2004;95:9–20.
- [77] Hoover-Plow J, Gong Y. Challenges for heart disease stem cell therapy. *Vasc Health Risk Manag* 2012;8:99–113.
- [78] Baek SJ, Kang SK, Ra JC. In vitro migration capacity of human adipose tissue-derived mesenchymal stem cells reflects their expression of receptors for chemokines and growth factors. *Exp Mol Med* 2011;43:596–603.
- [79] Ma J, Zhang S, Sun A, Shen J, Chen L, Wang K, et al. Time course of myocardial stromal cell-derived factor 1 expression and beneficial effects of intravenously administered bone marrow stem cells in rats with experimental myocardial infarction. *Basic Res Cardiol* 2005;100:217–23.
- [80] Son BR, Marquez Curtis LA, Kucia M, Wysoczynski M, Turner AR, Ratajczak J, et al. Migration of bone marrow and cord blood mesenchymal stem cells in vitro is regulated by stromal-derived factor-1-CXCR4 and hepatocyte growth factor-c-met axes and involves matrix metalloproteinases. *Stem Cells* 2006;24:1254–64.
- [81] Bhakta S, Hong P, Koc O. The surface adhesion molecule CXCR4 stimulates mesenchymal stem cell migration to stromal cell-derived factor-1 in vitro but does not decrease apoptosis under serum deprivation. *Cardiovasc Res* 2006;7:19–24.
- [82] Ceradini DJ, Kulkarni AR, Callaghan MJ, Tepper OM, Bastidas N, Kleinman ME, et al. Progenitor cell trafficking is regulated by hypoxic gradients through HIF-1 induction of SDF-1. *Nat Med* 2004;10:858–64.
- [83] Wojakowski W, Tendera M, Michałowska A, Majka M, Kucia M, Maślankiewicz K, et al. Mobilization of CD34/CXCR4+, CD34/CD117+, c-met+ stem cells, and mononuclear cells expressing early cardiac, muscle, and endothelial markers into peripheral blood in patients with acute myocardial infarction. *Circulation* 2004;110:3213–20.
- [84] Kinnaird T, Stabile E, Burnett MS, Lee CW, Barr S, Fuchs S, et al. Marrow-derived stromal cells express genes encoding a broad spectrum of

- arteriogenic cytokines and promote in vitro and in vivo arteriogenesis through paracrine mechanisms. *Circ Res* 2004;94:678–85.
- [85] Kinnaird T, Stabile E, Burnett MS, Shou M, Lee CW, Barr S, et al. Local delivery of marrow-derived stromal cells augments collateral perfusion through paracrine mechanisms. *Circulation* 2004;109:1543–9.
- [86] Kobayashi T, Hamano K, Li T-S, Katoh T, Kobayashi S, Matsuzaki M, et al. Enhancement of angiogenesis by the implantation of self bone marrow cells in a rat ischemic heart model. *J Surg Res* 2000;89:189–95.
- [87] Liechty KW, MacKenzie TC, Shaaban AF, Radu A, Moseley AM, Deans R, et al. Human mesenchymal stem cells engraft and demonstrate site-specific differentiation after in utero transplantation in sheep. *Nat Med* 2000;6:1282–6.
- [88] Bartholomew A, Sturgeon C, Siatskas M, Ferrer K, McIntosh K, Patil S, et al. Mesenchymal stem cells suppress lymphocyte proliferation in vitro and prolong skin graft survival in vivo. *Exp Hematol* 2002;30:42–8.
- [89] Aggarwal S, Pittenger MF. Human mesenchymal stem cells modulate allogeneic immune cell responses. *Blood* 2005;105:1815–22.
- [90] McIntosh K, Zvonick S, Garrett S, Mitchell JB, Floyd ZE, Hammill L, et al. The immunogenicity of human adipose-derived cells: temporal changes in vitro. *Stem Cells* 2006;24:1246–53.
- [91] Yoon Y-S, Wecker A, Heyd L, Park J-S, Tkebuchava T, Kusano K, et al. Clonally expanded novel multipotent stem cells from human bone marrow regenerate myocardium after myocardial infarction. *J Clin Invest* 2005;115:326–38.
- [92] Makino S, Fukuda K, Miyoshi S, Konishi F, Kodama H, Pan J, et al. Cardiomyocytes can be generated from marrow stromal cells in vitro. *J Clin Invest* 1999;103:697–705.
- [93] Xu W, Zhang X, Qian H, Zhu W, Sun X, Hu J, et al. Mesenchymal stem cells from adult human bone marrow differentiate into a cardiomyocyte phenotype in vitro. *Exp Biol Med* (Maywood) 2004;229:623–31.
- [94] Shim WSN, Jiang S, Wong P, Tan J, Chua YL, Tan YS, et al. Ex vivo differentiation of human adult bone marrow stem cells into cardiomyocyte-like cells. *Biochem Biophys Res Commun* 2004;324:481–8.
- [95] Rangappa S, Fen C, Lee EH, Bongso A, Sim EKW, Wei EKS. Transformation of adult mesenchymal stem cells isolated from the fatty tissue into cardiomyocytes. *Ann Thorac Surg* 2003;75:775–9.
- [96] Dijk A, Niessen HWM, Zandieh Doulabi B, Visser FC, Milligen FJ. Differentiation of human adipose-derived stem cells towards cardiomyocytes is facilitated by laminin. *Cell Tissue Res* 2008;334:457–67.
- [97] Planat-Bénard V, Menard C, André M, Puceat M, Perez A, Garcia-Verdugo J-M, et al. Spontaneous cardiomyocyte differentiation from adipose tissue stroma cells. *Circ Res* 2004;94:223–9.
- [98] Sliwa A, Balwierz A, Kiec-Wilk B, Polus A, Knapp A, Dembinska-Kiec A. Differentiation of human adipose tissue SVF cells into cardiomyocytes. *Genes Nutr* 2009;4:195–8.
- [99] Oswald J, Boxberger S, Jørgensen B, Feldmann S, Ehninger G, Bornhäuser M, et al. Mesenchymal stem cells can be differentiated into endothelial cells in vitro. *Stem Cells* 2004;22:377–84.
- [100] Chen M-Y, Lie P-C, Li Z-L, Wei X. Endothelial differentiation of Wharton's jelly-derived mesenchymal stem cells in comparison with bone marrow-derived mesenchymal stem cells. *Exp Hematol* 2009;37:629–40.
- [101] Colazzo F, Chester AH, Taylor PM, Yacoub MH. Induction of mesenchymal to endothelial transformation of adipose-derived stem cells. *J Heart Valve Dis* 2010;19:736–44.
- [102] Ning H, Liu G, Lin G, Yang R, Lue TF, Lin C-S. Fibroblast growth factor 2 promotes endothelial differentiation of adipose tissue-derived stem cells. *J Sex Med* 2009;6:967–79.
- [103] Kinner B, Zaleskas JM, Spector M. Regulation of smooth muscle actin expression and contraction in adult human mesenchymal stem cells. *Exp Cell Res* 2002;278:72–83.
- [104] Rodriguez LV, Alfonso Z, Zhang R, Leung J, Wu B, Ignarro LJ. Clonogenic multipotent stem cells in human adipose tissue differentiate into functional smooth muscle cells. *Proc Natl Acad Sci U S A* 2006;103:12167–72.
- [105] Yang J, Song T, Wu P, Chen Y, Fan X, Chen H, et al. Differentiation potential of human mesenchymal stem cells derived from adipose tissue and bone marrow to sinus node-like cells. *Mol Med Rep* 2012;5:108–13.
- [106] Liu Y, Song J, Liu W, Wan Y, Chen X, Hu C. Growth and differentiation of rat bone marrow stromal cells: does 5-azacytidine trigger their cardiomyogenic differentiation? *Cardiovasc Res* 2003;58:460–8.
- [107] Rangappa S, Entwistle JWC, Wechsler AS, Kresh JY. Cardiomyocyte-mediated contact programs human mesenchymal stem cells to express cardiogenic phenotype. *J Thorac Cardiovasc Surg* 2003;126:124–32.
- [108] Zhu Y, Liu T, Song K, Ning R, Ma X, Cui Z. ADSCs differentiated into cardiomyocytes in cardiac microenvironment. *Mol Cell Biochem* 2008;324:117–29.
- [109] Plotnikov EY, Khryapenkova TG, Vasileva AK, Marey MV, Galkina SI, Isaev NK, et al. Cell-to-cell cross-talk between mesenchymal stem cells and cardiomyocytes in co-culture. *J Cell Mol Med* 2008;12:1622–31.
- [110] Metzeler R, Alt C, Bai X, Yan Y, Zhang Z, Pan Z, et al. Human adipose tissue-derived stem cells exhibit proliferation potential and spontaneous rhythmic contraction after fusion with neonatal rat cardiomyocytes. *FASEB J* 2011;25:830–9.
- [111] Perán M, Marchal JA, López E, Jiménez-Navarro M, Boulaiz H, Rodríguez-Serrano F, et al. Human cardiac tissue induces transdifferentiation of adult stem cells towards cardiomyocytes. *Cytotherapy* 2010;12:332–7.
- [112] Ramesh B, Bishi DK, Rallapalli S, Arumugam S, Cherian KM, Guhathakurta S. Ischemic cardiac tissue conditioned media induced differentiation of human mesenchymal stem cells into early stage cardiomyocytes. *Cytotechnology* 2012;64:563–75.
- [113] Alvarez-Dolado M, Pardo R, Garcia-Verdugo JM, Fike JR, Lee HO, Pfeffer K, et al. Fusion of bone-marrow-derived cells with Purkinje neurons, cardiomyocytes and hepatocytes. *Nature* 2003;425:968–73.
- [114] Scherschel JA, Soonpaa MH, Srouf EF, Field LJ, Rubart M. Adult bone marrow-derived cells do not acquire functional attributes of cardiomyocytes when transplanted into peri-infarct myocardium. *Mol Ther* 2008;16:1129–37.
- [115] Yoon C-H, Koyanagi M, Iekushi K, Seeger F, Urbich C, Zeiher AM, et al. Mechanism of improved cardiac function after bone marrow mononuclear cell therapy: role of cardiovascular lineage commitment. *Circulation* 2010;121:2001–11.
- [116] Sadat S, Gehmert S, Song Y-H, Yen Y, Bai X, Gaiser S, et al. The cardioprotective effect of mesenchymal stem cells is mediated by IGF-I and VEGF. *Biochem Biophys Res Commun* 2007;363:674–9.
- [117] Rehman J, Traktuev D, Li J, Merfeld-Clauss S, Temm-Grove CJ, Bovenkerk JE, et al. Secretion of angiogenic and antiapoptotic factors by human adipose stromal cells. *Circulation* 2004;109:1292–8.
- [118] Uemura R, Xu M, Ahmad N, Ashraf M. Bone marrow stem cells prevent left ventricular remodeling of ischemic heart through paracrine signaling. *Circ Res* 2006;98:1414–21.
- [119] Nakagami H, Maeda K, Morishita R, Iguchi S, Nishikawa T, Takami Y, et al. Novel autologous cell therapy in ischemic limb disease through growth factor secretion by cultured adipose tissue-derived stromal cells. *Arterioscler Thromb Vasc Biol* 2005;25:2542–7.
- [120] Fazel S, Cimini M, Chen L, Li S, Angoulvant D, Fedak P, et al. Cardioprotective c-kit+ cells are from the bone marrow and regulate the myocardial balance of angiogenic cytokines. *J Clin Invest* 2006;116:1865–77.
- [121] Moon MH, Kim SY, Kim YJ, Kim SJ, Lee JB, Bae YC, et al. Human adipose tissue-derived mesenchymal stem cells improve postnatal neovascularization in a mouse model of hindlimb ischemia. *Cell Physiol Biochem* 2006;17:279–90.
- [122] Majumdar MK, Thiede MA, Mosca JD, Moorman M, Gerson SL. Phenotypic and functional comparison of cultures of marrow-derived mesenchymal stem cells (MSCs) and stromal cells. *J Cell Physiol* 1998;176:57–66.
- [123] Weimar IS, Miranda N, Muller EJ, Hekman A, Kerst JM, de Gast GC, et al. Hepatocyte growth factor/scatter factor (HGF/SF) is produced by human bone marrow stromal cells and promotes proliferation, adhesion and survival of human hematopoietic progenitor cells (CD34+). *Exp Hematol* 1998;26:885–94.
- [124] Anderson CD, Heydarkhan-Hagvall S, Schenke-Layland K, Yang JQ, Jordan MC, Kim JK, et al. The role of cytoprotective cytokines in cardiac ischemia/reperfusion injury. *J Surg Res* 2008;148:164–71.
- [125] Cheng SL, Zhang SF, Mohan S, Lecanda F, Fausto A, Hunt AH, et al. Regulation of insulin-like growth factors I and II and their binding proteins in human bone marrow stromal cells by dexamethasone. *J Cell Biochem* 1998;71:449–58.
- [126] Markel TA, Wang Y, Herrmann JL, Crisostomo PR, Wang M, Novotny NM, et al. VEGF is critical for stem cell-mediated cardioprotection and a crucial paracrine factor for defining the age threshold in adult and neonatal stem cell function. *Am J Physiol Heart Circ Physiol* 2008;295:H2308–14.
- [127] Bobis-Wozowicz S, Miekus K, Wybieralska E, Jarocha D, Zawisz A, Madeja Z, et al. Genetically modified adipose tissue-derived mesenchymal stem cells overexpressing CXCR4 display increased motility, invasiveness, and homing to bone marrow of NOD/SCID mice. *Exp Hematol* 2011;39: 686–696.e4.
- [128] Cheng Z, Ou L, Zhou X, Li F, Jia X, Zhang Y, et al. Targeted migration of mesenchymal stem cells modified with cxcr4 gene to infarcted myocardium improves cardiac performance. *Mol Ther* 2008;16:571–9.
- [129] Haider HK, Jiang S, Idris NM, Ashraf M. IGF-1—overexpressing mesenchymal stem cells accelerate bone marrow stem cell mobilization via paracrine activation of SDF-1 α /CXCR4 signaling to promote myocardial repair. *Circ Res* 2008;103:1300–8.
- [130] Huang J, Zhang Z, Guo J, Ni A, Deb A, Zhang L, et al. Genetic modification of mesenchymal stem cells overexpressing CCR1 increases cell viability, migration, engraftment, and capillary density in the injured myocardium. *Circ Res* 2010;106:1753–62.
- [131] Wang WE, Yang D, Li L, Wang W, Peng Y, Chen C, et al. Prolyl hydroxylase domain protein 2 silencing enhances the survival and paracrine function of transplanted adipose-derived stem cells in infarcted myocardium. *Circ Res* 2013;113:288–300.
- [132] Wood SM, Wiesener MS, Yeates KM, Okada N, Pugh CW, Maxwell PH, et al. Selection and analysis of a mutant cell line defective in the hypoxia-inducible factor-1 α -subunit (HIF-1 α). Characterization of hif-1 α -dependent and -independent hypoxia-inducible gene expression. *J Biol Chem* 1998;273:8360–8.
- [133] Iyer NV, Kotch LE, Agani F, Leung SW, Laughner E, Wenger RH, et al. Cellular and developmental control of O₂ homeostasis by hypoxia-inducible factor 1 α . *Genes Dev* 1998;12:149–62.
- [134] Tazuke SI, Mazure NM, Sugawara J, Carland G, Faessen GH, Suen LF, et al. Hypoxia stimulates insulin-like growth factor binding protein 1 (IGFBP-1) gene expression in HepG2 cells: a possible model for IGFBP-1 expression in fetal hypoxia. *Proc Natl Acad Sci U S A* 1998;95:10188–93.
- [135] Feldser D, Agani F, Iyer NV, Pak B, Ferreira G, Semenza GL. Reciprocal positive regulation of hypoxia-inducible factor 1 α and insulin-like growth factor 2. *Cancer Res* 1999;59:3915–8.

- [136] Oh JS, Ha Y, An SS, Khan M, Pennant WA, Kim HJ, et al. Hypoxia-preconditioned adipose tissue-derived mesenchymal stem cell increase the survival and gene expression of engineered neural stem cells in a spinal cord injury model. *Neurosci Lett* 2010;472:215–9.
- [137] Carmeliet P, Dor Y, Herbert JM, Fukumura D, Brusselmans K, Dewerchin M, et al. Role of HIF-1 α in hypoxia-mediated apoptosis, cell proliferation and tumour angiogenesis. *Nature* 1998;394:485–90.
- [138] Marti HJ, Bernaudin M, Bellail A, Schoch H, Euler M, Petit E, et al. Hypoxia-induced vascular endothelial growth factor expression precedes neovascularization after cerebral ischemia. *Am J Pathol* 2000;156:965–76.
- [139] Gerber HP, Condorelli F, Park J, Ferrara N. Differential transcriptional regulation of the two vascular endothelial growth factor receptor genes. Flt-1, but not Flk-1/KDR, is up-regulated by hypoxia. *J Biol Chem* 1997;272:23659–67.
- [140] Ryan HE, Lo J, Johnson RS. HIF-1 α is required for solid tumor formation and embryonic vascularization. *EMBO J* 1998;17:3005–15.
- [141] Lee EY, Xia Y, Kim W-S, Kim MH, Kim TH, Kim KJ, et al. Hypoxia-enhanced wound-healing function of adipose-derived stem cells: increase in stem cell proliferation and up-regulation of VEGF and bFGF. *Wound Repair Regen* 2009;17:540–7.
- [142] Kubo M, Li T-S, Kurazumi H, Takemoto Y, Ohshima M, Murata T, et al. Hypoxic preconditioning enhances angiogenic potential of bone marrow cells with aging-related functional impairment. *Circ J* 2012;76:986–94.
- [143] Kubo M, Li TS, Suzuki R, Shirasawa B, Morikage N, Ohshima M, et al. Hypoxic preconditioning increases survival and angiogenic potency of peripheral blood mononuclear cells via oxidative stress resistance. *Am J Physiol Heart Circ Physiol* 2008;294:H590–5.
- [144] Chacko SM, Ahmed S, Selvendiran K, Kuppusamy ML, Khan M, Kuppusamy P. Hypoxic preconditioning induces the expression of pro-survival and pro-angiogenic markers in mesenchymal stem cells. *Am J Physiol Cell Physiol* 2010;299:C1562–70.
- [145] Stubbs SL, Hsiao ST-F, Peshavariya HM, Lim SY, Disting GJ, Dilley RJ. Hypoxic preconditioning enhances survival of human adipose-derived stem cells and conditions endothelial cells in vitro. *Stem Cells Dev* 2012;21:1887–96.
- [146] Rosová I, Dao M, Capoccia B, Link D, Nolte JA. Hypoxic preconditioning results in increased motility and improved therapeutic potential of human mesenchymal stem cells. *Stem Cells* 2008;26:2173–82.
- [147] Tang YL, Zhu W, Cheng M, Chen L, Zhang J, Sun T, et al. Hypoxic preconditioning enhances the benefit of cardiac progenitor cell therapy for treatment of myocardial infarction by inducing CXCR4 expression. *Circ Res* 2009;104:1209–16.
- [148] Hu X, Yu SP, Fraser JL, Lu Z, Ogle ME, Wang J-A, et al. Transplantation of hypoxia-preconditioned mesenchymal stem cells improves infarcted heart function via enhanced survival of implanted cells and angiogenesis. *J Thorac Cardiovasc Surg* 2008;135:799–808.
- [149] Strauer BE, Brehm M, Zeus T, Köstering M, Hernandez A, Sorg RV, et al. Repair of infarcted myocardium by autologous intracoronary mononuclear bone marrow cell transplantation in humans. *Circulation* 2002;106:1913–8.
- [150] Schächinger V, Erbs S, Elsässer A, Haberbosch W, Hambrecht R, Holschermann H, et al. Intracoronary bone marrow-derived progenitor cells in acute myocardial infarction. *N Engl J Med* 2006;355:1210–21.
- [151] Meyer GP, Wollert KC, Lotz J, Steffens J, Lippolt P, Fichtner S, et al. Intracoronary bone marrow cell transfer after myocardial infarction: eighteen months' follow-up data from the randomized, controlled BOOST (BOne marrow transfer to enhance ST-elevation infarct regeneration) trial. *Circulation* 2006;113:1287–94.
- [152] Assmus B, Schächinger V, Teupe C, Britten M, Lehmann R, Döbert N, et al. Transplantation of progenitor cells and regeneration enhancement in acute myocardial infarction (TOPCARE-AMI). *Circulation* 2002;106:3009–17.
- [153] Assmus B, Honold J, Schächinger V, Britten MB, Fischer-Rasokat U, Lehmann R, et al. Transcatheter transplantation of progenitor cells after myocardial infarction. *N Engl J Med* 2006;355:1222–32.
- [154] Janssens S, Dubois C, Bogaert J, Theunissen K, Deroose C, Desmet W, et al. Autologous bone marrow-derived stem-cell transfer in patients with ST-segment elevation myocardial infarction: double-blind, randomised controlled trial. *Lancet* 2006;367:113–21.
- [155] Roncalli J, Mouquet F, Piot C, Trochu JN, Le Corvoisier P, Neuder Y, et al. Intracoronary autologous mononucleated bone marrow cell infusion for acute myocardial infarction: results of the randomized multicenter BONAMI trial. *Eur Heart J* 2011;32:1748–57.
- [156] Lunde K, Solheim S, Aakhus S, Arnesen H, Abdelnoor M, Egeland T, et al. Intracoronary injection of mononuclear bone marrow cells in acute myocardial infarction. *N Engl J Med* 2006;355:1199–209.
- [157] Meyer GP, Wollert KC, Lotz J, Pirr J, Rager U, Lippolt P, et al. Intracoronary bone marrow cell transfer after myocardial infarction: 5-year follow-up from the randomized-controlled BOOST trial. *Eur Heart J* 2009;30:2978–84.
- [158] Leistner DM, Fischer-Rasokat U, Honold J, Seeger FH, Schächinger V, Lehmann R, et al. Transplantation of progenitor cells and regeneration enhancement in acute myocardial infarction (TOPCARE-AMI): final 5-year results suggest long-term safety and efficacy. *Clin Res Cardiol* 2011;100:925–34.
- [159] Houtgraaf JH, Dekker den WK, van Dalen BM, Springeling T, de Jong R, van Geuns RJ, et al. First experience in humans using adipose tissue-derived regenerative cells in the treatment of patients with ST-segment elevation myocardial infarction. *J Am Coll Cardiol* 2012;59:539–40.
- [160] Perin EC, Sanchez PL, Ruiz RS, Perez-Cano R, Lasso J, Alonso-Farto JC, et al. First in man transendocardial injection of autologous adipose-derived stem cells in patients with non revascularizable ischemic myocardium (PRECISE). *Circulation* 2010;112:A17966.
- [161] Qayyum AA, Haack-Sørensen M, Mathiasen AB, Jørgensen E, Eklund A, Kastrup J. Adipose-derived mesenchymal stromal cells for chronic myocardial ischemia (MyStromalCell Trial): study design. *Regen Med* 2012;7:421–8.
- [162] Laflamme MA, Murry CE. Heart regeneration. *Nature* 2011;473:326–35.
- [163] Wollert KC, Meyer GP, Lotz J, Ringes-Lichtenberg S, Lippolt P, Breidenbach C, et al. Intracoronary autologous bone-marrow cell transfer after myocardial infarction: the BOOST randomised controlled clinical trial. *Lancet* 2004;364:141–8.
- [164] Hou D, Youssef EA-S, Brinton TJ, Zhang P, Rogers P, Price ET, et al. Radio-labeled cell distribution after intramyocardial, intracoronary, and interstitial retrograde coronary venous delivery: implications for current clinical trials. *Circulation* 2005;112:1150–6.
- [165] Pijnappels DA, Gregoire S, Wu SM. The integrative aspects of cardiac physiology and their implications for cell-based therapy. *Ann N Y Acad Sci* 2010;1188:7–14.
- [166] van der Bogt KEA, Schrepfer S, Yu J, Sheikh AY, Hoyt G, Govaert JA, et al. Comparison of transplantation of adipose tissue- and bone marrow-derived mesenchymal stem cells in the infarcted heart. *Transplantation* 2009;87:642–52.
- [167] Tang YL, Tang Y, Zhang YC, Qian K, Shen L, Phillips MI. Improved graft mesenchymal stem cell survival in ischemic heart with a hypoxia-regulated heme oxygenase-1 vector. *J Am Coll Cardiol* 2005;46:1339–50.
- [168] Frangogiannis NG, Entman ML. Chemokines in myocardial ischemia. *Trends Cardiovasc Med* 2005;15:163–9.
- [169] Wang F, Guan J. Cellular cardiomyoplasty and cardiac tissue engineering for myocardial therapy. *Adv Drug Deliv Rev* 2010;62:784–97.
- [170] Reeve JLV, Duffy AM, O'Brien T, Samali A. Don't lose heart—therapeutic value of apoptosis prevention in the treatment of cardiovascular disease. *J Cell Mol Med* 2005;9:609–22.
- [171] Hu X, Wang J, Chen J, Luo R, He A, Xie X, et al. Optimal temporal delivery of bone marrow mesenchymal stem cells in rats with myocardial infarction. *Eur J Cardiothorac Surg* 2007;31:438–43.
- [172] Van Linthout S, Savvatis K, Miteva K, Peng J, Ringe J, Warstat K, et al. Mesenchymal stem cells improve murine acute coxsackievirus B3-induced myocarditis. *Eur Heart J* 2011;32:2168–78.
- [173] Landa N, Miller L, Feinberg MS, Holbova R, Shachar M, Freeman I, et al. Effect of injectable alginate implant on cardiac remodeling and function after recent and old infarcts in rat. *Circulation* 2008;117:1388–96.
- [174] Leor J, Tuvia S, Guetta V, Manczur F, Castel D, Willenz U, et al. Intracoronary injection of in situ forming alginate hydrogel reverses left ventricular remodeling after myocardial infarction in swine. *J Am Coll Cardiol* 2009;54:1014–23.
- [175] Tsur-Gang O, Ruvinov E, Landa N, Holbova R, Feinberg MS, Leor J, et al. The effects of peptide-based modification of alginate on left ventricular remodeling and function after myocardial infarction. *Biomaterials* 2009;30:189–95.
- [176] Yu J, Du KT, Fang Q, Gu Y, Mihardja SS, Sievers RE, et al. The use of human mesenchymal stem cells encapsulated in RGD modified alginate microspheres in the repair of myocardial infarction in the rat. *Biomaterials* 2010;31:7012–20.
- [177] Lu W-N, Lü S-H, Wang H-B, Li D-X, Duan C-M, Liu Z-Q, et al. Functional improvement of infarcted heart by co-injection of embryonic stem cells with temperature-responsive chitosan hydrogel. *Tissue Eng Part A* 2009;15:1437–47.
- [178] Liu Z, Wang H, Wang Y, Lin Q, Yao A, Cao F, et al. The influence of chitosan hydrogel on stem cell engraftment, survival and homing in the ischemic myocardial microenvironment. *Biomaterials* 2012;33:3093–106.
- [179] Dai W, Wold LE, Dow JS, Kloner RA. Thickening of the infarcted wall by collagen injection improves left ventricular function in rats. *J Am Coll Cardiol* 2005;46:714–9.
- [180] Dai W, Hale SL, Kay GL, Jyrala AJ, Kloner RA. Delivering stem cells to the heart in a collagen matrix reduces relocation of cells to other organs as assessed by nanoparticle technology. *Regen Med* 2009;4:387–95.
- [181] Christman KL, Fok HH, Sievers RE, Fang Q, Lee RJ. Fibrin glue alone and skeletal myoblasts in a fibrin scaffold preserve cardiac function after myocardial infarction. *Tissue Eng* 2004;10:403–9.
- [182] Zhang G, Hu Q, Braunlin EA, Suggs LJ, Zhang J. Enhancing efficacy of stem cell transplantation to the heart with a PEGylated fibrin biomatrix. *Tissue Eng Part A* 2008;14:1025–36.
- [183] Zhang X, Wang H, Ma X, Adila A, Wang B, Liu F, et al. Preservation of the cardiac function in infarcted rat hearts by the transplantation of adipose-derived stem cells with injectable fibrin scaffolds. *Exp Biol Med* 2010;235:1505–15.
- [184] Ifkovits JL, Tous E, Minakawa M, Morita M, Robb JD, Koomalsingh KJ, et al. Injectable hydrogel properties influence infarct expansion and extent of postinfarction left ventricular remodeling in an ovine model. *Proc Natl Acad Sci U S A* 2010;107:11507–12.
- [185] Tous E, Ifkovits JL, Koomalsingh KJ, Shuto T, Soeda T, Kondo N, et al. Influence of injectable hyaluronic acid hydrogel degradation behavior on infarction-induced ventricular remodeling. *Biomacromolecules* 2011;12:4127–35.
- [186] Kofidis T, Lebl DR, Martinez EC, Hoyt G, Tanaka M, Robbins RC. Novel injectable bioartificial tissue facilitates targeted, less invasive, large-scale tissue restoration on the beating heart after myocardial injury. *Circulation* 2005;112:1173–7.

- [187] Zhang P, Zhang H, Wang H, Wei Y, Hu S. Artificial matrix helps neonatal cardiomyocytes restore injured myocardium in rats. *Artif Organs* 2006;30:86–93.
- [188] Laflamme MA, Chen KY, Naumova AV, Muskheli V, Fugate JA, Dupras SK, et al. Cardiomyocytes derived from human embryonic stem cells in pro-survival factors enhance function of infarcted rat hearts. *Nat Biotechnol* 2007;25:1015–24.
- [189] Padin-Iruegas ME, Misao Y, Davis ME, Segers VFM, Esposito G, Tokunou T, et al. Cardiac progenitor cells and biotinylated insulin-like growth factor-1 nanofibers improve endogenous and exogenous myocardial regeneration after infarction. *Circulation* 2009;120:876–87.
- [190] Lin YD, Yeh ML, Yang YJ, Tsai DC, Chu TY, Shih YY, et al. Intramyocardial peptide nanofiber injection improves postinfarction ventricular remodeling and efficacy of bone marrow cell therapy in pigs. *Circulation* 2010;122: S132–41.
- [191] Wang T, Jiang X-J, Tang Q-Z, Li X-Y, Lin T, Wu D-Q, et al. Bone marrow stem cells implantation with a-cyclodextrin/MPEG–PCL–MPEG hydrogel improves cardiac function after myocardial infarction. *Acta Biomater* 2009;5:2939–44.
- [192] Kraehenbuehl TP, Ferreira LS, Hayward AM, Nahrendorf M, Van der Vlies AJ, Vasile E, et al. Human embryonic stem cell-derived microvascular grafts for cardiac tissue preservation after myocardial infarction. *Biomaterials* 2011;32:1102–9.
- [193] Wang H, Liu Z, Li D, Guo X, Kasper FK, Duan C, et al. Injectable biodegradable hydrogels for embryonic stem cell transplantation: improved cardiac remodelling and function of myocardial infarction. *J Cell Mol Med* 2012;16: 1310–20.
- [194] Wu J, Zeng F, Huang X-P, Chung JCY, Konecny F, Weisel RD, et al. Infarct stabilization and cardiac repair with a VEGF-conjugated, injectable hydrogel. *Biomaterials* 2011;32:579–86.
- [195] Wang T, Wu DQ, Jiang XJ, Zhang XZ, Li XY, Zhang JF, et al. Novel thermo-sensitive hydrogel injection inhibits post-infarct ventricle remodelling. *Eur J Heart Fail* 2009;11:14–9.
- [196] Fujimoto KL, Ma Z, Nelson DM, Hashizume R, Guan J, Tobita K, et al. Synthesis, characterization and therapeutic efficacy of a biodegradable, thermoresponsive hydrogel designed for application in chronic infarcted myocardium. *Biomaterials* 2009;30:4357–68.
- [197] Wall ST, Yeh C-C, Tu RYK, Mann MJ, Healy KE. Biomimetic matrices for myocardial stabilization and stem cell transplantation. *J Biomed Mater Res A* 2010;95:1055–66.
- [198] Chachques JC, Azarine A, Mousseaux E, Serafi El M, Cortes-Morichetti M, Carpenter A. MRI evaluation of local myocardial treatments: epicardial versus endocardial (cell-fix catheter) injections. *J Interv Cardiol* 2007;20:188–96.
- [199] Dib N, Campbell A, Jacoby DB, Zawadzka A, Ratliff J, Miedzybrocki BM, et al. Safety and feasibility of percutaneous autologous skeletal myoblast transplantation in the coil-infarcted swine myocardium. *J Pharmacol Toxicol Methods* 2006;54:71–7.
- [200] Wall ST, Walker JC, Healy KE, Ratcliffe MB, Guccione JM. Theoretical impact of the injection of material into the myocardium: a finite element model simulation. *Circulation* 2006;114:2627–35.
- [201] Jackson BM, Gorman JH, Salgo IS, Moainie SL, Plappert T, St John-Sutton M, et al. Border zone geometry increases wall stress after myocardial infarction: contrast echocardiographic assessment. *Am J Physiol Heart Circ Physiol* 2003;284:H475–9.
- [202] Cornwell KG, Lei P, Andreadis ST, Pins GD. Crosslinking of discrete self-assembled collagen threads: effects on mechanical strength and cell–matrix interactions. *J Biomed Mater Res* 2006;80A:362–71.
- [203] Bhana B, Iyer RK, Chen WLK, Zhao R, Sider KL, Likhitpanichkul M, et al. Influence of substrate stiffness on the phenotype of heart cells. *Biotechnol Bioeng* 2010;105:1148–60.
- [204] Engler AJ, Sen S, Sweeney HL, Discher DE. Matrix elasticity directs stem cell lineage specification. *Cell* 2006;126:677–89.
- [205] Sharifi S, Blanquer SBC, van Kooten TG, Grijpma DW. Biodegradable nanocomposite hydrogel structures with enhanced mechanical properties prepared by photo-crosslinking solutions of poly(trimethylene carbonate)-poly(ethylene glycol)-poly(trimethylene carbonate) macromonomers and nanoclay particles. *Acta Biomater* 2012;8:4233–43.
- [206] Murphy CM, Haugh MG, O'Brien FJ. The effect of mean pore size on cell attachment, proliferation and migration in collagen-glycosaminoglycan scaffolds for bone tissue engineering. *Biomaterials* 2010;31:461–6.
- [207] Al-Shamkhani A, Duncan R. Radioiodination of alginate via covalently-bound tyrosinamide allows monitoring of its fate in vivo. *J Bioact Compat Polym* 1995;10:4–13.
- [208] Kong H-J, Alsberg E, Kaigler D, Lee K-Y, Mooney DJ. Controlling degradation of hydrogels via the size of crosslinked junctions. *Adv Mater* 2004;16:1917–21.
- [209] Bouhadir KH, Lee KY, Alsberg E, Damm KL, Anderson KW, Mooney DJ. Degradation of partially oxidized alginate and its potential application for tissue engineering. *Biotechnol Prog* 2001;17:945–50.
- [210] Yu J, Gu Y, Du KT, Mihaardja S, Sievers RE, Lee RJ. The effect of injected RGD modified alginate on angiogenesis and left ventricular function in a chronic rat infarct model. *Biomaterials* 2009;30:751–6.
- [211] Yu J, Christman KL, Chin E, Sievers RE, Saeed M, Lee RJ. Restoration of left ventricular geometry and improvement of left ventricular function in a rodent model of chronic ischemic cardiomyopathy. *J Thorac Cardiovasc Surg* 2009;137:180–7.
- [212] Kim I-Y, Seo S-J, Moon H-S, Yoo M-K, Park I-Y, Kim B-C, et al. Chitosan and its derivatives for tissue engineering applications. *Biotechnol Adv* 2008;26:1–21.
- [213] Hirano S, Tsuchida H, Nagao N. N-acetylation in chitosan and the rate of its enzymic hydrolysis. *Biomaterials* 1989;10:574–6.
- [214] Ishihara M, Nakanishi K, Ono K, Sato M, Kikuchi M, Saito Y, et al. Photocrosslinkable chitosan as a dressing for wound occlusion and accelerator in healing process. *Biomaterials* 2002;23:833–40.
- [215] Chenite A, Chaput C, Wang D, Combes C, Buschmann MD, Hoemann CD, et al. Novel injectable neutral solutions of chitosan form biodegradable gels in situ. *Biomaterials* 2000;21:2155–61.
- [216] Rosenblatt J, Devereux B, Wallace DG. Injectable collagen as a pH-sensitive hydrogel. *Biomaterials* 1994;15:985–95.
- [217] Williams BR, Gelman RA, Poppke DC, Piez KA. Collagen fibril formation. *J Biol Chem* 1978;253:6578–85.
- [218] Achilli M, Mantovani D. Tailoring mechanical properties of collagen-based scaffolds for vascular tissue engineering: the effects of pH, temperature and ionic strength on gelation. *Polymers* 2010;2:664–80.
- [219] Breen A, O'Brien T, Pandit A. Fibrin as a delivery system for therapeutic drugs and biomolecules. *Tissue Eng Part B Rev* 2009;15:201–14.
- [220] Kubota K, Kogure H, Masuda Y, Toyama Y, Kita R, Takahashi A, et al. Gelation dynamics and gel structure of fibrinogen. *Colloids Surf B Biointerfaces* 2004;38:103–9.
- [221] Rowe SL, Lee S, Stegemann JP. Influence of thrombin concentration on the mechanical and morphological properties of cell-seeded fibrin hydrogels. *Acta Biomater* 2007;3:59–67.
- [222] Bootle-Wilbraham CA, Tazzyman S, Thompson WD, Stirk CM, Lewis CE. Fibrin fragment E stimulates the proliferation, migration and differentiation of human microvascular endothelial cells in vitro. *Angiogenesis* 2001;4:269–75.
- [223] Guo H-D, Wang H-J, Tan Y-Z, Wu J-H. Transplantation of marrow-derived cardiac stem cells carried in fibrin improves cardiac function after myocardial infarction. *Tissue Eng Part A* 2010;17:45–58.
- [224] Christman KL, Vardanian AJ, Fang Q, Sievers RE, Fok HH, Lee RJ. Injectable fibrin scaffold improves cell transplant survival, reduces infarct expansion, and induces neovasculation formation in ischemic myocardium. *J Am Coll Cardiol* 2004;44:654–60.
- [225] Burdick JA, Prestwich GD. Hyaluronic acid hydrogels for biomedical applications. *Adv Mater* 2011;23:H41–56.
- [226] Stern R, Asari AA, Sugahara KN. Hyaluronan fragments: an information-rich system. *Eur J Cell Biol* 2006;85:699–715.
- [227] Kim J, Kim IS, Cho TH, Lee KB, Hwang SJ, Tae G, et al. Bone regeneration using hyaluronic acid-based hydrogel with bone morphogenic protein-2 and human mesenchymal stem cells. *Biomaterials* 2007;28:1830–7.
- [228] Yoon SJ, Fang YH, Lim CH, Kim BS, Son HS, Park Y, et al. Regeneration of ischemic heart using hyaluronic acid-based injectable hydrogel. *J Biomed Mater Res* 2009;91B:163–71.
- [229] van Dijk-Wolthuis W, Hooijboom J, Van Steenberghe MJ, Tsang S, Hennink WE. Degradation and release behavior of dextran-based hydrogels. *Macromolecules* 1997;30:4639–45.
- [230] Laflamme MA, Gold J, Xu C, Hassanipour M, Rosler E, Police S, et al. Formation of human myocardium in the rat heart from human embryonic stem cells. *Am J Pathol* 2005;167:663–71.
- [231] Hughes CS, Postovit LM, Lajoie GA. Matrigel: a complex protein mixture required for optimal growth of cell culture. *Proteomics* 2010;10:1886–90.
- [232] Zhang S, Holmes TC, DiPersio CM, Hynes RO, Su X, Rich A. Self-complementary oligopeptide matrices support mammalian cell attachment. *Biomaterials* 1995;16:1385–93.
- [233] Hsieh PCH, Davis ME, Gannon J, MacGillivray C, Lee RT. Controlled delivery of PDGF-BB for myocardial protection using injectable self-assembling peptide nanofibers. *J Clin Invest* 2006;116:237–48.
- [234] Davis ME, Hsieh PC, Takahashi T, Song Q, Zhang S, Kamm RD, et al. Local myocardial insulin-like growth factor 1 (IGF-1) delivery with biotinylated peptide nanofibers improves cell therapy for myocardial infarction. *Proc Natl Acad Sci U S A* 2006;103:8155–60.
- [235] Li Z, Guan J. Hydrogels for cardiac tissue engineering. *Polymers* 2011;3:740–61.
- [236] Wu D-Q, Wang T, Lu B, Xu X-D, Cheng S-X, Jiang X-J, et al. Fabrication of supramolecular hydrogels for drug delivery and stem cell encapsulation. *Langmuir* 2008;24:10306–12.
- [237] Jiang X-J, Wang T, Li X-Y, Wu D-Q, Zheng Z-B, Zhang J-F, et al. Injection of a novel synthetic hydrogel preserves left ventricle function after myocardial infarction. *J Biomed Mater Res A* 2009;90:472–7.
- [238] Kraehenbuehl TP, Zammaretti P, Van der Vlies AJ, Schoenmakers RG, Lutolf MP, Jacini ME, et al. Three-dimensional extracellular matrix-directed cardioprogenitor differentiation: systematic modulation of a synthetic cell-responsive PEG-hydrogel. *Biomaterials* 2008;29:2757–66.
- [239] Lee BH, Vernon B. In situ-gelling, erodible N-isopropylacrylamide copolymers. *Macromol Biosci* 2005;5:629–35.
- [240] Li X-Y, Wang T, Jiang X-J, Lin T, Wu D-Q, Zhang X-Z, et al. Injectable hydrogel helps bone marrow-derived mononuclear cells restore infarcted myocardium. *Cardiology* 2010;115:194–9.
- [241] Iacobellis G. Epicardial and pericardial fat: close, but very different. *Obes (Silver Spring)* 2009;17:625. authorreply626–627.
- [242] Dinsmore JH, Dib N. Stem cell therapy for the treatment of acute myocardial infarction. *Cardiol Clin* 2010;28:127–38.