See discussions, stats, and author profiles for this publication at: https://www.researchgate.net/publication/257659649

Reply to the "Comment on the paper 'The role of HBF4 in electro-catalysis: Arsenic contamination and anion adsorption' by A.L. Santos, R. Nagao, C.P. Oliveira, R.B. de Lima, H. Var...

ARTICLE in JOURNAL OF ELECTROANALYTICAL CHEMISTRY · NOVEMBER 2012

Impact Factor: 2.73 · DOI: 10.1016/j.jelechem.2012.09.031

READS

69

5 AUTHORS, INCLUDING:



Raphael Nagao

Saint Louis University

23 PUBLICATIONS 159 CITATIONS

SEE PROFILE



R.B. de Lima

Universidade Federal do Maranhão

14 PUBLICATIONS 463 CITATIONS

SEE PROFILE



Hamilton Varela

University of São Paulo

105 PUBLICATIONS 1,317 CITATIONS

SEE PROFILE

FISEVIER

Contents lists available at SciVerse ScienceDirect

Journal of Electroanalytical Chemistry

journal homepage: www.elsevier.com/locate/jelechem



Short Communication

Reply to the "Comment on the paper 'The role of HBF4 in electro-catalysis: Arsenic contamination and anion adsorption' by A.L. Santos, R. Nagao, C.P. Oliveira, R.B. de Lima, H. Varela [J. Electroanal. Chem. 660 (2011) 147–152]"

A.L. Santos^a, R. Nagao^a, C.P. Oliveira^a, R.B. de Lima^b, H. Varela^{a,*}

ARTICLE INFO

Article history: Available online 13 October 2012

Keywords: Electro-catalysis Electrochemical quartz crystal nanobalance Formic acid Tetrafluoroboric acid Anion adsorption

ABSTRACT

This is a Reply to recent comment by M. Schell to our previously published article: Reply to the "Comment on the paper 'The role of HBF4 in electro-catalysis: arsenic contamination and anion adsorption' by A.L. Santos, R. Nagao, C.P. Oliveira, R.B. de Lima, H. Varela [J. Electroanal. Chem. 660 (2011) 147–152]" © 2012 Elsevier B.V. All rights reserved.

We have recently reported an experimental investigation on the influence of different amounts of HBF4 on the Pt|HClO4(aq) and Pt|HClO₄(aq), HCOOH systems.[1] Following the previous work by Feliu and co-workers [2] on the presence of arsenic contamination in as-received HBF₄, we used this chemical from two different suppliers and characterized it in terms of the amount of arsenic con-Hvdride Generation Atomic Absorption tamination by Spectroscopy (HGAAS). The systems were studied by means of in situ nanogravimetry with the electrochemical quartz crystal nanobalance (EQCN), and in spite of the similar voltammetric profiles, the presence of arsenic was reflected in dramatic variations in the mass change vs. potential profiles, due to the deposition/dissolution of arsenic prior to the surface oxidation. Finally, the effect of As-contaminated HBF₄ on the electro-oxidation of formic acid was rationalized in terms of: (a) the enhancement of the electro-oxidation current at low potentials, and (b) the inhibition at high potentials. The former effect was unambiguously attributed to the arsenic-assisted electro-oxidation of formic acid, whereas the current decrease at high potential was suggested to result from anion adsorption.

In his comment [3] on our paper [1], Schell argues that we have misinterpreted some of his arguments and thus wrongly referred to his ideas. In this reply we address these criticisms.

The discussion given in Santos et al. [1] is based on our interpretation of the previous reports by Schell and co-workers. Of course,

* Corresponding author. Tel.: +55 16 33738059. E-mail address: varela@iqsc.usp.br (H. Varela). if our interpretation does not strictly reflect the authors' ideas, this can cause misinterpretation of their papers. In some sense, it seems we extrapolated some of Schell's conjectures and conclusions to a broader context. Specifically, concerning the particular role *we attributed* to fluoroboric acid, it was deduced from the observation that the catalytic enhancement caused by fluoroboric acid was actually due to the presence of arsenic as an impurity.

Schell obtained similar behavior with fluoroborate in the oxidation of formic acid and the evidence is strong that it was caused for the same reason, arsenic poisoning [4,5]. If one was unaware of the presence of arsenic, it would be reasonable to believe that fluoroborate induced a catalytic mechanism. Schell has developed a theory for a catalytic mechanism induced by anions [6,7]. Besides small concentrations, we achieved effects with large concentrations of fluoroboric acid. These results and the interpretation of Schell and co-workers previous work led us to attribute to them the claim that fluoroborate anion primarily enhanced catalytic activity.

In spite of any misinterpretation, we emphasize that the main experimental findings presented in our paper [1] remain unchanged.

References

- A.L. Santos, R. Nagao, C.P. Oliveira, Roberto B. de Lima, H. Varela, J. Electroanal. Chem. 660 (2011) 147–152.
- [2] C.A. Angelucci, J. Souza-Garcia, E. Herrero, J.M. Feliu, J. Electroanal. Chem. 646 (2010) 100–106.

^a Institute of Chemistry of São Carlos, University of São Paulo, CP 780, CEP: 13560-970 São Carlos, SP, Brazil

^b Department of Chemistry, Federal University of Maranhão, Campus do Bacanga, CEP: 65085-580 São Luis, MA, Brazil

- [3] M. Schell, "Comment on the paper 'The role of HBF4 in electro-catalysis: arsenic contamination and anion adsorption' by A.L. Santos, R. Nagao, C.P. Oliveira, R.B. de Lima, H. Varela [J. Electroanal. Chem. 660 (2011) 147-152]", J. Electroanal. Chem. (2012).
- [4] S. Chen, D. Lee, M. Schell, Electrochim. Acta 46 (2001) 3481–3492.
 [5] S. Chen, D. Lee, M. Schell, Electrochem. Commun. 3 (2001) 81–85.
 [6] M. Schell, B.E.K. Swamy, J. Electroanal. Chem. 584 (2005) 157–166.

- [7] D.M. Zeng, M. Schell, Electrochim. Acta 55 (2011) 4703–4710.