



ELSEVIER

Contents lists available at ScienceDirect

## American Journal of Infection Control

journal homepage: [www.ajicjournal.org](http://www.ajicjournal.org)

AJIC  
American Journal of  
Infection Control

## Letter to the Editor

Absence of *Clostridium difficile* in asymptomatic hospital staff

## To the Editor:

*Clostridium difficile* is considered a leading cause of hospital-acquired diarrhea. To evaluate the prevalence of asymptomatic *C difficile* stool carriage among hospital staff, we tested stool samples from 112 volunteers for toxigenic *C difficile*.

Usually, there is no significant risk for healthy adults in acquiring a *C difficile* infection (CDI) because their mature colonic bacterial flora will resist colonization with *C difficile*.<sup>1</sup> CDI in humans is generally linked to procedures that alter the normal enteric flora (antibiotic treatment or antineoplastic chemotherapy). Several outbreaks via a contaminated hospital environment or the contaminated hands of hospital personnel have shown the risk of nosocomial acquisition of CDI.<sup>2</sup> We chose to study the prevalence of *C difficile* stool carriage in asymptomatic hospital staff in Austria to assess the risk of healthy hospital staff acquiring *C difficile*.

Probandes were restricted to persons without a history of diarrhea or antibiotic usage for 3 months among hospital staff prior to testing (Table 1). Stool samples (n = 112) were gained from April to July 2010 at a 1,200 bed tertiary care university hospital. The male to female ratio was 1:5.3 (Table 2). The study was approved by the local ethics committee. Microbiological diagnostics were performed by direct plating onto *C difficile* selective agar (cycloserine/cefoxitin agar; bioMérieux, Marcy l'Etoile, France) plus broth enrichment as described elsewhere.<sup>3</sup>

One hundred twelve healthy and asymptomatic staff members from 9 departments and the medical school were screened for *C difficile*. All stool samples tested negative by primary inoculation of the selective plate. Enough material ( $\geq 1$  g) to perform testing by broth enrichment technique was available for 45 of the 112 samples (40.2%).

Broth enrichment technique yielded negative results for all stool samples tested. Also, 67 stool samples provided by administrative staff of a major Austrian groceries chain proved negative when tested (parallel to our study) using both methods (data not shown).

In medical literature, it is widely supposed that *C difficile* stool carriage is frequently found in healthy adults, including hospital staff. Van Nood et al reported 13% of asymptomatic hospital staff (75% of these being physicians) in The Netherlands to show *C difficile* stool carriage.<sup>4</sup> Kato et al demonstrated that 4.3% of hospital personnel in Japan carried *C difficile*.<sup>5</sup> Our finding of absence of fecal carriage in Austrian hospital staff was surprising but in accordance with results reported by Carmeli et al, who showed absence of intestinal carriage of *C difficile* among 55 hospital staff in Israel.<sup>6</sup> In addition, our study has the limitation of a relatively low sample

Table 1

Study group by affiliation

	Absolute number	% Of total
Gastroenterology	9	8.0
Hematology and oncology	19	16.9
Pediatrics	21	18.8
General surgery	19	16.9
Pediatric surgery	10	8.9
Vascular surgery	11	9.8
Medical school students	12	10.7
Anaesthesiology	7	6.3
Intensive care unit	1	0.9
Physical therapy	3	2.7
Total	112	100

Table 2

Study group by occupation

Study group	Age, yr		Sex	
	Absolute number	% Of total	Median	Female
Clerical staff	2	1.8	35	100%
Janitorial staff	3	2.7	53	100%
Physiotherapists	3	2.7	32	100%
Nurses	77	68.8	39	94%
Medical school Students	11	9.8	24	82%
Physicians	16	14.3	47	44%
Total	112	100	39	86%

size, but it guarantees an upper confidence bound of 2.6% for zero findings (95% confidence level).

Because specimens from symptomatic patients—tested for *C difficile* at the same time by the same method (direct plating)—yielded toxigenic *C difficile*, we abnegate the occurrence of false-negative results.<sup>7</sup> Similar discrepant findings have previously been reported for occurrence of *C difficile* in food (ground meat): whereas Rodriguez-Palacios et al in Canada found 20% of tested samples positive for *C difficile*,<sup>8</sup> Jöbstl et al reported a contamination rate of only 3% *C difficile* in ground meat in Austria.<sup>9</sup> This different contamination rate of ground meat might be just one of many factors explaining the discrepant results concerning *C difficile* carriage in healthy adults from different countries. In Austria, *C difficile* is the most frequently diagnosed bacterial pathogen in patients with community-acquired gastroenteritis, accounting for 18.7% of positive results, second only to norovirus (36.0%).<sup>10</sup> Our finding of absence of fecal carriage in healthy hospital staff and in healthy administrative staff of a food trade company underlines the diagnostic relevance of a positive stool result in patients with acute community acquired gastroenteritis. It also indicates that there is only low risk for healthy hospital staff for acquiring *C difficile*. Our findings also emphasize the diagnostic relevance of *C difficile* detection in symptomatic health care workers with diarrhea.

Conflicts of interest: None to report.

## Acknowledgment

We would like to thank Professor Christian Datz for providing us with stool samples of the Non-hospital-staff and also Dr. David Schein for his support in proofreading.

## References

1. Reeves AE, Theriot CM, Bergin IL, Huffnagle GB, Schloss PD, Young VB. The interplay between microbiome dynamics and pathogen dynamics in a murine model of *Clostridium difficile* infection. *Gut Microbes* 2011;2:145-58.
2. McFarland LV, Mulligan ME, Kwok RY, Stamm WE. Nosocomial Acquisition of *Clostridium difficile* infection. *N Engl J Med* 1989;320:204-10.
3. Mac Faddin JD. Media for isolation-cultivation-identification-maintenance of medical bacteria. Vol. 1. Baltimore [MD]: Williams & Wilkins; 1985. p. 240-246.
4. van Nood E, van Dijk K, Hegeman Z, Speelman P, Visser CE. Asymptomatic carriage of *Clostridium difficile* among HCWs: do we disregard the doctor? *Infect Control Hosp Epidemiol* 2009;30:924-5.
5. Kato H, Kita H, Karasawa T, Maegawa T, Koino Y, Takakuwa H, et al. Colonisation and transmission of *Clostridium difficile* in healthy individuals examined by PCR ribotyping and pulsed-field gel electrophoresis. *J Med Microbiol* 2001; 50:720-7.
6. Carmeli Y, Venkataraman L, DeGirolami PC, Lichtenberg DA, Karchmer AW, Samore MH. Stool colonization of healthcare workers with selected resistant bacteria. *Infect Control Hosp Epidemiol* 1998;19:38-40.
7. Hell M, Permoser M, Chmelizek G, Kern JM, Maass M, Huhulescu S, et al. *Clostridium difficile* infection: monoclonal or polyclonal genesis? *Infection* 2011;39:461-5.
8. Rodriguez-Palacios A, Staempfli HR, Duffield T, Weese JS. *Clostridium difficile* in retail ground meat, Canada. *Emerg Infect Dis* 2007;13:485-7.
9. Jöbstl M, Heuberger S, Indra A, Nepf R, Köfer J, Wagner M. *Clostridium difficile* in raw products of animal origin. *Int J Food Microbiol* 2010;138:172-5.
10. Huhulescu S, Kiss R, Brettlecker M, Cerny RJ, Hess C, Wewalka G, et al. Etiology of acute gastroenteritis in three sentinel general practices, Austria 2007. *Infection* 2009;37:103-8.

Markus Hell, MD\*

Kathrin Sickau, MD

Department of Hospital Epidemiology and Infection Control,  
University Hospital Salzburg, Austria

Gregor Chmelizek, MD

Department of Hospital Epidemiology and Infection Control,  
University Hospital Salzburg, Austria

Institute of Microbiology, Hygiene and Infectious Diseases, University  
Hospital Salzburg, Salzburg, Austria

Jan Marco Kern, MD

Matthias Maass, MD, Prof.

Institute of Microbiology, Hygiene and Infectious Diseases, University  
Hospital Salzburg, Salzburg, Austria

Steliana Huhulescu, MD

National Reference Laboratory for *Clostridium difficile*, Austrian  
Agency for Health and Food Safety (AGES), Vienna, Austria

Franz Allerberger, MD, Prof.

Institute of Microbiology, Hygiene and Infectious Diseases, University  
Hospital Salzburg, Salzburg, Austria

National Reference Laboratory for *Clostridium difficile*, Austrian

Agency for Health and Food Safety (AGES), Vienna, Austria

\* Address correspondence to Markus Hell, MD, University Hospital  
Salzburg, Department of Hospital Epidemiology and Infection  
Control, Muellner Hauptstr.48, 5020 Salzburg, Austria.  
E-mail address: [m.hell@salk.at](mailto:m.hell@salk.at) (M. Hell)