

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/257738909>

# Corrigendum to “Theoretical study of the electronic (hyper)polarizabilities of amino acids in gaseous and aqueous phases” [Comput. Theor. Chem. 976 (2011) 188–190]

**ARTICLE** *in* COMPUTATIONAL AND THEORETICAL CHEMISTRY · JANUARY 2012

Impact Factor: 1.55 · DOI: 10.1016/j.comptc.2011.11.033

---

CITATIONS

6

---

READS

23

## 1 AUTHOR:



[Andrea Alparone](#)

University of Catania

67 PUBLICATIONS 791 CITATIONS

SEE PROFILE



## Corrigendum


## Corrigendum to "Theoretical study of the electronic (hyper)polarizabilities of amino acids in gaseous and aqueous phases" [Comput. Theor. Chem. 976 (2011) 188–190]

Andrea Alparone\*

Department of Chemistry, University of Catania, Viale A. Doria 6, Catania 95125, Italy

The author regrets that the range of  $\langle\alpha\rangle$  values in Fig. 4 was incorrectly presented in the published article. Fig. 4 is correctly

reproduced below (with the range for the  $\langle\alpha\rangle$  values at 40–130 a.u. instead of 0–100 a.u.).



**Fig. 4.** Relationship between  $\langle\gamma\rangle$  and  $\langle\alpha\rangle$  values of the aliphatic amino acids computed at the MP2/aug-cc-pVDZ level. Gas phase  $\langle\alpha\rangle$  values are taken from Ref. [1]. Gas phase:  $\langle\gamma\rangle = -0.186 + 0.020 \cdot \langle\alpha\rangle$ ,  $r = 0.99$ . Water solution:  $\langle\gamma\rangle = -0.594 + 0.034 \cdot \langle\alpha\rangle$ ,  $r = 0.99$ .

## Reference

- [1] S. Millefiori, A. Alparone, A. Millefiori, A. Vanella, Electronic and vibrational polarizabilities of the twenty naturally occurring amino acids, *Biophys. Chem.* 132 (2008) 139–147 (and references therein).