



NEW DRUGS

DT-diaphorase: a target for new anticancer drugs

S. Danson^{a,b,*}, T.H. Ward^a, J. Butler^c, M. Ranson^{a,b}^a Paterson Institute for Cancer Research, Manchester, UK^b Department of Medical Oncology, Manchester, UK^c Salford University, Manchester, UK

KEYWORDS

DT-diaphorase;
Bioreductive;
Diaziridinylquinones;
RH1;
Resistance;
Induction

Summary DT-diaphorase (DTD) is an obligate two-electron reductase which bioactivates chemotherapeutic quinones. DTD levels are elevated in a number of tumour types, including non-small cell lung carcinoma, colorectal carcinoma, liver cancers and breast carcinomas, when compared to the surrounding normal tissue. The differential in DTD between tumour and normal tissue should allow targeted activation of chemotherapeutic quinones in the tumour whilst minimising normal tissue toxicity.

The prototypical bioreductive drug is Mitomycin C (MMC) which is widely used in clinical practice. However, MMC is actually a relatively poor substrate for DTD and its metabolism is pH-dependent. Other bioreductive drugs have failed because of poor solubility and inability to surpass other agents in use. RH1, a novel diaziridinylbenzoquinone, is a more efficient substrate for DTD. It has been demonstrated to have anti-tumour effects both in vitro and in vivo and demonstrates a relationship between DTD expression levels and drug response. RH1 has recently entered a phase I clinical trial in solid tumours under the auspices of Cancer Research UK.

Recent work has demonstrated that DTD is present in the nucleus and is associated with both p53 and the heat shock protein, HSP-70. Furthermore, DTD is inducible by several non-toxic compounds and therefore much interest has focussed on increasing the differential in DTD levels between tumour and normal tissues.

© 2004 Elsevier Ltd. All rights reserved.

Introduction

The enzyme DTD was first described in 1958 by Ernster and Navazio.¹ DTD was isolated from soluble rat liver homogenates and was shown to catalyse the oxidation of the co-factors nicotinamide adenine dinucleotide (NADH) and nicotinamide adenine dinucleotide phosphate (NADPH) equally

well. Subsequently the structure, mechanism of action and interactions of DTD have been investigated. During this time DTD has been referred to by a variety of names including NAD(P)H:quinone oxidoreductase (NQO) (EC 1.6.99.2), vitamin K reductase, phyloquinone reductase, menadione reductase, azo dye reductase, X-ray inducible transcript 3 (Xip3) and nicotinamide menadione oxidoreductase but, in keeping with current literature, the term "DT-Diaphorase" will be used throughout this review.

DTD bioactivates chemotherapeutic quinones, which are referred to as bioreductive agents. There

* Corresponding author. Present address: Department of Medical Oncology, Christie Hospital NHS Trust, Wilmslow Road, Withington, Manchester M20 4BX, UK. Fax: +44-161-446-3299.
E-mail address: sdanson@fsmail.net (S. Danson).

is often significant over-expression of DTD in neoplastic tissue compared to the surrounding normal tissues. As a result of this, DTD has attracted considerable attention as a potential candidate for targeted anticancer therapy.

Many comprehensive reviews have already been published about DTD and bioreductive drugs.²⁻⁷ This review seeks to discuss the general background and include more recent developments.

Structure, distribution and mechanism of action

DTD is a flavoprotein which exists as a dimer with each unit having a molecular weight of 32,000 kD and being associated with a flavin adenine dinucleotide (FAD) group, noncovalently attached to the protein. DTD is predominantly cytosolic with over 90% being found in the cytoplasm.⁸ However, earlier studies showed DTD in the endoplasmic reticulum,⁹ mitochondria¹⁰ and Golgi body¹¹ and recent work has shown that DTD is present in the nucleus.¹²

DTD is an obligatory two electron reductase enzyme. This is in contrast to the one electron reductase enzymes such as NADH:cytochrome *b* reductase, NADPH:cytochrome P450 reductase and xanthine oxidase. DTD functions via a "ping-pong" kinetic mechanism (cofactor binds on and then off followed by substrate on and then off) utilising the co-factors NADH and NADPH equally well as the electron donor.¹³

DTD has multiple cellular roles, many of which are poorly understood. In this review, we are most interested in the reduction of benzoquinones to hydroquinones, especially cytotoxic quinones, but it is also involved in reduction of dietary and environmental quinones. DTD also acts as a Phase II detoxification enzyme with the detoxifying step bypassing the formation of free radicals and so protecting tissues against mutagens, carcinogens and cytotoxics. DTD can recycle the membrane antioxidants ubiquinone and vitamin E.^{14,15} DTD is induced in a "stress response", for example in hypoxic conditions, along with other enzymes, including the glutathione S-transferases which conjugate hydrophobic electrophiles and reactive oxygen species, UDP-transferases which catalyse the conjugation and thus excretion of xenobiotics, epoxide hydrolase which inactivates epoxides and γ -glutamylcysteine synthetase which regulates glutathione metabolism.¹⁶

DTD is widely distributed in animals, plants and bacteria. In humans, DTD is distributed throughout

the body with highest levels in epithelial and endothelial cells, especially of the kidney and gastrointestinal tract.^{17,18} The amino acid sequence has been determined for rat, mouse and human DTD.¹⁹ The amino acid sequence of mouse DTD is more homologous to the rat enzyme than the human enzyme. However, rat DTD has faster reductive activation than mouse DTD due to a base difference at residue 104 which is critical for catalytic activity.¹⁹ In addition, DTD has high levels of activity in extrahepatic tissues in humans but is most abundant in the liver of other mammals such as rat, monkey and dog. Mouse has less DTD in the liver than other animals and this, in combination with the similar reductive rate to humans, means that mouse models may be more comparable to humans than those of rats or other species.

Gene

DTD activity is encoded by four genetic loci known as NQO1, NQO2, NQO3 and NQO4.²⁰ The majority of DTD is coded for by the NQO1 gene but as yet the roles of the other three are poorly understood. NQO1 is a single copy gene located on chromosome 16q22.1 and encodes for a protein of 274 residues.²¹ NQO1 consists of 6 exons and 5 introns and spans a 20 kilobase region. Exon 1 codes for the 5'UT region whilst exons 2-6 code for the remainder of the gene and the 3'UT region. The sixth exon has four potential polyadenylation signal sequences, three of which are used. A single copy of human Alu repetitive sequence is sited between the second and third polyadenylation sites. Jaiswal et al.²² have sequenced the region upstream of the protein coding sequence for DTD. They have identified several *cis*-activating transcriptional activators, including an antioxidant response element (ARE), within which a perfect AP1 site resides along with a xenobiotic response element (XRE), an AP2 element, a CAT box, a TATA box and a NF- κ B binding site.²²⁻²⁴ Jun, fos and other novel proteins from the nucleus, including oestrogen receptors,²⁵ bind to the ARE and mediate signal transduction. The molecular biology of DTD expression has been extensively reviewed by Jaiswal.¹⁶

A second gene product, NQO2 (NRH:quinone oxidoreductase) has been described. The role of NQO2 in the detoxification of quinones is unknown. The NQO2 gene is located on chromosome 6p25 and the protein product has 231 residues (43 amino acids shorter than NQO1 at the carboxyl terminal end). The initiation codon of the NQO2 cDNA aligns perfectly with the initiation codons of human and rat NQO1 cDNAs. NQO1 and NQO2 are homologous

for 54% of cDNA and 49% of protein. NQO2 is expressed in fewer tissues (including heart, lung, brain and skeletal muscle) than NQO1. NQO2 and NQO1 differ in their cofactor requirement with NQO2 using dihydronicotinamide riboside (NRH) rather than NADH or NADPH as the electron donor. NQO2 is resistant to typical inhibitors of DTD such as dicumarol. Readers are referred to the reviews by Jaiswal, Chen and Long for more detailed discussion of NQO2.^{26–28}

Polymorphism

Traver et al.²⁹ have demonstrated a single point nucleotide polymorphism (SNP), known as NQO1*2, in the NQO1 sequence of the BE colorectal cell line, which has no detectable DTD activity. A homologous base substitution (cytosine to thymidine) at position 609 on the cDNA sequence results in a proline to serine substitution in the protein. The inactive DTD expressed is much less stable than wild-type DTD. Wild-type DTD persists in cells for over 18 h whilst the mutant protein is broken down via the ubiquitin proteasomal pathway (UPP) in just 1.2 h.³⁰ This SNP appears to cause a change in the enzyme conformation so that there is a decrease in the FAD binding affinity and thus a loss of enzyme activity.³¹ There is ethnic variation in the occurrence of this SNP with 4.4% of the Caucasian population and 20.3% of Asians having the homologous genotype.³² Heterozygotes for this SNP have intermediate activity compared with people with the wild type.³³ Multiple further aberrations of DTD have been reported but the physiological significance of these have not yet been determined. These include the 465C>T variant (NQO1*3) which is associated with increased alternative splicing and decreased expression of DTD.^{34,35}

Localisation

Winski et al.¹² recently confirmed the presence of DTD in the nucleus. This is of interest when we consider the bioreduction of cytotoxic agents as it would allow the drug to be activated in the nucleus itself. The process of nuclear localisation can occur by either passive diffusion through nuclear pores, active transport by nuclear localisation signals or binding to chaperone proteins. The DTD molecule is too large for passive diffusion, its diameter is 60–70 nm and would need to be less than 10 nm.³⁶ The classic nuclear localisation signal required for active transport is not present in the DTD molecule. Therefore DTD is thought to bind to molecular chaperone proteins, such as HSP-70. HSPs are in-

volved in protein synthesis and folding, vesicular trafficking, and antigen processing and presentation. HSP-70 binds to hydrophobic regions of unfolded proteins which go through ATP-dependent binding cycles and the nascent protein is manipulated into an active conformation. Using co-immunoprecipitation, it has been shown that HSP-70 associates with wild-type DTD, and thus stabilises it, but not with the mutant DTD.³⁷ Anwar et al.³⁷ have generated a plasmid containing a DTD coding region with a mutated HSP-70 binding site. This plasmid did not associate with HSP-70, was catalytically inactive and was degraded by the UPP. DTD itself has been implicated in the stabilisation of the tumour suppressor gene, p53, which is responsible for regulating cell cycle G1 arrest and for activating apoptosis. Wild-type DTD does interact with p53 but the cellular role of this interaction is not yet known. DTD-mediated reduction of quinones induces p53³⁸ and this reduction, and subsequent p53 induction, can be suppressed using the specific DTD inhibitor, dicumarol.^{39,40} Comparison of inhibitors of DTD and inhibitors of another HSP, HSP-90 indicates that DTD and HSP-90 act through different mechanisms to induce p53 degradation and thus suppress apoptosis.⁴⁰

DTD and cancer

DTD is over-expressed in many cancerous tissues compared to normal tissue, especially non-small cell lung carcinoma (NSCLC) (Table 1).^{18,41–53} Since DTD acts to detoxify and protect the cell from toxins and mutagens, it may be that upregulation of DTD is part of the early carcinogenic process. This may involve epigenetic factors such as hypomethylation of the NQO1 gene,⁵⁴ elevation of the jun and fos proto-oncogenes,⁵⁵ altered promoter function, altered repressor function or post-transcriptional modifications.⁵

All four histological sub-types of NSCLC exhibit an increase in enzyme expression and activity, especially tumours that are well-differentiated, stage I and II and epidermoid.⁵³ Small cell lung carcinomas (SCLC) do not have raised DTD activity.⁵⁰ However, when SCLC cell lines are transfected with the v-Ha-ras proto-oncogene, thus causing a transition to the NSCLC phenotype through an increase in AP1 activity,⁵⁶ DTD activity is increased.⁵⁰ Some workers have reported that DTD activity is highest in the lung tumours of those who have never smoked or who have stopped smoking. This trend is not replicated for colon tumours.¹⁸ Conversely, those lung cancer patients

Table 1 Ratio of DTD activity in tumour tissue compared to normal tissue

Cells	Ratio	References
Rat leydig cell tumours	2.5–7.7	41
Rat hepatomas	2–2.4	42
Hyperplastic rat liver nodules	6.0	43
Mouse liver tumours	1.5–2.2	44
Rat brain tumours	10.8	45
Rat ascites hepatoma	4.7	46
Human colon carcinoma primary	2.5–3.9	18,47,52
Human colon carcinoma metastases	4.7	52
Human breast carcinoma	NS–9.5	18,48,53
Human NSCLC	8.2–19.2	18,50,52,53
Human SCLC	0.25	50
Human liver carcinoma	3.8–50	18,49
Human stomach carcinoma	0.39	18
Human kidney carcinoma	0.1–0.3	18,51
Human head/neck (squamous)	1.07	52

NS: not significant.

who are alcohol users have significantly higher DTD activity in their tumours than those who did not; again this trend has not been replicated for those with colon tumours.¹⁸ These differences in the lung tissue of smokers were not observed in the work of Marin et al.⁵³

Other tumours which over-express DTD include colorectal, liver and breast carcinomas but increased expression is not seen with all tumour types. For example, stomach and kidney tumours have lower DTD activity than the surrounding normal tissue, and head and neck tumours have the same DTD activity as the surrounding normal tissue.^{51,52} Some tissue types have high baseline activity levels, such as stomach and kidney, and it may not be possible to increase the DTD activity further. There may also be marked variation of DTD activity within a specific tumour type⁵⁰ and possibly within an individual tumour. Advanced metastatic tissue has been found to have greater expression and activity of DTD than tumours without metastases and poorly differentiated tumours have higher activity than well-differentiated ones.^{41,57} However, others have found lower absolute DTD activity in lymph node and liver metastases than in the primary tumour.⁵² High expression and activity of DTD are typically found in "normal" hepatic tissue surrounding hepatic tumours.^{46,49} It should be noted that DTD protein levels do not necessarily correlate with DTD activity.

DTD may play a role in cancer prevention so, conversely, it would be expected that individuals with the NQO1*2 SNP, and thus no active DTD,

would have a higher susceptibility to developing cancer. Several studies have shown a correlation between NQO1 polymorphism and cancer incidence. Kidney tumour, lung cancer, colorectal carcinoma and oesophageal and gastric cancer patients all have a higher percentage of the NQO1 polymorphism than the normal population.^{58–61} This correlation has not been shown in other studies with kidney, lung and colorectal cancer patients^{62–65} or in patients with adult glioma,⁶⁶ Hodgkins Disease,⁶⁷ non-Hodgkins lymphoma⁶⁷ or prostate cancer.⁶⁸ Lack of DTD is associated with an increased risk of developing adult leukaemia⁶⁹ and paediatric leukaemia.^{70,71} In cancer patients, the NQO1 polymorphism may be associated with an increased risk of chemotherapy-related myeloid leukaemia, including those leukaemias associated with abnormalities in chromosomes 5 and 7⁷² but others disagree.⁷³ In addition, individuals with absent DTD activity do have increased susceptibility to bone marrow suppression after environmental exposure to benzene and benzene-like compounds.⁷⁴ In NQO1 knock-out mice, no detectable change in phenotype is observed and they appear indistinguishable from wild-type mice, however they are more susceptible to quinone toxicity⁷⁵ and have increased sensitivity to the development of skin and visceral tumours.¹⁶

Role in anticancer therapy

Paradoxically, the detoxifying enzyme DTD can activate certain xenobiotics, such as MMC, ortho-naphthoquinones and aziridinybenzoquinones.^{76–78} This is of interest since drugs that are effectively bioactivated by DTD may allow tumour cytotoxicity without corresponding high levels of toxicity to normal tissues (see review by Workman,⁵⁵). Reduction of quinones by DTD may lead to either production of a reactive alkylating species by rearrangement, or hydroquinones may auto-oxidise leading to the production of reactive oxygen species and toxicity. The reductive cycle can continue along both pathways unless the system becomes anaerobic at which time oxygen radical production decreases and the semiquinone can accumulate.² A correlation exists between the cytotoxicity of these bioreductive compounds, DTD activity, an increase in the pro-apoptotic BAD protein and the induction of apoptosis.⁷⁹ Although, bioreductive quinones are more active in tumour cells that over-express DTD, there is still some effect on cells without DTD, presumably due to single electron enzymatic reduction.⁸⁰

Mitomycin C

Mitomycin C (6-Amino-8-{[(aminocarbonyl)oxy]methyl}-1,1a,2,8,8a,8b-hexahydro-8-methoxy-5-methyl[1aR-(1a α ,8 β ,8a α ,8b α)]azirino(2',3':3,4)-pyrrolo-(1,2- α)indole-4,7-dione) is a naturally occurring anti-tumour quinone derived from *Streptomyces caespitosus*, a strain of actinomyces. It has been used as a cytotoxic since the 1960s and has activity against lung, stomach, head and neck, prostate, breast and bladder tumours. The dose-limiting toxicity of MMC is myelosuppression. MMC is the prototypical bioreductive agent as it is inactive until reduced by one-electron reductases or DTD.⁸¹ Reduction allows DNA cross-links and strand breaks to form, thus inhibiting DNA replication. MMC sensitivity has been shown to be associated with high DTD levels by some workers but others have demonstrated discrepancies between in vitro and in vivo cytotoxicity.^{29,50,78,82-87} Relating MMC cytotoxicity to DTD levels is complicated by the fact that MMC is a relatively poor substrate for DTD, is also bioactivated by one-electron reductases, activation is pH-dependent and may be influenced by hypoxia.^{87,88} It has become evident that superior substrates for DTD exist and there has been much effort to produce novel agents that will target DTD more effectively.

E09

The indoloquinone E09 (3-hydroxy-5-aziridinyl-1-methyl-2(1H-indole-4,7-dione)-prop-beta-en-alpha-ol) is a synthetic analogue of MMC. E09 is a moderately improved substrate for DTD when compared to MMC as it can redox cycle more effectively and is not pH-dependent. A strong association has been described between elevated DTD activity and E09 sensitivity.^{81,83, 89} E09 entered clinical trials on the basis of its distinct anti-tumour profile and lack of myelosuppression. However, E09 failed to produce any clinical responses and this has been attributed to a combination of rapid plasma clearance and dose-limiting kidney toxicity.⁹⁰

CB1954

CB1954 (5-(aziridin-1-yl)-2,4-dinitrobenzamide) is a nitrophenylaziridine which is cytotoxic to rat carcinoma cells which over-express DTD.⁹¹ However, CB1954 is bioactivated much more slowly by human DTD than by rat DTD which has profound implications on its clinical utilisation.⁹² Human tumour cells require the NQO2 gene, which is

normally latent, to be activated by the synthetic co-factor NRH before CB1954 can be used in a cytotoxic manner.⁹³

AZQ

AZQ (2,5-bis(carboethoxyamino)-3,6-diaziridinyl-1,4-benzoquinone) is an aziridinylbenzoquinone with good solubility. AZQ does not readily alkylate DNA at physiologically relevant pH but has a relatively high reduction potential and is easily reduced by both one electron reductases and DTD to form DNA alkylating species.⁹⁴ AZQ activation is inhibited by the DTD inhibitor, dicumarol.⁹⁴ A correlation is seen between DTD levels and cytotoxicity with AZQ.⁹⁵ AZQ underwent clinical trials, activity was demonstrated against progressive gliomas⁹⁶ and refractory lymphomas⁹⁷ but it was not able to surpass agents already in use.

BZQ

BZQ (2,5-bis(2-hydroxyethylamino)-3,6-diaziridinyl-1,4-benzoquinone) is a powerful cross-linking aziridinylbenzoquinone which does not easily undergo reduction by either one or two electron reductases. BZQ is much less stable than AZQ and can readily alkylate in the absence of reduction⁹⁴ and its cytotoxicity is unaffected by the addition of dicumarol.⁸⁰

MeDZQ

MeDZQ (2,5-diaziridinyl-3,6-dimethyl-1,4-benzoquinone) is the lead compound in the aziridinylbenzoquinone series. MeDZQ is an excellent substrate for DTD^{95,98} and, after reduction by DTD, forms cross-links at GNC sequences.⁹⁹ There is a strong correlation between DTD levels in NSCLC cell lines and the cytotoxicity of MeDZQ.⁸⁷ However, the development of MeDZQ was limited by its poor solubility.

RH1

The novel diaziridinylbenzoquinone RH1 (2,5-diaziridinyl-3-hydroxymethyl-6-methyl-1,4-benzoquinone) (Fig. 1) may have advantages over existing bioreductive agents as it has good solubility, is an excellent substrate for DTD and anti-tumour effect has been demonstrated both in vitro and in vivo. RH1 is more soluble in water




Figure 1 RH1 molecule, with two aziridine rings and hydroxyl group.

than MeDZQ due to its hydroxyl moiety. RH1 is rapidly taken up by cells and undergoes aziridine ring opening, which is enhanced by the presence of DTD, and causes formation of DNA adducts, particularly at GCC sequences.¹⁰⁰ The products of reduction by RH1 are more stable, less likely to be involved in redox cycling and less likely to be influenced by hypoxia than those of MMC.¹⁰¹ RH1 is cleared ten-fold slower from the plasma than EO9.¹⁰² In keeping with its mechanism of action RH1 has been shown to be significantly more active in cells that over-express DTD, naturally or through transfection, than in cell types with low DTD.^{102,103} RH1 is an excellent substrate for DTD and is 50% more active than MeDZQ and 225 times better than MMC in cell lines.¹⁰³ RH1 has undergone in vivo efficacy testing by the Developmental Therapeutics Program of the National Cancer Institute and has demonstrated significant activity in both NSCLC and ovarian xenograft models.¹⁰⁴ Based upon the entire toxicology data in mice, rats and dogs, a starting dose level of 40 $\mu\text{g}/\text{m}^2$ daily for five days was suggested for human trials based upon this being 1/10th of the maximum tolerated dose in the most sensitive species, the dog. The mouse may be a better model as mouse and human liver have low levels as DTD whereas rat, monkey and dog have substantial amounts. Toxicity seen has included myelosuppression and local injection site inflammation. The lower renal metabolic rate seen with RH1, compared with EO9, suggests that RH1 may cause less renal toxicity.¹⁰² RH1 formulated in 20% hydroxypropyl- β -cyclodextrin is associated with a reduction in injection site toxicity compared to its formulation in water without a detrimental effect on stability and this has been used as the drug delivery vehicle in subsequent studies.¹⁰⁴ RH1, diluted in cyclodextrin and given once a day for five days, has entered a phase I dose escalation study in human patients with solid tumours, under the auspices of Cancer Research UK.

Manipulation of DTD

Much is known about the inhibition of DTD by dicumarol. Of more interest are the selective induction of DTD in tumour tissue to increase tumour cell kill whilst minimising normal tissue toxicity and potential mechanisms of resistance to RH1 and whether they can be overcome.

Inhibition

DTD is very sensitive to the inhibitory effects of the anti-coagulants, dicumarol and warfarin.¹⁰⁵ Dicumarol binds with the FAD group and competitively inhibits DTD with respect to the nicotinamide co-enzymes.¹⁰⁶ DTD is also competitively inhibited by Cibacron blue, chrysin, 7,8-dihydroxyflavone and phenidone. Recently, a potent inhibitor of DTD, ES936 (5-methoxy-1,2-dimethyl-3-[(4-nitrophenoxy)methyl]indole-4,7-dione) has been described.¹⁰⁷

Induction

DTD expression in tumour cell lines can be induced by a variety of structurally different agents, including the thiol, 1,2-dithiol-3-thione (D3T); oltipraz (5-(2-pyrazinyl)-4-methyl-1,2-dithiole-3-thione), a synthetic D3T analogue that was used to treat schistosomiasis; extracts of cruciferous vegetables (broccoli and green onions) such as fisetin and quercetin, two flavenols; dietary antioxidants, such as 2(3)-tert-butyl-4-hydroxyanisole (BHA); benzenes; azo dyes; aspirin-like drugs (ALD); dioxin; diphenols, anti-oestrogens, glucocorticoids and isothiocyanates^{108–126} (Table 2). DTD is also induced by hypoxia,¹²⁷ light¹²⁸ and the addition of some chemotherapeutic agents, including MMC and doxorubicin.^{125,129} There are several comprehensive reviews on induction including that by Begleiter et al.¹³⁰

Induction of DTD is mediated via an ARE, which is essential for the co-ordinated induction and expression of DTD. The AP1 site in the ARE is sensitive to induction by a number of agents and hypoxia, however AP1 binding proteins may not be necessary in all instances for ARE activation. Over-expression of c-fos or FRA1 represses binding to the ARE. Disruption of c-fos can increase DTD activity suggesting that c-fos is negatively regulating DTD.^{3,16} It has been hypothesised¹⁶ that Nrf2 is bound to Keap1, a repressor protein, and is retained in the cytoplasm. Dissociation of the Nrf2-keap1 dimer allows free Nrf2 to enter the nucleus where it heterodimerises with c-jun and binds to ARE. In-

terestingly Kepa et al.¹³¹ have reported that in NSCLC, FRA1 and FRA2 along with jun-b bind to the AP1 site thus enabling induction of DTD. Conversely in SCLC, FRA1/FRA2 and jun-b appear to be absent which could explain why SCLC does not over-express DTD whilst NSCLC does. There is also evi-

dence of involvement in induction by the NF- κ B response element in the 5' flanking region of the DTD gene.²⁴ Others have shown that the extracellular signal regulated protein kinase (ERK) pathway negatively regulates ARE induction.¹³² It is felt that unknown cytosolic factors are involved in the

Table 2 Levels of DTD induction (induced value/baseline value)

Inducer	Target	Ratio	Reference
D3T	Murine lymphoma	22	108
	Murine bone marrow	2–3	109
	Human leukaemia	2–10	110–112
	Human head & neck cancer	1.3	111,112
	Human lung cancer	1.1–8.9	111,112
	Human colon cancer	NS–2.5	111,112
	Human breast cancer	NS–9.9	111,112
	Human ovarian cancer	1.5–2.7	111
	Human prostate cancer	NS–2.6	111,112
	Human skin cancer	1.2–1.4	111
	Human liver cancer	NS	111,112
	Human stomach cancer	1.7–3.3	111
	Human bone marrow	10.5	111
	Human kidney	3.5	111
	Human lung	2.4	111
D3T/MMC	Human NSCLC	2.5	113
	Human breast cancer	1.8–3.6	113
	Human colon cancer	2.3	113
	Human melanoma	2.3	113
	Human gastric cancer	1.9	113
Oltipraz	Human marrow	6.7	113
	Murine cell lines	1.5–3	114
	Rat liver post-aflatoxin	2.2–7.1	115
	Human colorectal cancer	2.0	116
Fisetin	Murine hepatoma	2.6	121
Quercetin	Human breast cancer	2.0	123
Retinoic acid	Human breast cancer	2.2	113
DMM	Human breast cancer	4.4	113
DMM/MMC	Human breast cancer	2.9	113
DMF	Human breast cancer	4.6	113
	Human lymphocytes	6.5	118
PG	Human breast cancer	2.9	113
PG/MMC	Human breast cancer	1.8	113
Sulforaphone	Murine hepatoma	6.0	117
	Murine liver and gut	1.5–3.1	119,124
	Human hepatoma	5.9	117
	Human breast cancer	0.4–9.4	113,117
	Human prostate cancer	1.8	117
	Human cervical cancer	3.9	117
	Human colon cancer	10.3	117
BHA	Murine liver	10.0	17
	Murine kidney	4.2	17
	Murine lung	3.6	17
	Murine small intestine	2.7–6.2	17,124
tBHQ	Murine hepatoma	5.3	117
	Human hepatoma	6.4	117
	Human breast cancer	0.6–10.0	117
	Human prostate cancer	3.0	117
	Human cervical cancer	4.3	117
	Human colon cancer	10.8	117

Table 2 (continued)

Inducer	Target	Ratio	Reference
β -naphthoflavone	Murine hepatoma	8.5	117
	Human hepatoma	6.3	117
	Human breast cancer	0.5–12.5	117
	Human prostate cancer	1.4	117
	Human cervical cancer	4.0	117
	Human colon cancer	9.5	117
	Human lymphocytes	1.5	118
Methyl vinyl sulfone	Human lymphocytes	3.9	118
Aspirin	Human breast cancer	1.5	113
Ibuprofen	Human breast cancer	NS	113
ALD	Human T lymphocytes	2.0–4.0	120
Anti-oestrogens	Murine hepatoma	1–2.9	122
Doxorubicin	Murine mammary	1.4	125
	Murine bone marrow	NS	125

ALD: aspirin like drugs, BHA: 2(3)-tert-butyl-4-hydroxyanisole, DMF: dimethyl fumarate, DMM: dimethyl maleate, MMC: mitomycin C, NS: not significant, PG: propyl gallate, tBHQ: tert-butylhydroquinone.

stimulation of Nrf and Jun binding to the ARE and that these may be kinases, phosphatases or redox proteins.¹⁶

Most work has been undertaken on D3T which is an effective monofunctional inducer (it elevates phase II enzymes selectively), both in vitro and in vivo.¹¹⁵ Oltipraz is an analogue of D3T that has been shown to act as a chemoprotective agent in vivo and is now being assessed in clinical trials.¹³³ In this instance, we are interested in oltipraz as an inducing agent in established tumours. Dose–response curves of D3T and oltipraz have indicated that D3T was 15-fold more potent as an enzyme inducer than oltipraz,¹³⁴ and indeed D3T is superior to all the other analogues against which it has been compared. However, in practice, there are differences between the effects of inducers on murine and human cell lines¹¹¹ and oltipraz may have less side-effects in patients than D3T although this remains to be tested.

The differential between normal and neoplastic tissue is maintained, or even accentuated, when DTD is induced by D3T. In murine lymphoma cells, D3T increases the DTD activity by 22-fold but has little or no effect on normal mouse bone marrow cells.¹⁰⁸ Transcription is upregulated and increased levels of DTD, with increased catalytic activity, are observed in the cells 24–48 h after exposure to the inducing agent, oltipraz in this case, but enzyme activity returns to control levels after 72 h in drug free media, although mRNA levels remain elevated.¹¹⁶ Induction occurs in a dose-dependent manner.¹²¹ The optimal duration of drug exposure is 24 h and more prolonged exposure is associated with lower levels of induction of detoxification enzymes, including DTD. Oltipraz given twice-

weekly for 12 weeks did not modulate DTD suggesting that chronic continuous dosing is not useful in induction.¹³⁵ These data support intermittent schedules of inducing agents.¹¹⁶

An increased DTD level is associated with increased sensitivity to MMC and EO9. Combination therapy with D3T and MMC or EO9 produces a 2- or 7-fold enhancement respectively of the cytotoxic effects of these antitumour agents in murine lymphoma cells.¹⁰⁸ This antitumour activity can be inhibited by dicumarol, supporting the proposal that the enhanced tumour cell kill is due to elevated levels of DTD.¹³⁰ D3T has been shown to induce DTD activity in 28 of 38 human tumour cell lines, representing 10 tissue types, and significantly increased the cytotoxicity of EO9 in these cells. DTD levels were also induced in normal bone marrow (from low baseline levels), kidney (from low levels) and lung (from intermediate levels) without an increase in toxicity to normal kidney cells.¹¹¹ The addition of MMC to D3T did not enhance toxicity in marrow cells whereas D3T with EO9 produced a small increase in marrow toxicity.¹⁰⁸ A further study¹¹³ demonstrated that D3T increased DTD levels (but not the levels of other detoxification enzymes) in all six tumour cell lines studied and this resulted in increased cytotoxicity of MMC in four of the six lines by 1.4 to threefold. In this study, the combination of D3T and MMC was associated with a 50% increase in toxicity in normal human bone marrow cells but this was small in comparison with the enhanced cytotoxicity.¹¹³ Induction of DTD with D3T and ALDs has been shown to reduce the cytotoxicity of benzene and its metabolites on bone marrow stromal cells and lymphocytes.^{109,120} The absolute DTD level appears

to be important for induction: DTD activity cannot be induced if there is no baseline DTD activity when cells are homozygous for the NQO1*2 polymorphism¹³⁰ and it cannot be further induced if the baseline levels are very high. In the study by Wang et al.¹¹³ the two cell lines that did not exhibit an increase in MMC cytotoxicity had relatively high baseline DTD levels.

Resistance

The clinical role of bioreductive agents may be limited by both inherent and developed drug resistance. MMC resistance may be mediated by increased drug efflux, the repair of DNA lesions or a deficiency in bioreductive enzymes, such as DTD.¹³⁶ Down-regulation of DTD is associated with MMC resistance in human gastric, bladder and colon cancer cell lines^{29,137–140} and this is not necessarily associated with downregulation of P450 reductase.²⁹ Resistance by this mechanism can be reversed by transfection of a resistant cell line with DTD and P450 reductase or by placing it under hypoxic conditions.¹⁴¹ Cell lines often show broad cross-resistance to other cytotoxic agents; this may be due to a "molecular switch" controlling the expression of several detoxification enzymes, some involved in primary resistance and others in cross-resistance. Cisplatin-resistant ovarian cell lines demonstrate increased DTD levels with increased resistance to cisplatin and cross-resistance to MMC.¹⁴² Multidrug resistance involving a decrease in DTD activity has been demonstrated in rat ascites hepatoma cells treated with doxorubicin which were cross-resistant to MMC and the MMC analogue, porfiromycin.¹⁴³ Serial exposure of human erythroleukaemic cells to the diaziridinylbenzoquinones, AZQ and BZQ, resulted in resistant cell lines with decreased levels of both DTD and P450 reductase and increased levels of glutathione and superoxide dismutase compared to the parent cell line.¹⁴⁴ These resistant cell lines also demonstrated cross-resistance to doxorubicin and MMC. Resistance to RH1 has been developed in cancer cell lines and studies into the mechanisms of resistance in these lines are ongoing.¹⁴⁵

Summary

DTD has the potential to be a very important target in oncology. The differential expression of DTD between neoplastic tissue and normal tissue should allow targeting of bioreductive agents to the tumour cells whilst minimising normal tissue toxicity.

However, no cell system can predict for the multitude of other factors influencing cytotoxicity and normal tissue toxicity, including tumour heterogeneity, effect of other bioreductive enzymes, degree of hypoxia, cell nutrients, type of damage produced and effectiveness of DNA repair. In addition, none of the bioreductive agents developed to date has been specific for activation by a single reductive enzyme. Previous agents have provided disappointing results, especially EO9 which failed in clinical practice due to a combination of rapid plasma clearance and dose-limiting renal toxicity. Theoretically, RH1 is superior to other bioreductive drugs. It is soluble, has anti-tumour activity in vitro and in vivo and may have a high enough specificity for DTD to allow increased effect in the tumour cells whilst sparing normal tissue. The phase I clinical trial of RH1 in human solid tumours is ongoing. Other areas of interest include manipulation and localisation of DTD and investigation of the mechanisms of drug resistance.

References

1. Ernster L, Navazio F. Soluble diaphorase in animal tissues. *Acta Chemica Scandinavica* 1958;12:595–602.
2. Gutierrez PL. The role of NAD(P)H oxidoreductase (DT-diaphorase) in the bioactivation of quinone-containing antitumour agents: a review. *Free Rad Biol Med* 2000;29:263–75.
3. Ross D, Kepa J, Winski S, Beall HD, Anwar A, Siegel D. NAD(P)H:quinone oxidoreductase 1 (NQO1): chemoprotection, gene regulation and genetic polymorphisms. *Chem Biol Int* 2000;129:77–97.
4. Rauth AM, Goldberg Z, Misra V. DT-diaphorase: possible roles in cancer chemotherapy and carcinogenesis. *Oncol Res* 1997;9:339–49.
5. Ross D, Beall H, Traver R, Siegel D, Phillips RM, Gibson NW. Bioactivation of quinones by DT-diaphorase, molecular, biochemical and chemical studies. *Oncol Res* 1994;6:493–500.
6. Ross D, Siegel D, Beall H, Prakash AS, Mulcahy RT, Gibson NW. DT-diaphorase in activation and detoxification of quinones. Bioreductive activation of mitomycin C. *Cancer Metastasis Rev* 1993;12:83–101.
7. Riley RJ, Workman P. DT-diaphorase and cancer chemotherapy. *Biochem Pharmacol* 1992;43:1657–69.
8. Eliasson M, Bostrom M, Depierre JW. Levels and subcellular distributions of detoxifying enzymes in the ovarian corpus luteum of the pregnant and non-pregnant pig. *Biochem Pharmacol* 1999;58:1287–92.
9. Danielson L, Ernster L, Ljunggren M. Selective extraction of DT-diaphorase from mitochondria and microsomes. *Acta Chem Scand* 1960;14:1837–8.
10. Conover TE, Ernster L. DT-diaphorase IV. Coupling of extramitochondrial reduced pyridine nucleotide oxidation to mitochondrial respiratory chain. *Biochem Biophys Acta* 1968;67:268–80.
11. Edlund C, Elhrammer A, Dallner G. Distribution of newly synthesised DT-diaphorase in rat liver. *Biosci Rep* 1982;2:861–5.

12. Winski SL, Koutalos Y, Bentley DL, Ross D. Subcellular localisation of NAD(P)H:quinone oxidoreductase 1 in human cancer cells. *Cancer Res* 2002;**62**:1420–4.
13. Siegel D, Gibson NW, Preusch PC, Ross D. Metabolism of NAD(P)H: (quinone acceptor) oxidoreductase (DT-diaphorase): role in diaziquone-induced DNA damage and cytotoxicity in human colon carcinoma cells. *Cancer Res* 1990;**50**:7293–300.
14. Beyer RE, Segura-Aguilar J, Di Bernardo S, et al. The role of DT-diaphorase in the maintenance of the reduced antioxidant form of coenzyme Q in membrane systems. *PNAS* 1996;**93**:2528–32.
15. Siegel D, Bolton EM, Burr JA, Liebler DC, Ross D. The reduction of alpha-tocopherolquinone by human NAD(P)H:quinone oxidoreductase: the role of alpha-tocopherolhydroquinone as a cellular antioxidant. *Mol Pharmacol* 1997;**52**:300–5.
16. Jaiswal AK. Regulation of genes encoding NAD(P)H:quinone oxidoreductases. *Free Rad Biol Med* 2000;**29**:254–62.
17. Benson AN, Hunkler MJ, Talalay P. Increase of NAD(P)H:quinone reductase by dietary antioxidants: possible role in protection against carcinogenesis and toxicity. *PNAS* 1980;**77**:5216–20.
18. Schlager JJ, Powis G. Cytosolic NAD(P)H: (quinone-acceptor) oxidoreductase in human and normal tissue: effects of cigarette smoking and alcohol. *Int J Cancer* 1990;**45**:403–9.
19. Chen S, Knox R, Wu K, Deng PSK, Zhou D. Molecular basis of the catalytic differences among DT-diaphorase of human, rat, and mouse. *J Biol Chem* 1997;**272**:1437–9.
20. Edwards YH, Potter J, Hopkinson DA. Human FAD-dependent NAD(P)H diaphorase. *Biochem J* 1980;**187**:429–36.
21. Jaiswal AK, McBride OW, Adesnik M, Nebert DW. Human dioxin-inducible cytosolic NAD(P)H:menadione oxidoreductase cDNA sequence and localization of gene to chromosome 16. *J Biol Chem* 1988;**263**:13572–8.
22. Jaiswal AK. Human NAD(P)H:quinone oxidoreductase (NQO1) gene structure and induction by dioxin. *Biochemistry* 1991;**30**:10647–53.
23. Joseph P, Xie T, Xu Y, Jaiswal AK. NAD(P)H:quinone oxidoreductase 1 (DT-diaphorase): expression, regulation and role in cancer. *Oncol Res* 1994;**6**:525–32.
24. Yao K, O'Dwyer PJ. Involvement of NF- κ B in the induction of NAD(P)H:quinone oxidoreductase (DT-diaphorase) by hypoxia, oltpiraz and mitomycin C. *Biochem Pharmacol* 1995;**49**:275–82.
25. Montano MM, Jaiswal AK, Katzenellenbogen BS. Transcriptional regulation of the human quinone reductase gene by antioestrogen-liganded oestrogen receptor- α and oestrogen receptor- β . *Oncogene* 1998;**273**:25443–9.
26. Jaiswal AK, Burnett P, Adesnik M, McBride OW. Nucleotide and deduced amino acid sequence of a human cDNA (NQO2) corresponding to a second member of the NAD(P)H:quinone oxidoreductase gene family. Extensive polymorphism at the NQO2 gene locus on chromosome 6. *Biochemistry* 1990;**29**: 1899–906.
27. Chen S, Wu K, Knox R. Structure-function studies of DT-diaphorase (NQO1) and NRH:quinone oxidoreductase (NQO2). *Free Rad Biol Med* 2000;**29**:276–84.
28. Long II DJ, Jaiswal AK. NRH:quinone oxidoreductase 2 (NQO2). *Chem Biol Int* 2000;**129**:99–112.
29. Traver RD, Horikoshi T, Danenberg KD, et al. NAD(P)H:quinone oxidoreductase gene expression in human colon carcinoma cells: characterisation of a mutation which modulates DT-diaphorase activity and mitomycin sensitivity. *Cancer Res* 1992;**52**:797–802.
30. Siegel D, Anwar A, Winski SL, Kepa JK, Zolman KL, Ross D. Rapid polyubiquitination and proteosomal degradation of a mutant form of NAD(P)H:quinone oxidoreductase 1. *Mol Pharmacol* 2001;**59**:263–8.
31. Li R, Bianchet MA, Talalay P, Amzel LM. The three-dimensional structure of NAD(P)H:quinone reductase, a flavoprotein involved in cancer chemoprotection and chemotherapy: mechanism of the two-electron reduction. *PNAS* 1995;**92**:8846–50.
32. Kelsey KT, Ross D, Traver RD, et al. Ethnic variation in the prevalence of a common NAD(P)H quinone oxidoreductase polymorphism and its implications for anticancer chemotherapy. *Br J Cancer* 1997;**76**:852–4.
33. Misra V, Grondin A, Klamut HJ, Rauth AM. Assessment of the relationship between genotypic status of a DT-diaphorase point mutation and enzymatic activity. *Br J Cancer* 2000;**83**:998–1002.
34. Gaedigk A, Tyndale RF, Jurima-Romet M, Sellers EM, Grant DM, Leeder JS. NAD(P)H:quinone oxidoreductase: polymorphisms and allele frequencies in Caucasian, Chinese and Canadian Native Indian and Inuit populations. *Pharmacogenetics* 1998;**8**:305–13.
35. Pan SS, Han Y, Farabaugh P, Xia H. Implication of alternative splicing for expression of a variant NAD(P)H:quinone oxidoreductase-1 with a single nucleotide polymorphism at 465C>T. *Pharmacogenetics* 2002;**12**:479–88.
36. Yoneda Y, Hieda M, Nagoshi E, Miyamoto Y. Nucleocytoplasmic protein transport and recycling of Ran. *Cell Struct Funct* 1999;**24**:425–33.
37. Anwar A, Siegel D, Kepa JK, Ross D. Interaction of the molecular chaperone HSP70 with human NAD(P)H:quinone oxidoreductase 1. *J Biol Chem* 2002;**277**:14060–7.
38. Qiu X, Schonthal AH, Cadenas E. Anticancer quinones induce pRb-preventable G2/M cell cycle arrest and apoptosis. *Free Rad Biol Med* 1998;**24**:848–54.
39. Ngo EO, Nutter LM, Sura T, Gutierrez PL. Induction of p53 by the concerted actions of aziridine and quinone moieties of diaziquone. *Chem Res Toxicol* 1988;**11**:360–8.
40. Asher G, Lotem J, Kama R, Sachs L, Shaul Y. NQO1 stabilises p53 through a distinct pathway. *PNAS* 2002;**99**:3099–104.
41. Schor NA, Rice BF, Huseby RA. Dehydrogenation of reduced pyridine nucleotides by leydig cell tumours of the rat testis. *Proc Soc Exp Biol Med* 1976;**151**:418–21.
42. Schor NA, Morris HP. The activity of DT-diaphorase in experimental hepatomas. *Biochem Biophys Acta* 1977;**2**: 5–9.
43. Schor NA, Ogawa K, Lee G, Farber E. The use of the DT-diaphorase for the detection of foci of early neoplastic transformation in rat liver. *Cancer Lett* 1978;**5**:167–71.
44. Becker FF, Stout DL. A constitutive deficiency in the monooxygenase system of spontaneous mouse liver tumours. *Carcinogenesis* 1984;**5**:785–8.
45. Berger MS, Talcott RE, Rosenblum ML. Use of quinones in brain tumour therapy: preliminary results of preclinical laboratory investigations. *J Toxicol Env Health* 1985;**16**: 713–9.
46. Beyer RE, Segura-Aguilar JE, Ernster L. The anticancer enzyme DT-diaphorase is induced selectively in the liver during ascites hepatoma growth. *Anticancer Res* 1988;**8**:233–8.
47. Schor NA, Cornelisse CJ. Biochemical and quantitative histochemical study of reduced pyridine nucleotide dehydrogenation by human colon carcinomas. *Cancer Res* 1983;**43**:4850–5.
48. Batist K, Cowan KH, Curt G, Kathi AG. Increased glutathione S-transferase activity (GST) in drug-treated human breast cancer cells. *Proc Am Assoc Cancer Res* 1985;**26**:1362a.

49. Cresteil T, Jaiswal AK. High levels of expression of the NAD(P)H:quinone oxidoreductase (NQO1) gene in tumour cells compared to normal cells of the same origin. *Biochem Pharm* 1991;42:1021–7.
50. Malkinson AM, Siegel D, Forrest GL, et al. Elevated DT-diaphorase activity and messenger RNA content in human non-small cell lung carcinomas – relationship to the response of human tumour xenografts to mitomycin C. *Cancer Res* 1992;52:4752–7.
51. Eickelmann P, Ebert T, Warskulat U, Schulz WA, Sies H. Expression of NAD(P)H:quinone oxidoreductase and glutathione S-transferase alpha and pi in human renal cell carcinoma and in kidney cancer-derived cell lines. *Carcinogenesis* 1994;15:219–25.
52. Smitskamp-Wilms E, Giaccone G, Pinedo HM, van der Laan BFAM, Peters GJ. DT-diaphorase activity in normal and neoplastic human tissues; an indicator for sensitivity to bioreductive agents. *Br J Cancer* 1995;72:917–21.
53. Marin A, Lopez de Cerain A, Hamilton E, et al. DT-diaphorase and cytochrome B5 reductase in human lung and breast tumours. *Br J Cancer* 1997;76:923–9.
54. Williams JB, Lu AYH, Cameron RG, Pickett CB. Rat liver NAD(P)H:quinone reductase. Construction of a quinone reductase cDNA clone and regulation of quinone reductase mRNA by 3-methylcholanthrene and in persistent hepatocyte nodules induced by chemical carcinogens. *J Biol Chem* 1986;261:5524–8.
55. Workman P. Enzyme-directed bioreductive drug development revisited: a commentary on recent progress and future prospects with emphasis on quinone anticancer agents and quinone metabolising enzymes, particularly DT-diaphorase. *Oncol Res* 1994;6:461–75.
56. Sistonen L, Holtta E, Makela TP, Keski-Oje J, Alitalo K. The cellular response to induction of the p21 c-Ha-ras oncoprotein includes stimulation of jun gene expression. *EMBO* 1989;8:815–22.
57. Mikami K, Naito M, Ishiguro T, et al. Immunological quantification of DT-diaphorase in carcinoma cell lines and clinical colon cancer cases: advanced tumours express greater levels of DT-diaphorase. *Jpn J Cancer Res* 1998;89:910–5.
58. Schulz WA, Krummeck A, Rosinger I, et al. Increased frequency of a null-allele for NAD(P)H:quinone oxidoreductase in patients with urological malignancies. *Pharmacogenetics* 1977;7:235–9.
59. Rosvold EA, Mcglynn KA, Lustbader ED, Buetow KH. Identification of an NAD(P)H:quinone oxidoreductase polymorphism and its association with lung cancer and smoking. *Pharmacogenetics* 1995;5:199–206.
60. Lafuente MJ, Casterad X, Trias M, et al. NAD(P)H:quinone oxidoreductase-dependent risk for colorectal cancer and its association with the presence of k-ras mutations in tumours. *Carcinogenesis* 2000;21:1813–9.
61. Zhang J-H, Li Y, Wang R, et al. NQO1 C609T polymorphism associated with esophageal and gastric cardiac carcinoma in North China. *World J Gastroenterol* 2003;9:1390–3.
62. Longuemaux S, Delomenie C, Gallon C, et al. Candidate genetic modifiers of individual susceptibility to renal cell carcinoma: a study of polymorphic human xenobiotic-metabolizing enzymes. *Cancer Res* 1999;59:2903–8.
63. Yin L, Pu Y, Liu T-Y, Tung Y-H, Chen K-W, Lin P. Genetic polymorphisms of NAD(P)H quinone oxidoreductase, CYP1A1 and microsomal epoxide hydrolase and lung cancer risk in Nanjing, China. *Lung Cancer* 2001;33:133–41.
64. Hamajima N, Matsuo K, Iwato H, et al. NAD(P)H:quinone oxidoreductase 1 (NQO1) C609T polymorphism and the risk of eight cancers for Japanese. *Int J Clin Oncol* 2002;7:103–8.
65. Harth V, Donat S, Ko Y, Abel J, Vetter H, Bruning T. NAD(P)H quinone oxidoreductase 1 codon 609 polymorphism and its association to colorectal cancer. *Arch Toxicol* 2000;73:528–31.
66. Peters ES, Kelsey KT, Wiencke JK, et al. NAT2 and NQO1 polymorphisms are not associated with adult glioma. *Cancer Epidemiol Biomark Prev* 2001;10:151–2.
67. Soucek P, Sarmanova J, Kristensen VN, Apltauerova M, Gut I. Genetic polymorphisms of biotransformation enzymes in patients with Hodgkin's and non-Hodgkin's lymphomas. *Int Arch Occup Environ Health* 2002;75:S86–92.
68. Steiner M, Hillenbrand M, Borkowski M, Seiter H, Schuff-Werner P. 609 C>T polymorphism in NAD(P)H:quinone oxidoreductase gene in patients with prostatic adenocarcinoma or benign prostatic hyperplasia. *Cancer Lett* 1999;135:67–71.
69. Smith MT, Wang Y, Kane R, et al. Low NAD(P)H:quinone oxidoreductase 1 activity is associated with increased risk of acute leukaemia in adults. *Blood* 2001;97:1422–6.
70. Wiemels JL, Pagnamenta A, Taylor GM, Eden OB, Alexander FE, Greaves MF. A lack of a functional NAD(P)H:quinone oxidoreductase allele is selectively associated with paediatric leukaemias that have MLL fusions. *Cancer Res* 1999;59:4095–9.
71. Krajcinovic M, Sinnett H, Richer C, Labuda D, Sinnett D. Role of NQO1, MPO and CYP2E1 genetic polymorphisms in the susceptibility to childhood acute lymphoblastic leukaemia. *Int J Cancer* 2002;97:230–6.
72. Larson RA, Wang Y, Banerjee M, et al. Prevalence of the inactivating 609C→T polymorphism in the NAD(P)H:quinone oxidoreductase (NQO1) gene in patients with primary and therapy-related myeloid leukemia. *Blood* 1999;94:803–7.
73. Blanco JG, Edick MJ, Hancock ML, et al. Genetic polymorphisms in CYP3A5 CYP3A4 and NQO1 in children who developed therapy-related malignancies. *Pharmacogenetics* 2002;12:605–11.
74. Rothman N, Smith MT, Hayes RB, et al. Benzene poisoning, a risk factor for haematological malignancy, is associated with the NQO1 609C>T mutation and rapid fractional excretion of chlorzoxazone. *Cancer Res* 1997;57:2839–42.
75. Radjendirane V, Joseph P, Lee YH, et al. Disruption of the DT-diaphorase (NQO1) gene in mice leads to increased menadione toxicity. *J Biol Chem* 1998;273:7382–9.
76. Walton MI, Smith PJ, Workman P. The role of NAD(P)H:quinone oxidoreductase (EC 1.6.99.2, DT-diaphorase) in the reductive bioactivation of the novel indoloquinone antitumour agent E09. *Cancer Commun* 1991;3:199–206.
77. Pardee AB, Li YZ, Li CJ. Cancer therapy with β -lapachone. *Curr Cancer Drug Targets* 2002;2:227–42.
78. Siegel D, Gibson NW, Preusch PC, Ross D. Metabolism of mitomycin C by DT-diaphorase: role in mitomycin C-induced DNA damage and cytotoxicity in human colon carcinoma cells. *Cancer Res* 1990;50:7483–9.
79. Tudor G, Gutierrez P, Aguilera-Gutierrez A, Sausville EA. Cytotoxicity and apoptosis of benzoquinones: redox cycling, cytochrome release, and BAD protein expression. *Biochem Pharmacol* 2003;65:1061–75.
80. Nemeikaite-Ceniene A, Sarlauskas J, Anusevicius Z, Nivinskas H, Cenas N. Cytotoxicity of RH1 and related aziridinylbenzoquinones: involvement of activation by NAD(P)H:quinone oxidoreductase (NQO1) and oxidative stress. *Arch Biochem Biophys* 2003;416:110–8.
81. Hoey BM, Butler JB, Swallow AJ. Reductive activation of mitomycin C. *Biochemistry* 1988;27:2608–14.

82. Fitzsimmons SA, Workman P, Grever M, Paull K, Camalier R, Lewis AD. Reductase enzyme expression across the national cancer institute tumour cell line panel: correlation with sensitivity to mitomycin C and E09. *J Natl Cancer Inst* 1996;**83**:259–69.
83. Gustafson DL, Beal HD, Bolton EM, Ross D, Waldren CA. Expression of human NAD(P)H: quinone oxoreductase (DT-diaphorase) in chinese hamster ovary cells; effects on the cytotoxicity of antitumour quinones. *Mol Pharmacol* 1996;**50**:728–35.
84. Nishiyama M, Saeki S, Aogi K, Hirabayashi N, Toge T. Relevance of DT-diaphorase activity to mitomycin C efficacy on human cancer cells: differences in vitro and in vivo systems. *Int J Cancer* 1993;**53**:1013–6.
85. Phillips RM, Burger AM, Loadman PM, Jarrett CM, Swaine DJ, Fiebig H-H. Predicting tumour responses to mitomycin C on the basis of DT-diaphorase activity or drug metabolism by tumour homogenates: implications for enzyme-directed bioreductive drug development. *Cancer Res* 2000;**60**:6384–90.
86. Marshall RS, Paterson MC, Rauth AM. DT-diaphorase activity and mitomycin C sensitivity in non-transformed cell strains derived from members of a cancer-prone family. *Biochem Pharmacol* 1991;**12**:1175–80.
87. Beall HD, Murphy AM, Siegel D, Hargreaves RH, Butler J, Ross D. Nicotinamide adenine dinucleotide (Phosphate):quinone oxidoreductase (DT-diaphorase) as a target for bioreductive antitumour quinones; quinone cytotoxicity and selectivity in human lung and breast cancer cell lines. *Mol Pharmacol* 1995;**48**:499–504.
88. Siegel D, Beall H, Kasai M, Atal H, Gibson NW, Ross D. pH-dependent inactivation of DT-diaphorase by mitomycin C and porfiromycin. *Mol Pharmacol* 1993;**44**:1128–34.
89. Robertson N, Stratford IJ, Houlbrook S, Carmichael J, Adams GE. The sensitivity of human tumour cells to quinone bioreductive drugs: what role for DT-diaphorase. *Biochem Pharm* 1992;**44**:409–12.
90. Schellens JHM, Planting AS, van Acker BAC, et al. Phase I and pharmacologic study of the novel indoloquinone bioreductive alkylating cytotoxic drug E09. *J Natl Cancer Inst* 1994;**86**:906–12.
91. Knox RJ, Friedlos F, Jarman M, Roberts JJ. A new cytotoxic DNA interstrand crosslinking agent 5-(aziridin-1-yl)-4-hydroxylamino-2-nitrobenzamide is formed from 5-(aziridin-1-yl)-2,4-dinitrobenzamide (CB1954) by a nitroreductase enzyme in walker carcinoma cells. *Biochem Pharmacol* 1988;**37**:4661–9.
92. Boland MP, Knox RJ, Roberts JJ. The differences in kinetics of rat and human DT-diaphorase result in a differential sensitivity of derived cell lines to CB1954 (5-(aziridin-1-yl)-2,4-dinitrobenzamide). *Biochem Pharmacol* 1991;**41**:867–75.
93. Knox RJ, Jenkins TC, Hobbs SM, Chen S, Melton RG, Burke PJ. Bioactivation of 5-(aziridin-yl)-2,4-dinitrobenzamide (CB1954) by human NAD(P)H:quinone oxidoreductase. 2. A novel co-substrate-mediated antitumour prodrug therapy. *Cancer Res* 2000;**60**:4179–86.
94. Butler J, Zielendziak A, Lea JS, Ward TH, Hoey BM. Contrasting cytotoxic mechanisms of similar antitumour diaziridinylbenzoquinones. *Free Rad Res Comms* 1990;**8**:231–9.
95. Gibson NW, Hartley JA, Butler J, Siegel D, Ross D. Relationship between DT-diaphorase mediated metabolism of a series of aziridinylbenzoquinones and DNA damage and cytotoxicity. *Mol Pharmacol* 1992;**42**:531–6.
96. Haid M, Khandekhar JD, Christ M, et al. Aziridinylbenzoquinone in recurrent, progressive glioma of the central nervous system. A Phase II study by the Illinois Cancer Council. *Cancer* 1985;**56**:1311–5.
97. Case Jr DC, Hayes DM. Phase II study of aziridinylbenzoquinone in refractory lymphoma. *Cancer Treat Rep* 1983;**67**:993–6.
98. Lee CS, Hartley JA, Berardini MD, et al. Alteration in DNA cross-linking and sequence selectivity of a series of aziridinylbenzoquinones after enzymatic reduction by DT-diaphorase. *Biochemistry* 1992;**31**:3019–25.
99. Berardini MD, Souhami RL, Lee CS, Gibson NW, Butler J, Hartley JA. Two structurally related diaziridinylbenzoquinones preferentially cross-link DNA at different sites upon reduction with DT-diaphorase. *Biochemistry* 1993;**32**:3306–12.
100. Naylor MA, Swann E, Everett J, et al. Indolequinone antitumor agents: reductive activation and elimination from (5-methoxy-1-methyl-4,7-dioxindol-3-yl)methyl derivatives and hypoxia-selective cytotoxicity in vitro. *J Med Chem* 1998;**41**:2720–31.
101. Xing C, Skibo EB. Sigmoidal reactions of the aziridinyl semiquinone species. Why aziridinyl benzoquinones are metabolically more stable than aziridinyl indoloquinones. *Biochemistry* 2000;**39**:10770–80.
102. Loadman PM, Phillips RM, Lim LE, Bibby MC. Pharmacological properties of a new aziridinylbenzoquinone, RH1 (2,5-diaziridinyl-3-(hydroxyl)-6-methyl-1,4-benzoquinone) in mice. *Biochem Pharmacol* 2000;**59**:831–7.
103. Winski SL, Hargreaves RH, Butler J, Ross D. A new screening system for NAD(P)H:quinone oxidoreductase (NQO1)-directed antitumour quinones: identification of a new aziridinylbenzoquinone, RH1, as a NQO1-directed antitumour agent. *Clin Cancer Res* 1998;**4**:3083–8.
104. Sausville E. NCI developmental therapeutics program (personal communication).
105. Hollander PM, Ernster L. Studies on the reaction mechanism of DT diaphorase. Action of dead-end inhibitors and effects of phospholipids. *Arch Biochem Biophys* 1975;**169**:560–7.
106. Chen S, Wu K, Zhang D, Sherman M, Knox R, Yang CS. Molecular characterisation of binding of substrates and inhibitors to DT-diaphorase: combined approach involving site-directed mutagenesis, inhibitor-binding analysis and computer modelling. *Mol Pharmacol* 1999;**56**:272–8.
107. Winski SL, Faig M, Bianchet MA, et al. Characterization of a mechanism-based inhibitor of NAD(P)H:quinone oxidoreductase 1 by biochemical, X-ray crystallographic and mass spectrometric approaches. *Biochemistry* 2001;**40**:15135–42.
108. Begleiter A, Leith MK, Curphey TJ. Induction of DT-diaphorase by 1,2-dithiole-3-thione and increase of anticancer activity of bioreductive agents. *Br J Cancer* 1996;**27**:S9–S14.
109. Twerdok LE, Rembish SJ, Trush MA. Induction of quinone reductase and glutathione in bone marrow cells by 1,2-dithiole-3-thione: effect on hydroquinone-induced cytotoxicity. *Toxicol Appl Pharmacol* 1992;**112**:273–81.
110. Li Y, Lafuente A, Trush MA. Characterisation of quinone oxidoreductase, glutathione and glutathione S-transferase in human myeloid cell lines: induction by 1,2-dithiole-3-thione and effects on hydroquinone-induced cytotoxicity. *Life Sci* 1994;**54**:901–16.
111. Doherty GP, Leith MK, Wang X, Curphey TJ, Begleiter A. Induction of DT-diaphorase by 1,2-dithiole-3-thiones in human tumour and normal cells and effect on anti-tumour activity of bioreductive agents. *Br J Cancer* 1998;**77**:1241–52.
112. Begleiter A, Leith MK, Doherty GP, Digby TJ, Pan S. Factors influencing the induction of DT-diaphorase activity by 1,2-

- dithiole-3-thione in human tumour cell lines. *Biochem Pharmacol* 2001;**61**:955–64.
113. Wang X, Doherty GP, Leith MK, Curphey TJ, Begleiter A. Enhanced cytotoxicity of mitomycin C in human tumour cells with inducers of DT-diaphorase. *Br J Cancer* 1999;**80**:1223–30.
114. Carr BA, Franklin MR. Induction of drug metabolising enzymes by 1,7-phenanthroline and oltipraz in mice is unrelated to Ah-responsiveness. *J Biochem Toxicol* 1999;**13**:77–82.
115. Kensler TW, Egner PA, Dolan PM, Groopman JD, Roebuck BD. Mechanism of protection against aflatoxin tumorigenicity in rats fed 5-(2-pyrazinyl)-4-methyl-1,2-dithiol-3-thione (oltipraz) and related 1,2-dithiol-3-thiones and 1,2-dithiol-3-ones. *Cancer Res* 1987;**47**:4271–7.
116. O'Dwyer PJ, Clayton M, Halbherr T, Myers CB, Yao K. Cellular kinetics of induction by oltipraz and its keto derivative of detoxification enzymes in human colon adenocarcinoma cells. *Clin Cancer Res* 1997;**3**:783–91.
117. Jiang Z-Q, Chen C, Yang B, Habbar V, Kong A-NT. Differential responses from seven mammalian cell lines to the treatments of detoxifying enzyme inducers. *Life Sci* 2003;**72**:2243–53.
118. Gordon GB, Prochaska HJ, Yang LY-S. Induction of NAD(P)H:quinone reductase in human peripheral blood lymphocytes. *Carcinogenesis* 1991;**12**:2393–6.
119. Zhang Y, Talalay P, Cho CG, Posner GH. A major inducer of anticarcinogenic protective enzymes from broccoli: isolation and elucidation of structure. *PNAS* 1992;**89**:2399–403.
120. Flescher E, Snyder CA. Aspirin-like drugs can protect human T lymphocytes against benzoquinone cytotoxicity. *Arch Toxicol* 1995;**69**:684–9.
121. Hou D-X, Fukuda M, Johnson JA, Miyamori K, Ushikai M, Fujii M. Fisetin induces transcription of NADPH:quinone oxidoreductase gene through an antioxidant responsive element-involved activation. *Int J Oncol* 2001;**18**:1175–9.
122. Heo YH, Kim S, Park JE, Jeong LS, Lee SK. Induction of quinone reductase activity by stilbene analogs in mouse Hepa 1c1c7 cells. *Arch Pharm Res* 2001;**24**:597–600.
123. Valerio Jr LG, Kepa JK, Pickwell GV, Quattrochi LC. Induction of human NAD(P)H:quinone oxidoreductase (NQO1) gene expression by the flavenol quercetin. *Toxicol Lett* 2001;**119**:49–57.
124. McMahon M, Itoh K, Yamamoto M, et al. The Cap'n'Collar basic leucine zipper transcription factor Nrf2 (NF-E2 p45-related factor2) controls both constitutive and inducible expression of intestinal detoxification and glutathione biosynthetic enzymes. *Cancer Res* 2001;**61**:3299–307.
125. Begleiter A, Leith MK. Induction of DT-diaphorase by doxorubicin and combination therapy with mitomycin C in vitro. *Biochem Pharmacol* 1995;**50**:1281–6.
126. Gonzalez S, Labombarda F, Deniselle MCG, Saravia FE, Roig P, De Nicola AF. Glucocorticoid effects on fos immunoreactivity and NADPH-diaphorase histochemical staining following spinal cord injury. *Brain Res* 2001;**912**:144–53.
127. Yao KS, Clayton M, O'Dwyer PJ. Interaction of heat and hypoxia in modulating transcription of DT diaphorase in human colon adenocarcinoma cells. *Cell Growth Differ* 1994;**5**:125–31.
128. Alvarez S, Boveria A. Induction of antioxidant enzymes and DT-diaphorase in human blood mononuclear cells by light stress. *Arch Biochem Biophys* 1993;**305**:247–51.
129. Kolesar J, Miller J, Duan W, et al. T1 induction in PBMC by mitomycin C predicts response to irinotecan. *Proc ASCO* 2002;**22**:456a.
130. Begleiter A, Leith MK, Curphey TJ, Doherty GP. Induction of DT-diaphorase in cancer chemoprevention and chemotherapy. *Oncol Res* 1997;**9**:371–82.
131. Kepa JK, Ross D. DT-diaphorase activity in NSCLC and SCLC cell lines: a role for fos/jun regulation. *Br J Cancer* 1999;**79**:1679–84.
132. Yu R, Lei W, Mandlekar S, et al. Role of a mitogen-activated protein kinase pathway in the induction of phase II detoxifying enzymes by chemicals. *J Biol Chem* 1999;**274**: 27545–52.
133. Wattenberg LW, Bueding E. Inhibitory effects of 5-(2-pyrazinyl)-4-methyl-1,2-dithiol-3-thione (Oltipraz) on carcinogenesis induced by benzo[a]pyrene, diethylnitrosamine and uracil mustard. *Carcinogenesis* 1986;**7**:1379–81.
134. Egner PA, Kensler TW, Prestera T, et al. Regulation of phase 2 enzyme induction by oltipraz and other dithiol-ethiones. *Carcinogenesis* 1994;**15**:177–81.
135. Szarka CE, Yao KS, Pfeiffer GR, et al. Chronic dosing of oltipraz in people at increased risk for colorectal cancer. *Cancer Detect Prev* 2001;**25**:352–61.
136. Morrow CS, Cowan KH. Mechanisms and clinical significance of multidrug resistance. *Oncology* 1988;**2**:55–63.
137. Mikami K, Naito M, Tomida A, Yamada M, Sirakusa T, Tsuruo T. DT-diaphorase as a critical determinant of sensitivity to mitomycin C in human colon and gastric carcinoma cell lines. *Cancer Res* 1996;**56**:2823–6.
138. Sagara N, Katoh M. Mitomycin C resistance is induced by TCF-3 overexpression in gastric cancer cell line MKN28 is associated with DT-diaphorase down-regulation. *Cancer Res* 2000;**60**:5959–62.
139. Lambert PA, Kang Y, Greaves B, Perry RR. The importance of DT-diaphorase in mitomycin C resistance in human colon cancer cell lines. *J Surg Res* 1998;**80**:177–81.
140. Singh SV, Scalapogna D, Xia H, et al. Biochemical characterisation of a mitomycin C resistant human bladder cancer cell line. *Int J Cancer* 1996;**65**:852–7.
141. Baumann P, Hodnick WF, Scow HA, et al. Reversal of mitomycin C resistance by overexpression of bioreductive enzymes in Chinese hamster ovary cells. *Cancer Res* 2001;**61**:7770–6.
142. O'Dwyer PJ, Perez RP, Yao KS, Godwin AK, Hamilton TC. Increased DT-diaphorase expression and cross-resistance to mitomycin C in a series of cisplatin-resistant human ovarian cancer cell lines. *Biochem Pharmacol* 1996;**52**: 21–7.
143. Wakusawa S, Nalamura S, Miyamoto K. Establishment by adriamycin exposure of multidrug-resistant rat ascites hepatoma AH130 cells showing low DT-diaphorase activity and high cross resistance to mitomycins. *Jpn J Cancer Res* 1997;**88**:88–96.
144. Ward TH, Haran MS, Whittaker D, Watson AJ, Howard TD, Butler J. Cross-resistance studies on two K562 sublines resistant to diaziridinylbenzoquinones. *Biochem Pharmacol* 1995;**50**:459–64.
145. Danson S, Ranson M, Denny O, Jones N, Ward TH. Drug resistance studies on two paired cell lines resistant to the diaziridinylbenzoquinone, RH1. British Oncological Association Annual Meeting, 2003.