

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/7258230>

Albumin cobalt binding and ischemia modified albumin generation: An endogenous response to ischemia?

ARTICLE *in* INTERNATIONAL JOURNAL OF CARDIOLOGY · MAY 2006

Impact Factor: 4.04 · DOI: 10.1016/j.ijcard.2005.03.040 · Source: PubMed

CITATIONS

70

READS

31

3 AUTHORS:



Giuseppe Lippi

University of Verona

1,316 PUBLICATIONS **15,901** CITATIONS

SEE PROFILE



Martina Montagnana

University of Verona

333 PUBLICATIONS **5,218** CITATIONS

SEE PROFILE



Gian Cesare Guidi

University of Verona

419 PUBLICATIONS **5,641** CITATIONS

SEE PROFILE

Letter to the Editor

Albumin cobalt binding and ischemia modified albumin generation:
An endogenous response to ischemia?

Giuseppe Lippi*, Martina Montagnana, Gian Cesare Guidi

*Istituto di Chimica e Microscopia Clinica, Dipartimento di Scienze Morfologico-Biomediche, Università degli Studi di Verona,
Ospedale Policlinico G.B. Rossi, Piazzale Scuro, 10, 37121-Verona, Italy*

Received 8 March 2005; accepted 12 March 2005

Available online 27 April 2005

Keywords: Myocardial ischemia; Acute coronary syndrome; Ischemia modified albumin; Albumin cobalt binding; Hypoxia-induced factor

Myocardial ischemia has become the leading cause of mortality in western countries and this unfavorable trend is unlikely to reverse in the near future. The events progression leading to acute coronary syndrome involves a physiological continuum that traditionally spans from plaque instability to rupture, intracoronary thrombus, reduced blood flow, myocardial ischemia, reversible damage and necrosis [1]. From this biological perspective, myocardial necrosis is time-dependent and occurs when the action of the endogenous mechanisms of response to ischemia is finally overwhelmed. During acute ischemic conditions, the metal binding capacity of albumin for transition metals, like copper, nickel and cobalt is reduced, generating a metabolic variant of the protein, commonly known as ischemia modified albumin (IMA). IMA measurement has recently been proposed as the first US FDA-cleared sensitive marker for the diagnosis of myocardial ischemia in patients presenting with typical acute chest pain [2]. The precise mechanism for IMA generation is yet unknown, though it appears that reactive oxygen species, produced during ischemia, might generate highly reactive hydroxyl free radicals, resulting in site-specific modification to the N-terminus of the albumin moiety, especially at the *N*-Asp-Ala-His-Lys sequence [3,4].

It has been recently reported that cobalt chloride administration might promote delayed cardiac precondition-

ing through selective activation of the HIF-1, AP-1 and iNOS signaling [5]. Cobalt is a relatively rare transition metal with properties similar to iron, chromium and nickel. Cobalt chloride, a water-soluble compound traditionally employed to treat anemia in pregnant women, infants and chronic anemia patients undergoing long-term haemodialysis [6], is a well-established chemical inducer of hypoxia-like responses, such as erythropoiesis and angiogenesis in vivo, likely involving an increased DNA binding activity of hypoxia-inducible factor-1 α (HIF-1 α) to its target genomic sequences. Genes modulated by the HIF-1 α binding include those encoding for angiogenic growth factors (angiopoietin, erythropoietin, vascular endothelial growth factor, insulin growth factor-2, placental growth factor, platelet-derived growth factor β), cytokines (transforming growth factors α and β), survival factors and proapoptotic proteins (RTP801, NIX) [7]. It has been speculated that cobalt might stabilize HIF-1 α through generation of reactive oxygen species (ROS) by a nonenzymatic, nonmitochondrial mechanism. Under normoxic conditions, the main mediator HIF-1 α is rapidly degraded by the proteasome. However, at lower oxygen conditions, HIF-1 α undergoes a stabilization process and induces activation of genetic sequences that promote efficient adaptations to hypoxia. On the basis of the evidence provided by the article of Xi et al. [5], the in vivo generation of IMA might thus be interpreted as an efficient endogenous mechanism of response to ischemia, preventing myocardial damage or limiting the extent of myocyte necrosis. The decreased albumin affinity for transition metals, as it occurs during acute ischemia, generates IMA and increases the concentration of biological active free

* Corresponding author. Tel.: +39 045 8074516; fax: +39 045 8201889.*E-mail addresses:* ulippi@tin.it, giuseppe.lippi@mail.univr.it
(G. Lippi).

cobalt in plasma, which might finally exert a beneficial effect, limiting the extent of the ischemic injury, by modulating the HIF pathway. In this perspective, we believe that this evidence might provide a reliable explanation for the in vivo generation of IMA during acute ischemia.

References

- [1] Schachter M. The pathogenesis of atherosclerosis. *Int J Cardiol* 1997;62(Suppl 2):S3.
- [2] Roy D, Quiles J, Aldama G, Sinha M, Avanzas P, Arroyo-Espliguero R, et al. Ischemia modified albumin for the assessment of patients presenting to the emergency department with acute chest pain but normal or non-diagnostic 12-lead electrocardiograms and negative cardiac troponin T. *Int J Cardiol* 2004;97:297–330.
- [3] Marx G, Chevion M. Site-specific modification of albumin by free radicals. *Biochem J* 1985;236:397–400.
- [4] Gidenne S, Ceppa F, Fontan E, Perrier F, Burnat P. Analytical performance of the Albumin Cobalt Binding (ACB) test on the cobas MIRA plus analyzer. *Clin Chem Lab Med* 2004;42:455–61.
- [5] Xi L, Tahler M, Yin C, Salloum F, Kukreja RC. Cobalt chloride induces delayed cardiac preconditioning in mice through selective activation of HIF-1 α and AP-1 and iNOS signaling. *Am J Physiol Heart Circ Physiol* 2004;287:H2369–75.
- [6] Barceloux DG. Cobalt. *Toxicol Clin Toxicol* 1999;37:201–6.
- [7] Dery MA, Michaud MD, Richard DE. Hypoxia-inducible factor-1: regulation by hypoxic and non-hypoxic activators. *Int J Biochem Cell Biol* 2005;37:535–40.