

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/7418420>

# Niles, J.C., Wishnok, J.S. & Tannenbaum, S.R. Peroxynitrite-induced oxidation and nitration products of guanine and 8-oxoguanine: structures and mechanisms of product formation. Ni...

ARTICLE *in* NITRIC OXIDE · APRIL 2006

Impact Factor: 3.52 · DOI: 10.1016/j.niox.2005.11.001 · Source: PubMed

---

CITATIONS

119

---

READS

55

3 AUTHORS, INCLUDING:



Jacquín C Niles

Massachusetts Institute of Technology

32 PUBLICATIONS 1,379 CITATIONS

SEE PROFILE



John S Wishnok

Massachusetts Institute of Technology

169 PUBLICATIONS 15,827 CITATIONS

SEE PROFILE

# Peroxynitrite-induced oxidation and nitration products of guanine and 8-oxoguanine: Structures and mechanisms of product formation <sup>☆</sup>

Jacquin C. Niles <sup>a</sup>, John S. Wishnok <sup>a</sup>, Steven R. Tannenbaum <sup>a,b,\*</sup>

<sup>a</sup> *Biological Engineering Division, Massachusetts Institute of Technology, Cambridge, MA 02139, USA*

<sup>b</sup> *Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA 02139, USA*

Received 8 July 2005; revised 3 November 2005

Available online 13 December 2005

## Abstract

Peroxynitrite induces DNA base damage predominantly at guanine (**G**) and 8-oxoguanine (**8-oxoG**) nucleobases via oxidation reactions. Nitration products are also observed, consistent with the generation of radical intermediates that can recombine with the  $\cdot\text{NO}_2$  formed during peroxynitrite degradation. The neutral **G** radical, **G** $\cdot$ , reacts with  $\cdot\text{NO}_2$  to yield 8-nitroguanine (8-nitroG) and 5-nitro-4-guanidinohydantoin (**NI**), while for **8-oxoG** we have proposed a reactive guanidinyldene radical intermediate. The products generated during peroxynitrite-mediated 8-oxoG oxidation depend on oxidant flux, with dehydroguanidinohydantoin (**DGh**), 2,4,6-trioxo-[1,3,5]triazinane-1-carboxamide (**CAC**) and **NO<sub>2</sub>-DGh** predominating at high fluxes and spiroiminodihydantoin (**Sp**), guanidinohydantoin (**Gh**) and 4-hydroxy-2,5-dioxo-imidazolidine-4-carboxylic acid (**HICA**) predominating at low fluxes. Both product sets are observed at intermediate fluxes. It is therefore important in model systems to ensure that the relative concentrations are well controlled to minimize competing reactions that may not be relevant in vivo. Increasingly sophisticated systems for modeling peroxynitrite production in vivo are being developed and these should help with predicting the products most likely to be formed in vivo. Together with the emerging information on the genotoxic and mutational characteristics of the individual oxidation products, it may be found that the extent of tissue damage, mutational spectra and, hence, cancer risk may change as a function of peroxynitrite fluxes as different product combinations predominate.

© 2005 Elsevier Inc. All rights reserved.

**Keywords:** Peroxynitrite; Guanine; 8-Oxoguanine; 8-Nitroguanine; Dehydroguanidinohydantoin; 2,4,6-Trioxo- [1,3,5]triazinane-1-carboxamide; Spiroiminodihydantoin; 4-Hydroxy-2,5-dioxo-imidazolidine-4-carboxylic acid; Nitric oxide; DNA damage; Oxidative damage

Nitric oxide ( $\cdot\text{NO}$ ) plays a pivotal role in many biological processes, and is produced primarily by the enzyme nitric oxide synthase (NOS), of which there are three isoforms, namely, endothelial (eNOS), neuronal (nNOS), and inducible (iNOS). The expression, regulation, synthesis, and role of NOS and  $\cdot\text{NO}$  in physiologic processes such as vasodilation, neurotransmission, inhibition of platelet aggregation, and in proper immunologic function, have been extensively studied and previously reviewed [1–8].  $\cdot\text{NO}$

plays a dual role in human physiology: both eNOS and nNOS produce relatively low  $\cdot\text{NO}$  levels which modulate processes such as vasodilation and platelet aggregation in the cardiovascular system, and long term potentiation and neurotransmission in the nervous system. In the immune system,  $\cdot\text{NO}$  is produced by iNOS in higher amounts and is used by macrophages as a cytotoxic agent against engulfed pathogens. More recently, nitric oxide has been shown also to arise under certain conditions via reduction of nitrite by deoxygenated hemoglobin [9,10].

Since  $\cdot\text{NO}$  is toxic at levels higher than normal physiological levels, its production must be tightly controlled both spatially and temporally to minimize undesired effects. This balance, however, can be disrupted in certain pathophysiological states. During chronic gastrointestinal inflammatory

<sup>☆</sup> Peroxynitrite is used to denote the sum of the conjugate acid (HOONO) and base (ONOO<sup>−</sup>) of oxoperoxonitrate (1<sup>−</sup>). Specific reference to either species is indicated by use of HOONO or ONOO<sup>−</sup>.

\* Corresponding author. Fax: +1 617 252 1787.

E-mail address: [srt@mit.edu](mailto:srt@mit.edu) (S.R. Tannenbaum).

conditions, such as the inflammatory bowel diseases [11–13], and chronic *Helicobacter pylori* infection [14,15], exuberant NO production by activated macrophages is believed to be an important tissue-damage mediator, and to potentiate the increased cancer risk associated with these conditions. In fact, significant increases in the NO-derived 8-nitroguanine (**8-NO<sub>2</sub>G**) and 8-oxoguanine (**8-oxoG**) products (vide infra), have been demonstrated in gastric tissue from *H. pylori* infected individuals, and these mutagenic lesions may play an important role in *H. pylori* related carcinogenesis. Thus, understanding the connection between NO and these outcomes requires consideration of the fate of NO in biological systems, and the capacity of the associated intermediates to react with various cellular components, especially DNA, since damage to this informational molecule can lead to mutations and ultimately carcinogenesis.

NO itself is relatively unreactive toward DNA, but it can be converted into several highly reactive species. These include nitrous anhydride (N<sub>2</sub>O<sub>3</sub>), nitrogen dioxide (NO<sub>2</sub>), nitryl chloride (NO<sub>2</sub>Cl), and peroxynitrite (ONOO<sup>−</sup>),<sup>1</sup> and the chemistry underlying the formation and reactivity of these species has been recently reviewed [16]. In particular, our group has been interested in understanding the contribution of peroxynitrite-induced DNA damage to the cytotoxicity and carcinogenesis associated with pathological states in which NO is overproduced. Toward this objective, we have been characterizing the peroxynitrite-induced DNA-damage products and, through collaborative efforts, determining how they interact with the replication and repair machinery in vitro and in vivo to establish their individual genotoxic and mutagenic profiles. We have also been supplementing these biological studies with experiments aimed at understanding the mechanism by which the various products form, and have gained some key chemical insights that are discussed later. Of particular relevance toward connecting the chemistry of formation with the biological significance of the products, we have established that both product distribution and identity are dramatically affected by peroxynitrite fluxes. Specifically, two distinct product sets dominate at low and high peroxynitrite fluxes, respectively, raising the intriguing possibility that peroxynitrite may induce significantly different mutational spectra and genotoxic profiles at different production rates in chronically inflamed tissues.

This review summarizes the progress in identifying the major peroxynitrite-induced DNA-damage products, with emphasis on the nucleobases guanine (**G**) and 8-oxoguanine (**8-oxoG**). The quantitative aspects related to these substances are also of considerable interest [17], but treatment of these is beyond the scope of this review. The chemical insights gained from our studies with peroxynitrite,

together with those of other groups using different systems to study oxidation of these bases, have been synthesized to highlight how seemingly-different oxidants ultimately result in similar or identical intermediates and final products. Finally, some biological aspects of these reaction products are also discussed.

### A brief review of peroxynitrite decomposition chemistry

Peroxynitrite chemistry is complex due to dependence on a combination of pH and carbon dioxide concentrations. In the absence of CO<sub>2</sub>, peroxynitrite will decay ( $t_{1/2} \sim 1$  s) via an acid-catalyzed pathway. At physiologic pH, both ONOO<sup>−</sup> and its conjugate acid, ONOOH ( $pK_a = 6.8$ ) exist, and the latter decomposes via homolysis to give the hydroxyl (HO<sup>•</sup>) and nitrogen dioxide (NO<sub>2</sub>) free radicals, each in approximately 25–28% yield [18–20]. In the presence of physiologic CO<sub>2</sub> levels ( $\sim 1$ – $1.3$  mM), decay occurs almost exclusively via a CO<sub>2</sub>-catalyzed pathway. ONOO<sup>−</sup> reacts with CO<sub>2</sub> ( $k_2 = 3$ – $5 \times 10^4$  M<sup>−1</sup> s<sup>−1</sup>) to give peroxy-nitrosocarbonate, which undergoes homolysis to give the CO<sub>3</sub><sup>•−</sup> and NO<sub>2</sub> free radicals, each in  $\sim 30$ – $33\%$  yields [21–25]. Peroxynitrite reactivity with various substrates can be induced by these radical species, but both ONOO<sup>−</sup> and ONOOH are reactive in their own right. ONOO<sup>−</sup> is known to act as a nucleophile in reactions with CO<sub>2</sub> [23,24], aldehydes [26] and ketones [27], while ONOOH mediates ebselen oxidation [28]. Overall, in solutions buffered at physiologic pH and containing CO<sub>2</sub>, the major reactive species are expected to be CO<sub>3</sub><sup>•−</sup>, NO<sub>2</sub>, and ONOO<sup>−</sup>, along with minor amounts of ONOOH and HO<sup>•</sup>. Together, these reactive intermediates determine the spectrum of oxidation products observed when peroxynitrite reacts with DNA.

### Peroxynitrite-induced DNA oxidation products

Peroxynitrite induces both sugar-phosphate backbone damage and base damage. Direct backbone damage at reasonably low peroxynitrite concentrations is manifested as single strand breaks ([29,30]; for detailed reviews, see [31,32]), and H-atom abstraction from the deoxyribose moiety by the peroxynitrite-derived radicals outlined above is believed to be the initiating event [31]. The formation of base propenals when deoxynucleotides are treated with peroxynitrite supports this hypothesis [33]. Interestingly, CO<sub>2</sub> dramatically suppresses peroxynitrite-induced sugar damage, both at the nucleoside and DNA levels [33,34]. In DNA, while the total damage (sugar and base) remains unchanged, independent of CO<sub>2</sub>, proportionately more base damage occurs in the presence of CO<sub>2</sub> [34].

In this review, we focus exclusively on base damage, and specifically on **G** and its oxidation product, **8-oxoG**, since these are the most reactive toward peroxynitrite [35], and possibly the major contributors to peroxynitrite-derived genotoxic and mutagenic lesions.

<sup>1</sup> Abbreviations used: SIN-1, 3-morpholinosydnonimine; MnTMPyP, manganese (III) 5,10,15,20-tetrakis(*N*-methylpyridyl)porphyrin; **Iz**, 2,5-diamino-4H-imidazol-4-one; **Oz**, 2,2,4-triamino-5-(2H)-oxazolone; **HICA**, 4-hydroxy-2,5-dioxo-imidazolidine-4-carboxylic acid.

### Peroxynitrite-mediated guanine oxidation

**G** has the lowest reduction potential of the four normal DNA bases ( $E^\circ = 1.29$  V [36]), and its reaction with the peroxynitrite-derived  $\text{HO}^\cdot$  ( $E^\circ = 1.9\text{--}2.1$  V [37]) and  $\text{CO}_3^{\cdot-}$  ( $E^\circ = 1.5$  V [38]) radicals peroxynitrite is thermodynamically favorable. Furthermore, the bimolecular rate constants for reaction of these radicals with **G** ( $7.8 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  for  $\text{HO}^\cdot$  [39]) and  $7 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$  for  $\text{CO}_3^{\cdot-}$  [40] dictate extremely fast kinetics for the initial oxidation event. The **G** radical cation, **G** $^+$ , is initially produced and subsequently converted into several nitration and oxidation products that have been structurally characterized by us and other groups, as summarized in Fig. 1.

### Nitration products

The first nitration product to be identified was 8-nitro-guanine, **8-NO<sub>2</sub>G** [41]. This modified base, when formed in the context of oligonucleotides [34] or double stranded DNA [42], depurinates with a half-life on the order of hours under simulated physiologic conditions. Thus, **8-NO<sub>2</sub>G** formation in vivo may be an important source of apurinic sites arising from peroxynitrite production.

Later, we identified 5-nitro-4-guanidinohydantoin, **NI**, as a novel peroxynitrite-induced **G** nitration product [43]. Unlike **8-NO<sub>2</sub>G**, this compound has a stable glycosidic bond, and persists either as the free nucleoside or nucleotide in DNA. **NI** represents the only peroxynitrite-induced guanine nitration product characterized to date in which the pyrimidine ring has been disrupted.

There is a single report of a peroxynitrite addition product, 4,5-dihydro-5-hydroxy-4-(nitrosooxy)guanine [44]. Apparently, this compound does not spontaneously depurinate, and appears to be stable despite the favorable elimination of nitrous acid to extend the  $\pi$ -conjugation in the pyrimidine ring. Furthermore, several C8-substituted 5-hydroxypurines are known to be hydrolytically unstable, readily undergoing hydration at the C6 position to yield hydantoin products [45–48]. The extent to which the C8-substitution impacts the rate of this reaction has not been

fully explored, but the above issues need to be reconciled before acceptance of this proposed structure.

### Oxidation products

Two main oxidation products are formed during the peroxynitrite-induced oxidation of **G**, namely, **Iz** [49] and **8-oxoG** [30]. **Iz**, which had previously been identified as a product of Type I **G** photooxidation, undergoes hydrolysis to **Oz** at physiologic pH [50]. Thus, in tissues under oxidative stress in which **Iz** is initially formed, **Oz** is the product that is expected to accumulate.

The proposed formation of **8-oxoG** from the reaction of **G** with peroxynitrite was initially controversial, with some investigators using fairly high ratios of peroxynitrite to **G** claiming no formation of **8-oxoG** [42,49] and others using lower ratios reporting its formation [30]. This apparent discrepancy was finally resolved when Pryor and co-workers demonstrated that peroxynitrite reacted preferentially with **8-oxoG** over **G** [51]. When relatively high concentrations of peroxynitrite are used, the resulting **8-oxoG** formed is rapidly consumed in subsequent reactions with excess peroxynitrite. Thus, a compound previously identified as 4-hydroxy-7,8-dihydro-8-oxoG was isolated from the reaction of **G** with peroxynitrite [49,52]. However, further work by our group and others has since established that, first, this product is in fact spiroiminodihydantoin, **Sp** [47,53,54] and, second, that it most likely arises from further oxidation of **8-oxoG**.

### Peroxynitrite-mediated 8-oxoG oxidation

In addition to the inherently interesting chemistry underlying peroxynitrite-mediated **G** oxidation, we have also been interested in elucidating the potential biological importance of these reactions. Several findings coincided to influence our decision to focus on the reaction between peroxynitrite and **8-oxoG** as a potentially important source of biologically relevant DNA lesions.

First, **8-oxoG** has been detected, in amounts ranging from 4 to 11 in  $10^7$  bases, in genomic DNA isolated from tissues not under chronically elevated levels of oxidative stress [55]. Reliable measurements of levels in chronically inflamed tissues are lacking, but presumably these will be higher than the levels mentioned above, thus making **8-oxoG** an even more likely target for oxidative damage in DNA. Second, **8-oxoG** is inherently more reactive than **G** with peroxynitrite, and in a study by Pryor's group a  $10^3$ -fold excess of **G** over **8-oxoG** protected only half of the **8-oxoG** from reacting with peroxynitrite [51]. Thermodynamically, **8-oxoG** ( $E^\circ = 0.74$  V [56]) is more susceptible to oxidation than **G**. Recent studies have also highlighted that the peroxynitrite-derived radicals will rapidly react with **8-oxoG**, and bimolecular rate constants of  $8 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$  [40] and  $5 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$  [57] have been reported for  $\text{CO}_3^{\cdot-}$  and  $\cdot\text{NO}_2$ , respectively. Furthermore, direct electron transfer from **8-oxoG** to the oxidizing **G** $^\cdot$  readily occurs with a bimolecular rate constant of




Fig. 1. Summary of the peroxynitrite-induced **G** oxidation and nitration products.


$4.6 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$  [56]. These observations indicate that even with **G** in large excess over **8-oxoG**, oxidation of the latter can still dominate because of (i) the inherent thermodynamic and kinetic favorability of the reaction of **8-oxoG** with peroxynitrite, and (ii) the high propensity for damage initiated on **G** to be transferred to **8-oxoG**. Third, the ability of double stranded DNA to support charge transfer over long distances to sites remote from the initial oxidation site has become well established [58,59], and ultimately sites with the lowest reduction potentials serve as sinks for the migrating positive hole. Thus, in oligonucleotides containing **8-oxoG** in addition to the other normal bases, oxidation occurs preferentially at the **8-oxoG** even when the initial oxidizing event is well removed from this residue [60]. Taken all together, these facts suggest that oxidative damage may preferentially accumulate at **8-oxoG** residues in DNA.

#### Studying **8-oxoG** oxidation using authentic peroxynitrite introduced by bolus addition

In early experiments, bolus addition of preformed peroxynitrite to solutions containing acetylated 8-oxo-(2'-deoxy)guanosine was used as a model system to characterize several reaction products, shown in Fig. 2. The nature of the sugar moiety did not affect the identity of the nucleobase oxidation products, and so both 3',5'-di-*O*-acetyl- $\beta$ -D-erythropentofuranosyl- and 2',3',5'-tri-*O*-acetyl- $\beta$ -D-erythropentofuranosyl-**8-oxoG** derivatives were used. Without acetylation, the reaction products are extremely polar, so this modification proved useful by improving their chromatographic properties, and facilitated purification by standard reversed-phase HPLC.

Although the structure of the major product formed over a range of peroxynitrite concentrations was initially assigned tentatively [61], subsequent detailed NMR study of this compound produced by MnTMPyP/KHSO<sub>5</sub>-mediated oxidation of d(GpT) established it as dehydroguanidinohydantoin, **DGh** [62]. In buffered phosphate/bicarbonate solutions (pH 7.4) it readily hydrolyzes to give oxaluric acid (**OA**) as a relatively stable product [61,62]. Interestingly, in a more physiologic buffer containing  $\text{Mg}^{2+}$  and bicarbonate, **OA** decomposes to give the urea (**UA**) [63]. This reaction occurs in the context of both nucleoside and oligonucleotide, and suggests that the initial **DGh** lesion can give rise in vivo to several further hydrolysis products that may be of biological significance.

The second most abundant product was identified as 2,4,6-trioxo-[1,3,5]triazinane-1-carboxamide (**CAC**). This is identical to the major singlet oxygen-derived **8-oxoG** oxidation product [64], which will hydrolyze to cyanuric acid (**CA**) [61,64]. There appear to be two routes to **CAC**, one involving an unstable, uncharacterized intermediate (**CAP**) and the other the further oxidation of **DGh** by peroxynitrite. Interestingly,  $\text{CO}_2$ , which is known to alter the reactive species produced during peroxynitrite decomposition [23–25], has a profound impact on the product profile with respect to **DGh** and **CAC** formation. When reactions are carried out in the absence of  $\text{CO}_2$ , **DGh** is the major product up to  $\sim 1 \text{ mM}$  peroxynitrite, above which **CAC** becomes the major product. We have shown that further oxidation of **DGh** to **CAC** is at least one factor accounting for this phenomenon [61]. In the presence of  $\text{CO}_2$ , however, **DGh** remains the major product up to  $2 \text{ mM}$  peroxynitrite, suggesting that the oxidants produced under these conditions





do not as effectively yield **CAC** (J.C. Niles, J.S. Wishnok, S.R. Tannenbaum, unpublished results).<sup>2</sup>

Two relatively minor products identified in this reaction are parabanic acid (**PA**) and *N*-nitro-dehydroguanosine (NO<sub>2</sub>-**DGh**) [61,65]. The former is likely an intermediate in the hydrolysis of **DGh** to **OA**, but a direct route to its formation cannot be excluded. NO<sub>2</sub>-**DGh** is the only peroxynitrite-induced **8-oxoG**-derived nitration product identified to date, and provides some important insights into the mechanism via which the various reaction products are formed. Like **DGh**, both **PA** and NO<sub>2</sub>-**DGh** are hydrolytically unstable and will decompose to **OA**.

The final oxidation product of this reaction is **Iz**, the same product that is formed during peroxynitrite-induced **G** oxidation [49]. It is therefore tempting to conclude that **Iz** formed during the reaction with **G** is arising from secondary oxidation of **8-oxoG**. However, there is a distinct difference in the pH-dependence of **Iz** formation when starting from **G** versus **8-oxoG**. With **G**, **Iz** formation is prominent at pH 7.2, while with **8-oxoG**, **Iz** formation becomes prominent only at pH > 7.5, and increases up to pH 8.5 (J.C. Niles, J.S. Wishnok, S.R. Tannenbaum, unpublished results). Therefore, in the oxidation of **G**, any contribution to **Iz** formation from secondary **8-oxoG** oxidation at pH 7.2 must be minor, indicating that there are at least two paths to **Iz** that are, respectively, **8-oxoG** dependent and independent.

Additionally, it is important to note that **DGh** is not a precursor to **Iz**, as was suggested in earlier literature [66,67]. It is possible under basic conditions in the presence of ammonium hydroxide to convert **DGh** into the *base moiety* of **Iz**, but not the intact nucleoside [61]. Taken together with the fact that **DGh** hydrolyzes exclusively to **OA** [61,62,68] and not **Iz** at near neutral pH, the most logical conclusion is that **DGh** and **Iz** represent distinct fates of an earlier reaction intermediate(s).

In an effort to gain further insights into the mechanism underlying product formation, we attempted reductive trapping of reaction intermediates. Using thiol reductants (glutathione and cysteine), in addition to the expected decrease in overall product yield, a new compound identified as **Sp** was formed [53]. While this strategy failed to trap compounds that could be unambiguously assigned as product precursors, it served together with other work [47,54] to establish that the oxidation product previously identified as 4-hydroxy-7,8-dihydro-8-oxoG [52] is in fact **Sp**.

#### *Studying 8-oxoG oxidation using infused authentic and in situ generated peroxynitrite*

While bolus addition of preformed peroxynitrite to reaction mixtures served as a convenient initial model, it clearly does not simulate peroxynitrite formation in vivo. Thus, the possibility that the chemistry and hence the products identified during bolus addition experiments

may not accurately reflect the in vivo situation could not be excluded. To test this, we reduced the instantaneous peroxynitrite concentrations in reaction mixtures using two strategies, namely: (i) infusion of preformed peroxynitrite and (ii) in situ peroxynitrite generation using SIN-1, a compound that decomposes in aqueous solution to produce stoichiometric amounts of O<sub>2</sub><sup>•−</sup> and NO that combine to produce peroxynitrite at very low fluxes [69–71]. The products formed in these studies are also summarized in Fig. 2.

When peroxynitrite was infused into buffered solutions containing **8-oxoG**, three additional compounds, i.e., **Sp**, guanosine (Gh) and **HICA** became quantitatively important [48,71]. Delivering the same total amount of peroxynitrite over increasing time periods led to **Sp/Gh** becoming the major reaction product [48]. It is important to note that while the **Sp** formed is structurally identical to that isolated during our attempted thiol trapping experiments [53], it arises here via a distinct mechanism, and this is discussed later. Additionally, an inverse, pH-dependent relationship exists between **Sp** and **Gh** formation, with **Sp** being produced predominately at neutral pH and **Gh** being produced predominantly at low pH [47,48].

Interestingly, using SIN-1 to generate lower peroxynitrite fluxes than achievable by infusion leads predominantly to **Sp/Gh** and **HICA** formation and the complete absence of bolus addition products [53]. Thus, two distinctly different product sets are obtained depending on whether peroxynitrite is added quickly to **8-oxoG** at high (bolus addition) or slowly at low (SIN-1) fluxes. At intermediate fluxes (infusion) these two pathways compete, yielding a combination of products. These intriguing findings motivated us to explore in greater detail the mechanism(s) of product formation during the peroxynitrite-mediated **G** and **8-oxoG** oxidation, and our studies, together with work from several other groups, have provided additional insights into these complex reactions.

### **Mechanistic insights into peroxynitrite-induced oxidation of guanine and 8-oxoguanine**

#### *Guanine oxidation*

Oxidation of **G** has been extensively studied, and it is well established that the initial event, using a variety of oxidant systems, is a one-electron oxidation to produce the **G** radical cation, (**G**<sup>•+</sup>) [39]. As previously mentioned, this is carried out efficiently by the peroxynitrite-derived HO<sup>•</sup> and CO<sub>3</sub><sup>•−</sup> radicals. NO<sub>2</sub> (*E*<sup>o</sup> = 1.04 V [37]) the other peroxynitrite-derived radical, however, has been shown experimentally not to induce **G** oxidation [57]. Thus, the initial step in the reaction between peroxynitrite and **G** is one-electron oxidation of **G** to **G**<sup>•+</sup> by either HO<sup>•</sup> or CO<sub>3</sub><sup>•−</sup>. In our experiments, ~1–1.3 mM CO<sub>2</sub> was used, making CO<sub>3</sub><sup>•−</sup> the quantitatively more important initiating radical.

Once formed, **G**<sup>•+</sup> (p*K*<sub>a</sub> = 3.9) deprotonates at physiological pH to give the neutral radical, **G**<sup>•</sup> [39], which is

<sup>2</sup> **CAP** is formed under these conditions, but we are unable to accurately quantitate it. Therefore, it has not been included in the total **CAC** formed.




Fig. 3. Proposed mechanism for the peroxynitrite-induced formation of G-derived products.

expected to have significant unpaired electron density at O6, C5, and C8 [72]. Formation of 8-NO<sub>2</sub>G and NI can be mechanistically rationalized by radical recombination of <sup>•</sup>NO<sub>2</sub> at C8 and C5 of G<sup>•</sup>, respectively (Fig. 3) [43,73]. For 8-NO<sub>2</sub>G, the aromatic purine ring is restored by deprotonation at C8, while for NI, the purine ring is disrupted to give the aromatic imidazole nucleus [43]. Supporting this proposal is the recent demonstration that G<sup>•</sup> generated independently of peroxynitrite will react with <sup>•</sup>NO<sub>2</sub> to give both 8-NO<sub>2</sub>G and NI [73,74].

The mechanism of Iz formation is an intriguing aspect of G oxidation chemistry, since the same product forms via a variety of oxidants, including HO<sup>•</sup> [50], MnTMPyP/KHSO<sub>5</sub> [67], and peroxynitrite [43,49], yet the source of the exogenous [O] atom incorporated at C4 varies among these oxidants. In HO<sup>•</sup>-mediated oxidations, molecular oxygen provides the C4 [O] atom [50], while for MnTMPyP/KHSO<sub>5</sub> and peroxynitrite, HSO<sub>5</sub><sup>-</sup> [67] and ONOO<sup>-</sup> (J.C. Niles, J.S. Wishnok, S.R. Tannenbaum, unpublished results) are the sources, respectively. These findings highlight two important facts. First, there are multiple, indepen-

dent paths leading to formation of common intermediates that ultimately converge on the same final product(s). This is true for both G and 8-oxoG oxidation, and applies to all the purely oxidative products. Second, involvement of HSO<sub>5</sub><sup>-</sup> and ONOO<sup>-</sup> in direct nucleophilic reactions suggests that electrophilic intermediates are produced during the reaction.

We propose that the C5-alkoxide radical intermediate, 2, is important in Iz formation. With peroxynitrite, this intermediate is generated when ONOO<sup>-</sup> attacks an electrophilic intermediate, which is likely to be the guanine cation, G<sup>+</sup> [67] at C5. The weak peroxo bond in the resulting ROONO species, 1, will undergo homolysis to give the C5-alkoxide radical and <sup>•</sup>NO<sub>2</sub>. Indeed, precedent exists for the favorable peroxo bond homolysis in the analogous HOONO [15,16] and ONOOCO<sub>2</sub><sup>-</sup> [75,76] species. Pyrimidine ring opening of the C5-alkoxide is proposed to lead to formation of a putative guanidinyldene radical intermediate, 3, which is in equilibrium with 4. H-atom abstraction by the latter, followed by hydrolysis, then produces formamide [67] and Iz.

The guanidinyldene radical has been proposed as a reactive intermediate to rationalize a critical deficiency in the mechanisms proposed to explain **Iz** formation. First, as outlined earlier and contrary to previously proposed mechanisms, **DGh** is not a direct precursor to **Iz**. Second, despite the fact that the guanidino functional group is a poor nucleophile, requiring elevated temperatures, alkaline pH ( $pK_a \sim 12$ ) and prolonged reaction times (hours to days) for reasonable reaction yields, this chemistry is invoked to account for the rapid formation of **Iz** on the timescale of seconds (for peroxynitrite) to minutes (singlet oxygen, MnTMPyP/KHSO<sub>5</sub>, and riboflavin) at ambient temperatures and pH  $\sim 7$ . This short timescale for **Iz** formation indicates the involvement of an intermediate with very high reactivity, and the guanidinyldene radical, which is expected to have reactivity more comparable to that of aminyl, iminyl, and amidyl radicals [77,78], is a putative candidate. While there is no direct evidence to date for this intermediate, **NO<sub>2</sub>-DGh** formation during the oxidation of **8-oxoG** suggests its existence during these reactions (vide infra).

We have also determined that peroxynitrite is the major source of the oxygen atom incorporated into **8-oxoG** during the peroxynitrite-mediated oxidation of **G** (J.C. Niles, J.S. Wishnok, S.R. Tannenbaum, unpublished results). **8-OxoG** was not directly detected under our reaction conditions, so the **8-oxoG**-derived product **CAC** was used as a surrogate marker. We established that **CAC** derived from the reaction of <sup>18</sup>O-labeled peroxynitrite with unlabeled **G** and **8-oxoG** contained three and two peroxynitrite-derived [O] atoms, respectively, indicating that the extra label in the **G**-derived product must have been present in the precursor **8-oxoG**. In rationalizing this fact mechanistically, we have proposed that while ONOO<sup>−</sup> reacts with **G**<sup>+</sup> at C5 to give **Iz**, reaction at C8 gives the adduct, **5**, that decomposes via **6** and **7** into **8-oxoG**. Alternatively, elimination of nitrous acid from **5** can directly yield **8-oxoG**. Here again, for the same final product, the pattern of exogenous [O] atom incorporation is dependent on the oxidizing system used. Thus, for riboflavin- (Type I) and singlet oxygen- (Type II) mediated photooxidation, H<sub>2</sub>O [79] and O<sub>2</sub> [52] are the sources of this [O] atom, respectively. Furthermore, these findings—along with the formation of **8-NO<sub>2</sub>G** and **NI**—add experimental support to C5 and C8 being primary sites of unpaired electron density and/or electrophilicity.

### 8-Oxoguanine oxidation

As with **G**, the initial step in **8-oxoG** oxidation is the one-electron oxidation by the peroxynitrite-derived radicals to produce the radical cation, **8-oxoG**<sup>•+</sup> (Fig. 4) [40,56,57]. This reaction is both thermodynamically and kinetically more favorable than **G** oxidation, as discussed earlier. The  $pK_a$  of **8-oxoG**<sup>•+</sup> is 6.6 [56], and thus, at physiologic pH, both the neutral radical and radical cation exist, making it difficult to readily distinguish the reacting species. In addition to the **8-oxoG**<sup>•+</sup>, electrochemical studies indicate that oxidation to

a reactive quinonoid diimine, **8-oxoG**<sup>ox</sup> intermediate readily occurs [45,80]. We have included this species as a key reactive intermediate in our proposed model of peroxynitrite-induced **8-oxoG** oxidation. However, we cannot exclude the possibility that **8-oxoG**/**8-oxoG**<sup>•+</sup> are relevant reactive intermediates in the chemistry proposed below.

For the reaction of **8-oxoG** with peroxynitrite, it is helpful to consider the mechanistic details in the context of high and low peroxynitrite fluxes. The products formed at high fluxes, namely, **DGh**, **NO<sub>2</sub>-DGh**, and **CAC**, all incorporate peroxynitrite-derived [O] atoms, while those formed at low fluxes, namely, **Sp**, **Gh**, and **HICA**, all incorporate H<sub>2</sub>O-derived [O] atoms [48]. Based on these observations, we have proposed that nucleophiles in solution compete for reaction with **8-oxoG**<sup>ox</sup>. At high and low peroxynitrite fluxes, respectively, reaction of **8-oxoG**<sup>ox</sup> with ONOO<sup>−</sup> and H<sub>2</sub>O will occur exclusively. It is only when peroxynitrite is infused that both pathways directly compete, leading to formation of both product sets.

### Mechanism of high flux peroxynitrite product formation

In our model, at sufficiently high peroxynitrite concentrations ONOO<sup>−</sup> reacts with **8-oxoG**<sup>ox</sup> at C5 to give **8**, which rapidly undergoes homolytic peroxo bond cleavage to give the C5-alkoxide, **9**. Just as proposed for the analogous intermediate formed during **G** oxidation, pyrimidine ring opening leads to the guanidinyldene radical intermediate, **10**, which has multiple fates, namely: (i) H-abstraction to yield **DGh**, the major reaction product; (ii) combination with **NO<sub>2</sub>** to give **NO<sub>2</sub>-DGh** and, (iii) cyclization via **11** and **12** to give **Iz**. As before, the guanidinyldene radical is proposed as an intermediate to rationalize the formation of several key products. However, the intriguing challenge—of directly observing or specifically trapping this intermediate in both **G** and **8-oxoG** oxidation reactions—remains.

Another issue for further study is the mechanism of **CAC** formation. As mentioned earlier, there are at least two pathways to this product, namely: (i) an indirect route involving further oxidation of **DGh** [61], and (ii) a “direct” route involving formation of a hydrolytically unstable intermediate, **CAP**, that has eluded detailed structural characterization (J.C. Niles, J.S. Wishnok, S.R. Tannenbaum, unpublished results). Based upon a putative [M-H]<sup>−</sup> ion produced in ESI-MS along with the fact that it hydrolyzes to **CAC**, we speculate that this product may be **CAP**. Several attempts to trap this species with various amines to isolate a stable urea derivative have proven unsuccessful. Full structural elucidation of this product will be essential for improving our mechanistic understanding of **CAC** formation.

### Mechanism of low flux peroxynitrite product formation

At low peroxynitrite fluxes, H<sub>2</sub>O dominates as the nucleophile, and attacks **8-oxoG**<sup>ox</sup> at C5 to give 5-hydroxy-8-oxoG. This is an important intermediate in





to alcohols, are present [53]. Thus, **Sp/Gh** are the expected final products in our system. Second, **8-oxoG<sup>+</sup>/8'-oxoG** reacts very slowly with  $O_2$  ( $k_2 < 10^2 M^{-1} s^{-1}$  [82]) and hence this reaction cannot effectively compete with  $O_2^{\cdot-}$  for **8-oxoG<sup>+</sup>/8-oxoG**. Lastly, it is likely that aerobic peroxynitrite decomposition in the presence of thiols leads to  $O_2^{\cdot-}$  production [83], thus ensuring that reaction of  $O_2^{\cdot-}$  with **8-oxoG<sup>+</sup>/8-oxoG** effectively out-competes the slow reaction with  $O_2$ . Thus, mechanism (ii) is an attractive model for explaining **Sp** formation with incorporation of an  $O_2$ -derived [O] atom.

Recognition of **HICA** as a product of peroxynitrite-mediated **8-oxoG** oxidation came only after performing low flux experiments. Isotope labeling indicates that it incorporates three water-derived [O] atoms, and we have proposed that it is formed via hydration of **8-oxoG<sup>ox</sup>** at both C4 and C5 to give **12**, with subsequent cleavage of the N3–C4 bond and elimination of guanidine to yield **HICA** [71]. Chemically, **HICA** is a minor reaction product in comparison to **Gh/Sp**, and this may in part reflect preferential hydration at C5 over the relatively more hindered C4. However, it may still be important biologically, even if formed at very low levels, if it is a highly persistent mutagenic or toxic lesion.

#### *Biological implications of peroxynitrite-mediated G and 8-oxoG oxidation*

Thus far, we have emphasized the identity and chemistry of peroxynitrite-induced **G** and **8-oxoG** products at the nucleoside level. However, several studies have shown that they are also produced in single and double stranded oligodeoxynucleotides and calf thymus DNA treated with bolus peroxynitrite. Both **G**-derived nitration products **8-NO<sub>2</sub>G** [34,42] and **NI** [84] have been reported, albeit the latter at lower levels in double relative to single stranded DNA. The oxidation product **8-oxoG** has also been detected in peroxynitrite-treated plasmid DNA [30], and its further oxidation products identified in oligodeoxynucleotides [85,86]. Taken together, these observations demonstrate that the same products formed at the nucleoside level are produced in DNA, and hence may be important in understanding peroxynitrite's mutagenicity.

Indeed, a strong correlation exists between the preferential reactivity of peroxynitrite with **G** and **8-oxoG** and its ability to induce DNA mutations predominantly at **G** residues. The mutation types and frequencies in initial studies consisted of **G:C** → **T:A** (75%), **G:C** → **C:G** (12%), and **G:C** → **A:T** (14%) and were determined using shuttle vectors treated with high peroxynitrite fluxes [87]. Our studies demonstrating a dramatic shift in the products obtained at high and low peroxynitrite fluxes, prompted further investigations comparing mutational profiles at high (bolus addition), intermediate (infusion), and low (SIN-1 generated) peroxynitrite fluxes. While mutation frequencies were significantly reduced for both infusion and SIN-1 reactions relative to bolus addition, **G:C** → **T:A** transversions were still the predominant

mutations observed [88]. These results suggest that the DNA-damage products formed at low and high peroxynitrite fluxes may differ, but there is significant overlap in their propensity to induce similar mutations originating at damaged **G** residues.

This conclusion is supported by studies examining the mutational properties of the individual oxidation products introduced site-specifically into oligonucleotides or viral genomes for in vitro and in vivo study, respectively. The sufficiently stable major products of **G** and **8-oxoG** oxidation have been studied, and their mutational spectra are summarized below. **8-OxoG** is known to induce **G:C** → **T:A** mutations [89–93]. Of the other **G**-derived products, **Iz** gives rise mainly to **G:C** → **C:G** mutations [94], in agreement with a previous preliminary prediction made based on molecular orbital calculations [95]. However, its hydrolysis product **Oz** causes **G:C** → **T:A** mutations [86,96]. **NI** gives rise to **G:C** → **T:A** (22%), **G:C** → **A:T** (19%), and **G:C** → **C:G** (9%) mutations [94]. **8-NO<sub>2</sub>G** depurinates to leave an abasic site, and in systems that conform to the A-rule, is expected to lead to **G:C** → **T:A** mutation [97,98]. Recently, this product has also been shown to be a miscoding lesion, directly inducing **G:C** → **T:A** mutations predominantly, and **G:C** → **C:G** and **G:C** → **A:T** mutations and deletions to a lesser extent [99]. Therefore, **8-NO<sub>2</sub>G** induces mutations directly and indirectly, but the principal mechanism cannot easily be distinguished by examining mutational spectra, as both pathways lead to the same predominant mutation. Of the **8-oxoG** products, **OA** and **UA** are expected to be the major stable products in bolus addition studies. **OA** has been shown to induce **G:C** → **T:A** mutations at high frequency [86,100], and this is also true for **UA** [63]. Similarly, **CA** causes predominantly **G:C** → **T:A** mutations [86,101]. On the other hand, the major products of low peroxynitrite fluxes, **Gh** and **Sp** will induce both **G:C** → **C:G** and **G:C** → **T:A** transversions [102,103], with the former as the predominant mutation (98% for **Gh** and 65% for **Sp**, respectively) [103]. Unfortunately, **DGh** and **NO<sub>2</sub>-DGh** are too unstable for determination of mutational spectra, while the more recently elucidated **HICA** has not been studied. Overall, the strong concordance between the types of mutations induced by the individual products with those observed in aggregate studies lend further support to the conclusion that oxidation of **G** and **8-oxoG** can explain the majority of peroxynitrite-induced mutations.

#### **Conclusions**

Peroxynitrite induces DNA base damage predominantly at **G** and **8-oxoG** nucleobases via oxidation reactions that are thermodynamically and kinetically favorable. In addition to purely 'oxidative' products, nitration products result in both cases, consistent with the generation of radical intermediates that can recombine with the  $NO_2$  formed during peroxynitrite degradation. The neutral **G** radical, **G<sup>•</sup>**, is the intermediate

that reacts with  $\text{NO}_2$  to yield **8-NO<sub>2</sub>G** and **NI**, while for **8-oxoG** we have proposed a reactive guanidynylidene radical intermediate. In addition to rationalizing the formation of **NO<sub>2</sub>-DGh**, the putative guanidynylidene radical generated during **G** and **8-oxoG** oxidation also reconciles the formation of **Iz** on the very rapid timescale of peroxynitrite-mediated reactions, and may also be important in other **G** and **8-oxoG** oxidation models.

The products generated during peroxynitrite-mediated **8-oxoG** oxidation are dependent on oxidant flux. At high fluxes, **DGh**, **CAC**, and **NO<sub>2</sub>-DGh** are the predominant products, and result from reactions involving attack by the peroxynitrite anion,  $\text{ONOO}^-$ , on an electrophilic intermediate. At low fluxes, **Sp**, **Gh**, and **HICA** are formed, and these arise from  $\text{H}_2\text{O}$  attacking the electrophilic intermediate. At intermediate fluxes, these two pathways compete and both product sets result.

The dramatic shift in product distribution as a function of peroxynitrite flux illustrates the importance of using in vitro model systems that, as closely as possible, recapitulate the physiologic system. Given that peroxynitrite and its derived radicals are all chemically reactive, it is crucial to ensure that the relative concentrations are well controlled to minimize competing reactions that may not be relevant in vivo. In addition to the reaction of **8-oxoG** with peroxynitrite, a similar phenomenon has been observed in experiments with tyrosine, where 3-nitrotyrosine and 3,3'-dityrosine formation dominates at high and low peroxynitrite fluxes, respectively [104–108]. Increasingly sophisticated systems for modeling in vivo peroxynitrite production are being developed [104–108] and these should help with predicting more precisely the most likely in vivo products.

Nevertheless, this product shift with changing peroxynitrite flux is intriguing. Together with the emerging information on the genotoxic and mutational characteristics of the individual oxidation products, it suggests that the extent of tissue damage, mutational spectra and, hence, cancer risk may change as a function of peroxynitrite fluxes as different product combinations predominate. This will require further work to establish how and whether the individual products are repaired, and their frequency in DNA from tissues under oxidative stress. The former goal is readily achievable, but the latter presents an enormous challenge since some of these products are unstable under the conditions required to digest DNA, and the resulting nucleosides and aglycones are highly polar, making chromatographic separation difficult. However, with the rapid improvements in sensitive and specific tandem chromatographic and mass spectrometric methods, this problem too, should be tractable.

## Acknowledgments

This work was supported by NIH Grants 5F31-HG00144 (J.C.N), CA 26731, and ES02109-24 (MIT Center for Environmental Health Sciences).

## References

- [1] J. Garthwaite, Glutamate, nitric oxide and cell-cell signaling in the nervous system, *Trends Neurosci.* 14 (2) (1991) 60–67.
- [2] M.A. Marletta, M.A. Tayeh, J.M. Hevel, Unraveling the biological significance of nitric oxide, *Biofactors* 2 (4) (1990) 219–225.
- [3] L.J. Ignarro, Biosynthesis and metabolism of endothelium-derived nitric oxide, *Annu. Rev. Pharmacol. Toxicol.* (1990) 30535–30560.
- [4] S. Moncada, R.M. Palmer, E.A. Higgs, Nitric oxide: physiology, pathophysiology, and pharmacology, *Pharmacol. Rev.* 43 (2) (1991) 109–142.
- [5] C. Nathan, Nitric oxide as a secretory product of mammalian cells, *FASEB J.* 6 (12) (1992) 3051–3064.
- [6] J. MacMicking, Q.W. Xie, C. Nathan, Nitric oxide and macrophage function, *Annu. Rev. Immunol.* (1997) 15323–15350.
- [7] C. Nathan, Q.W. Xie, Regulation of biosynthesis of nitric oxide, *J. Biol. Chem.* 269 (19) (1994) 13725–13728.
- [8] M.A. Marletta, Nitric oxide synthase: aspects concerning structure and catalysis, *Cell* 78 (6) (1994) 927–930.
- [9] M.T. Gladwin, J.H. Crawford, R.P. Patel, The biochemistry of nitric oxide, nitrite, and hemoglobin: role in blood flow regulation, *Free Radic. Biol. Med.* 36 (6) (2003) 707–717.
- [10] Z. Huang, R. Shiva, D.B. Kim-Shapiro, R.P. Patel, L.A. Ringwood, C.E. Irby, K.T. Huang, C. Ho, N. Hogg, A.N. Schechter, M.T. Gladwin, Enzymatic function of hemoglobin as a nitrite reductase that produces NO under allosteric control, *J. Clin. Invest.* 115 (8) (2005) 2099–2107.
- [11] I.I. Singer, D.W. Kawka, S. Scott, J.R. Weidner, R.A. Mumford, T.E. Riehl, W.F. Stenson, Expression of inducible nitric oxide synthase and nitrotyrosine in colonic epithelium in inflammatory bowel disease, *Gastroenterology* 111 (4) (1996) 871–885.
- [12] G. Dijkstra, H. Moshage, H.M. van Dullemen, A. de Jager-Krikken, A.T. Tiebosch, J.H. Kleibeuker, P.L. Jansen, H. van Goor, Expression of nitric oxide synthases and formation of nitrotyrosine and reactive oxygen species in inflammatory bowel disease, *J. Pathol.* 186 (4) (1998) 416–421.
- [13] D.M. McCafferty, Peroxynitrite and inflammatory bowel disease, *Gut* 46 (3) (2000) 436–439.
- [14] H. Ohshima, H. Bartsch, Chronic infections and inflammatory processes as cancer risk factors: possible role of nitric oxide in carcinogenesis, *Mutat. Res.* 305 (2) (1994) 253–264.
- [15] N. Ma, Y. Adachi, Y. Hiraku, N. Horiki, S. Horiike, I. Imoto, S. Pinlaor, M. Murata, R. Semba, S. Kawanishi, Accumulation of 8-nitroguanine in human gastric epithelium induced by *Helicobacter pylori* infection, *Biochem. Biophys. Res. Commun.* 319 (2) (2004) 506–510.
- [16] P.C. Dedon, S.R. Tannenbaum, Reactive nitrogen species in the chemical biology of inflammation, *Arch. Biochem. Biophys.* 423 (1) (2004) 12–22.
- [17] H.B. Yu, L. Venkatarangan, J.S. Wishnok, S.R. Tannenbaum, Quantitation of four guanine oxidation products from reaction of DNA with varying doses of peroxynitrite, *Chem. Res. Toxicol.* (2005), in Press.
- [18] G. Merenyi, J. Lind, Free radical formation in the peroxynitrous acid ( $\text{ONOOH}$ )/peroxynitrite ( $\text{ONOO}^-$ ) system, *Chem. Res. Toxicol.* 11 (4) (1998) 243–246.
- [19] C.E. Richeson, P. Mulder, V.W. Bowry, K.U. Ingold, The complex chemistry of peroxynitrite decomposition: new insights, *J. Am. Chem. Soc.* 120 (29) (1998) 7211–7219.
- [20] J.W. Coddington, J.K. Hurst, S.V. Lymar, Hydroxyl radical formation during peroxynitrous acid decomposition, *J. Am. Chem. Soc.* 121 (11) (1999) 2438–2443.
- [21] S.V. Lymar, J.K. Hurst, Rapid reaction between peroxynitrite ion and carbon dioxide: implications for biological activity, *J. Am. Chem. Soc.* 117 (34) (1995) 8867–8868.
- [22] A. Denicola, B.A. Freeman, M. Trujillo, R. Radi, Peroxynitrite reaction with carbon dioxide/bicarbonate: kinetics and influence on peroxynitrite-mediated oxidations, *Arch. Biochem. Biophys.* 333 (1) (1996) 49–58.

- [23] R.M. Uppu, G.L. Squadrito, W.A. Pryor, Acceleration of peroxynitrite oxidations by carbon dioxide, *Arch. Biochem. Biophys.* 327 (2) (1996) 335–343.
- [24] S.V. Lyman, J.K. Hurst, CO<sub>2</sub>-catalyzed one-electron oxidations by peroxynitrite: properties of the reactive intermediate, *Inorg. Chem.* 37 (2) (1998) 294–301.
- [25] S. Goldstein, G. Czapski, Formation of peroxynitrate from the reaction of peroxynitrite with CO<sub>2</sub>: evidence for carbonate radical production, *J. Am. Chem. Soc.* 120 (14) (1998) 3458–3463.
- [26] R.M. Uppu, G.W. Winston, W.A. Pryor, Reactions of peroxynitrite with aldehydes as probes for the reactive intermediates responsible for biological nitration, *Chem. Res. Toxicol.* 10 (12) (1997) 1331–1337.
- [27] D. Yang, Y. Tang, J. Chen, X. Wang, M.D. Bartberger, K.N. Houk, L. Olson, Ketone-catalyzed decomposition of peroxynitrite via dioxirane intermediates, *J. Am. Chem. Soc.* 121 (51) (1999) 11976–11983.
- [28] H. Masumoto, H. Sies, The reaction of ebselen with peroxynitrite, *Chem. Res. Toxicol.* 9 (1) (1996) 262–267.
- [29] M.G. Salgo, K. Stone, G.L. Squadrito, J.R. Battista, W.A. Pryor, Peroxynitrite causes DNA nicks in plasmid pBR322, *Biochem. Biophys. Res. Commun.* 210 (3) (1995) 1025–1030.
- [30] L.J. Kennedy, K. Moore Jr., J.L. Caulfield, S.R. Tannenbaum, P.C. Dedon, Quantitation of 8-oxoguanine and strand breaks produced by four oxidizing agents, *Chem. Res. Toxicol.* 10 (4) (1997) 386–392.
- [31] C. Szabo, H. Ohshima, DNA damage induced by peroxynitrite: subsequent biological effects, *Nitric Oxide* 1 (5) (1997) 373–385.
- [32] S. Burney, J.L. Caulfield, J.C. Niles, J.S. Wishnok, S.R. Tannenbaum, The chemistry of DNA damage from nitric oxide and peroxynitrite, *Mutat. Res.* 424 (1999) 37–49.
- [33] V. Yermilov, Y. Yoshie, J. Rubio, H. Ohshima, Effects of carbon dioxide/bicarbonate on induction of DNA single-strand breaks and formation of 8-nitroguanine, 8-oxoguanine and base propenal mediated by peroxynitrite, *FEBS Lett.* 399 (1996) 67–39970.
- [34] N.Y. Tretyakova, S. Burney, B. Pamir, J.S. Wishnok, P.C. Dedon, G.N. Wogan, S.R. Tannenbaum, Peroxynitrite-induced DNA damage in the supF gene: correlation with the mutational spectrum, *Mutat. Res. Fundam. Mol. Mech. Mutagen.* 447 (2) (2000) 287–303.
- [35] S. Burney, J.C. Niles, P.C. Dedon, S.R. Tannenbaum, DNA damage in deoxynucleosides and oligonucleotides treated with peroxynitrite, *Chem. Res. Toxicol.* 12 (6) (1999) 513–520.
- [36] S. Steenken, S.V. Jovanovic, How easily oxidized is DNA? One-electron reduction potentials of adenosine and guanosine radicals in aqueous solution, *J. Am. Chem. Soc.* 119 (3) (1997) 617–618.
- [37] D.M. Stanbury, Reduction potentials involving inorganic free radicals in aqueous solution, *Adv. Inorg. Chem.* 33 (1989) 69–138.
- [38] R.E. Huie, C.L. Clifton, P. Neta, Electron transfer reaction rates and equilibria of the carbonate and sulfate radical anions, *Radiat. Phys. Chem.* 38 (5) (1991) 477–481.
- [39] S. Steenken, Purine-bases, nucleosides and nucleotides-aqueous-solution redox chemistry and transformation reactions of their radical cations and e- and OH adducts, *Chem. Rev.* 89 (3) (1989) 503–520.
- [40] V. Shafirovich, A. Dourandin, W.D. Huang, N.E. Geacintov, The carbonate radical is a site-selective oxidizing agent of guanine in double-stranded oligonucleotides, *J. Biol. Chem.* 276 (27) (2001) 24621–24626.
- [41] V. Yermilov, J. Rubio, M. Becchi, M.D. Friesen, B. Pignatelli, H. Ohshima, Formation of 8-nitroguanine by the reaction of guanine with peroxynitrite in vitro, *Carcinogenesis* 16 (9) (1995) 2045–2050.
- [42] V. Yermilov, J. Rubio, H. Ohshima, Formation of 8-nitroguanine in DNA treated with peroxynitrite in vitro and its rapid removal from DNA by depurination, *FEBS Lett.* 376 (1995) 207–210.
- [43] J.C. Niles, J.S. Wishnok, S.R. Tannenbaum, A novel nitroimidazole compound formed during the reaction of peroxynitrite with 2',3',5'-tri-O-acetyl-guanosine, *J. Am. Chem. Soc.* 123 (49) (2001) 12147–12151.
- [44] T. Douki, J. Cadet, B.N. Ames, An adduct between peroxynitrite and 2'-deoxyguanosine: 4,5-dihydro-5-hydroxy-4-(nitrosooxy)-2'-deoxyguanosine, *Chem. Res. Toxicol.* 9 (1) (1996) 3–7.
- [45] R.N. Goyal, G. Dryhurst, Redox chemistry of guanine and 8-oxyguanine and a comparison of the peroxidase-catalyzed and electrochemical oxidation of 8-oxyguanine, *J. Electroanal. Chem.* 135 (1982) 75–91.
- [46] K. Kahn, P. Serfozo, P.A. Tipton, Identification of the true product of the urate oxidase reaction, *J. Am. Chem. Soc.* 119 (23) (1997) 5435–5442.
- [47] W. Luo, J.G. Muller, E.M. Rachlin, C.J. Burrows, Characterization of spiroiminodihydantoin as a product of one-electron oxidation of 8-oxo-7,8-dihydroguanosine, *Org. Lett.* 2 (5) (2000) 613–616.
- [48] J.C. Niles, J.S. Wishnok, S.R. Tannenbaum, Spiroiminodihydantoin and guanidinohydantoin are the dominant products of 8-oxoguanosine oxidation at low fluxes of peroxynitrite: mechanistic studies with O-18, *Chem. Res. Toxicol.* 17 (11) (2004) 1510–1519.
- [49] T. Douki, J. Cadet, Peroxynitrite mediated oxidation of purine bases of nucleosides and isolated DNA, *Free Radic. Res.* 24 (5) (1996) 369–380.
- [50] J. Cadet, M. Berger, G.W. Buchko, P.C. Joshi, S. Raoul, J.-L. Ravanat, 2,2-Diamino-4-[(3,5-di-O-acetyl-2-deoxy-β-D-erythro-pentofuransyl)amino-5-(2H)-oxazolone]: a novel and predominant radical oxidation product of 3',5'-di-O-acetyl-2'-deoxyguanosine, *J. Am. Chem. Soc.* 116 (1994) 7403–7404.
- [51] R.M. Uppu, R. Cueto, G.L. Squadrito, M.G. Salgo, W.A. Pryor, Competitive reactions of peroxynitrite with 2'-deoxyguanosine and 7,8-dihydro-8-oxo-2'-deoxyguanosine (8-oxo-dG), relevance to the formation of 8-oxo-dG in DNA exposed to peroxynitrite, *Free Radic. Biol. Med.* 21 (3) (1996) 407–411.
- [52] J. Ravanat, J. Cadet, Reaction of singlet oxygen with 2'-deoxyguanosine and DNA. Isolation and characterization of the main oxidation products, *Chem. Res. Toxicol.* 8 (3) (1995) 379–388.
- [53] J.C. Niles, J.S. Wishnok, S.R. Tannenbaum, Spiroiminodihydantoin is the major product of the 8-oxo-7,8-dihydroguanosine reaction with peroxynitrite in the presence of thiols and guanosine photooxidation by methylene blue, *Org. Lett.* 3 (7) (2001) 963–966.
- [54] W. Adam, M.A. Arnold, M. Grune, W.M. Nau, U. Pischel, C.R. Saha-Moller, Spiroiminodihydantoin is a major product in the photooxidation of 2'-deoxyguanosine by the triplet states and oxyl radicals generated from hydroxyacetophenone photolysis and dioxetane thermolysis, *Org. Lett.* 4 (4) (2002) 537–540.
- [55] H.J. Helbock, K.B. Beckman, M.K. Shigenaga, P.B. Walter, A.A. Woodall, H.C. Yeo, B.N. Ames, DNA oxidation matters: the HPLC-electrochemical detection assay of 8-oxo-deoxyguanosine and 8-oxo-guanine, *Proc. Natl. Acad. Sci. USA* 95 (1) (1998) 288–293.
- [56] S. Steenken, S.V. Jovanovic, M. Bietti, K. Bernhard, The trap depth (in DNA) of 8-oxo-7,8-dihydro-2'-deoxyguanosine as derived from electron-transfer equilibria in aqueous solution, *J. Am. Chem. Soc.* 122 (10) (2000) 2373–2374.
- [57] V. Shafirovich, J. Cadet, D. Gasparutto, A. Dourandin, N.E. Geacintov, Nitrogen dioxide as an oxidizing agent of 8-oxo-7,8-dihydro-2'-deoxyguanosine but not of 2'-deoxyguanosine, *Chem. Res. Toxicol.* 14 (2) (2001) 233–241.
- [58] S.O. Kelley, J.K. Barton, Radical migration through the DNA helix: chemistry at a distance, in: *Metal Ions in Biological Systems*, vol. 36, Marcel Dekker, New York, 1999, pp. 211–249.
- [59] C.J. Burrows, J.G. Muller, Oxidative nucleobase modifications leading to strand scission, *Chem. Rev.* 98 (3) (1998) 1109–1151.
- [60] G.B. Schuster, Long-range charge transfer in DNA: Transient structural distortions control the distance dependence, *Acc. Chem. Res.* 33 (4) (2000) 253–260.
- [61] J.C. Niles, S. Burney, S.P. Singh, J.S. Wishnok, S.R. Tannenbaum, Peroxynitrite reaction products of 3',5'-di-O-acetyl-8-oxo-7,8-dihydro-2'-deoxyguanosine, *Proc. Natl. Acad. Sci. USA* 96 (12) (1999) 11729–11734.
- [62] A. Chworos, Y. Coppel, I. Dubey, G. Pratviel, B. Meunier, Guanine oxidation: NMR characterization of a dehydro-guanidinohydantoin residue generated by a 2e-oxidation of d(GpT), *J. Am. Chem. Soc.* 123 (25) (2001) 5867–5877.

- [63] P.T. Henderson, W.L. Neeley, J.C. Delaney, F. Gu, J.C. Niles, S.S. Hah, S.R. Tannenbaum, J.M. Essigmann, Urea lesion formation in DNA as a consequence of 7,8-dihydro-8-oxoguanine oxidation and hydrolysis provides a potent source of point mutations, *Chem. Res. Toxicol.* 18 (1) (2005) 12–18.
- [64] S. Raoul, J. Cadet, Photosensitized reaction of 8-oxo-7,8-dihydro-2'-deoxyguanosine: identification of 1-(2-deoxy- $\beta$ -D-erythro-pentofuranosyl)cyanoic acid as the major singlet oxygen oxidation product, *J. Am. Chem. Soc.* 118 (8) (1996) 1892–1898.
- [65] J.C. Niles, J.S. Wishnok, S.R. Tannenbaum, A novel nitration product formed during the reaction of peroxynitrite with 2',3',5'-tri-O-acetyl-7,8-dihydro-8-oxoguanosine: N-nitro-N'-[1-(2,3,5-tri-O-acetyl- $\beta$ -D-erythro-pentofuranosyl)-2,4-dioximidazolidin-5-ylidene]guanidine, *Chem. Res. Toxicol.* 13 (5) (2000) 390–396.
- [66] W. Adam, C.R. Saha-Moller, A. Schonberger, Photooxidation of 8-oxo-7,8-dihydro-2'-deoxyguanosine by thermally generated triplet-excited ketones from 3-(hydroxymethyl)-3,4,4-trimethyl-1,2-dioxetane and comparison with type I and type II photosensitizers, *J. Am. Chem. Soc.* 118 (39) (1996) 9233–9238.
- [67] C. Vialas, G. Pratviel, C. Claparols, B. Meunier, Efficient oxidation of 2'-deoxyguanosine by Mn-TMPyP/KHSO<sub>5</sub> to imidazolone dIz without formation of 8-oxo-dG, *J. Am. Chem. Soc.* 120 (45) (1998) 11548–11553.
- [68] V. Duarte, D. Gasparutto, L.F. Yamaguchi, J.L. Ravanat, G.R. Martinez, M.H.G. Medeiros, P. Di Mascio, J. Cadet, Oxaluric acid as the major product of singlet oxygen-mediated oxidation of 8-oxo-7,8-dihydroguanine in DNA, *J. Am. Chem. Soc.* 122 (51) (2000) 12622–12628.
- [69] M. Kelm, R. Dahmann, D. Wink, M. Feelisch, The nitric oxide superoxide assay—insights into the biological chemistry of the NO/O<sub>2</sub><sup>•-</sup> interaction, *J. Biol. Chem.* 272 (15) (1997) 9922–9932.
- [70] R.J. Singh, N. Hogg, J. Joseph, E. Konorev, B. Kalyanaram, The peroxynitrite generator, SIN-1, becomes a nitric oxide donor in the presence of electron acceptors, *Arch. Biochem. Biophys.* 361 (2) (1999) 331–339.
- [71] J.C. Niles, J.S. Wishnok, S.R. Tannenbaum, Mass spectrometric identification of 4-hydroxy-2,5-dioxoimidazolidine-4-carboxylic acid during oxidation of 8-oxoguanosine by peroxynitrite and KHSO<sub>5</sub>/CoCl<sub>2</sub>, *Chem. Res. Toxicol.* 17 (11) (2004) 1501–1509.
- [72] E.O. Hole, W.H. Nelson, D.M. Close, E. Sagstuen, ESR and ENDOR study of the guanine cation: secondary product in 5'-dGMP, *J. Chem. Phys.* 86 (9) (1987) 5218–5219.
- [73] R. Misiaszek, C. Crean, N.E. Geacintov, V. Shafirovich, Combination of nitrogen dioxide radicals with 8-oxo-7,8-dihydroguanine and guanine radicals in DNA: oxidation and nitration end-products, *J. Am. Chem. Soc.* 127 (7) (2005) 2191–2200.
- [74] J. Heo, K.C. Prutzman, V. Mocanu, S.L. Campbell, Mechanism of free radical nitric oxide-mediated Ras guanine nucleotide dissociation, *J. Mol. Biol.* 346 (5) (2005) 1423–1440.
- [75] K.N. Houk, K.R. Condroski, W. Pryor, Radical and concerted mechanisms in the oxidations of amines, sulfides, and alkenes by peroxynitrite, peroxynitrous acid and the peroxynitrite-CO<sub>2</sub> adduct: density functional theory transition structures and energetics, *J. Am. Chem. Soc.* 118 (51) (1996) 13002–13006.
- [76] G. Merenyi, J. Lind, Thermodynamics of peroxynitrite and its CO<sub>2</sub> adduct, *Chem. Res. Toxicol.* 10 (11) (1997) 1216–1220.
- [77] J.L. Esker, M. Newcomb, The generation of nitrogen radicals and their cyclizations for the construction of the pyrrolidine nucleus, in: A.R. Katritzky (Ed.), *Advances in Heterocyclic Chemistry*, 58, Academic Press, San Diego, 1993, pp. 1–45.
- [78] A.G. Fallis, I.M. Brinza, Free radical cyclizations involving nitrogen, *Tetrahedron* 53 (52) (1997) 17543–17594.
- [79] H. Kasai, Z. Yamaizumi, M. Berger, J. Cadet, Photosensitized formation of 7,8-dihydro-8-oxo-2'-deoxyguanosine (8-hydroxy-2'-deoxyguanosine) in DNA by riboflavin: a non singlet oxygen mediated reaction, *J. Am. Chem. Soc.* 114 (24) (1992) 9692–9694.
- [80] R.N. Goyal, N. Jain, D.K. Garg, Electrochemical and enzymatic oxidation of guanosine and 8-hydroxyguanosine and the effects of oxidation products in mice, *Bioelectrochem. Bioenerg.* 43 (1) (1997) 105–114.
- [81] W.C. Luo, J.G. Muller, C.J. Burrows, The pH-dependent role of superoxide in riboflavin-catalyzed photooxidation of 8-oxo-7,8-dihydroguanosine, *Org. Lett.* 3 (18) (2001) 2801–2804.
- [82] R. Misiaszek, Y. Uvaydov, C. Crean, N.E. Geacintov, V. Shafirovich, Combination reactions of superoxide with 8-oxo-7,8-dihydroguanine radicals in DNA—kinetics and end products, *J. Biol. Chem.* 280 (8) (2005) 6293–6300.
- [83] C. Quijano, B. Alvarez, R.M. Gatti, O. Augusto, R. Radi, Pathways of peroxynitrite oxidation of thiol groups, *Biochem. J.* 322 (1997) 167–173.
- [84] F. Gu, W.G. Stillwell, J.S. Wishnok, A.J. Shallop, R.A. Jones, S.R. Tannenbaum, Peroxynitrite-induced reactions of synthetic oligo 2'-deoxynucleotides and DNA containing guanine: formation and stability of a 5-guanidino-4-nitroimidazole lesion, *Biochemistry* 41 (23) (2002) 7508–7518.
- [85] N.Y. Tretyakova, J.C. Niles, S. Burney, J.S. Wishnok, S.R. Tannenbaum, Peroxynitrite-induced reactions of synthetic oligonucleotides containing 8-oxoguanine, *Chem. Res. Toxicol.* 12 (5) (1999) 459–466.
- [86] P.T. Henderson, J.C. Delaney, F. Gu, S.R. Tannenbaum, J.M. Essigmann, Oxidation of 7,8-dihydro-8-oxoguanine affords lesions that are potent sources of replication errors in vivo, *Biochemistry* 41 (3) (2002) 914–921.
- [87] M.J. Juedes, G.N. Wogan, Peroxynitrite-induced mutation spectra of pSP189 following replication in bacteria and in human cells, *Mutat. Res. Fundam. Mol. Mech. Mutagen.* 349 (1) (1996) 51–61.
- [88] M.Y. Kim, M. Dong, P.C. Dedon, G.N. Wogan, Effects of peroxynitrite dose and dose rate on DNA damage and mutation in the supF shuttle vector, *Chem. Res. Toxicol.* 18 (1) (2005) 76–86.
- [89] M.L. Wood, M. Dizdaroglu, E. Gajewski, J.M. Essigmann, Mechanistic studies of ionizing radiation and oxidative mutagenesis: genetic effects of a single 8-hydroxyguanine (7-hydro-8-oxoguanine) residue inserted at a unique site in a viral genome, *Biochemistry* 29 (30) (1990) 7024–7032.
- [90] S. Shibutani, M. Takeshita, A.P. Grollman, Insertion of specific bases during DNA synthesis past the oxidation-damaged base 8-oxodG, *Nature* 349 (6308) (1991) 431–434.
- [91] K.C. Cheng, D.S. Cahill, H. Kasai, S. Nishimura, L.A. Loeb, 8-Hydroxyguanine, an abundant form of oxidative DNA damage, causes G → T and A → C substitutions, *J. Biol. Chem.* 267 (1) (1992) 166–172.
- [92] L.G. Lowe, F.P. Guengerich, Steady-state and pre-steady-state kinetic analysis of dNTP insertion opposite 8-oxo-7,8-dihydroguanine by *Escherichia coli* polymerases I exo- and II exo, *Biochemistry* 35 (30) (1996) 9840–9849.
- [93] V. Duarte, J.G. Muller, C.J. Burrows, Insertion of dGMP and dAMP during in vitro DNA synthesis opposite an oxidized form of 7,8-dihydro-8-oxoguanine, *Nucleic Acids Res.* 27 (2) (1999) 496–502.
- [94] W.L. Neeley, J.C. Delaney, P.T. Henderson, J.M. Essigmann, In vivo bypass efficiencies and mutational signatures of the guanine oxidation products 2-aminoimidazolone and 5-guanidino-4-nitroimidazole, *J. Biol. Chem.* 279 (42) (2004) 43568–43573.
- [95] K. Kino, I. Saito, H. Sugiyama, Product analysis of GG-specific photooxidation of DNA via electron transfer: 2-aminoimidazolone as a major guanine oxidation product, *J. Am. Chem. Soc.* 120 (29) (1998) 7373–7374.
- [96] V. Duarte, D. Gasparutto, M. Jaquinod, J. Cadet, In vitro DNA synthesis opposite oxazolone and repair of this DNA damage using modified oligonucleotides, *Nucleic Acids Res.* 28 (7) (2000) 1555–1563.
- [97] L.A. Loeb, B.D. Preston, Mutagenesis by apurinic apyrimidinic sites, *Annu. Rev. Genet.* 20 (1986) 201–230.
- [98] S. Shibutani, M. Takeshita, A.P. Grollman, Translesional synthesis on DNA templates containing a single abasic site—a mechanistic study of the “A rule”, *J. Biol. Chem.* 272 (21) (1997) 13916–13922.
- [99] N. Suzuki, M. Yasui, N.E. Geacintov, V. Shafirovich, S. Shibutani, Miscoding events during DNA synthesis past the nitration-damaged base 8-nitroguanine, *Biochemistry* 44 (25) (2005) 9238–9245.



- [100] V. Duarte, D. Gasparutto, M. Jaquinod, J.L. Ravanat, J. Cadet, Repair and mutagenic potential of oxaluric acid, a major product of singlet oxygen-mediated oxidation of 8-oxo-7,8-dihydroguanine, *Chem. Res. Toxicol.* 14 (1) (2001) 46–53.
- [101] D. Gasparutto, S. Da Cruz, A. Bourdat, M. Jaquinod, J. Cadet, Synthesis and biochemical properties of cyanuric acid nucleoside-containing DNA oligomers, *Chem. Res. Toxicol.* 12 (7) (1999) 630–638.
- [102] O. Korniyushyna, A.M. Berges, J.G. Muller, C.J. Burrows, In vitro nucleotide misinsertion opposite the oxidized guanosine lesions spiroiminodihydantoin and guanidinohydantoin and DNA synthesis past the lesions using *Escherichia coli* DNA polymerase I (Klenow fragment), *Biochemistry* 41 (51) (2002) 15304–15314.
- [103] P.T. Henderson, J.C. Delaney, J.G. Muller, W.L. Neeley, S.R. Tanenbaum, C.J. Burrows, J.M. Essigmann, The hydantoin lesions formed from oxidation of 7,8-dihydro-8-oxoguanine are potent sources of replication errors in vivo, *Biochemistry* 42 (31) (2003) 9257–9262.
- [104] G.R. Hodges, J. Marwaha, T. Paul, K.U. Ingold, A novel procedure for generating both nitric oxide and superoxide in situ from chemical sources at any chosen mole ratio. First application: tyrosine oxidation and a comparison with preformed peroxynitrite, *Chem. Res. Toxicol.* 13 (12) (2000) 1287–1293.
- [105] S. Pfeiffer, B. Mayer, Lack of tyrosine nitration by peroxynitrite generated at physiological pH, *J. Biol. Chem.* 273 (42) (1998) 27280–27285.
- [106] S. Pfeiffer, K. Schmidt, B. Mayer, Dityrosine formation outcompetes tyrosine nitration at low steady-state concentrations of peroxynitrite. Implications for tyrosine modification by nitric oxide/superoxide in vivo, *J. Biol. Chem.* 275 (9) (2000) 6346–6352.
- [107] S. Goldstein, G. Czapski, J. Lind, G. Merenyi, Tyrosine nitration by simultaneous generation of NO and  $O_2^{\cdot-}$  under physiological conditions. How the radicals do the job, *J. Biol. Chem.* 275 (5) (2000) 3031–3036.
- [108] C. Wang, W.M. Deen, Peroxynitrite delivery methods for toxicity studies, *Chem. Res. Toxicol.* 17 (1) (2004) 32–44.