

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/236251776>

Glucocorticoids and their receptors: Insights into specific roles in mitochondria

ARTICLE *in* PROGRESS IN BIOPHYSICS AND MOLECULAR BIOLOGY · APRIL 2013

Impact Factor: 2.27 · DOI: 10.1016/j.pbiomolbio.2013.04.001 · Source: PubMed

CITATIONS

11

READS

90

12 AUTHORS, INCLUDING:



In-Sung Song

Inje University

213 PUBLICATIONS 3,152 CITATIONS

SEE PROFILE



Jae Boum Youm

58 PUBLICATIONS 895 CITATIONS

SEE PROFILE



Seung Hun Jeong

Inje University

32 PUBLICATIONS 262 CITATIONS

SEE PROFILE




Jin Han

College of Medicine, Cardiovascular and Me...

147 PUBLICATIONS 2,127 CITATIONS

SEE PROFILE



Review

Glucocorticoids and their receptors: Insights into specific roles in mitochondria

Sung-Ryul Lee, Hyoung-Kyu Kim, In-Sung Song, Jaeboum Youm, Louise Anne Dizon, Seung-Hun Jeong, Tae-Hee Ko, Hye-Jin Heo, Kyoung Soo Ko, Byoung Doo Rhee, Nari Kim, Jin Han*

National Research Laboratory for Mitochondrial Signaling, Department of Physiology, College of Medicine, Cardiovascular and Metabolic Disease Center, Inje University, 633-165 Gaegeum-Dong, Busanjin-Gu, 613-735 Busan, Republic of Korea

ARTICLE INFO

Article history:

Available online 16 April 2013

Keywords:

Glucocorticoid
Glucocorticoid receptor
Glucocorticoid response element
Mitochondria
Genomic
Non-genomic

ABSTRACT

Glucocorticoids (GCs) affect most physiological systems and are the most frequently used drugs for multiple disorders and organ transplantation. GC functions depend on a balance between circulating GC and cytoplasmic glucocorticoid receptor II (GR). Mitochondria individually enclose circular, double-stranded DNA that is expressed and replicated in response to nuclear-encoded factors imported from the cytoplasm. Fine-tuning and response to cellular demands should be coordinately regulated by the nucleus and mitochondria; thus mitochondrial–nuclear interaction is vital to optimal mitochondrial function. Elucidation of the direct and indirect effects of steroids, including GCs, on mitochondria is an important and emerging field of research. Mitochondria may also be under GC control because GRs are present in mitochondria, and glucocorticoid response elements (GREs) reside in the mitochondrial genome. Therefore, mitochondrial gene expression can be regulated by GCs via at least two different mechanisms: direct action on mitochondrial DNA and oxidative phosphorylation (OXPHOS) genes, or by an indirect effect through interaction with nuclear genes. In this review, we outline possible mechanisms of regulation of mitochondrial genes in response to GCs in view of translocation of the GR into mitochondria and the possible regulation of OXPHOS genes by GREs in the mitochondrial genome.

© 2013 Elsevier Ltd. All rights reserved.

Contents

1. Introduction	45
2. Glucocorticoid receptor	45
3. Mitochondria	46
4. GCs and mitochondrial biogenesis	47
5. GCs, the GR, and mitochondria	48
5.1. Transcriptional control of mitochondrial proteins encoded by nDNA: nucleus to mitochondria	49
5.2. Transcriptional control of nDNA-encoded transcription factors that affect nDNA-encoded mitochondrial proteins	49
5.3. Transcriptional control of mitochondrial proteins encoded by mtDNA	50
5.4. Activation of cytosolic signaling proteins by plasma membrane or cytosolic steroid receptors	50
5.5. Non-genomic effect of GCs on mitochondria	50
6. Concluding remarks	51
Conflict of Interest	51
Acknowledgments	51
References	51

Abbreviations: AF, activation function (domain); AMP, adenosine monophosphate; ATP, adenosine triphosphate; COX, cytochrome c oxidase; D-loop, displacement loop (of mitochondrial genome); DBD, DNA-binding domain; LBD, (C-terminal) ligand-binding domain; GC, glucocorticoid; GR, glucocorticoid receptor II; GRE, glucocorticoid response element; mtDNA, mitochondrial DNA; mRNA, messenger RNA; nDNA, nuclear DNA; NLS, nuclear localizing sequence; NO, nitric oxide; NOS, nitric oxide synthase; NRF, nuclear respiratory factors; NTD, N-terminal domain; OXPHOS, (mitochondrial) oxidative phosphorylation; PGC-1, peroxisome proliferator-activated receptor (PPAR)- γ coactivator 1; PPAR, peroxisome proliferator-activated receptor; rRNA, ribosomal RNA; TFAM, mitochondrial transcription factor A; tRNA, transfer RNA; UCP-1, (mitochondrial) uncoupling protein-1.

* Corresponding author. Tel.: +82 51 890 6727; fax: +82 51 894 5714.

E-mail address: phyhanj@inje.ac.kr (J. Han).

1. Introduction

Our understanding of the multiple effects of steroid hormones continues to advance (Hammes and Levin, 2011). Glucocorticoids (GCs; cortisol in humans and corticosterone in rodents) control approximately 20% of the expressed human genome, targeting almost all organs and tissues (Chrousos and Kino, 2005). GCs are crucial to life because they regulate and support an array of vital cardiovascular, metabolic, immunologic, and homeostatic functions. Although cortisol secretion varies diurnally (Dallman et al., 1993; Walker et al., 2012), its release also fluctuates depending on traumatic stress, as well as other types of stress, including illness, surgery, extreme temperatures, and psychological stress (Barnes, 2011; Biddie et al., 2012). GC participates in the establishment of a hypermetabolic state linked to physical or mental stress and trauma, and accompanied by an increased metabolic rate, skeletal muscle protein metabolism, liver gluconeogenesis, and hyperglycemia with insulin resistance (Brillon et al., 1995). GC deficiency or insufficiency is uncommon; for example, cortisol deficiency due to adrenocortical insufficiency rarely occurs, but may lead to hypotension, weight loss, asthenia, hypoglycemia, and secondary endocrine effects, including GC-reversible hypothyroidism, hyperprolactinemia, and hypercalcemia (Burke, 1985). Clinically, under- or over-production of GCs, as seen in Addison's (cortisol deficiency) or Cushing's (cortisol excess) syndrome, may result in adverse reactions or even death (Burke, 1985; Whitworth et al., 2000). As in the case of adrenalectomy, although GC insufficiency leads to upregulation of the glucocorticoid receptor II (GR), the GR does not function properly in the absence of GC (Gregory et al., 1976). Similarly, GC alone does not have significant effects on the cardiovascular or other organ systems. Therefore, the actions of GCs and the functional receptor system, the GR, are tightly coupled, and over- or under-stimulation of GC signaling may evoke pathological changes.

Mitochondria are integral to adenosine triphosphate (ATP) production, apoptosis (Fosslien, 2001; Green and Kroemer, 2004; Jeong and Seol, 2008), and production of biomolecules such as amino acids, lipids, hemes, purines, and steroid hormones (Miller, 2011). Intracellular cholesterol is ultimately converted to GC in the mitochondria (Miller, 2011) and the secretion of synthesized GC is tightly controlled by a specific mechanism in the adrenal gland (Rosol et al., 2001). Mitochondria also participate in intracellular calcium homeostasis; however, exposure to excessive Ca^{2+} levels causes massive swelling of the mitochondrial matrix, leading to necrotic cell death (Green and Kroemer, 2004). Furthermore, mitochondria are first-line responders to different stresses (Ryan and Hoogenraad, 2007) and can adjust bioenergetics, thermogenesis, and oxidative stress and/or apoptotic responses in response to perturbed homeostasis (Lesnefsky et al., 2001; Manoli et al., 2007). Consequences of mitochondrial damage include various diseases associated with multiple organ failure, which often precipitates the death of critical care patients (Kozlov et al., 2011). The concept that the nucleus is the domain of steroid receptor action has become controversial with the discovery of steroid hormone receptors and the GR in mitochondria and the identification of glucocorticoid response elements (GREs) in the mitochondrial genome (Demonacos et al., 1995, 1996; Martinus et al., 1996; Tsiriyotis et al., 1997). The consensus sequence of a positive GRE is 5'-GGTA-CAnnnTGTCT-3' (Gruber et al., 2004), whereas a negative GRE has a more variable sequence (Adcock, 2000). Based on these findings, exploration of the direct and rapid effects of steroid hormones on mitochondria is a novel field worthy of investigation (Gavrilova-Jordan and Price, 2007).

The dual localization of proteins in mitochondria and the nucleus, and the involvement of these proteins in control of gene expression or post-transcriptional processes, is also a common

phenomenon among plants and higher eukaryotes (Duchene and Giege, 2012). It has been strongly postulated that the significance of the GR and GRE in mitochondria is crucial to understanding the effect of GCs on this organelle and the possibility of genomic interplay between the nucleus and mitochondria (Ioannou et al., 1988). Cytoplasmic GRs, which share no sequence homology with other known mitochondrial localization signals, can translocate into the liver (Demonacos et al., 1993; Psarra and Sekeris, 2011) and lymphoid cell mitochondria (Boldizar et al., 2010), or move out of brain mitochondria (Demonacos et al., 1995), in response to dexamethasone (DEX) treatment or acute stress, implying that the mitochondrial GR is not static but is dynamically regulated in response to GCs. Simoes et al. (2012) also demonstrated that asthma, an inflammatory disease, reduces mitochondrial GR and OXPHOS enzyme biosynthesis in mouse lung epithelial cell monolayers. Although mitochondrial gene expression has been less well characterized than that of nuclear genes, there is a strong possibility that the GR also plays a role in mitochondrial gene expression. Because mitochondria also possess functional GREs, these organelles are not simply regulated by the nucleus; rather, the regulation of expression of some genes is finely and rapidly coordinated by translocation of the GR between the nucleus and mitochondria. Thus, the regulatory role of GC and GR on mitochondrial gene expression may take place through at least two different mechanisms: a direct effect on mitochondrial DNA (mtDNA) or an indirect effect via interaction with nuclear genes (e.g., those encoding mitochondrial RNA-processing enzymes, mitochondrial transcription factors, and nuclear respiratory factors). In this review, we outline the possible effects of GCs on regulation of mitochondrial gene expression in view of the localization of the GR in mitochondria and possible impacts on regulation of mitochondrial proteins.

2. Glucocorticoid receptor

The action of GCs is dependent on the expression of the GR, and, in turn, the functions of the GR in gene regulation are affected by chromatin structure, epigenetic factors, genetic variation, and the pattern of GC secretion (Biddie et al., 2012; John et al., 2011; Nicolaides et al., 2010). In addition, the GR can be also influenced by ligand binding, chemical compounds, post-translational modification, chaperones (or co-chaperones), GR isoforms, transcriptional co-regulators, transcription factors, and other proteins (Chrousos and Kino, 2005, 2007; Lu and Cidlowski, 2004). Therefore, complex GC–GR signaling activity contributes to the intricate nature of human physiology and pathophysiology.

The human GR is encoded by nine exons of a single gene located on chromosome 5 (Fig. 1). The protein consists of a variable N-terminal domain (NTD), two hormone-independent activation function domains (AF; AF1 and 2), a DNA-binding domain (DBD) with two zinc finger motifs, a hinge (H) region, and a C-terminal ligand-binding domain (LBD). The human GR also contains two nuclear localization sequences (Savory et al., 1999). Sumoylation sites are present in AF1. In the absence of ligand, newly synthesized GR is phosphorylated at three sites (S203, S221, and S226) and then rapidly dephosphorylated by protein phosphatase(s) associated with the LBD. Several post-translational modifications of GR- α , including phosphorylation, sumoylation, and ubiquitination, can modify aspects of signaling, such as subcellular localization, interaction with other proteins, and degradation (Anbalagan et al., 2012; Ismaili and Garabedian, 2004). The GR- α is a classical glucocorticoid receptor and has eight isoforms, with NTDs of different lengths, that are produced by alternative translation initiation from a single GR mRNA species. GR- α isoforms have different cellular distributions and gene regulatory profiles (Lu and Cidlowski, 2005; Oakley




Fig. 1. Structure of the human GR. A single GR gene resides on human chromosome 5. Alternative splicing of nine exons produces the GR- α and GR- β isoforms. GR- α has an NTD, DBD with two zinc finger motifs, H region, and C-terminal LBD. GR- β has no LBD and acts as a dominant-negative form of GR- α (Yudt et al., 2003). The NLS and domains AF1 and AF2 are shown. Multiple GR- α isoforms are produced by initiation of translation of the NTD at alternative sites. Other GR variants, including GR- γ (arginine insertion between the two zinc fingers of the DBD), GR- δ (deletion of the proximal 185 amino acids of the LBD, encoded by exons 5–7), and GR-P (a truncated receptor mutant lacking the distal half of the LBD) have been reported in cancer, but exact functions of these variants are still unknown (McMaster and Ray, 2007). AF, hormone-independent activation function domain; DBD, DNA-binding domain; GR, glucocorticoid receptor II; LBD, ligand-binding domain; H, hinge; NLS, nuclear localizing sequence; NTD, N-terminal domain.

and Cidlowski, 2011). The splice variant GR- β is expressed at a lower level than GR- α (Pujols et al., 2002), and exerts a dominant-negative effect on the GR- α isoforms due to the absence of the LBD (Yudt et al., 2003). Other GR splice variants, including GR- γ , GR- δ , and GR-P, have been reported in some types of cancer (Fig. 1). GR- γ , an integral splice variant of GR- α , has an extra arginine in the DBD and shows reduced transcriptional activity compared with GR- α (Krett et al., 1995; Moalli et al., 1993; Rivers et al., 1999). However, the exact functions of these variants are still under investigation (McMaster and Ray, 2007). Mutations in the human GR gene can impair GC signal transduction by altering tissue reactivity to the GR, and can have various outcomes, ranging from asymptomatic to extreme cases of hyperandrogenism, fatigue, and/or mineralocorticoid excess (Charmandari and Kino, 2007). The GR is primarily metabolized via the ubiquitin/proteasome-dependent protein degradation pathway (Wallace and Cidlowski, 2001; Weber et al., 2002).

Insight into the pathophysiological role of the GR was gained in research using transgenic animal models (Cole et al., 2001; Goodwin et al., 2011; Sainte-Marie et al., 2007). The events of embryonic development were less affected in GR-deficient mice, but neonatal mortality was attributed to retarded lung maturation and lung collapse (Cole et al., 1995). In addition, these mice had decreased ability to induce expression of major gluconeogenic enzymes (i.e., those that affect mitochondrial function indirectly in the liver) (Cole et al., 1995), but showed markedly elevated plasma adrenocorticotrophic hormone and corticosterone (Cole et al., 1995, 2001). Additionally, knockout of GR genes can increase secretion of corticosteroid-binding globulin (CBG), a 50–60-kDa glycoprotein with a single steroid-binding site. CBG is a primary glucocorticoid transporter in the blood stream (Gagliardi et al., 2010; Perogamvros et al., 2010) and, thereby, suppresses GC availability in the circulation (Cole et al., 1999). In mice with a partially deleted GR gene,

both fetuses and adults showed profound resistance to DEX-mediated functional activation or repression (Cole et al., 2001). Surprisingly, when the point mutation A458T was introduced into the GR D-loop region, resulting in defective DNA binding, this mutant had normal local and systemic anti-inflammatory responses (Reichardt et al., 2001). This implies that the anti-inflammatory effect of GCs is not solely dependent on GC-mediated genomic effects.

3. Mitochondria

Mitochondria individually encapsulate their own genome (Fig. 2), which is expressed and replicated by the action of nuclear-encoded factors exported to the cytoplasm. Although there are large differences between tissues, a cell may contain several thousand mitochondria, and each mitochondrion contains 1–10 copies of mtDNA (Bogenhagen and Clayton, 1974; Garesse and Vallejo, 2001). Mitochondrial morphology is not static, but is dynamically maintained through the processes of fission and fusion (Frazier et al., 2006). In humans, the maternally-inherited mtDNA consists of circular, double-stranded DNA containing 37 genes. The genes encode 13 proteins of the mitochondrial respiratory chain, two ribosomal RNAs (rRNAs; 12S and 16S), and 22 transfer RNAs (tRNAs) (Bonawitz et al., 2006). Mitochondrial integrity is essential for the function and survival of cells. Surprisingly, in rho-0 mitochondria, which have little or no mtDNA, the formation and function of mitochondrial respiratory complexes is inhibited; however, these mitochondria have a proton gradient required for importing protein and preventing apoptosis (Zamzami and Kroemer, 2001). Moreover, the content of mitochondrial proteins, including respiratory complexes, catabolic proteins, transport proteins, and the mitochondrial translational system, are dramatically changed in rho(0) mitochondria (Chevallet et al., 2006; Jeon et al., 2011). This implies that mitochondrial protein functional networks can be reorganized to respond to the physiological status of the cell, and that alterations in mtDNA can lead to pleiotropic effects on the mitochondrial architecture.

A noncoding sequence of mtDNA, the D-loop or hypervariable region, serves as a control region for both DNA transcription and replication (Scarpulla, 2012; Scarpulla et al., 2012). Unlike the nuclear genome, all mitochondrial genes are expressed together, primarily from three promoters, the L-strand promoter (LSP), and H-strand promoters (HSPs) 1 and 2. These promoters are located in the regulatory D-loop (Montoya et al., 1982) and are recognized by the mitochondrial basal transcriptional machinery, which is comprised of the mitochondrial RNA polymerase and mitochondrial transcription factors (mtTFAs) (Bonawitz et al., 2006; Scarpulla, 2008; Scarpulla et al., 2012). Mitochondrial RNAs are produced from polycistronic transcripts and undergo processing and modification to produce mature mRNA (polyadenylation), rRNA (nucleotide modification), and tRNA (addition of CCA). However, the abundance of each mRNA, rRNA, and tRNA is highly variable, possibly due to further post-transcriptional modifications that affect transcript processing, maturation, stability, and degradation (Rackham et al., 2012). Although mitochondrial transcription has been intensively studied, the mechanism of mitochondrial translation remains, for the most part, unknown (Bonawitz et al., 2006; Rackham et al., 2012).

The basal metabolic rate of resting hepatocytes is controlled by energy from non-mitochondrial processes (10%) and mitochondrial proton leak (20%); the greatest proportion of the metabolic rate is dependent on mitochondrial ATP production (70%) (Hulbert and Else, 2000). The maintenance of mitochondrial function requires a delicate balance between continuous protein synthesis and degradation, i.e., protein turnover. Recent metabolic labeling




Fig. 2. Location of putative GREs in the mitochondrial genome. The mitochondrial genome is circular and composed of heavy (H) and light (L) strands. The displacement loop (D-loop) is the major noncoding region of the genome and is a stable, triple-stranded structure that harbors the L-strand promoter (LSP), one of the H-strand promoters (HSPs), and the origin of replication of the H-strand (O_H) found in many mtDNA molecules *in vivo*. The major L-strand origin (O_L) is located in the “WANCY” cluster of tRNAs. The consensus GRE is 5'-GGTACAnnnTGTCT-3', wherein “n” indicates any base. The putative GREs in the mitochondrial genome are located in the D-loop and within the structural genes, and their role could be in transcription initiation. In the mouse mitochondrial genome, there are at least eight GRE-like elements. GRE 1 and 2 are found in the D-loop region and contain the mitochondrial promoters. These two GREs are involved in the regulation of mitochondrial transcription, and the same transcription factor (GR) regulates nuclear and mitochondrial gene transcription in a similar manner (Demonacos et al., 1996). The 12S rRNA gene and the tRNA^{Leu} gene contain GRE 3 and 4, respectively. Three putative GREs (GRE 5, 6, and 7) are located within the COX1 subunit, and GRE 8 is present within the COX3 gene (Demonacos et al., 1996; Sekeris, 1990). The putative GREs were arbitrarily named for the purpose of this review. tRNAs are denoted with capital letters for specific amino acids according to the standard one-letter nomenclature code. There are two tRNAs for serine and leucine, which are labeled S1 and S2 or L1 and L2, respectively (Bonawitz et al., 2006). COX, cytochrome c oxidase; GRE, glucocorticoid response element; ND, NADH dehydrogenase; TERM, specific termination site.

studies determined the turnover rates of at least 458 proteins in mouse cardiac and hepatic mitochondria; respective median rates of 0.0402 and 0.163 per day were calculated, corresponding to median half-lives of 17.2 and 4.26 days (Kim et al., 2012). The mitochondrial proteome shows significant heterogeneity among tissues, based on mitochondrial functional requirements (Johnson et al., 2007; Mootha et al., 2003); examples include enhancements of fatty acid oxidation for ATP production in the heart, specific aspects of the urea cycle and biosynthesis in liver, uncoupled respiration in adipose tissue, coupled and uncoupled respiration in skeletal muscle, metabolism of reabsorbed amino acids (e.g., tryptophan) in kidney, and specialized neurotransmitter metabolism and some aspects of reactive oxygen species (ROS) inactivation in brain. Thus, individual tissues have specific protein requirements that are provided by nuclear-encoded proteins.

4. GCs and mitochondrial biogenesis

Mitochondria are not produced *de novo*, but duplicate by a mechanism involving recruitment of proteins and replication of mtDNAs, which are added to preexisting subcompartments. Biogenesis and proliferation of mitochondria are influenced by external factors, such as nutrients, hormones, temperature, exercise, hypoxia, and aging (Ryan and Hoogenraad, 2007), which modify mitochondrial bioenergetics, thermogenesis, and oxidative stress and/or apoptotic responses (Manoli et al., 2007). Mitochondrial biogenesis and metabolism require the coordinated expression of multiple genes in two organelles, the nucleus and the mitochondrion. It has been proposed that, in the mitochondrial

respiratory chain, mtDNA-encoded proteins serve constitutive functions and nuclear-encoded proteins have regulatory roles (Kelly and Scarpulla, 2004; Scarpulla, 2012). However, changes in mitochondrial activity can alter nuclear gene expression, a process known as retrograde communication (Finley and Haigis, 2009). Depletion of mtDNA sends signals to the nucleus, which, in turn, compensates by modifying expression of multiple genes (Liu and Butow, 2006; Park et al., 2006). For example, cardiomyocytes with low mtDNA content and/or decreased mitochondrial membrane potential generate a complex nuclear response, possibly through Ca^{2+} signaling, which results in the alteration of signaling pathways involving the transcription factors calcineurin-dependent nuclear factor of activated T-cells cytoplasmic 1, c-Jun N-terminal kinase (JNK)-dependent ATF2, and nuclear factor-kappa B (NF- κ B), and enhances transcription of the gene encoding cytochrome c oxidase (COX) Vb (Biswas et al., 1999). In addition, cell proliferation and differentiation are closely related in terms of the status of mitochondria in myoblasts (Herzberg et al., 1993; Rochard et al., 2000). The regulation of mtDNA copy number, which is believed to be influenced, at least, by nuclear-encoded mtTFAs, is multifaceted, with different pathways influencing mtDNA replication and stability; and together, these pathways determine the specific amount and dynamic regulation of mtDNA *in vivo*. However, genetic mutation in mtDNA does not eventually induce mitochondrial biogenesis (Moraes et al., 1992). The contribution of mutations in mtDNA to carcinogenesis has been scrutinized using various approaches, but the conclusions remain controversial.

It is assumed that the main effect of GCs is the activation of nuclear-encoded genes, leading to enhanced mitochondrial

biogenesis and increased mitochondrial mass and respiration (Goffart and Wiesner, 2003). The impairment of GC production by adrenalectomy does not interfere *per se* with mitochondrial oxidation; rather, it alters the oxidative phosphorylation process in mitochondria and decreases ADP/O coefficient values and respiratory control (Kittas et al., 1984). In fetal adrenalectomized rats, COX mRNA level and mtDNA content were unchanged by GCs in heart, whereas mitochondrial maturation and differentiation were subject to the regulatory effect of GCs (Prieur et al., 1998). Mitochondrial biogenesis is sensitive to ROS (Scarpulla, 2012) and is precisely orchestrated by nuclear respiratory factors (NRFs) 1 and 2, mitochondrial transcription factors (TFAM A and B), and the integrative activity of coactivators, such as the peroxisome proliferator-activated receptor (PPAR)- γ coactivator 1 (PGC-1) family (PGC-1 α , PGC-1 β , and PRC) (Finley and Haigis, 2009; Scarpulla et al., 2012). GCs are essential for the rise in expression of oxidative phosphorylation enzymes and the development of the inner mitochondrial membrane, but do not regulate postnatal changes in mitochondrial density and mtDNA content in kidney or other tissues (Djouadi et al., 1994), with the exception of skeletal muscle (Weber et al., 2002).

5. GCs, the GR, and mitochondria

Circulating GC levels are regulated by CBG (Perogramvros et al., 2010). Tissue sensitivity can also be controlled by the enzyme

11 β -hydroxysteroid dehydrogenase 1 (converts inactive cortisone into active cortisol), 11 β -hydroxysteroid dehydrogenase 2 (a cortisol-degrading enzyme), or 5 α -reductase (converts GC into tetrahydrocortisol or alldihydrocortisol). It should be mentioned that 11-ketometabolites (cortisone, 11-dehydrocorticosterone) produced in degradative pathways have some additional effects, such as reducing the response to aldosterone (Odermatt et al., 2001), and some products of GC metabolism have the potential for activation (McInnes et al., 2004). The GR is a ligand-regulated transcription factor, located primarily in the cytoplasm in its inactive form as a multimeric complex (Fig. 3). The GR is also a target of multiple kinases, and post-translational modifications, including sumoylation and ubiquitination, can affect subcellular localization, interaction with other proteins, and degradation (Anbalagan et al., 2012; Ismaili and Garabedian, 2004). Several protein kinases phosphorylate the GR at specific sites. Phosphorylation of the GR may affect its transcriptional activation in response to hormone stability, receptor stability, subcellular localization, regulation of transcription, and interactions with co-regulators (Ismaili and Garabedian, 2004; Weigel and Moore, 2007). Mitogen-activated protein kinase (MAPK) phosphorylates the GR residues T171 and S246, whereas multiple cyclin-dependent kinase complexes modify S224 and S232 (Krstic et al., 1997). The GR undergoes cyclic phosphorylation by one of these kinases and dephosphorylation, by protein phosphatase 5, at three sites (S203, S221, and S226), which serves to maintain steady-state receptor phosphorylation at a low




Fig. 3. Functional modulation of the mitochondrion by GCs and the GR. GC binding to the GR releases GC–GR from the cytoplasmic inactive GR complex. ① Active GR translocates into the nucleus and alters expression (transactivation or transrepression) of nuclear genes encoding OXPHOS and other mitochondrial proteins. ② The GR influences the expression of transcription factors for control of nDNA-encoded mitochondrial proteins. ③ Mitochondrial regulatory factors from the nucleus control transcription and mitochondrial gene expression. After binding GC, the GR translocates into the mitochondrion and regulates mitochondrial gene expression via mGREs. Because the mitochondrial genome contains putative GREs, transcription of mitochondrial genes encoding the OXPHOS proteins and other proteins required for protein synthesis can be initiated or suppressed. This kind of direct action of the GR on the mitochondrial genome may be faster than that of nDNA-encoded regulatory proteins due to the time lag associated with expression and transport into the mitochondrion. ④ Components released from the cytoplasmic GR complex, a non-specific intracellular receptor, and the GR and a GC transporter in the plasma membrane rapidly initiate cytosolic signaling, leading to non-genomic and genomic activity in both nucleus and mitochondrion. ⑤ The GR can affect OXPHOS activity directly or indirectly through cellular signaling in mitochondria. In addition, ⑥ The mitochondrial products or other cellular messengers, including ADP, Ca²⁺, and NAD⁺, derived from mitochondria, can regulate nuclear gene expression (Liu and Butow, 2006). These changes eventually affect mitochondrial functions. Therefore, non-classical GC activities evoked in mitochondria, together with classical activities in the nucleus, add complexity to the physiological phenomenon. “+” and “–” indicate transactivation and transrepression of genes, respectively. Solid arrows and dashed arrows indicate possible genomic and non-genomic actions, respectively. GC, glucocorticoid; GR, glucocorticoid receptor; GRE, glucocorticoid response element; mGR, membrane-bound GR; mGRE, mitochondrial GRE; mOXPHOS, mitochondrial DNA-encoded OXPHOS; nDNA, nuclear DNA; nOXPHOS, nuclear DNA-encoded OXPHOS; nGRE, nuclear glucocorticoid response element; OXPHOS, oxidative phosphorylation system.

basal level in the absence of ligand. Inhibition of protein phosphatase affects retention of the GR in the nucleus and transcriptional activation (Wang et al., 2007; Zuo et al., 1999).

Detection of the GR in mitochondria stimulated research to elucidate its specific role in this organelle and broaden the understanding of the actions of GCs in mitochondria. The activity of the respiratory chain depends on the expression of hundreds of nuclear-encoded genes and 13 genes encoded by the mitochondrial genome. In addition, nuclear DNA (nDNA)-encoded mitochondrial proteins for protein import and assembly, such as the transporter inner membrane and transporter outer membrane complexes, are important factors in mitochondrial biogenesis or homeostasis (Tome et al., 2012). GCs have diverse genomic effects in the nucleus through regulation of transcription factors. Apart from nuclear-encoded mitochondrial transcriptional regulators, it was recently discovered that human and rat mitochondrial genomes contain sequences similar to that of the nuclear GREs (Demonacos et al., 1995; Gregory et al., 1976; Gruber et al., 2004). Among mtDNA genes, those encoding COX subunits I and III, 12S and 16S rRNA, NADH dehydrogenase subunit I, and tRNA^{Leu}, as well as the D-loop (promoter region) of the mitochondrial genome (Demonacos et al., 1995; Ioannou et al., 1988; Tsiriyotis et al., 1997), contain putative GREs (Fig. 2). Considering the presence of GREs in both the nucleus and mitochondria, it is reasonable to postulate that expression of mitochondrial components can be tightly controlled temporally. Possible mitochondrial functions influenced by the GR can be categorized as 1) transcriptional control of mitochondrial proteins encoded by nDNA, 2) transcriptional control of nuclear-encoded transcription factors affecting nDNA-encoded mitochondrial proteins, 3) transcriptional control of mitochondrial proteins encoded by mtDNA, 4) activation of cytosolic signaling proteins by plasma membrane receptors or cytosolic steroid receptors that subsequently affect the mitochondria, and 5) non-genomic effects of mitochondrial membrane-bound GR or GCs through cognate mitochondrial receptors and the interaction of these receptors with binding sites in the mitochondrial genome (Fig. 3). These categories are briefly discussed in a separate section. It should be mentioned that the activities of mitochondrial OXPHOS proteins are regulated by protein level, as well as by post-translational modifications; however, this aspect will not be considered here.

5.1. Transcriptional control of mitochondrial proteins encoded by nDNA: nucleus to mitochondria

In cardiomyocytes from chronically restraint-stressed rats, mitochondrial proteomic analysis revealed that differentially expressed proteins were all nuclear-encoded (i.e., carnitine palmitoyltransferase 2, mitochondrial acyl-CoA thioesterase 1, isocitrate dehydrogenase3 [NAD1] alpha, fumarate hydratase 1, pyruvate dehydrogenase beta, creatine kinase, and prohibitin) and, at the same time, mitochondrial respiration capacity was impaired (Liu et al., 2004). It has been reported that among the genes whose expression was increased (140 genes) or decreased (108 genes) by corticosterone treatment (Chen et al., 2005), there were 15 signaling molecules, eight transcription factors, three chromatin- or DNA-binding proteins, and 11 enzymes involved in energy production or nucleic acid, lipid, or steroid metabolism. Thus, these modulated genes can indirectly or directly influence mitochondrial function, but whether the effect enhances or suppresses mitochondrial function is not easily determined. The pyruvate dehydrogenase complex (PDC) is a key regulatory enzyme in the oxidation of glucose to acetyl-CoA. Pyruvate dehydrogenase kinase 4 (PDK4) phosphorylates PDC, thereby reducing its activity. GCs contribute to the initiation of PDK4 gene transcription in fasting and diabetes, reinforced in part by GREs in the distal promoter (Connaughton et al., 2010).

Expression of proteins in the respiratory apparatus relies largely on gene-specific regulatory mechanisms. Chromatin accessibility is a primary factor in GR binding to chromatin and may determine cell- and gene-specific activities of the GR (John et al., 2011). GC-mediated transactivation can be achieved by translocation of the GR into the nucleus followed by interaction with regulatory regions (e.g., GREs), whereas transrepression of target genes (Santos et al., 2011) can be accomplished by binding to a negative GRE, counteracting the transactivating effects of other general transcription factors, or repression of inflammatory responses without binding of GCs to the GR. It has been proposed that the GR interacts directly with NF- κ B or one of several AP-1 components (Schacke et al., 2002), and thereby suppresses the activity of these transcription factors. In addition, GRE tethering has been proposed, whereby, instead of binding directly to the DNA to achieve an effect, the GR binds to DNA-bound transcription factors in a regulatory complex to assist in transactivation or transrepression (Schoneveld et al., 2004). The action of GCs on RNA metabolism has been relatively well studied in the liver (Psarra and Sekeris, 2009), but little information is available for the heart or cardiovascular system. Regarding the inflammatory response, signaling pathways, including the extracellular signal-regulated kinase, JNK/SAPK, and P38/SAPK2 pathways, are involved in the control of mRNA turnover, and translation of mRNAs of proinflammatory proteins can be inhibited by GCs (Stellato, 2004). This raises the possibility that mitochondrial proteins can also be targets of GC–GR through modulation of mRNA stability or translation, although there is no direct evidence for this.

GCs may attenuate apoptosis caused by serum deprivation in rat hepatoma cells (Evans-Storms and Cidlowski, 2000). In addition, the GC agonist DEX can prevent cardiac injury caused by ischemia/reperfusion damage through transcriptional activation of the Bcl-xL gene, which prevents mitochondrial release of cytochrome c, in mouse heart (Xu et al., 2011). This GC-induced activation may be related to the presence of hormone response element-like sequences (e.g., GRE) in the promoter region of the Bcl-xL gene (Xu et al., 2011).

5.2. Transcriptional control of nDNA-encoded transcription factors that affect nDNA-encoded mitochondrial proteins

The ligand-activated GR- α is able to regulate gene expression without binding to GREs, perhaps by acting as a monomer to influence the capacity of other transcription factors to enhance or depress the transcription rates of target genes (Gottlicher et al., 1998; Nicolaides et al., 2010). The GR can also influence the expression of transcription factors controlling nDNA-encoded mitochondrial proteins. The general transcription factor Sp1 is crucial in the expression of a multitude of genes implicated in various constitutive and tissue-specific cellular activities (Suske, 1999). NRF-1 and NRF-2 play critical roles in transcription of nuclear-encoded mitochondrial proteins (Kelly and Scarpulla, 2004). NRF-1 is a 68-kD polypeptide with a C-terminal transcriptional activation domain comprised of glutamine-containing clusters of hydrophobic amino acid residues. NRF-1 participates in transcriptional control of genes encoding cytochrome c, proteins in five respiratory complexes, proteins involved in assembly of the respiratory apparatus, constituents of the mtDNA transcription and replication machinery, mitochondrial and cytosolic enzymes of the heme biosynthetic pathway, and proteins involved in mitochondrial protein import (Kelly and Scarpulla, 2004). However, regulation of most nuclear genes encoding mitochondrial enzymes upstream of the respiratory chain is not dependent on NRF-1/NRF-2 (Kelly and Scarpulla, 2004).

PGC-1 α , which serves as a master regulator in the transcriptional control of mitochondrial biogenesis and respiratory functions (Scarpulla, 2011), can be induced by GCs in liver tissue, potentiating expression of gluconeogenic genes. The actual interaction of mammalian PGC-1 α with the LBD of the GR serves as an effective stimulator of GR-mediated transcription in cells (Knutti et al., 2000). PGC-1 α also plays a role in thermogenesis by the activation of mitochondrial uncoupling protein-1 (UCP-1), which promotes mitochondrial uncoupled respiration in brown adipose tissue (Puigserver et al., 1998). In human tissues, expression of PGC-1 α is tissue-specific and at a high level in skeletal muscle, heart, kidney, and liver (Knutti et al., 2000). This suggests that the action of GCs can vary or is related to the differential responsiveness of GC–GR according to preexisting conditions in tissues. Nitric oxide (NO) can affect mitochondrial function directly or indirectly; PGC-1 α is modulated in a cGMP-dependent manner, as demonstrated by the observation that the mitochondrial thermogenic response is significantly altered in mice lacking endothelial nitric oxide synthase (eNOS) (Kelly and Scarpulla, 2004). GCs can rapidly induce NO production by activation of eNOS by a non-genomic mechanism (Hafezi-Moghadam et al., 2002); however, in the long term, NO production is decreased by reduction of levels of inducible NOS (Simmons et al., 1996) and eNOS (Akaike and Matsumoto, 2007; Duckles and Miller, 2010; Wallerath et al., 2004), eventually reducing the level of bioavailable NO. It should be noted that cellular and mitochondrial NO levels or antioxidative status (Psarra et al., 2009) can impair GC–GR action. Nitrosative stress, such as septic shock, can decrease binding of GC to its receptor as a result of S-nitrosylation of critical –SH groups, and thus suppresses GR-hsp90 heterocomplex dissociation (Galigniana et al., 1999).

5.3. Transcriptional control of mitochondrial proteins encoded by mtDNA

Mitochondrial gene expression is largely influenced by the abundance of mtDNA, transcriptional activity of mitochondrial RNA polymerase, activities of mitochondrial transcription factor B (TFB) and TFAM, and RNA stability and translation efficiency (Fernandez-Vizarra et al., 2008). The bulk of mitochondrial regulatory proteins is nuclear-encoded and recruited to the D-loop regulatory region of mtDNA. TFAM is an NRF-1 target gene and is required for mtDNA replication and maintenance. Increased NRF-1 and TFAM mRNAs in mtDNA-deficient cells or defective mitochondria due to lipopolysaccharide-induced oxidative damage might constitute a coping mechanism for cellular stress. PGC-1 α and TFAM are direct transcriptional coactivators of PPAR- γ in the mitochondrial biogenetic regulatory cascade (Puigserver and Spiegelman, 2003).

Recently, PPAR- γ and the heterodimeric transcription factor AP-1 has been identified in mammalian mitochondria. It is evident that binding of GR and AP-1 to the mitochondrial genome could potentially regulate mitochondrial gene expression (Leigh-Brown et al., 2010). For the GR to induce a genomic effect in mitochondria, translocation of the GR to mitochondria would be a prerequisite, but the underlying molecular mechanism remains unclear. It has been suggested that the GR may possess cryptic mitochondrial targeting signals in the C-terminal region, similar to the estrogen and androgen receptors (Psarra et al., 2005), although no known classical mitochondrial import sequences have been identified in the GR. It has also been noted that the GR/Bcl-2 complex is similar to estrogen receptors with regard to mitochondrial protein translocation via binding to heat shock protein 70/90 chaperone proteins (Du et al., 2009a). Direct activation of mitochondrial transcription by GCs has been proposed in various cell lines. In the presence of α -amanitin, a specific inhibitor of nuclear RNA polymerase II (Lindell et al., 1970), mitochondrial genes, such as those encoding COX I–III

and 12S and 18S rRNAs, but not the nuclear COX IV gene, are upregulated by GCs (Psarra and Sekeris, 2009). DNA–RNA hybridization analysis showed that hydrocortisone (cortisol) administration induces expression of all mitochondrial genes simultaneously in rat liver, and does not selectively enhance individual gene transcription (Minchenko, 1989). However, treatment with the GC analog DEX increased the transcript levels of COX II and III and 12S RNA in skeletal muscle, but not in the colon, liver, or kidney (Weber et al., 2002). In addition, several nuclear transcription factors, including AP-1, NF- κ B, cyclic-AMP response element-binding protein (CREB), T3 receptor p43 (activated by ligand binding), signal transducer and activator of transcription (STAT)3 (stimulated by growth hormone signaling), the estrogen receptor, and tumor suppressor p53 (activated in response to stress) were identified in mitochondria in various cell lines and tissues, with different localization (Psarra and Sekeris, 2009). Among these, mitochondrial STAT3 serves as a subunit of mitochondrial complex I, which directly regulates mitochondrial function via its effect on the electron transport chain (Wegrzyn et al., 2009). Mitochondrial p43 and CREB bind to mtDNA to regulate gene expression (Leigh-Brown et al., 2010). The presence of nuclear transcription factors in mitochondria and their interplay with GCs in regulation of mitochondrial gene expression should be further investigated to gain insight into the activities of GC and the GR.

5.4. Activation of cytosolic signaling proteins by plasma membrane or cytosolic steroid receptors

After GC binding, the multimeric GR complex undergoes a conformational change, generating a heteromeric complex and GR dimerization. Upon ligand activation, the GR is actively imported into the nucleus and mitochondria, where it influences gene regulation (Fig. 3). Apart from GC–GR, various components released from the cytosolic complex, including heat shock protein 90, heat shock protein 70, Src tyrosine kinase, serine/threonine protein phosphatase type 5 (Hinds and Sanchez, 2008), and heat shock protein 56 (Buttgereit et al., 2004; McMaster and Ray, 2007) can independently induce cell signaling (Fig. 3). The GR interacts with diverse cytosolic proteins, such as tubulin, actin, and other proteins involved in various signal transduction pathways (i.e., AP-1, NF- κ B, Stat5, 14-3-3, HOP, p23, FKBP51, FKBP52, and Raf-1) (Hedman et al., 2006). Although further investigation is required, 27 putative GR-interacting proteins were newly identified, including PP63 (a protease inhibitor and tyrosine kinase inhibitor), major vault protein, TATA-binding interacting protein 49a, transcription elongation factor B, and hypoxanthine-guanine phosphoribosyltransferase. Activation of cellular signaling by these proteins can produce rapid non-genomic action. In HepG2 hepatoma cells, DEX significantly suppressed the activities of mitochondrial complexes I and II, whereas complex III activity was enhanced by a p38 MAPK-dependent mechanism (Desquiret et al., 2008). These cell signaling events initiated by the GR can eventually impact mitochondrial function.

5.5. Non-genomic effect of GCs on mitochondria

In the cell, complete synthesis and post-translational modification of a protein requires 15 min to hours; nuclear translocation takes 10–30 min and transcription, 5–120 min (Haller et al., 2008). Thus, achievement of the GC-mediated genomic effect is a more time-consuming process than that required to effect non-genomic changes, which occur within several minutes in the absence of transcriptional changes (Hafezi-Moghadam et al., 2002; Haller et al., 2008). These rapidly-occurring non-genomic effects of steroids are established and have been extensively described (Billing

et al., 2012; Haller et al., 2008; Lee et al., 2012; Samarasinghe et al., 2011; Song and Buttgerit, 2006; Zhang et al., 2006; Zhou et al., 2011). Non-genomic effects include hormone–ligand activity involving nuclear receptor proteins in the cytoplasm, which are stimulated either at the cell surface within mitochondria or by binding to other cytoplasmic receptors. In isolated kidney and brain mitochondria, low doses of GCs cause a decrease in state 3 respiration, which may affect the activity of COX, F_0F_1 -ATPase, or complex IV (Katyare et al., 2003; Morin et al., 2000; Simon et al., 1998). Similar to what has been observed on the plasma membrane and nuclear pores (Losel and Wehling, 2003), GCs may directly modulate mitochondrial function by interpolating in the mitochondrial membrane, which may result in compromised ATP production due to increased proton leakage from the mitochondrion (Song and Buttgerit, 2006). In isolated rat brain mitochondria, a physiological dose of corticosterone non-specifically inhibits mitochondrial ATP production by suppressing electron transfer from NADH to the electron transfer chain through complex I (Fujita et al., 2009). Long-term administration of varying doses (100 nM–1 μ M) of corticosterone dynamically modulates major functional indicators of mitochondria, including mitochondrial oxidation, membrane potential, and mitochondrial Ca^{2+} level, in cortical neurons (Du et al., 2009b). These findings imply that GC can directly modulate mitochondrial functions, such as ATP production, but the possibility of GC action through the GR cannot be completely excluded. The synthetic GC DEX can directly bind and activate both conventional PKCs and atypical PKC isoforms, and this activation is not blocked by the GR antagonist RU38486 (Alzamora and Harvey, 2008). This direct and non-genomic activation of PKC signaling by GC binding may further influence mitochondrial activity.

As the result of a variety of apoptotic stimuli, mitochondrial membrane potential can be changed by genomic and non-genomic actions of the GR (Kfir-Erenfeld et al., 2010; Sionov et al., 2006). Recently, it has been suggested that novel ligand-induced translocation of GC into mitochondria can initiate the apoptotic cascade in thymocytes by decreasing mitochondrial transmembrane potential (Buttgerit et al., 2000; Palinkas et al., 2008) or interacting with mitochondrial proteins that regulate apoptosis (Boldizar et al., 2010; Psarra and Sekeris, 2009; Sionov et al., 2006). However, the exact mechanism underlying induction of the apoptotic cascade by the translocated GC–GR complex in mitochondria is still under investigation. The proposed mechanisms underlying GR-mediated apoptosis are 1) interaction with Bcl-2 family members (Thompson and Winoto, 2008), 2) promotion of Bax/Bak apoptotic oligomer formation by the GR (Brunelle and Letai, 2009), and 3) regulation of the activity of members of the pro-apoptotic Bcl-2 family of proteins via a genomic effect on Src tyrosine kinase (Herr et al., 2007).

GCs and the GR are responsible for multiple complex actions that are initiated both in the nucleus and mitochondria. The physiological status of cells and mitochondria play integral roles in determining the actions of GCs, rendering a complex and dynamic environment in which GC activity takes place (Pandya et al., 2007). In addition, the interactions of GR signaling pathways with other signaling pathways, and the existence of multiple coactivators contribute to an intricate, tightly regulated network, and represent phenomena central to gaining an understanding of hormone responsiveness and resistance.

6. Concluding remarks

GC–GR interactions modulate expression of OXPHOS and mitochondrial proteins directly and indirectly. In the present review, we have briefly outlined the specific roles of GCs and their receptors in mitochondria. Mitochondrial genes are under the tight

control of nuclear-encoded structural and transcriptional regulatory proteins. However, the translocation of the GR into mitochondria and the existence of GRE-like elements in the mitochondrial genome provide evidence of highly ordered behavior of GCs and the GR in these organelles. In addition, the biological effects of GC–GR interactions vary, but are highly dependent on ligand specificity and availability, which, in turn, are influenced by the physiological status of the cell and mitochondria. For appropriate regulation of mitochondrial activities, the mitochondria should send signals to the nucleus indicating the nature of the stimuli so that the cell can adapt and respond accordingly. Understanding the mechanisms underlying retrograde signaling between mitochondria and the nucleus following the GC–GR response and the regulatory roles of GCs in mitochondria will be valuable for understanding GC activity and necessary for basic and clinical application of GCs.

Conflict of Interest

The authors declare no conflicts of interest.

Acknowledgments

This work was supported by the Priority Research Centers Program (2010-0020224) and the Basic Science Research Program (2010-0025855 and 2012R1A2A1A03007595) through the National Research Foundation of Korea (NRF), funded by the Ministry of Education, Science, and Technology.

References

- Adcock, I.M., 2000. Molecular mechanisms of glucocorticosteroid actions. *Pulm. Pharmacol. Ther.* 13, 115–126.
- Akaike, M., Matsumoto, T., 2007. [Glucocorticoid-induced reduction in NO bioavailability and vascular endothelial dysfunction]. *Clin. Calcium* 17, 864–870.
- Alzamora, R., Harvey, B.J., 2008. Direct binding and activation of protein kinase C isoforms by steroid hormones. *Steroids* 73, 885–888.
- Anbalagan, M., Huderson, B., Murphy, L., Rowan, B.G., 2012. Post-translational modifications of nuclear receptors and human disease. *Nucl. Recept. Signal.* 10, e001.
- Barnes, P.J., 2011. Glucocorticosteroids: current and future directions. *Br. J. Pharmacol.* 163, 29–43.
- Biddie, S.C., Conway-Campbell, B.L., Lightman, S.L., 2012. Dynamic regulation of glucocorticoid signalling in health and disease. *Rheumatology (Oxf.)* 51, 403–412.
- Billing, A.M., Revets, D., Hoffmann, C., Turner, J.D., Vernocchi, S., Muller, C.P., 2012. Proteomic profiling of rapid non-genomic and concomitant genomic effects of acute restraint stress on rat thymocytes. *J. Proteomics* 75, 2064–2079.
- Biswas, G., Adebajo, O.A., Freedman, B.D., Anandatheerthavarada, H.K., Vijayarath, C., Zaidi, M., Kotlikoff, M., Avadhani, N.G., 1999. Retrograde Ca^{2+} signaling in C_2C_{12} skeletal myocytes in response to mitochondrial genetic and metabolic stress: a novel mode of inter-organelle crosstalk. *EMBO J.* 18, 522–533.
- Bogenhagen, D., Clayton, D.A., 1974. The number of mitochondrial deoxyribonucleic acid genomes in mouse L and human HeLa cells. Quantitative isolation of mitochondrial deoxyribonucleic acid. *J. Biol. Chem.* 249, 7991–7995.
- Boldizar, F., Talaber, G., Szabo, M., Bartis, D., Palinkas, L., Nemeth, P., Berki, T., 2010. Emerging pathways of non-genomic glucocorticoid (GC) signalling in T cells. *Immunobiology* 215, 521–526.
- Bonawitz, N.D., Clayton, D.A., Shadel, G.S., 2006. Initiation and beyond: multiple functions of the human mitochondrial transcription machinery. *Mol. Cell* 24, 813–825.
- Brillon, D.J., Zheng, B., Campbell, R.G., Matthews, D.E., 1995. Effect of cortisol on energy expenditure and amino acid metabolism in humans. *Am. J. Physiol.* 268, E501–E513.
- Brunelle, J.K., Letai, A., 2009. Control of mitochondrial apoptosis by the Bcl-2 family. *J. Cell Sci.* 122, 437–441.
- Burke, C.W., 1985. Adrenocortical insufficiency. *Clin. Endocrinol. Metab.* 14, 947–976.
- Buttgerit, F., Burmester, G.R., Brand, M.D., 2000. Bioenergetics of immune functions: fundamental and therapeutic aspects. *Immunol. Today* 21, 192–199.
- Buttgerit, F., Straub, R.H., Wehling, M., Burmester, G.R., 2004. Glucocorticoids in the treatment of rheumatic diseases: an update on the mechanisms of action. *Arthritis Rheum.* 50, 3408–3417.
- Charmandari, E., Kino, T., 2007. Novel causes of generalized glucocorticoid resistance. *Horm. Metab. Res.* 39, 445–450.

- Chen, Q.M., Alexander, D., Sun, H., Xie, L., Lin, Y., Terrand, J., Morrissey, S., Purdom, S., 2005. Corticosteroids inhibit cell death induced by doxorubicin in cardiomyocytes: induction of antiapoptosis, antioxidant, and detoxification genes. *Mol. Pharmacol.* 67, 1861–1873.
- Chevallet, M., Lescuyer, P., Diemer, H., van Dorsselaer, A., Leize-Wagner, E., Rabilloud, T., 2006. Alterations of the mitochondrial proteome caused by the absence of mitochondrial DNA: a proteomic view. *Electrophoresis* 27, 1574–1583.
- Chrousos, G.P., Kino, T., 2005. Intracellular glucocorticoid signaling: a formerly simple system turns stochastic. *Sci. STKE* 2005, pe48.
- Chrousos, G.P., Kino, T., 2007. Glucocorticoid action networks and complex psychiatric and/or somatic disorders. *Stress* 10, 213–219.
- Cole, T.J., Blendy, J.A., Monaghan, A.P., Kriegstein, K., Schmid, W., Aguzzi, A., Fantuzzi, G., Hummler, E., Unsicker, K., Schutz, G., 1995. Targeted disruption of the glucocorticoid receptor gene blocks adrenergic chromaffin cell development and severely retards lung maturation. *Genes Dev.* 9, 1608–1621.
- Cole, T.J., Harris, H.J., Hoong, I., Solomon, N., Smith, R., Krozowski, Z., Fullerton, M.J., 1999. The glucocorticoid receptor is essential for maintaining basal and dexamethasone-induced repression of the murine corticosteroid-binding globulin gene. *Mol. Cell. Endocrinol.* 154, 29–36.
- Cole, T.J., Myles, K., Purton, J.F., Brereton, P.S., Solomon, N.M., Godfrey, D.I., Funder, J.W., 2001. GRKO mice express an aberrant dexamethasone-binding glucocorticoid receptor, but are profoundly glucocorticoid resistant. *Mol. Cell. Endocrinol.* 173, 193–202.
- Connaughton, S., Chowdhury, F., Attia, R.R., Song, S., Zhang, Y., Elam, M.B., Cook, G.A., Park, E.A., 2010. Regulation of pyruvate dehydrogenase kinase isoform 4 (PDK4) gene expression by glucocorticoids and insulin. *Mol. Cell. Endocrinol.* 315, 159–167.
- Dallman, M.F., Strack, A.M., Akana, S.F., Bradbury, M.J., Hanson, E.S., Scribner, K.A., Smith, M., 1993. Feast and famine: critical role of glucocorticoids with insulin in daily energy flow. *Front. Neuroendocrinol.* 14, 303–347.
- Demonacos, C., Djordjevic-Markovic, R., Tsawdaroglou, N., Sekeris, C.E., 1995. The mitochondrion as a primary site of action of glucocorticoids: the interaction of the glucocorticoid receptor with mitochondrial DNA sequences showing partial similarity to the nuclear glucocorticoid responsive elements. *J. Steroid Biochem. Mol. Biol.* 55, 43–55.
- Demonacos, C., Tsawdaroglou, N.C., Djordjevic-Markovic, R., Papalopoulou, M., Galanopoulos, V., Papadogeorgaki, S., Sekeris, C.E., 1993. Import of the glucocorticoid receptor into rat liver mitochondria in vivo and in vitro. *J. Steroid Biochem. Mol. Biol.* 46, 401–413.
- Demonacos, C.V., Karayanni, N., Hatzoglou, E., Tsiroyiotis, C., Spandidos, D.A., Sekeris, C.E., 1996. Mitochondrial genes as sites of primary action of steroid hormones. *Steroids* 61, 226–232.
- Desquiret, V., Gueguen, N., Malthiery, Y., Ritz, P., Simard, G., 2008. Mitochondrial effects of dexamethasone imply both membrane and cytosolic-initiated pathways in HepG2 cells. *Int. J. Biochem. Cell Biol.* 40, 1629–1641.
- Djouadi, F., Bastin, J., Gilbert, T., Rotig, A., Rustin, P., Merlet-Benichou, C., 1994. Mitochondrial biogenesis and development of respiratory chain enzymes in kidney cells: role of glucocorticoids. *Am. J. Physiol.* 267, C245–C254.
- Du, J., McEwen, B., Manji, H.K., 2009a. Glucocorticoid receptors modulate mitochondrial function: a novel mechanism for neuroprotection. *Commun. Integr. Biol.* 2, 350–352.
- Du, J., Wang, Y., Hunter, R., Wei, Y., Blumenthal, R., Falke, C., Khairova, R., Zhou, R., Yuan, P., Machado-Vieira, R., McEwen, B.S., Manji, H.K., 2009b. Dynamic regulation of mitochondrial function by glucocorticoids. *Proc. Natl. Acad. Sci. U.S.A.* 106, 3543–3548.
- Duchene, A.M., Giege, P., 2012. Dual localized mitochondrial and nuclear proteins as gene expression regulators in plants? *Front. Plant Sci.* 3, 221.
- Duckles, S.P., Miller, V.M., 2010. Hormonal modulation of endothelial NO production. *Pflügers Arch.* 459, 841–851.
- Evans-Storrs, R.B., Cidlowski, J.A., 2000. Delineation of an antiapoptotic action of glucocorticoids in hepatoma cells: the role of nuclear factor-kappaB. *Endocrinology* 141, 1854–1862.
- Fernandez-Vizcarra, E., Enriquez, J.A., Perez-Martos, A., Montoya, J., Fernandez-Silva, P., 2008. Mitochondrial gene expression is regulated at multiple levels and differentially in the heart and liver by thyroid hormones. *Curr. Genet.* 54, 13–22.
- Finley, L.W., Haigis, M.C., 2009. The coordination of nuclear and mitochondrial communication during aging and calorie restriction. *Ageing Res. Rev.* 8, 173–188.
- Fosslien, E., 2001. Mitochondrial medicine—molecular pathology of defective oxidative phosphorylation. *Ann. Clin. Lab. Sci.* 31, 25–67.
- Frazier, A.E., Kiu, C., Stojanovski, D., Hoogenraad, N.J., Ryan, M.T., 2006. Mitochondrial morphology and distribution in mammalian cells. *Biol. Chem.* 387, 1551–1558.
- Fujita, C., Ichikawa, F., Teratani, T., Murakami, G., Okada, T., Shinohara, M., Kawato, S., Ohta, Y., 2009. Direct effects of corticosterone on ATP production by mitochondria from immortalized hypothalamic GT1-7 neurons. *J. Steroid Biochem. Mol. Biol.* 117, 50–55.
- Gagliardi, L., Ho, J.T., Torpy, D.J., 2010. Corticosteroid-binding globulin: the clinical significance of altered levels and heritable mutations. *Mol. Cell. Endocrinol.* 316, 24–34.
- Galigniana, M.D., Piwien-Pilipuk, G., Assreuy, J., 1999. Inhibition of glucocorticoid receptor binding by nitric oxide. *Mol. Pharmacol.* 55, 317–323.
- Garesse, R., Vallejo, C.G., 2001. Animal mitochondrial biogenesis and function: a regulatory cross-talk between two genomes. *Gene* 263, 1–16.
- Gavrilova-Jordan, L.P., Price, T.M., 2007. Actions of steroids in mitochondria. *Semin. Reprod. Med.* 25, 154–164.
- Goffart, S., Wiesner, R.J., 2003. Regulation and co-ordination of nuclear gene expression during mitochondrial biogenesis. *Exp. Physiol.* 88, 33–40.
- Goodwin, J.E., Zhang, J., Gonzalez, D., Albinsson, S., Geller, D.S., 2011. Knockout of the vascular endothelial glucocorticoid receptor abrogates dexamethasone-induced hypertension. *J. Hypertens.* 29, 1347–1356.
- Gottlicher, M., Heck, S., Herrlich, P., 1998. Transcriptional cross-talk, the second mode of steroid hormone receptor action. *J. Mol. Med. (Berl.)* 76, 480–489.
- Green, D.R., Kroemer, G., 2004. The pathophysiology of mitochondrial cell death. *Science* 305, 626–629.
- Gregory, M.C., Duval, D., Meyer, P., 1976. Changes in cardiac and hepatic glucocorticoid receptors after adrenalectomy. *Clin. Sci. Mol. Med.* 51, 487–493.
- Gruber, C.J., Gruber, D.M., Gruber, I.M., Wieser, F., Huber, J.C., 2004. Anatomy of the estrogen response element. *Trends Endocrinol. Metab.* 15, 73–78.
- Hafezi-Moghadam, A., Simoncini, T., Yang, Z., Limbourg, F.P., Plumier, J.C., Rebsamen, M.C., Hsieh, C.M., Chui, D.S., Thomas, K.L., Prorock, A.J., Laubach, V.E., Moskowitz, M.A., French, B.A., Ley, K., Liao, J.K., 2002. Acute cardiovascular protective effects of corticosteroids are mediated by non-transcriptional activation of endothelial nitric oxide synthase. *Nat. Med.* 8, 473–479.
- Haller, J., Mikics, E., Makara, G.B., 2008. The effects of non-genomic glucocorticoid mechanisms on bodily functions and the central neural system. A critical evaluation of findings. *Front. Neuroendocrinol.* 29, 273–291.
- Hammes, S.R., Levin, E.R., 2011. Minireview: Recent advances in extranuclear steroid receptor actions. *Endocrinology* 152, 4489–4495.
- Hedman, E., Widen, C., Asadi, A., Dinnetz, I., Schroder, W.P., Gustafsson, J.A., Wikstrom, A.C., 2006. Proteomic identification of glucocorticoid receptor interacting proteins. *Proteomics* 6, 3114–3126.
- Herr, I., Gassler, N., Friess, H., Buchler, M.W., 2007. Regulation of differential pro- and anti-apoptotic signaling by glucocorticoids. *Apoptosis* 12, 271–291.
- Herzberg, N.H., Zwart, R., Wolterman, R.A., Ruiter, J.P., Wanders, R.J., Bolhuis, P.A., van den Bogert, C., 1993. Differentiation and proliferation of respiration-deficient human myoblasts. *Biochim. Biophys. Acta* 1181, 63–67.
- Hinds Jr., T.D., Sanchez, E.R., 2008. Protein phosphatase 5. *Int. J. Biochem. Cell Biol.* 40, 2358–2362.
- Hulbert, A.J., Else, P.L., 2000. Mechanisms underlying the cost of living in animals. *Annu. Rev. Physiol.* 62, 207–235.
- Ioannou, I.M., Tsawdaroglou, N., Sekeris, C.E., 1988. Presence of glucocorticoid responsive elements in the mitochondrial genome. *Anticancer Res.* 8, 1405–1409.
- Ismaili, N., Garabedian, M.J., 2004. Modulation of glucocorticoid receptor function via phosphorylation. *Ann. N. Y. Acad. Sci.* 1024, 86–101.
- Jeon, J., Jeong, J.H., Baek, J.H., Koo, H.J., Park, W.H., Yang, J.S., Yu, M.H., Kim, S., Pak, Y.K., 2011. Network clustering revealed the systemic alterations of mitochondrial protein expression. *PLoS Comput. Biol.* 7, e1002093.
- Jeong, S.Y., Seol, D.W., 2008. The role of mitochondria in apoptosis. *BMB Rep.* 41, 11–22.
- John, S., Sabo, P.J., Thurman, R.E., Sung, M.H., Biddie, S.C., Johnson, T.A., Hager, G.L., Stamatoyannopoulos, J.A., 2011. Chromatin accessibility pre-determines glucocorticoid receptor binding patterns. *Nat. Genet.* 43, 264–268.
- Johnson, D.T., Harris, R.A., French, S., Blair, P.V., You, J., Bemis, K.G., Wang, M., Balaban, R.S., 2007. Tissue heterogeneity of the mammalian mitochondrial proteome. *Am. J. Physiol. Cell Physiol.* 292, C689–C697.
- Katyare, S.S., Balasubramanian, S., Parmar, D.V., 2003. Effect of corticosterone treatment on mitochondrial oxidative energy metabolism in developing rat brain. *Exp. Neurol.* 183, 241–248.
- Kelly, D.P., Scarpulla, R.C., 2004. Transcriptional regulatory circuits controlling mitochondrial biogenesis and function. *Genes Dev.* 18, 357–368.
- Kfir-Erenfeld, S., Sionov, R.V., Spokoini, R., Cohen, O., Yefenof, E., 2010. Protein kinase networks regulating glucocorticoid-induced apoptosis of hematopoietic cancer cells: fundamental aspects and practical considerations. *Leuk. Lymphoma* 51, 1968–2005.
- Kim, T.Y., Wang, D., Kim, A.K., Lau, E., Lin, A.J., Liem, D.A., Zhang, J., Zong, N.C., Lam, M.P., Ping, P., 2012. Metabolic labeling reveals proteome dynamics of mouse mitochondria. *Mol. Cell. Proteomics* 11, 1586–1594.
- Kittas, C., Chaniotis, F.J., Antonakopoulos, G., Papacharalampous, N.X., 1984. The effects of bilateral adrenalectomy on the fine structure of the rat myocardium. *Zentralbl. Allg. Pathol.* 129, 225–229.
- Knutti, D., Kaul, A., Kralli, A., 2000. A tissue-specific coactivator of steroid receptors, identified in a functional genetic screen. *Mol. Cell. Biol.* 20, 2411–2422.
- Kozlov, A.V., Bahrami, S., Calzia, E., Dangel, P., Gille, L., Kuznetsov, A.V., Troppmair, J., 2011. Mitochondrial dysfunction and biogenesis: do ICU patients die from mitochondrial failure? *Ann. Intensive Care* 1, 41.
- Krett, N.L., Pillay, S., Moalli, P.A., Greipp, P.R., Rosen, S.T., 1995. A variant glucocorticoid receptor messenger RNA is expressed in multiple myeloma patients. *Cancer Res.* 55, 2727–2729.
- Krstic, M.D., Rogatsky, I., Yamamoto, K.R., Garabedian, M.J., 1997. Mitogen-activated and cyclin-dependent protein kinases selectively and differentially modulate transcriptional enhancement by the glucocorticoid receptor. *Mol. Cell. Biol.* 17, 3947–3954.
- Lee, S.R., Kim, H.K., Youm, J.B., Dizon, L.A., Song, I.S., Jeong, S.H., Seo, D.Y., Ko, K.S., Rhee, B.D., Kim, N., Han, J., 2012. Non-genomic effect of glucocorticoids on cardiovascular system. *Pflügers Arch.* 464, 549–559.
- Leigh-Brown, S., Enriquez, J.A., Odom, D.T., 2010. Nuclear transcription factors in mammalian mitochondria. *Genome Biol.* 11, 215.
- Lesnefsky, E.J., Moghaddas, S., Tandler, B., Kerner, J., Hoppel, C.L., 2001. Mitochondrial dysfunction in cardiac disease: ischemia–reperfusion, aging, and heart failure. *J. Mol. Cell. Cardiol.* 33, 1065–1089.
- Lindell, T.J., Weinberg, F., Morris, P.W., Roeder, R.G., Rutter, W.J., 1970. Specific inhibition of nuclear RNA polymerase II by alpha-amanitin. *Science* 170, 447–449.

- Liu, Z., Butow, R.A., 2006. Mitochondrial retrograde signaling. *Annu. Rev. Genet.* 40, 159–185.
- Liu, X.H., Qian, L.J., Gong, J.B., Shen, J., Zhang, X.M., Qian, X.H., 2004. Proteomic analysis of mitochondrial proteins in cardiomyocytes from chronic stressed rat. *Proteomics* 4, 3167–3176.
- Losel, R., Wehling, M., 2003. Nongenomic actions of steroid hormones. *Nat. Rev. Mol. Cell Biol.* 4, 46–56.
- Lu, N.Z., Cidlowski, J.A., 2004. The origin and functions of multiple human glucocorticoid receptor isoforms. *Ann. N. Y. Acad. Sci.* 1024, 102–123.
- Lu, N.Z., Cidlowski, J.A., 2005. Translational regulatory mechanisms generate N-terminal glucocorticoid receptor isoforms with unique transcriptional target genes. *Mol. Cell Biol.* 18, 331–342.
- Manoli, I., Alesci, S., Blackman, M.R., Su, Y.A., Rennert, O.M., Chrousos, G.P., 2007. Mitochondria as key components of the stress response. *Trends Endocrinol. Metab.* 18, 190–198.
- Martinus, R.D., Garth, G.P., Webster, T.L., Cartwright, P., Naylor, D.J., Hoj, P.B., Hoogenraad, N.J., 1996. Selective induction of mitochondrial chaperones in response to loss of the mitochondrial genome. *Eur. J. Biochem.* 240, 98–103.
- McInnes, K.J., Kenyon, C.J., Chapman, K.E., Livingstone, D.E., Macdonald, L.J., Walker, B.R., Andrew, R., 2004. 5 α -reduced glucocorticoids, novel endogenous activators of the glucocorticoid receptor. *J. Biol. Chem.* 279, 22908–22912.
- McMaster, A., Ray, D.W., 2007. Modelling the glucocorticoid receptor and producing therapeutic agents with anti-inflammatory effects but reduced side-effects. *Exp. Physiol.* 92, 299–309.
- Miller, W.L., 2011. Role of mitochondria in steroidogenesis. *Endocr. Dev.* 20, 1–19.
- Minchenko, A.G., 1989. The time-dependent stimulating action of hydrocortisone on mitochondrial RNA biosynthesis in the rat liver. *Probl. Endocrinol. (Mosk.)* 35, 72–77.
- Moalli, P.A., Pillay, S., Krett, N.L., Rosen, S.T., 1993. Alternatively spliced glucocorticoid receptor messenger RNAs in glucocorticoid-resistant human multiple myeloma cells. *Cancer Res.* 53, 3877–3879.
- Montoya, J., Christianson, T., Levens, D., Rabinowitz, M., Attardi, G., 1982. Identification of initiation sites for heavy-strand and light-strand transcription in human mitochondrial DNA. *Proc. Natl. Acad. Sci. U.S.A.* 79, 7195–7199.
- Mootha, V.K., Bunkenborg, J., Olsen, J.V., Hjerrild, M., Wisniewski, J.R., Stahl, E., Bolouri, M.S., Ray, H.N., Sihag, S., Kamal, M., Patterson, N., Lander, E.S., Mann, M., 2003. Integrated analysis of protein composition, tissue diversity, and gene regulation in mouse mitochondria. *Cell* 115, 629–640.
- Moraes, C.T., Ricci, E., Bonilla, E., DiMauro, S., Schon, E.A., 1992. The mitochondrial tRNA(Leu)(UUR) mutation in mitochondrial encephalomyopathy, lactic acidosis, and stroke-like episodes (MELAS): genetic, biochemical, and morphological correlations in skeletal muscle. *Am. J. Hum. Genet.* 50, 934–949.
- Morin, C., Zini, R., Simon, N., Charbonnier, P., Tillement, J.P., Le Louet, H., 2000. Low glucocorticoid concentrations decrease oxidative phosphorylation of isolated rat brain mitochondria: an additional effect of dexamethasone. *Fundam. Clin. Pharmacol.* 14, 493–500.
- Nicolaides, N.C., Galata, Z., Kino, T., Chrousos, G.P., Charmandari, E., 2010. The human glucocorticoid receptor: molecular basis of biologic function. *Steroids* 75, 1–12.
- Oakley, R.H., Cidlowski, J.A., 2011. Cellular processing of the glucocorticoid receptor gene and protein: new mechanisms for generating tissue-specific actions of glucocorticoids. *J. Biol. Chem.* 286, 3177–3184.
- Odermatt, A., Arnold, P., Frey, F.J., 2001. The intracellular localization of the mineralocorticoid receptor is regulated by 11 β -hydroxysteroid dehydrogenase type 2. *J. Biol. Chem.* 276, 28484–28492.
- Palinkas, L., Talaber, G., Boldizsar, F., Bartis, D., Nemeth, P., Berki, T., 2008. Developmental shift in TCR-mediated rescue of thymocytes from glucocorticoid-induced apoptosis. *Immunobiology* 213, 39–50.
- Pandya, J.D., Agarwal, N.A., Katiyar, S.S., 2007. Dexamethasone treatment differentially affects the oxidative energy metabolism of rat brain mitochondria in developing and adult animals. *Int. J. Dev. Neurosci.* 25, 309–316.
- Park, S.Y., Lee, S., Park, K.S., Lee, H.K., Lee, W., 2006. Proteomic analysis of cellular change involved in mitochondria-to-nucleus communication in L6 GLUT4myc myocytes. *Proteomics* 6, 1210–1222.
- Perogamvros, I., Underhill, C., Henley, D.E., Hadfield, K.D., Newman, W.G., Ray, D.W., Lightman, S.L., Hammond, G.L., Trainer, P.J., 2010. Novel corticosteroid-binding globulin variant that lacks steroid binding activity. *J. Clin. Endocrinol. Metab.* 95, E142–E150.
- Prieur, B., Bismuth, J., Delaval, E., 1998. Effects of adrenal steroid hormones on mitochondrial maturation during the late fetal period. *Eur. J. Biochem.* 252, 194–199.
- Psarra, A.M., Hermann, S., Panayotou, G., Spyrou, G., 2009. Interaction of mitochondrial thioredoxin with glucocorticoid receptor and NF- κ B modulates glucocorticoid receptor and NF- κ B signalling in HEK-293 cells. *Biochem. J.* 422, 521–531.
- Psarra, A.M., Sekeris, C.E., 2009. Glucocorticoid receptors and other nuclear transcription factors in mitochondria and possible functions. *Biochim. Biophys. Acta* 1787, 431–436.
- Psarra, A.M., Sekeris, C.E., 2011. Glucocorticoids induce mitochondrial gene transcription in HepG2 cells: role of the mitochondrial glucocorticoid receptor. *Biochim. Biophys. Acta* 1813, 1814–1821.
- Psarra, A.M., Solakidi, S., Trougakos, I.P., Margaritis, L.H., Spyrou, G., Sekeris, C.E., 2005. Glucocorticoid receptor isoforms in human hepatocarcinoma HepG2 and SaOS-2 osteosarcoma cells: presence of glucocorticoid receptor alpha in mitochondria and of glucocorticoid receptor beta in nucleoli. *Int. J. Biochem. Cell Biol.* 37, 2544–2558.
- Puigserver, P., Spiegelman, B.M., 2003. Peroxisome proliferator-activated receptor-gamma coactivator 1 alpha (PGC-1 alpha): transcriptional coactivator and metabolic regulator. *Endocr. Rev.* 24, 78–90.
- Puigserver, P., Wu, Z., Park, C.W., Graves, R., Wright, M., Spiegelman, B.M., 1998. A cold-inducible coactivator of nuclear receptors linked to adaptive thermogenesis. *Cell* 92, 829–839.
- Pujols, L., Mullol, J., Roca-Ferrer, J., Torrego, A., Xaubet, A., Cidlowski, J.A., Picado, C., 2002. Expression of glucocorticoid receptor alpha- and beta-isoforms in human cells and tissues. *Am. J. Physiol. Cell Physiol.* 283, C1324–C1331.
- Rackham, O., Mercer, T.R., Filipovska, A., 2012. The human mitochondrial transcriptome and the RNA-binding proteins that regulate its expression. *Wiley Interdiscip. Rev. RNA* 3, 675–695.
- Reichardt, H.M., Tuckermann, J.P., Gottlicher, M., Vujic, M., Weih, F., Angel, P., Herrlich, P., Schütz, G., 2001. Repression of inflammatory responses in the absence of DNA binding by the glucocorticoid receptor. *EMBO J.* 20, 7168–7173.
- Rivers, C., Levy, A., Hancock, J., Lightman, S., Norman, M., 1999. Insertion of an amino acid in the DNA-binding domain of the glucocorticoid receptor as a result of alternative splicing. *J. Clin. Endocrinol. Metab.* 84, 4283–4286.
- Rochard, P., Rodier, A., Casas, F., Cassar-Malek, I., Marchal-Victorion, S., Daury, L., Wrutniak, C., Cabello, G., 2000. Mitochondrial activity is involved in the regulation of myoblast differentiation through myogenin expression and activity of myogenic factors. *J. Biol. Chem.* 275, 2733–2744.
- Rosol, T.J., Yarrington, J.T., Latendresse, J., Capen, C.C., 2001. Adrenal gland: structure, function, and mechanisms of toxicity. *Toxicol. Pathol.* 29, 41–48.
- Ryan, M.T., Hoogenraad, N.J., 2007. Mitochondrial-nuclear communications. *Annu. Rev. Biochem.* 76, 701–722.
- Sainte-Marie, Y., Nguyen Dinh Cat, A., Perrier, R., Mangin, L., Soukaseum, C., Peuchmaur, M., Tronche, F., Farman, N., Escoubet, B., Benitah, J.P., Jaissier, F., 2007. Conditional glucocorticoid receptor expression in the heart induces atrioventricular block. *FASEB J.* 21, 3133–3141.
- Samarasinghe, R.A., Di Maio, R., Volonte, D., Galbiati, F., Lewis, M., Romero, G., DeFranco, D.B., 2011. Nongenomic glucocorticoid receptor action regulates gap junction intercellular communication and neural progenitor cell proliferation. *Proc. Natl. Acad. Sci. U.S.A.* 108, 16657–16662.
- Santos, G.M., Fairall, L., Schwabe, J.W., 2011. Negative regulation by nuclear receptors: a plethora of mechanisms. *Trends Endocrinol. Metab.* 22, 87–93.
- Savory, J.G., Hsu, B., Laquian, I.R., Giffin, W., Reich, T., Hache, R.J., Lefebvre, Y.A., 1999. Discrimination between NL1- and NL2-mediated nuclear localization of the glucocorticoid receptor. *Mol. Cell Biol.* 19, 1025–1037.
- Scarpulla, R.C., 2008. Transcriptional paradigms in mammalian mitochondrial biogenesis and function. *Physiol. Rev.* 88, 611–638.
- Scarpulla, R.C., 2011. Metabolic control of mitochondrial biogenesis through the PGC-1 family regulatory network. *Biochim. Biophys. Acta* 1813, 1269–1278.
- Scarpulla, R.C., 2012. Nucleus-encoded regulators of mitochondrial function: integration of respiratory chain expression, nutrient sensing and metabolic stress. *Biochim. Biophys. Acta* 1819, 1088–1097.
- Scarpulla, R.C., Vega, R.B., Kelly, D.P., 2012. Transcriptional integration of mitochondrial biogenesis. *Trends Endocrinol. Metab.* 23, 459–466.
- Schacke, H., Docke, W.D., Asadullah, K., 2002. Mechanisms involved in the side effects of glucocorticoids. *Pharmacol. Ther.* 96, 23–43.
- Schoneveld, O.J., Gaemers, I.C., Lamers, W.H., 2004. Mechanisms of glucocorticoid signalling. *Biochim. Biophys. Acta* 1680, 114–128.
- Sekeris, C.E., 1990. The mitochondrial genome: a possible primary site of action of steroid hormones. *In Vivo* 4, 317–320.
- Simmons, W.W., Ungureanu-Longrois, D., Smith, G.K., Smith, T.W., Kelly, R.A., 1996. Glucocorticoids regulate inducible nitric oxide synthase by inhibiting tetrahydrobiopterin synthesis and L-arginine transport. *J. Biol. Chem.* 271, 23928–23937.
- Simoes, D.C.M., Psarra, A.-M.G., Mauad, T., Pantou, I., Roussos, C., Sekeris, C.E., Gratziau, C., 2012. Glucocorticoid and estrogen receptors are reduced in mitochondria of lung epithelial cells in asthma. *PLoS ONE* 7, e39183.
- Simon, N., Jolliet, P., Morin, C., Zini, R., Urien, S., Tillement, J.P., 1998. Glucocorticoids decrease cytochrome c oxidase activity of isolated rat kidney mitochondria. *FEBS Lett.* 435, 25–28.
- Sionov, R.V., Kfir, S., Zafir, E., Cohen, O., Zilberman, Y., Yefenof, E., 2006. Glucocorticoid-induced apoptosis revisited: a novel role for glucocorticoid receptor translocation to the mitochondria. *Cell Cycle* 5, 1017–1026.
- Song, I.H., Buttgeriet, F., 2006. Non-genomic glucocorticoid effects to provide the basis for new drug developments. *Mol. Cell. Endocrinol.* 246, 142–146.
- Stellato, C., 2004. Post-transcriptional and nongenomic effects of glucocorticoids. *Proc. Am. Thorac. Soc.* 1, 255–263.
- Suske, G., 1999. The Sp-family of transcription factors. *Gene* 238, 291–300.
- Thompson, J., Winoto, A., 2008. During negative selection, Nur77 family proteins translocate to mitochondria where they associate with Bcl-2 and expose its proapoptotic BH3 domain. *J. Exp. Med.* 205, 1029–1036.
- Tome, M.E., Lee, K., Jaramillo, M.C., Briehl, M.M., 2012. Mitochondria are the primary source of the H(2)O(2) signal for glucocorticoid-induced apoptosis of lymphoma cells. *Exp. Ther. Med.* 4, 237–242.
- Tsiriou, C., Spandidos, D.A., Sekeris, C.E., 1997. The mitochondrion as a primary site of action of glucocorticoids: mitochondrial nucleotide sequences, showing similarity to hormone response elements, confer dexamethasone inducibility to chimaeric genes transfected in LTK-cells. *Biochem. Biophys. Res. Commun.* 235, 349–354.

- Walker, J.J., Spiga, F., Waite, E., Zhao, Z., Kershaw, Y., Terry, J.R., Lightman, S.L., 2012. The origin of glucocorticoid hormone oscillations. *PLoS Biol.* 10, e1001341.
- Wallace, A.D., Cidlowski, J.A., 2001. Proteasome-mediated glucocorticoid receptor degradation restricts transcriptional signaling by glucocorticoids. *J. Biol. Chem.* 276, 42714–42721.
- Wallerath, T., Godecke, A., Molojavyy, A., Li, H., Schrader, J., Forstermann, U., 2004. Dexamethasone lacks effect on blood pressure in mice with a disrupted endothelial NO synthase gene. *Nitric Oxide* 10, 36–41.
- Wang, Z., Chen, W., Kono, E., Dang, T., Garabedian, M.J., 2007. Modulation of glucocorticoid receptor phosphorylation and transcriptional activity by a C-terminal-associated protein phosphatase. *Mol. Endocrinol.* 21, 625–634.
- Weber, K., Bruck, P., Mikes, Z., Kupper, J.H., Klingenspor, M., Wiesner, R.J., 2002. Glucocorticoid hormone stimulates mitochondrial biogenesis specifically in skeletal muscle. *Endocrinology* 143, 177–184.
- Wegrzyn, J., Potla, R., Chwae, Y.J., Sepuri, N.B., Zhang, Q., Koeck, T., Derecka, M., Szczepanek, K., Szelag, M., Gornicka, A., Moh, A., Moghaddas, S., Chen, Q., Bobbili, S., Cichy, J., Dulak, J., Baker, D.P., Wolfman, A., Stuehr, D., Hassan, M.O., Fu, X.Y., Avadhani, N., Drake, J.L., Fawcett, P., Lesniewski, E.J., Lerner, A.C., 2009. Function of mitochondrial Stat3 in cellular respiration. *Science* 323, 793–797.
- Weigel, N.L., Moore, N.L., 2007. Steroid receptor phosphorylation: a key modulator of multiple receptor functions. *Mol. Endocrinol.* 21, 2311–2319.
- Whitworth, J.A., Mangos, G.J., Kelly, J.J., 2000. Cushing, cortisol, and cardiovascular disease. *Hypertension* 36, 912–916.
- Xu, B., Strom, J., Chen, Q.M., 2011. Dexamethasone induces transcriptional activation of Bcl-xL gene and inhibits cardiac injury by myocardial ischemia. *Eur. J. Pharmacol.* 668, 194–200.
- Yudt, M.R., Jewell, C.M., Bienstock, R.J., Cidlowski, J.A., 2003. Molecular origins for the dominant negative function of human glucocorticoid receptor beta. *Mol. Cell. Biol.* 23, 4319–4330.
- Zamzami, N., Kroemer, G., 2001. The mitochondrion in apoptosis: how Pandora's box opens. *Nat. Rev. Mol. Cell Biol.* 2, 67–71.
- Zhang, L., Zhou, R., Li, X., Ursano, R.J., Li, H., 2006. Stress-induced change of mitochondria membrane potential regulated by genomic and non-genomic GR signaling: a possible mechanism for hippocampus atrophy in PTSD. *Med. Hypotheses* 66, 1205–1208.
- Zhou, J., Li, M., Sheng, C.Q., Liu, L., Li, Z., Wang, Y., Zhou, J.R., Jing, Z.P., Chen, Y.Z., Jiang, C.L., 2011. A novel strategy for development of glucocorticoids through non-genomic mechanism. *Cell. Mol. Life Sci.* 68, 1405–1414.
- Zuo, Z., Urban, G., Scammell, J.G., Dean, N.M., McLean, T.K., Aragon, I., Honkanen, R.E., 1999. Ser/Thr protein phosphatase type 5 (PP5) is a negative regulator of glucocorticoid receptor-mediated growth arrest. *Biochemistry* 38, 8849–8857.