

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/8466014>

# Interaction of the disintegrin and cysteine-rich domains of ADAM12 with integrin $\alpha 7 \beta 1$

ARTICLE *in* EXPERIMENTAL CELL RESEARCH · SEPTEMBER 2004

Impact Factor: 3.25 · DOI: 10.1016/j.yexcr.2004.04.005 · Source: PubMed

CITATIONS

35

READS

36

8 AUTHORS, INCLUDING:



[Joanna Gruszczynska-Biegala](#)

International Institute of Molecular and Cel...

16 PUBLICATIONS 303 CITATIONS

[SEE PROFILE](#)



[Tracey Cheuvront](#)

University of Kansas

7 PUBLICATIONS 101 CITATIONS

[SEE PROFILE](#)



[Klaus von der Mark](#)

Friedrich-Alexander-University of Erlangen-...

270 PUBLICATIONS 14,386 CITATIONS

[SEE PROFILE](#)



[Stephen J Kaufman](#)

University of Illinois, Urbana-Champaign

77 PUBLICATIONS 3,742 CITATIONS

[SEE PROFILE](#)

## Interaction of the disintegrin and cysteine-rich domains of ADAM12 with integrin $\alpha 7 \beta 1$

Zhefeng Zhao,<sup>a</sup> Joanna Gruszczynska-Biegala,<sup>a</sup> Tracey Cheuvront,<sup>a</sup> Haiqing Yi,<sup>a</sup> Helga von der Mark,<sup>b</sup> Klaus von der Mark,<sup>b</sup> Stephen J. Kaufman,<sup>c</sup> and Anna Zolkiewska<sup>a,\*</sup>

<sup>a</sup>Department of Biochemistry, Kansas State University, Manhattan, KS 66506, USA

<sup>b</sup>Institute of Experimental Medicine, Friedrich-Alexander-University Erlangen-Nürnberg, 91054 Erlangen, Germany

<sup>c</sup>Department of Cell and Structural Biology, University of Illinois, Urbana, IL 61801, USA

Received 7 May 2003, revised version received 2 April 2004

Available online 6 May 2004

### Abstract

We describe a novel interaction between the disintegrin and cysteine-rich (DC) domains of ADAM12 and the integrin  $\alpha 7 \beta 1$ . Integrin  $\alpha 7 \beta 1$  extracted from human embryonic kidney 293 cells transfected with  $\alpha 7$  cDNA was retained on an affinity column containing immobilized DC domain of ADAM12. 293 cells stably transfected with  $\alpha 7$  cDNA adhered to DC-coated wells, and this adhesion was partially inhibited by 6A11 integrin  $\alpha 7$  function-blocking antibody. The X1 and the X2 extracellular splice variants of integrin  $\alpha 7$  supported equally well adhesion to the DC protein. Integrin  $\alpha 7 \beta 1$ -mediated cell adhesion to DC had different requirements for  $Mn^{2+}$  than adhesion to laminin. Furthermore, integrin  $\alpha 7 \beta 1$ -mediated cell adhesion to laminin, but not to DC, resulted in efficient cell spreading and phosphorylation of focal adhesion kinase (FAK) at Tyr397. We also show that adhesion of L6 myoblasts to DC is mediated in part by the endogenous integrin  $\alpha 7 \beta 1$  expressed in these cells. Since integrin  $\alpha 7$  plays an important role in muscle cell growth, stability, and survival, and since ADAM12 has been implicated in muscle development and regeneration, we postulate that the interaction between ADAM12 and integrin  $\alpha 7 \beta 1$  may be relevant to muscle development, function, and disease. We also conclude that laminin and the DC domain of ADAM12 represent two functional ligands for integrin  $\alpha 7 \beta 1$ , and adhesion to each of these two ligands via integrin  $\alpha 7 \beta 1$  triggers different cellular responses.

© 2004 Elsevier Inc. All rights reserved.

**Keywords:** Myoblast; Muscle; Laminin; Cell–cell interaction

### Introduction

ADAM proteins constitute a family of transmembrane or secreted glycoproteins that play important roles in cell surface proteolysis, cell–cell or cell–matrix interactions, cell migration, and differentiation [1–3]. The extracellular portion of a typical ADAM protein contains an N-terminal secretion signal, prodomain, metalloprotease, disintegrin, cysteine-rich, and epidermal growth factor-like domains. The prodomain, metalloprotease, disintegrin, and cysteine-rich domains of ADAMs are related to the P-III class of soluble snake venom metalloproteases (SVMPs) [4]. By analogy with the interactions mediated by SVMPs, the

disintegrins and/or the cysteine-rich domains of ADAMs have been proposed to bind to integrins and to mediate cell–cell or cell–matrix interactions [5]. The disintegrin domains of ADAM2 (fertilin  $\beta$ ) and ADAM9 have been shown to bind to integrin  $\alpha_6 \beta_1$  [6–9], the disintegrin domain of ADAM15 interacts with integrins  $\alpha_v \beta_3$  [10,11] and  $\alpha_5 \beta_1$  [11], the disintegrin domain of ADAM23 associates with integrin  $\alpha_v \beta_3$  [12], and the disintegrin domain of ADAM28 is a ligand for integrin  $\alpha_4 \beta_1$  [13]. The disintegrin and cysteine-rich (DC) domains of ADAM13, in addition to interacting with  $\beta 1$  integrins, bind to fibronectin [14], an extracellular matrix protein that associates with several integrins containing  $\beta 1$  chains.

ADAM12 is highly expressed in developing and regenerating skeletal muscle [15–19]. It is an active metalloprotease that cleaves insulin-like growth factor binding protein-3 and -5 [20,21], and heparin binding-epidermal growth factor (HB-EGF) [22]. ADAM12 cytoplasmic domain inter-

\* Corresponding author. Department of Biochemistry, Kansas State University, 104 Willard Hall, Manhattan, KS 66506. Fax: +1-785-532-7278.

E-mail address: [zolkiea@ksu.edu](mailto:zolkiea@ksu.edu) (A. Zolkiewska).

acts with protein tyrosine kinase Src [23], phosphatidylinositol 3-kinase [24], and cytoskeletal proteins  $\alpha$ -actinin-2 [18] and  $\alpha$ -actinin-1 [25]. Transgenic expression of ADAM12 in differentiated myofibers of skeletal muscle using muscle creatine kinase promoter results in a pronounced accumulation of adipocytes in skeletal muscle [26]. Overexpression of ADAM12 in myofibers of the dystrophin-deficient *mdx* mice alleviates the skeletal muscle pathology [27].

The disintegrin and cysteine-rich domains of ADAM12, similar to the corresponding domains in other ADAM proteins, appear to interact with cell adhesion receptors. The cysteine-rich domain of ADAM12 binds to syndecan-4, a member of the heparan sulfate proteoglycan family [28–30], and the disintegrin domain of ADAM12 interacts with integrin  $\alpha 9 \beta 1$  [31,32]. To identify integrin  $\alpha 9 \beta 1$  as a binding partner for the disintegrin domain of ADAM12, a series of CHO cells stably transfected with individual integrin  $\alpha$  chains ( $\alpha 2$ ,  $\alpha 3$ ,  $\alpha 4$ ,  $\alpha 5$ ,  $\alpha 6$ , or  $\alpha 9$ ) was tested for adhesion to the recombinant disintegrin domain of ADAM12 [31]. The list of integrin  $\alpha$  chains transfected into CHO cells did not include integrin  $\alpha 7$ . Integrin  $\alpha 7$ , which forms a dimer with the  $\beta 1$  chain, is the major laminin-binding integrin in skeletal muscle and it plays an important role in muscle development, disease, and regeneration [33]. Since ADAM12 has been implicated in muscle development and regeneration as well [18,19,34], we asked whether integrin  $\alpha 7 \beta 1$  can represent a binding partner for ADAM12. The  $\alpha 7$  chain can exist as one of the three cytoplasmic tail splice variants (A, B, or C) and two extracellular domain splice variants (X1 and X2) [33]. In this report, we demonstrate that the disintegrin and cysteine-rich domains of ADAM12 bind to the  $\alpha 7 \beta 1$  dimer (containing either the X1B or X2B variant of  $\alpha 7$ ) in vitro and in intact 293 human embryonic kidney cells and L6 myoblasts. However, we also provide evidence that integrin  $\alpha 7 \beta 1$ -mediated adhesion to disintegrin and cysteine-rich domains, unlike adhesion to laminin, does not efficiently induce cell spreading and phosphorylation of focal adhesion kinase (FAK) at Tyr397. These results point to important differences between integrin  $\alpha 7 \beta 1$ -mediated signaling initiated by two different ligands: laminin and ADAM12.

## Materials and methods

### Cells

Rat L6 myoblasts were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS) in the presence of 5% CO<sub>2</sub> at 37°C. 293EBNA cells [human embryonic kidney 293 cells stably expressing the Epstein Barr Virus (EBV) EBNA-1 gene] were cultured in DMEM supplemented with 250  $\mu$ g/ml G418. 293-X1 and 293-X2 cells (293EBNA cells stably

transfected with mouse integrin  $\alpha 7 X1B$  or  $\alpha 7 X2B$  isoforms, respectively, in pCEP4, an episomal mammalian expression vector containing the EBV replication origin, Ref. [35]) were cultured in DMEM/medium F12 (1:1) supplemented with 5% FBS, 250  $\mu$ g/ml G418, and 300  $\mu$ g/ml hygromycin.

### Affinity chromatography using recombinant disintegrin and cysteine-rich domains of ADAM12

Recombinant DC-6xHis protein (amino acids 425–657, CGNGYV ... VFGVHK, from mouse ADAM12, containing a C-terminal 6xHis tag) was produced in *Spodoptera frugiperda* Sf21 cell line using baculovirus expression system, and was purified from cell-associated material, as described [37]. DC-6xHis was immobilized on a fresh Ni-NTA affinity column (1.5 mg/ml resin, 0.1 ml bed volume). In parallel, a control column containing Ni-NTA resin and no DC-6xHis was subjected to the same treatment. 293-X1 or 293-X2 cells were extracted with 150 mM NaCl, 50 mM Tris-HCl (pH 7.5), 50 mM octylglucopyranoside, 1 mM MnCl<sub>2</sub>, 1 mM AEBSF, 5  $\mu$ g/ml aprotinin, 5  $\mu$ g/ml pepstatin A, 5  $\mu$ g/ml leupeptin; cell extracts were supplemented with 10 mM imidazole, divided into two equal parts, loaded onto the DC-6xHis column and the control column, and incubated for 1 h at 4°C. Columns were then washed with 10 ml of 150 mM NaCl, 50 mM Tris-HCl (pH 7.5), 1% Triton X-100 (washing buffer), then with washing buffer supplemented with 0.1 M imidazole, and then eluted with 0.3 ml of washing buffer containing 0.5 M imidazole.

### Cell adhesion assays

Wells of 96-microtiter plates were coated with indicated concentrations of EHS laminin (Invitrogen) or DC-6xHis in NaHCO<sub>3</sub>, pH 9.0, overnight at 4°C, and then blocked with 2.5% (w/v) BSA in DPBS for 1 h at 37°C. Adherent cells were detached in DPBS containing 1 mM EDTA and 0.05% (w/v) trypsin, washed with DPBS containing 1 mg/ml of trypsin inhibitor, suspended at 10<sup>6</sup> cells/ml in Tyrode's buffer (5 mM HEPES-KOH, pH 7.4, 12 mM NaHCO<sub>3</sub>, 150 mM NaCl, 2.6 mM KCl, 5 mM glucose, 0.5 mM MgCl<sub>2</sub>, 1 mM CaCl<sub>2</sub>) containing 1% BSA. Cells were then pre-incubated for 15 min with or without the following function-blocking antibodies: anti- $\alpha 7$  (clone 6A11, Ref. [35]; 10  $\mu$ g/ml), anti- $\beta 1$  (clone Ha2/5, Pharmingen; 20  $\mu$ g/ml), anti- $\beta 1$  (clone P4C10), anti- $\alpha 1$  (clone FB12), anti- $\alpha 2$  (clone P1E6), anti- $\alpha 3$  (clone P1B5), anti- $\alpha 5$  (clone P1D6), anti- $\alpha 6$  (clone NKI-GoH3), or anti- $\alpha 9$  (clone Y9A2) (obtained from Chemicon, used at 20  $\mu$ g/ml). Cells were then incubated in the wells of a 96-well plate (10<sup>5</sup> cells/well) for 1 h at 37°C. Unbound cells were removed by washing the wells three times with DMEM. Adherent cells were fixed for 15 min at room temperature with 3% glutaraldehyde in DPBS and stained for 5 min with 0.04% crystal

violet in 20% methanol. After washing the wells and adding 1% SDS, absorbance was measured with a Vmax microplate reader (Molecular Devices) at 550 nm; the linear range of the plate reader was 0–3.5 O.D. Each assay point was derived from two to four wells in at least two independent experiments.

### Immunoblotting

To analyze FAK phosphorylation, cells were serum starved for 24 h; to analyze integrin  $\alpha 7$  binding to the DC protein, cells were used without starvation. Cellular proteins were extracted with 50 mM Tris–HCl (pH 7.4), 150 mM NaCl, 1% (v/v) Triton X-100, 1% sodium deoxycholate, 0.1% SDS, 1 mM AEBSF, 5  $\mu$ g/ml aprotinin, 5  $\mu$ g/ml leupeptin, 5  $\mu$ g/ml pepstatin A, 5 mM EDTA, 50 mM NaF, 2 mM  $\text{Na}_3\text{VO}_4$  and 10 mM  $\text{Na}_4\text{P}_2\text{O}_7$ ; 2 ml/100-mm plate. Samples were centrifuged for 30 min at  $21,000 \times g$ , supernatants were resolved by SDS-PAGE and transferred to a nitrocellulose membrane. Membranes were blocked in DPBS containing 3% (w/v) dry milk (for  $\alpha 7$ B and FAK immunoblotting) or 3% (w/v) BSA [for FAK (pY397) immunoblotting] and 0.3% (v/v) Tween-20, then incubated with a primary antibody in blocking buffer, followed by incubation with a horseradish peroxidase-labeled secondary antibody and development with a chemiluminescence detection method (West Pico, Pierce, Rockford, IL). The following dilutions of primary antibodies were used: rabbit anti-integrin  $\alpha 7$  antibody recognizing the C-terminus of the  $\alpha 7$ B isoform [36], 1:2000; anti-FAK (Santa Cruz, sc-558), 1:4000; anti-FAK (pY397) (BioSource), 1: 5000.

## Results

### *Integrin $\alpha 7\beta 1$ interacts with the DC domain of ADAM12 in vitro*

To examine whether integrin  $\alpha 7\beta 1$  can bind to the disintegrin and cysteine-rich (DC) domains of ADAM12 in vitro, protein extracts from cells expressing the  $\alpha 7\beta 1$  integrin were passed through Ni-NTA columns containing immobilized DC protein, and the eluate was analyzed by SDS-PAGE and Western blotting using an antibody specific for the C-terminus of integrin  $\alpha 7$ . As shown in Fig. 1, mouse integrin  $\alpha 7$ X1 and  $\alpha 7$ X2 chains stably expressed in 293 cells (293-X1 and 293-X2, respectively) were retained on the DC-containing columns but not on control Ni-NTA columns without DC. Immunoblotting with an antibody specific for integrin  $\beta 1$  further confirmed the presence of the  $\beta 1$  integrin chain in the eluates from DC but not from control columns (result not shown). This result indicates the  $\alpha 7\beta 1$  dimer containing either the X1 or X2 splice variant of the integrin  $\alpha 7$  interacts with the DC protein in vitro.




Fig. 1. Interaction of integrin  $\alpha 7\beta 1$  with the disintegrin-like and cysteine-rich (DC) domains of ADAM12 in vitro. (A) Total cell extracts from 293 cells stably transfected with mouse integrin  $\alpha 7$ X1B or  $\alpha 7$ X2B (293-X1 and 293-X2, respectively) were loaded on Ni-NTA columns containing immobilized DC-6xHis protein or on control Ni-NTA columns without DC-6xHis. Another DC-6xHis-containing column was incubated with cell extraction buffer only. Columns were washed and eluted with 0.5 M imidazole; samples of the eluates were subjected to SDS-PAGE under reducing conditions, followed by Western blotting with antibody raised against a C-terminal peptide from integrin  $\alpha 7$ B. The protein band of approximately 35 kDa detected by the antibody and corresponding to the C-terminal "light chain" of integrin  $\alpha 7$ B is shown. (B) Expression of integrin  $\alpha 7$ B in total cell lysates of 293-X1, 293-X2, but not in control mock-transfected 293 cells, was confirmed by Western blotting with anti-integrin  $\alpha 7$ B antibody.

### *Integrin $\alpha 7\beta 1$ mediates cell adhesion to the DC domain of ADAM12 in the presence of $\text{Mn}^{2+}$*

We next examined whether recombinant DC domain of ADAM12 can support cell adhesion via integrin  $\alpha 7\beta 1$ . In the presence of 1 mM  $\text{Mn}^{2+}$ , 1 mM  $\text{Ca}^{2+}$ , and 0.5 mM  $\text{Mg}^{2+}$ , both integrin  $\alpha 7$ -expressing cells (293-X1 and 293-X2) and mock-transfected 293 cells adhered well to DC-coated plates (Fig. 2A). Pre-incubation of 293-X1 and 293-X2 cells with integrin  $\alpha 7$ -blocking antibody 6A11 inhibited cell adhesion by approximately 50% (Fig. 2A). The same antibody did not have any effect on the adhesion of mock-transfected 293 cells to DC (Fig. 2A). This suggests that adhesion of 293-X1 and 293-X2 cells to DC under these conditions was mediated, at least in part, by integrin  $\alpha 7\beta 1$ . In contrast, in the presence of 1 mM  $\text{Ca}^{2+}$ , 0.5 mM  $\text{Mg}^{2+}$ , and in the absence of  $\text{Mn}^{2+}$ , all three cell lines (293-X1, 293-X2, and 293) adhered very poorly to DC-coated plates (Fig. 2B).

In the presence of  $\text{Mn}^{2+}$ , the attachment of 293 cells to DC was partially inhibited by integrin  $\beta 1$  blocking antibodies. The extent of this inhibition was similar to the




Fig. 2. Integrin  $\alpha 7 \beta 1$  supports adhesion of 293-X1 and 293-X2 cells to the DC domain of ADAM12 in the presence of  $Mn^{2+}$ . Wells of 96-well plates were coated with DC protein at indicated concentrations. 293 cells stably transfected with mouse integrin  $\alpha 7 X1B$ ,  $\alpha 7 X2B$ , or vector only (293-X1, 293-X2, and 293, respectively) were incubated for 15 min in Tyrode's buffer containing 1 mM  $Mn^{2+}$ , 1 mM  $Ca^{2+}$ , and 0.5 mM  $Mg^{2+}$  (A) or 1 mM  $Ca^{2+}$ , and 0.5 mM  $Mg^{2+}$  and no  $Mn^{2+}$  (B), without (closed symbols) or with 10  $\mu g/ml$  of 6A11 integrin  $\alpha 7$  function-blocking antibody (open symbols). Cells were then added to the wells ( $10^5$  cells/well) and incubated for 1 h at 37°C. After removing unbound cells, bound cells were fixed and stained with 0.04% crystal violet, as described in Materials and methods. Results are shown as the average from three measurements,  $\pm$  SE.

inhibition observed for 293-X1 and 293-X2 cells (Fig. 3). Thus, integrin  $\alpha 7 \beta 1$  is not the only integrin capable of binding to the DC domain of ADAM12, and other  $\beta 1$  integrins present in 293 cells may mediate the adhesion to DC as well. Previous studies demonstrated that 293 cells




Fig. 3. The effect of integrin  $\beta 1$  function-blocking antibodies on adhesion of 293-X1, 293-X2, and 293 cells to DC. 293-X1, 293-X2, or 293 cells in Tyrode's buffer/1% BSA/1 mM  $Mn^{2+}$  were incubated for 15 min without (black bars) or with integrin  $\beta 1$  function-blocking antibody Ha2/5 (20  $\mu g/ml$ ; gray bars) or P4C10 (20  $\mu g/ml$ ; white bars) and then transferred to wells coated with 10  $\mu g/ml$  of DC. The amounts of cells bound to the wells were determined as in Fig. 2.

express multiple integrin  $\alpha$  chains that can dimerize with the  $\beta 1$  chain [35,38]. We tried to determine which of these integrins might have interacted with DC. We reasoned that the best candidates were the integrins that were down-regulated after overexpression of integrin  $\alpha 7$ , namely  $\alpha 1$ ,  $\alpha 2$ ,  $\alpha 3$ ,  $\alpha 5$ , and  $\alpha 6$  [35]. Therefore, we tested the effect of function-blocking antibodies against these integrins on the adhesion of 293 cells to DC. In addition, since the disintegrin domain of ADAM12 was shown to bind to integrin  $\alpha 9 \beta 1$  [31,32] and since 293 cells express the  $\alpha 9$  chain [38], the effect of an anti-integrin  $\alpha 9$  antibody was also examined. None of the tested antibodies, however, added either individually or in combination, inhibited 293 cell adhesion to DC (data not shown). This suggests that either a different integrin  $\alpha$  subunit was involved in adhesion to DC or that the potency of one or more of the antibodies used in our studies was diminished in the presence of  $Mn^{2+}$ .

*Integrin  $\alpha 7 \beta 1$ -mediated cell adhesion to laminin has different requirements for  $Mn^{2+}$  than adhesion to DC*

Integrin  $\alpha 7 \beta 1$  is a laminin-binding protein [33,39,40] and it efficiently mediates adhesion to laminin-coated plates [35,36,41–43]. As integrin  $\alpha 7 \beta 1$  appears to bind to the DC domain of ADAM12 as well, we were interested in comparing integrin  $\alpha 7 \beta 1$ -mediated cell adhe-



sion to DC and to laminin. As shown in Fig. 2, the interaction of integrin  $\alpha 7 \beta 1$  with DC was critically dependent on  $Mn^{2+}$ . In contrast, adhesion of 293-X1 and 293-X2 cells to laminin was very efficient in the presence of 0.5 mM  $Mg^{2+}$  and 1 mM  $Ca^{2+}$ , and it was only weakly further stimulated by  $Mn^{2+}$  (Figs. 4A, B). Moreover, adhesion of 293-X1 and 293-X2, but not control 293 cells, to laminin in the presence of 0.5 mM  $Mg^{2+}$  and 1 mM  $Ca^{2+}$  was efficiently blocked by 6A11 antibody (Fig. 4B). Adhesion of 293-X1 and 293-X2 cells incubated in the presence of 6A11 antibody was lower than adhesion of 293 cells, which is consistent with decreased levels of several other laminin-binding integrins (mainly integrin  $\alpha 3 \beta 1$  and  $\alpha 6 \beta 1$ ) in 293-X1 and 293-X2 cells as a result of overexpression of integrin  $\alpha 7$  [35]. Thus, in the absence of  $Mn^{2+}$ , integrin  $\alpha 7 \beta 1$  is the major integrin in 293-X1 and 293-X2 cells that mediates adhesion to laminin.

In contrast, the 6A11 antibody had a much weaker effect on the adhesion of 293-X1 and 293-X2 cells to laminin when 1 mM  $Mn^{2+}$  was included in the incubation medium (Fig. 4A). This suggests that other laminin-binding integrins that are present in 293-X1 and 293-X2 cells contribute significantly to cell adhesion to laminin in the presence of 1 mM  $Mn^{2+}$ . Collectively, the results of the experiments shown in Figs. 2 and 4 indicate that integrin  $\alpha 7 \beta 1$  dimers expressed in 293-X1 and 293-X2 have different requirements for  $Mn^{2+}$  to mediate cell adhesion to DC or to laminin.

*293-X1 and 293-X2 cells spread well on laminin and poorly on DC*

We examined the ability of integrin  $\alpha 7 \beta 1$  to mediate cell spreading on laminin and on DC. 293-X1, 293-X2, and 293 cells plated on laminin-coated plates in the presence of 0.5 mM  $Mg^{2+}$  and 1 mM  $Ca^{2+}$  were well spread and assumed a flattened morphology (Fig. 5). Since in the absence of  $Mn^{2+}$ , adhesion of 293-X1 and 293-X2 cells to laminin is mediated predominantly via integrin  $\alpha 7 \beta 1$  (Fig. 4B); this result indicates that integrin  $\alpha 7 \beta 1$  binding to laminin-coated plates induces 293-X1 and 293-X2 cell spreading. In contrast, in the presence of 0.5 mM  $Mg^{2+}$ , 1 mM  $Ca^{2+}$ , and 1 mM  $Mn^{2+}$ , 293-X1 and 293-X2 cells plated on DC were mostly rounded and had few filopodia-like thin protrusions (Fig. 5). Since 293-X1 and 293-X2 cells in the presence of  $Mn^{2+}$  adhere to DC in large part via integrin  $\alpha 7 \beta 1$ , this result suggested that integrin  $\alpha 7 \beta 1$  binding to laminin and to DC may trigger different signaling pathways in these cells. The extent of spreading of control 293 cells plated on laminin or DC was similar and most cells appeared spread and flattened.

*Integrin  $\alpha 7 \beta 1$ -mediated adhesion to laminin leads to higher Tyr397 phosphorylation of FAK than adhesion to DC*

We next asked whether integrin  $\alpha 7 \beta 1$ -mediated cell adhesion to DC- and to laminin-coated plates results in the activation of the same intracellular signaling pathways.




Fig. 4. Integrin  $\alpha 7 \beta 1$ -mediated adhesion of 293-X1 and 293-X2 cells to laminin has a different requirements for  $Mn^{2+}$  than adhesion to DC. Wells of 96-well plates were coated with laminin at indicated concentrations. Adhesion of 293-X1, 293-X2, and 293 cells in the presence of 1 mM  $Mn^{2+}$  (A) or in the absence of  $Mn^{2+}$  (B), without (closed symbols) or with 10  $\mu$ g/ml of 6A11 integrin  $\alpha 7$  function-blocking antibody (open symbols), was determined as in Fig. 2. Results are shown as the average from three measurements,  $\pm$  SE.




Fig. 5. Integrin  $\alpha 7 \beta 1$ -mediated adhesion to DC induces less cell spreading than adhesion to laminin. Wells of 96-well plates were coated with laminin or DC at 20  $\mu\text{g}/\text{ml}$ . 293-X1, 293-X2, or 293 cells suspended in Tyrode's buffer containing 0.5 mM  $\text{Mg}^{2+}$ , 1 mM  $\text{Ca}^{2+}$  (for adhesion to laminin) or 0.5 mM  $\text{Mg}^{2+}$ , 1 mM  $\text{Ca}^{2+}$ , and 1 mM  $\text{Mn}^{2+}$  (for adhesion to DC) were added to the wells ( $10^4$  cells/well). After 1 h, cells were examined by phase contrast microscopy.

Autophosphorylation at Tyr397 and subsequent activation of focal adhesion kinase (FAK) are the early events that are observed in many cells after the engagement of integrins with their immobilized ligands [43,44]. To examine the level of FAK phosphorylation after cell adhesion to DC, 293-X1, 293-X2, and 293 cells were serum-starved for 24 h, detached from plates, and then allowed to attach to DC-coated wells in the presence of 1 mM  $\text{Mn}^{2+}$ . After 1 h incubation, the levels of FAK (pY397) and total FAK protein were analyzed by Western blotting. As shown in Figs. 6A and B, the level of FAK (pY397) in 293-X1 and in 293-X2 cells plated on DC protein was at least three times lower than in control 293 cells, whereas the amount of total FAK protein was the same. Since in the presence of 1 mM  $\text{Mn}^{2+}$  the adhesion of 293-X1 and 293-X2, but not 293 cells, to DC is mediated in large part by integrin  $\alpha 7 \beta 1$  (see also Fig. 2A), this result suggests that the interaction between  $\alpha 7 \beta 1$  and DC provides a poor stimulus for FAK phosphorylation.

In contrast, the interaction between integrin  $\alpha 7 \beta 1$  and laminin does lead to efficient phosphorylation of FAK at Tyr397. In 0.5 mM  $\text{Mg}^{2+}$  and 1 mM  $\text{Ca}^{2+}$ , the adhesion of 293-X1 and 293-X2 cells to laminin is mediated mostly by integrin  $\alpha 7 \beta 1$  (see Fig. 4B), whereas 293 cells adhere to laminin most likely via integrin  $\alpha 3 \beta 1$  and  $\alpha 6 \beta 1$  (Ref. [35]). The levels of FAK (pY397) in 293-X1, 293-X2, and 293 cells attached to laminin in the absence of  $\text{Mn}^{2+}$  were very similar (Fig. 6), indicating that integrin  $\alpha 7 \beta 1$  engagement (in 293-X1 and 293-X2 cells) was as efficient as the engagement of other laminin-binding integrins (in 293 cells) in inducing FAK phosphorylation. Collectively, these results suggest that although laminin and the DC domain of ADAM12 represent two ligands for integrin  $\alpha 7 \beta 1$ , adhesion to each of these two ligands via integrin  $\alpha 7 \beta 1$  triggers different cellular responses. Specifically,  $\alpha 7 \beta 1$ -mediated

adhesion to laminin leads to higher Tyr397 phosphorylation of FAK (Fig. 6), which is accompanied by more efficient cell spreading (Fig. 5).

#### *The endogenous integrin $\alpha 7 \beta 1$ expressed in myoblasts contributes to cell adhesion to DC*

We asked whether skeletal myoblasts expressing the endogenous  $\alpha 7$  chain utilize the  $\alpha 7 \beta 1$  integrin dimer during adhesion to the DC domain of ADAM12. As shown in Fig. 7A, L6 myoblasts incubated in the presence of 1 mM  $\text{Mn}^{2+}$ , 1 mM  $\text{Ca}^{2+}$ , and 0.5 mM  $\text{Mg}^{2+}$  adhered well to DC-coated plates. Similarly to 293-X1 and 293-X2 cells, the adhesion of L6 myoblasts to DC was critically dependent on the presence of  $\text{Mn}^{2+}$ . Pre-incubation with 6A11 integrin  $\alpha 7$  function-blocking antibody inhibited cell adhesion at lower DC-coating concentrations ( $<2 \mu\text{g}/\text{ml}$ ), but did not have any effect at higher DC concentrations (Fig. 7A), suggesting that other integrins may also play a role in L6 cell adhesion to




Fig. 6. FAK phosphorylation at Tyr397 in cells adhering to DC and to laminin. (A) Wells of 96-well plates were coated with DC protein or EHS laminin (L) at 20  $\mu\text{g}/\text{ml}$ . 293-X1, 293-X2, or 293 cells were suspended in Tyrode's buffer containing 0.5 mM  $\text{Mg}^{2+}$ , 1 mM  $\text{Ca}^{2+}$ , with or without 1 mM  $\text{Mn}^{2+}$ , as indicated. Cells were then added to the wells ( $10^5$  cells/well) and incubated for 1 h at 37°C. After removing unbound cells, bound cells were treated with lysis buffer containing protease and phosphatase inhibitors, centrifuged, and supernatants were analyzed by Western blotting using a phospho-specific antibody recognizing phosphorylated Tyr397 in FAK (top) or total FAK (bottom). (B) The intensities of the FAK (pY397) and total FAK signals shown in panel A were quantified by densitometry. The experiment was repeated three times, representative results from one experiment are shown in A, the average FAK (pY397)/FAK signals (in arbitrary units,  $\pm\text{SE}$ ) are shown in B.




Fig. 7. Adhesion of L6 myoblasts to DC is partially mediated by integrin  $\alpha 7 \beta 1$  and it requires the presence of  $Mn^{2+}$ . Wells of 96-well plates were coated with DC protein (A) or laminin (B) at indicated concentrations. 293-X1, 293-X2, and 293 cells were incubated for 15 min in Tyrode's buffer containing 1 mM  $Mn^{2+}$ , 1 mM  $Ca^{2+}$ , and 0.5 mM  $Mg^{2+}$  (left panels) or 1 mM  $Ca^{2+}$ , and 0.5 mM  $Mg^{2+}$  and no  $Mn^{2+}$  (right panels), without (closed symbols) or with 10  $\mu g/ml$  of 6A11 integrin  $\alpha 7$  function-blocking antibody (open symbols). Cells were then added to the wells ( $10^5$  cells/well) and incubated for 1 h at  $37^\circ C$ . The amount of bound cells was determined as in Fig. 2. Results are shown as the average from three measurements,  $\pm SE$ .

DC. Interestingly, the inhibitory effect of the antibody was more pronounced when L6 cells were plated on DC than on laminin (Fig. 7B). We conclude that the endogenous integrin  $\alpha 7 \beta 1$  expressed in L6 myoblasts does contribute to cell adhesion to DC, and its contribution is most evident at low concentrations of DC protein.

## Discussion

In this report, we describe a novel interaction between the disintegrin and cysteine-rich domains of ADAM12 and integrin  $\alpha 7 \beta 1$ . Using affinity columns containing immobilized DC domain of ADAM12, we showed that integrin  $\alpha 7 \beta 1$  bound to the DC protein *in vitro*. 293 cells stably transfected with  $\alpha 7$  cDNA, as well as L6 myoblasts expressing the endogenous  $\alpha 7$  integrin, adhered to DC-coated wells, and this adhesion was partially inhibited by 6A11 integrin  $\alpha 7$  function-blocking antibody.

Integrin  $\alpha 7$  chain can exist as one of the three cytoplasmic (A, B, or C) and two extracellular (X1 or X2) splice variants [33]. Our cell adhesion and column binding assays utilized 293 cells expressing the X1B or X2B forms of the  $\alpha 7$  chain, which are the major variants expressed in proliferating myoblasts [45,46]. Although the X1 and X2 splice

variants confer different specificities for laminin in cell adhesion [36,42], cell migration [35,36], acetylcholine receptor clustering [47], or solid phase binding assays [43], these two variants supported equally well adhesion to the DC protein.

Whereas laminin and the DC domain of ADAM12 represent two functional ligands for integrin  $\alpha 7 \beta 1$ , our results suggest that there are important differences between  $\alpha 7 \beta 1$  binding to these two ligands. While  $\alpha 7 \beta 1$ -mediated adhesion to laminin did not require activation of the integrin by  $Mn^{2+}$ ,  $Mn^{2+}$  was necessary to detect adhesion to the DC protein. In addition, cellular responses initiated upon integrin  $\alpha 7 \beta 1$  binding to laminin and to DC seem to be different as well, as  $\alpha 7 \beta 1$ -mediated adhesion to laminin results in better cell spreading and more potent Tyr397 phosphorylation than adhesion to DC. Interestingly, our results are similar to a previous report in which integrin  $\alpha 6 \beta 1$ -mediated adhesion of fibrosarcoma cells to the recombinant extracellular domain of ADAM9 led to a rounded cell morphology and higher cell motility than adhesion to laminin [9].

The different requirement for  $Mn^{2+}$  of  $\alpha 7 \beta 1$  binding to laminin and to DC is consistent with the recently proposed model of integrin activation [48–50]. According to this model, in the presence of  $Mn^{2+}$ , there is an equilibrium of the bent integrin conformer (with a low affinity for ligand), an extended conformer with closed headpiece (with an intermediate affinity for ligand), and an extended conformer with open headpiece (with a high affinity for ligand). In the absence of  $Mn^{2+}$ , the major conformational state of integrin is the bent conformer, with a very minor population of the extended conformers. Addition of a ligand shifts the equilibrium toward the extended conformer with an open headpiece [48]. Because the equilibria between conformational changes and ligand binding are thermodynamically linked, a ligand (such as laminin) that binds strongly to integrin  $\alpha 7 \beta 1$  may shift the equilibrium towards the extended conformer with an open headpiece even in the absence of  $Mn^{2+}$ . In the case of a ligand with weaker binding (such as DC),  $Mn^{2+}$  is required to destabilize the bent conformation and to induce the extended conformer with the open headpiece that can interact with the ligand. Our results suggest, therefore, that DC binding to integrin  $\alpha 7 \beta 1$  is significantly weaker than the binding of laminin. It should be noted, however, that the DC protein corresponds to a domain of ADAM12 and its integrin binding properties may be different from the properties of the intact transmembrane ADAM12.

Integrin  $\alpha 7 \beta 1$  is not the only protein capable of binding to the DC domain of ADAM12, as adhesion of 293-X1 and 293-X2 cells to DC was only partially inhibited by anti- $\alpha 7$  antibodies and parental 293 cells lacking integrin  $\alpha 7$  chain also adhered well to DC-coated plates. Indeed, it has been shown recently that multiple receptors present on the same cell can participate in cell binding to recombinant ADAM disintegrin domains [8]. In fact, two other proteins were reported to bind to the extracellular domain of ADAM12



previously: syndecan-4, a member of heparan sulfate family [28–30], and integrin  $\alpha 9\beta 1$  [31,32]. Syndecan-4 binds to the cysteine-rich fragment of ADAM12 located between amino acids 564 and 708 [28], a region that has a limited overlap (93 amino acids) with the DC construct used in our studies (amino acids 425–657). Syndecans, however, are not activated by  $Mn^{2+}$  and they seem unlikely to mediate cell adhesion observed in our studies. Integrin  $\alpha 9\beta 1$  binds to the  $RX_6DLPEF$  recognition motif that is present in the disintegrin domains of many ADAMs, including ADAM12 [32]. Thus, the  $\alpha 9\beta 1$  integrin is a receptor for many ADAM disintegrins. As both  $\alpha 9\beta 1$  and ADAMs are widely expressed,  $\alpha 9\beta 1$ -ADAM interactions are predicted to be of a broad biological importance [32]. Integrin  $\alpha 7\beta 1$ , on the other hand, is expressed predominantly in skeletal and cardiac muscle and therefore it might be involved in tissue-specific interactions with ADAM12. Interestingly, it was recently observed that transgenic overexpression of ADAM12 in mice using the muscle creatine kinase promoter leads to increased expression of integrin  $\alpha 7$  in skeletal muscle [51]. Clearly, an interaction between the DC domain of ADAM12 and integrin  $\alpha 7\beta 1$  is one of the possible mechanisms underlying this effect.

Integrin  $\alpha 7$  plays an important role in muscle development and disease [33]. Spatiotemporal pattern of expression of individual splice variants of integrin  $\alpha 7$  is developmentally regulated during myogenesis [52–54]. Mice lacking the integrin  $\alpha 7$  gene develop a specific form of muscular dystrophy that affects myotendinous junction [55], and mutations in the integrin  $\alpha 7$  gene in humans cause congenital myopathies [56]. Moreover, an altered expression of integrin  $\alpha 7$  has been observed in Duchenne muscular dystrophy and in *mdx* mice that lack dystrophin [57]. Enhanced expression of integrin  $\alpha 7$  reduced muscular dystrophy and stabilized muscle regeneration in *mdx/utr*–/– mice that lacked both dystrophin and utrophin [58]. Since ADAM12 has been implicated in muscle development and/or regeneration as well [18,19,34,51], we postulate that the interaction between ADAM12 and integrin  $\alpha 7\beta 1$  reported here may contribute to integrin  $\alpha 7\beta 1$  functions in muscle cell growth, stability, and/or survival.

## Acknowledgments

This work was supported by the NIH COBRE Award P20 RR17708 and matching support from the State of Kansas to AZ and by NIH and MDA awards to SJK. This is contribution 03-60-J from the Kansas Agricultural Experiment Station.

## References

- [1] J.M. White, ADAMs: modulators of cell–cell and cell–matrix interactions, *Curr. Opin. Cell Biol.* 15 (2003) 598–606.
- [2] J. Schlöndorff, C.P. Blobel, Metalloprotease-disintegrins: modular proteins capable of promoting cell–cell interactions and triggering signals by protein-ectodomain shedding, *J. Cell Sci.* 112 (1999) 3603–3617.
- [3] D.F. Seals, S.A. Courtneidge, The ADAMs family of metalloproteases: multidomain proteins with multiple functions, *Genes Dev.* 17 (2003) 7–30.
- [4] J.B. Bjarnason, J.W. Fox, Snake venom metalloendopeptidases: repolysins, *Methods Enzymol.* 248 (1995) 345–368.
- [5] J.P. Evans, Fertilin  $\beta$  and other ADAMs as integrin ligands: insights into cell adhesion and fertilization, *BioEssays* 23 (2001) 628–639.
- [6] D. Bigler, Y. Takahashi, M.S. Chen, E.A.C. Almeida, L. Osbourne, J.M. White, Sequence-specific interaction between the disintegrin domain of mouse ADAM 2 (fertilin  $\beta$ ) and murine eggs. Role of the  $\alpha 6$  integrin subunit, *J. Biol. Chem.* 275 (2000) 11576–11584.
- [7] Y. Takahashi, D. Bigler, Y. Ito, J.M. White, Sequence-specific interaction between the disintegrin domain of mouse ADAM 3 and murine eggs: role of  $\beta 1$  integrin-associated proteins CD9, CD81, and CD98, *Mol. Biol. Cell* 12 (2001) 809–820.
- [8] M. Tomczuk, Y. Takahashi, J. Huang, S. Murase, M. Mistretta, E. Klaffky, A. Sutherland, L. Bolling, S. Coonrod, C. Marcinkiewicz, D. Sheppard, M.A. Stepp, J.M. White, Role of multiple  $\beta 1$  integrins in cell adhesion to the disintegrin domains of ADAMs 2 and 3, *Exp. Cell Res.* 290 (2003) 68–81.
- [9] D. Nath, P.M. Slocombe, A. Webster, P.E. Stephens, A.J. Docherty, G. Murphy, Meltrin  $\gamma$  (ADAM-9) mediates cellular adhesion through  $\alpha 6\beta 1$  integrin, leading to a marked induction of fibroblast cell motility, *J. Cell Sci.* 113 (2000) 2319–2328.
- [10] X.-P. Zhang, T. Kamata, K. Yokoyama, W. Puzon-McLaughlin, Y. Takada, Specific interaction of the recombinant disintegrin-like domain of MDC-15 (metargidin, ADAM-15) with integrin  $\alpha v\beta 3$ , *J. Biol. Chem.* 273 (1998) 7345–7350.
- [11] D. Nath, P.M. Slocombe, P.E. Stephens, A. Warn, G.R. Hutchinson, K.M. Yamada, A.J.P. Docherty, G. Murphy, Interaction of metargidin (ADAM-15) with  $\alpha v\beta 3$  and  $\alpha 5\beta 1$  integrins on different haemopoietic cells, *J. Cell Sci.* 112 (1999) 579–587.
- [12] S. Cal, J.M. Freije, J.M. Lopez, Y. Takada, C. Lopez-Otin, ADAM 23/ MDC3, a human disintegrin that promotes cell adhesion via interaction with the  $\alpha v\beta 3$  integrin through an RGD-independent mechanism, *Mol. Biol. Cell* 11 (2000) 1457–1469.
- [13] L.C. Bridges, P.H. Tani, K.R. Hanson, C.M. Roberts, M.B. Judkins, R.D. Bowditch, The lymphocyte metalloprotease MDC-L (ADAM 28) is a ligand for the integrin  $\alpha 4\beta 1$ , *J. Biol. Chem.* 277 (2002) 3784–3792.
- [14] A. Gaultier, H. Cousin, T. Darriber, D. Alfandari, ADAM13 disintegrin and cysteine-rich domains bind to the second heparin-binding domain of fibronectin, *J. Biol. Chem.* 277 (2002) 23336–23344.
- [15] T. Yagami-Hiromasa, T. Sato, T. Kurisaki, K. Kamijo, Y. Nabeshima, A. Fujisawa-Sehara, A metalloprotease-disintegrin participating in myoblast fusion, *Nature* 377 (1995) 652–656.
- [16] T. Kurisaki, A. Masuda, N. Osumi, Y. Nabeshima, A. Fujisawa-Sehara, Spatially- and temporally-restricted expression of meltrin  $\alpha$  (ADAM12) and  $\beta$  (ADAM19) in mouse embryo, *Mech. Dev.* 73 (1998) 211–215.
- [17] B.J. Gilpin, F. Loechel, M.-G. Mattei, E. Engvall, R. Albrechtsen, U. Wewer, A novel, secreted form of human ADAM 12 (meltrin  $\alpha$ ) provokes myogenesis in vivo, *J. Biol. Chem.* 273 (1998) 157–166.
- [18] M.F. Galliano, C. Huet, J. Frygeli, A. Polgren, U.M. Wewer, E. Engvall, Binding of ADAM 12, a marker of skeletal muscle regeneration, to the muscle-specific actin-binding protein,  $\alpha$ -actinin-2, is required for myoblast fusion, *J. Biol. Chem.* 275 (2000) 13933–13939.
- [19] A. Bornemann, R. Kuschel, A. Fujisawa-Sehara, Analysis for transcript expression of meltrin  $\alpha$  in normal, regenerating, and denervated rat muscle, *J. Muscle Res. Cell Motil.* 21 (2000) 475–480.
- [20] Z. Shi, W. Xu, F. Loechel, U.M. Wewer, L.J. Murphy, ADAM12, a disintegrin metalloprotease, interacts with insulin-like growth factor-binding protein-3, *J. Biol. Chem.* 275 (2000) 18574–18580.

- [21] F. Loechel, J.W. Fox, G. Murphy, R. Albrechtsen, U.M. Wewer, ADAM12-S cleaves IGFBP-3 and IGFBP-5 and is inhibited by TIMP-3, *Biochem. Biophys. Res. Commun.* 278 (2000) 511–515.
- [22] M. Asakura, M. Kitakaze, S. Takashima, Y. Liao, F. Ishikura, T. Yoshinaka, H. Ohmoto, K. Node, K. Yoshino, H. Ishiguro, H. Asanuma, S. Sanada, Y. Matsumura, H. Takeda, S. Beppu, M. Tada, M. Hori, S. Higashiyama, Cardiac hypertrophy is inhibited by antagonism of ADAM12 processing of HB-EGF: metalloproteinase inhibitors as a new therapy, *Nat. Med.* 8 (2002) 35–40.
- [23] Q. Kang, Y. Cao, A. Zolkiewska, Metalloprotease-disintegrin ADAM12 binds to the SH3 domain of Src and activates Src tyrosine kinase in C2C12 cells, *Biochem. J.* 352 (2000) 883–892.
- [24] Q. Kang, Y. Cao, A. Zolkiewska, Direct interaction between the cytoplasmic tail of ADAM12 and the Src homology 3 domain of p85 $\alpha$  activates phosphatidylinositol 3-kinase in C2C12 cells, *J. Biol. Chem.* 276 (2001) 24466–24472.
- [25] Y. Cao, Q. Kang, A. Zolkiewska, Metalloprotease-disintegrin ADAM12 interacts with  $\alpha$ -actinin-1, *Biochem. J.* 357 (2001) 353–361.
- [26] N. Kawaguchi, X. Xu, R. Tajima, P. Kronqvist, C. Sundberg, F. Loechel, R. Albrechtsen, U.M. Wewer, ADAM12 protease induces adipogenesis in transgenic mice, *Am. J. Pathol.* 160 (2002) 1895–1903.
- [27] P. Kronqvist, N. Kawaguchi, R. Albrechtsen, X. Xu, H.D. Schroder, B. Moghadaszadeh, F.C. Nielsen, C. Frohlich, E. Engvall, U.M. Wewer, ADAM12 alleviates the skeletal muscle pathology in mdx dystrophic mice, *Am. J. Pathol.* 161 (2002) 1535–1540.
- [28] K. Iba, R. Albrechtsen, B.J. Gilpin, F. Loechel, U.M. Wewer, Cysteine-rich domain of human ADAM12 (meltrin  $\alpha$ ) supports tumor cell adhesion, *Am. J. Pathol.* 154 (1999) 1489–1501.
- [29] K. Iba, R. Albrechtsen, B. Gilpin, C. Fröhlich, F. Loechel, A. Zolkiewska, K. Ishihuro, T. Kojima, W. Liu, J.K. Langford, R.D. Sanderson, C. Brakebusch, R. Fässler, U.M. Wewer, The cysteine-rich supports cell adhesion through syndecans and triggers signaling events that lead to  $\beta$ 1 integrin-dependent cell spreading, *J. Cell Biol.* 149 (2000) 1143–1155.
- [30] C.K. Thodeti, R. Albrechtsen, M. Grauslund, M. Asmar, C. Larsson, Y. Takada, A.M. Mercurio, J.R. Couchman, U.M. Wewer, ADAM12/syndecan-4 signaling promotes  $\beta$ 1 integrin-dependent cell spreading through PKC $\alpha$  and RhoA, *J. Biol. Chem.* 278 (2003) 9576–9584.
- [31] K. Eto, W. Puzon-McLaughlin, D. Sheppard, A. Sehara-Fujisawa, X.P. Zhang, Y. Takada, RGD-independent binding of integrin  $\alpha$ 9 $\beta$ 1 to the ADAM-12 and -15 disintegrin domains mediates cell–cell interaction, *J. Biol. Chem.* 275 (2000) 34922–34930.
- [32] K. Eto, C. Huet, T. Tarui, S. Kupriyanov, H.-Z. Liu, W. Puzon-McLaughlin, X.-P. Zhang, D. Sheppard, E. Engvall, Y. Takada, Functional classification of ADAMs based on a conserved motif for binding to integrin  $\alpha$ 9 $\beta$ 1: implications for sperm-egg binding and other cell interactions, *J. Biol. Chem.* 277 (2002) 17804–17810.
- [33] D.J. Burkin, S.J. Kaufman, The  $\alpha$ 7 $\beta$ 1 integrin in muscle development and disease, *Cell Tissue Res.* 296 (1999) 183–190.
- [34] T. Kurisaki, A. Masuda, K. Sudo, J. Sakagami, S. Higashiyama, Y. Matsuda, A. Nagabukuro, A. Tsuji, Y. Nabeshima, M. Asano, Y. Iwakura, A. Sehara-Fujisawa, Phenotypic analysis of meltrin  $\alpha$  (ADAM12)-deficient mice: involvement of meltrin  $\alpha$  in adipogenesis and myogenesis, *Mol. Cell. Biol.* 23 (2003) 55–61.
- [35] S. Schöber, D. Mielenz, F. Echtermeyer, S. Hapke, E. Pöschl, H. von der Mark, H. Moch, K. von der Mark, The role of extracellular and cytoplasmic splice domains of  $\alpha$ 7-integrin in cell adhesion and migration on laminins, *Exp. Cell Res.* 255 (2000) 303–313.
- [36] S. Crawley, E.M. Farrell, W. Wang, M. Gu, H.-Y. Huang, V. Huynh, B.L. Hodges, D.N.W. Cooper, S.J. Kaufman, The  $\alpha$ 7 $\beta$ 1 integrin mediates adhesion and migration of skeletal myoblasts on laminin, *Exp. Cell Res.* 235 (1997) 274–286.
- [37] A. Zolkiewska, Disintegrin-like/cysteine-rich region of ADAM12 is an active cell adhesion domain, *Exp. Cell Res.* 252 (1999) 423–431.
- [38] S. Liu, M. Slepak, M.H. Ginsberg, Binding of paxillin to the  $\alpha$ 9 integrin cytoplasmic domain inhibits cell spreading, *J. Biol. Chem.* 276 (2001) 37086–37092.
- [39] H. von der Mark, J. Dürr, A. Sonnenberg, K. von der Mark, R. Deutzmann, S.L. Goodman, Skeletal myoblasts utilize a novel  $\beta$ 1-series integrin and not  $\alpha$ 6 $\beta$ 1 for binding to the E8 and T8 fragments of laminin, *J. Biol. Chem.* 266 (1991) 23593–23601.
- [40] W.K. Song, W. Wang, R.F. Foster, D.A. Bielser, S.J. Kaufman, H36- $\alpha$ 7 is a novel integrin  $\alpha$  chain that is developmentally regulated during skeletal myogenesis, *J. Cell Biol.* 117 (1992) 643–657.
- [41] C.-C. Yao, B.L. Ziober, A.E. Sutherland, D.L. Mendrick, R.H. Kramer, Laminins promote the locomotion of skeletal myoblasts via the  $\alpha$ 7 integrin receptor, *J. Cell Sci.* 109 (1996) 3139–3150.
- [42] B.L. Ziober, Y. Chen, R.H. Kramer, The laminin-binding activity of the (7) integrin receptor is defined by developmentally regulated splicing in the extracellular domain, *Mol. Biol. Cell* 8 (1997) 1723–1734.
- [43] H. von der Mark, I. Williams, O. Wendler, L. Sorokin, K. von der Mark, E. Pöschl, Alternative splice variants of  $\alpha$ 7 $\beta$ 1 integrin selectively recognize different laminin isoforms, *J. Biol. Chem.* 277 (2002) 6012–6016.
- [44] J.T. Parsons, K.H. Martin, J.K. Slack, J.M. Taylor, S.A. Weed, Focal adhesion kinase: a regulator of focal adhesion dynamics and cell movement, *Oncogene* 19 (2000) 5606–5613.
- [45] W.K. Song, W. Wang, H. Sato, D.A. Bielser, S.J. Kaufman, Expression of  $\alpha$ 7 integrin cytoplasmic domains during skeletal muscle development: alternate forms, conformational change, and homologies with serine/threonine kinases and tyrosine phosphatases, *J. Cell Sci.* 106 (1993) 1139–1152.
- [46] B.L. Ziober, M.P. Vu, N. Waleh, J. Crawford, C.-S. Lin, R.H. Kramer, Alternative extracellular and cytoplasmic domains of the integrin  $\alpha$ 7 subunit are differentially expressed during development, *J. Biol. Chem.* 268 (1993) 26773–26783.
- [47] D.J. Burkin, M. Gu, B.L. Hodges, J.T. Campanelli, S.J. Kaufman, A functional role for specific spliced variants of the  $\alpha$ 7 $\beta$ 1 integrin in acetylcholine receptor clustering, *J. Cell Biol.* 143 (1998) 1067–1075.
- [48] J. Takagi, B.M. Petre, T. Walz, T.A. Springer, Global conformational rearrangements in integrin extracellular domains in outside-in and inside-out signaling, *Cell* 110 (2002) 599–611.
- [49] R.O. Hynes, Integrins: bidirectional, allosteric signaling machines, *Cell* 110 (2002) 673–687.
- [50] M. Shimaoka, J. Takagi, T.A. Springer, Conformational regulation of integrin structure and function, *Annu. Rev. Biophys. Biomol. Struct.* 31 (2002) 485–516.
- [51] B. Moghadaszadeh, R. Albrechtsen, L.T. Guo, M. Zaik, N. Kawaguchi, R.H. Borup, P. Kronqvist, H.D. Schroder, K.E. Davies, T. Voit, F.C. Nielsen, E. Engvall, U.M. Wewer, Compensation for dystrophin-deficiency: ADAM12 overexpression in skeletal muscle results in increased  $\alpha$ 7 integrin, utrophin and associated glycoproteins, *Hum. Mol. Genet.* 12 (2003) 2467–2479.
- [52] Z.Z. Bao, M. Lakonishok, S.J. Kaufman, A.F. Horwitz,  $\alpha$ 7 $\beta$ 1 integrin is a component of the myotendinous junction on skeletal muscle, *J. Cell Sci.* 106 (1993) 579–590.
- [53] P.T. Martin, S.J. Kaufman, R.H. Kramer, J.R. Sanes, Synaptic integrins in developing, adult, and mutant muscle: selective association of  $\alpha$ 1,  $\alpha$ 7A, and  $\alpha$ 7B integrins with the neuromuscular junction, *Dev. Biol.* 174 (1996) 125–139.
- [54] T. Velling, G. Collo, L. Sorokin, M. Durbeek, H. Zhang, D. Gullberg, Distinct  $\alpha$ 7A $\beta$ 1 and  $\alpha$ 7B $\beta$ 1 integrin expression patterns during mouse development:  $\alpha$ 7A is restricted to skeletal muscle but  $\alpha$ 7B is expressed in striated muscle, vasculature, and nervous system, *Dev. Dyn.* 207 (1996) 355–371.
- [55] U. Mayer, G. Saher, R. Fässler, A. Bornemann, F. Echtermeyer, H. von der Mark, N. Miosge, E. Pöschl, K. von der Mark, Absence of integrin  $\alpha$ 7 causes a novel form of muscular dystrophy, *Nat. Genet.* 17 (1997) 318–323.

- [56] Y.K. Hayashi, F.L. Chou, E. Engvall, M. Ogawa, C. Matsuda, S. Hirabayashi, K. Yokochi, B.L. Ziober, R.H. Kramer, S.J. Kaufman, E. Ozawa, Y. Goto, I. Nonaka, T. Tsukahara, J.Z. Wang, E.P. Hoffman, K. Arahata, Mutations in the integrin  $\alpha 7$  gene cause congenital myopathy, *Nat. Genet.* 19 (1998) 94–97.
- [57] B.L. Hodges, Y.K. Hayashi, I. Nonaka, W. Wang, K. Arahata, S.J. Kaufman, Altered expression of the  $\alpha 7\beta 1$  integrin in human and murine muscular dystrophies, *J. Cell Sci.* 110 (1997) 2873–2881.
- [58] D.J. Burkin, G.Q. Wallace, K.J. Nicol, D.J. Kaufman, S.J. Kaufman, Enhanced expression of the  $\alpha 7\beta 1$  integrin reduces muscular dystrophy and restores viability in dystrophic mice, *J. Cell Biol.* 152 (2001) 1207–1218.