

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/257346074>

Corrigendum to “Conformational analysis and vibrational study of sulfanilamide” [Vib. Spectrosc. 58 (2012) 12–18]

ARTICLE *in* VIBRATIONAL SPECTROSCOPY · JANUARY 2013

Impact Factor: 2 · DOI: 10.1016/j.vibspec.2012.08.007

READS

23

2 AUTHORS:



Gulce Ildiz

T.C. Istanbul Kultur University

10 PUBLICATIONS 26 CITATIONS

SEE PROFILE




Sevim Akyuz

T.C. Istanbul Kultur University

170 PUBLICATIONS 1,535 CITATIONS

SEE PROFILE



Corrigendum

Corrigendum to “Conformational analysis and vibrational study of sulfanilamide”
[Vib. Spectrosc. 58 (2012) 12–18]

Gulce Ogruc Ildiz*, Sevim Akyuz

Istanbul Kultur University, Science and Letters Faculty, Physics Department, Atakoy Campus, 34156 Bakirkoy, Istanbul, Turkey

After the above paper was published, we found an incorrect statement in the text on p. 14, a few errors in Table 4, and a wrong version of Fig. 3. They should be corrected as follows:

1. On p. 14, middle section of the right hand column, this sentence starting with “While the NC stretching mode. . .” should be removed from the text and replaced with the following sentence: “While the OSCC torsion mode is observed at 563 and 564 cm^{-1} in IR and Raman spectra, respectively, and is calculated as 524 and 526 cm^{-1} for conformers I and II, respectively, this mode cannot be observed in the calculated spectra of conformers III and IV.”
2. In Table 4, the calculated “DFT (harmonic)” vibrational frequencies (cm^{-1}) for “ ν_{21} , ν_{SO_2} sym” mode in columns 6–9 should be corrected as 1155, 1155, 1157, and 1157, respectively. The corresponding scale factor of the calculated SO_2 stretching frequencies should be 1.023 (instead of 1.033) in footnote “b” under the table.
3. On p. 16, the simulated spectral plots presented in Fig. 3 (inserted there from a previous revision of the work by the editor’s mistake) should be replaced by the final version shown below (in accordance with the data of Table 4).

DOI of original article: <http://dx.doi.org/10.1016/j.vibspec.2011.10.005>.

* Corresponding author.

E-mail address: g.ogruc@iku.edu.tr (G.O. Ildiz).




Fig. 3. Scaled theoretical IR spectra of global minimum conformer (I), second (II), third (III) and fourth (IV) lowest energy conformers, from bottom to top, respectively.