

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/230851720>

In vitro antiplasmodial activity of marine sponge *Clathria indica* associated bacteria against *Plasmodium falciparum*

ARTICLE in ASIAN PACIFIC JOURNAL OF TROPICAL BIOMEDICINE · SEPTEMBER 2012

DOI: 10.1016/S2221-1691(12)60367-0

READS

79

2 AUTHORS, INCLUDING:



[S Jacob Inbaneson](#)

Premier Medical Corporation Limited, Dama...


30 PUBLICATIONS 398 CITATIONS

SEE PROFILE



Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtb

Document heading

In vitro antiplasmodial activity of marine sponge *Clathria indica* associated bacteria against *Plasmodium falciparum*

Samuel Jacob Inbaneson, Sundaram Ravikumar^{*},

School of Marine Sciences, Department of Oceanography and Coastal Area Studies, Alagappa University, Thondi Campus, Thondi–623 409, Ramanathapuram District, Tamil Nadu, India

ARTICLE INFO

Article history:

Received 25 June 2012

Received in revised form 5 July 2012

Accepted 7 August 2012

Available online 28 August 2012

Keywords:

Antiplasmodial compounds

Biochemical constituents

*Clathria indica*IC₅₀*Plasmodium falciparum*

Sponge bacteria

ABSTRACT

Objective: To identify the possible antiplasmodial drugs from bacteria associated with marine sponge *Clathria indica*. **Methods:** *Clathria indica* samples were collected from Thondi coast and subjected for enumeration and isolation of associated bacteria. Filter sterilized extracts (100, 50, 25, 12.5, 6.25 and 3.125 μ g.mL⁻¹) from isolated bacterial isolates were screened for antiplasmodial activity against *Plasmodium falciparum* and potential extracts were also screened for biochemical constituents. **Results:** The count of bacterial strains were maximum in November 2007 (19×10^4 CFU.g⁻¹) and the average count was maximum during the monsoon season (107×10^3 CFU.g⁻¹). Thirty one morphologically different bacterial isolates were isolated from *Clathria indica* and the ethyl acetate bacterial extracts were screened for antiplasmodial activity against *Plasmodium falciparum*. The antiplasmodial activity of a isolate THB23 (IC₅₀ 28.80 μ g.mL⁻¹) extract is highly comparable with the positive control chloroquine (IC₅₀ 19.59 μ g.mL⁻¹) and 17 bacterial extracts which showed IC₅₀ value of more than 100 μ g.mL⁻¹. Statistical analysis reveals that, significant *in vitro* antiplasmodial activity ($P < 0.05$) was observed between the concentrations and time of exposure. The chemical injury to erythrocytes showed no morphological changes in erythrocytes by the ethyl acetate extract of bacterial strains after 48 h of incubation. The *in vitro* antiplasmodial activity might be due to the presence of carbohydrates and alkaloids in the ethyl acetate extracts of bacterial isolates. **Conclusions:** The ethyl acetate extracts of THB23 possesses novel compounds for the development of antiplasmodial drugs.

1. Introduction

Marine microorganisms are emerging as an exciting resource for the discovery of novel bioactive compounds. Marine microorganisms exhibit a wide range of biological activity such as antibacterial, antifungal, anti-inflammatory, antitumor, Cytotoxic, algicidal, antiviral and immunosuppressive[1–12]. The microorganisms associated with marine animals produce pharmaceutically valuable secondary metabolites, since the marine microorganisms live in a significantly different environment from those of the terrestrial microorganisms. Sponges are well known to be hosts for a large community of microorganisms, such as bacteria and fungi and some

of these microbes are probably host specific[13–27]. That exceeds the number of bacteria in seawater by two–three orders of magnitude[28]. Earlier investigations reported that, the microbial communities associated with sponges produce novel bioactive natural compounds with host specificity[29,30]. As marine sponges are a rich source of diverse bacterial populations, the present investigation was carried out to explore the antiplasmodial activity of *Clathria indica* associated bacteria from Palk Strait region.

2. Materials and methods

2.1. Isolation of sponge associated bacteria

Marine sponge *Clathria indica* was collected by by-catch at Thondi (Lat. 9° 44' 11'' N and Lon. 79° 1' 13'' E) in the Palk Strait region of Tamil Nadu and was authenticated by Dr. S. Lazarus, Emeritus Fellow (Retired), Centre for Marine Science and Technology, Manonmaniam Sundaranar

^{*}Corresponding author: Sundaram Ravikumar, School of Marine Sciences, Department of Oceanography and Coastal Area Studies, Alagappa University, Thondi Campus, Thondi–623 409, Ramanathapuram District, Tamil Nadu, India.

Tel: +914561–243470, 0091–9003306959

E-mail: ravibiotech201321@gmail.com

Foundation Project: This work was financially supported by Indian Council of Medical Research, New Delhi (grant No. 59/6/2002/BMS/TRM).

University, Rajakkamangalam, Kanyakumari District, Tamil Nadu, India. All the collected samples were washed thrice with tap water and twice with distilled water to remove the adhering associated animals. One gram of sponge samples was cut into small pieces and serially diluted. Diluted sample was subjected for continuous shaking in a thermostat shaker and plated in triplicate on Zobell Marine agar 2216 medium (HiMedia Laboratories Pvt. Limited, Mumbai, India) using pour plate method. The plates were incubated in an inverted position for 24 h at $(28 \pm 2)^\circ\text{C}$ and the colonies were counted and recorded. Based on the morphological characteristics (forms, elevation, margin and colour of the colony), the colonies were selected and restreaked thrice in a nutrient agar medium (HiMedia Laboratories Pvt. Limited, Mumbai, India) and stored on nutrient agar slants.



Figure 1. Percentage of antiplasmodial activity IC_{50} values of *Clathria indica* associated bacterial strains against *Palsmodium falciparum*.

2.2. Mass cultivation of total heterotrophic bacteria

A loopful inoculum of isolated bacterial strains were further inoculated into 500 mL conical flask containing 100 mL of nutrient broth (pH 7.2) prepared with 50% of aged seawater and kept at $(28 \pm 2)^\circ\text{C}$ with continuous shaking. Twenty milliliter of the broth culture was then transformed to 1000 mL of nutrient broth prepared with 50% of aged seawater and incubated for 4–5 days under continuous shaking.

2.3. Extraction of bioactive principles from bacteria

The mass cultures of isolated bacterial stains were adjusted to pH 5.0 using 1N hydrochloric acid and centrifuged at 3000 rpm for 5 min to remove cells. The supernatant was collected and was mixed with equal volume of ethyl acetate in a separating funnel. After vigorous shaking, the flask was kept undisturbed until two separate layers obtained (aqueous and organic). The upper organic phase was concentrated in a vacuum evaporator at 40°C and the crude extract was obtained. This process was repeated three times to obtain complete extraction of active principles.

2.4. Parasite cultivation

The antiplasmodial activity of isolated bacterial extracts was assessed against *Palsmodium falciparum* obtained

from the Jawaharlal Nehru Centre for Advanced Scientific Research, Indian Institute of Science, Bangalore, India. *Palsmodium falciparum* are cultivated in human O Rh⁺ red blood cells using RPMI 1640 medium (HiMedia Laboratories Private Limited, Mumbai, India)^[31] supplemented with O Rh⁺ serum (10%), 5% sodium bicarbonate (HiMedia Laboratories Private Limited, Mumbai, India) and $40 \mu\text{g.mL}^{-1}$ of gentamycin sulphate (HiMedia Laboratories Private Limited, Mumbai, India). Hematocrits were adjusted at 5% and parasite cultures were used when they exhibited 2% parasitaemia^[32].

2.5. In vitro antiplasmodial assay

Filter sterilized extracts ($100, 50, 25, 12.5, 6.25$ and $3.125 \mu\text{g.mL}^{-1}$) from 31 bacterial isolates were incorporated into 96 well tissue culture plate containing $200 \mu\text{L}$ of *Palsmodium falciparum* culture with fresh red blood cells diluted to 2% hematocrit. Negative control was maintained with fresh red blood cells and 2% parasitized *Palsmodium falciparum* diluted to 2% hematocrit, positive control was maintained with parasitized blood cells culture treated with chloroquine and Artemether^[33]. Parasitaemia was evaluated after 48 h by Giemsa stain and the average percentage suppression of parasitaemia was calculated by the following formula: Average % suppression of parasitaemia = $\frac{\text{Average \% parasitaemia in control} - \text{Average \% parasitaemia in test}}{\text{Average \% parasitaemia in control}} \times 100$.

2.6. Antiplasmodial activity calculation and analysis

The antiplasmodial activities of isolated bacteria were expressed by the inhibitory concentrations (IC_{50}) of the drug that induced a 50% reduction in parasitaemia compared to the control (100% parasitaemia). The IC_{50} values were calculated (concentration of extract in X axis and percentage of inhibition in Y axis) using Office XP (SDAS) software with linear regression equation. This activity was analyzed in accordance with the norms of antiplasmodial activity of Rasoanaivo et al^[34]. According to this norms, an extract is very active if $\text{IC}_{50} < 5 \mu\text{g.mL}^{-1}$, active $5 \mu\text{g.mL}^{-1} < \text{IC}_{50} < 50 \mu\text{g.mL}^{-1}$, weakly active $50 \mu\text{g.mL}^{-1} < \text{IC}_{50} < 100 \mu\text{g.mL}^{-1}$ and inactive $\text{IC}_{50} > 100 \mu\text{g.mL}^{-1}$.

2.7. Chemical injury to erythrocytes

To assess any chemical injury to erythrocytes that might be attributed to the extract, $200 \mu\text{L}$ of erythrocytes were incubated with $100 \mu\text{g.mL}^{-1}$ of the extract at a dose equal to the highest used in the antiplasmodial assay. The conditions of the experiment were maintained as in the case of antiplasmodial assay. After 48 h of incubation, thin blood smears were stained with Giemsa stain and observed for morphological changes under high-power light microscopy. The morphological findings were compared with those in erythrocytes that were uninfected and not exposed to extract^[35].

Table 1Counts of associated bacterial isolates from *Clathria indica*.

Sample No.	Month of Collection	THB $\times 10^4$ CFU.g ⁻¹	Season	THB $\times 10^3$ CFU.g ⁻¹
1	August 2007	3	Pre monsoon	27
2	September 2007	2		
3	October 2007	3		
4	November 2007	19	Monsoon	107
5	December 2007	11		
6	January 2008	2		
7	February 2008	4	Post monsoon	23
8	March 2008	2		
9	April 2008	1		
10	May 2008	2	Summer	53
11	June 2008	7		
12	July 2008	7		

Table 2Morphological characteristics and antiplasmodial IC₅₀ values of isolated bacterial isolates.

Isolate No.	Form	Elevation	Margin	Colour of the colony	IC ₅₀ (μ g.mL ⁻¹)
THB-10	Irregular	Raised	Filamentous	White	51.08
THB-11	Circular	Flat	Entire	Brownish yellow	58.83
THB-21	Irregular	Raised	Undulate	Waxy	35.47
THB-22	Irregular	Raised	Undulate	Pale	42.84
THB-23	Circular	Raised	Entire	Transparent yellow	28.80
THB-36	Circular	Raised	Entire	Waxy	75.30
THB-37	Irregular	Raised	Undulate	Light yellow	34.61
THB-49	Circular	Raised	Entire	Waxy	>100
THB-50	Circular	Convex	Entire	Transparent waxy	>100
THB-51	Circular	Raised	Entire	Transparent waxy	>100
THB-61	Circular	Raised	Entire	Centre white, surrounding transparent yellow	>100
THB-62	Circular	Raised	Entire	White	69.00
THB-74	Circular	Raised	Entire	Light yellow	>100
THB-75	Irregular	Raised	Undulate	White	>100
THB-76	Circular	Flat	Entire	Lemon yellow	56.41
THB-88	Irregular	Flat	Undulate	White	>100
THB-89	Circular	Raised	Entire	Light yellow	>100
THB-90	Circular	Convex	Entire	Dull white	>100
THB-102	Circular	Raised	Entire	Waxy	>100
THB-103	Circular	Flat	Entire	Orange	92.58
THB-104	Circular	Raised	Entire	Yellow	>100
THB-116	Irregular	Raised	Undulate	Light yellow	>100
THB-117	Circular	Raised	Entire	Yellow	>100
THB-118	Circular	Raised	Entire	Light brown	88.61
THB-130	Circular	Raised	Entire	Light yellow	>100
THB-131	Circular	Raised	Entire	Whitish yellow	>100
THB-141	Irregular	Flat	Curled	White	>100
THB-142	Circular	Convex	Entire	Orange	86.92
THB-151	Irregular	Raised	Undulate	Waxy	96.37
THB-152	Circular	Raised	Entire	Light yellow	>100
THB-153	Circular	Raised	Entire	Transparent white	74.32
Positive control					
Chloroquine					19.59
Artemether					4.09

Values are found significant between concentrations and time of exposure ($P < 0.05$)

3. Results

The counts of isolated bacterial isolates from the marine sponge *Clathria indica* samples are represented in Table 1.

The bacterial count was maximum in the month of November 2007 (19×10^4 CFU.g⁻¹) and minimum in the month of April 2008 (1×10^4 CFU.g⁻¹). The average count was maximum during the monsoon season (November – January) (107×10^3

Table 3

Biochemical constituents in chosen sponge associated bacterial isolates extracts.

Biochemical Constituents	THB21	THB22	THB23	THB37
Reducing sugars	+	+	+	+
Amino acids	–	–	–	–
Proteins	–	+	–	–
Alkaloids	+	+	+	+
Steroids	–	–	–	–
Triterpenoids	–	–	–	–

“+” indicates positive; “–” indicates negative.

CFU.g⁻¹) and followed by summer season (May – July) (53 × 10³ CFU.g⁻¹). A total of 31 different bacterial isolates were isolated from *Clathria indica* based on the morphological characteristics (Table 2). The extract of THB23 (28.80 µg.mL⁻¹) showed minimum level of IC₅₀ value and followed by THB37 (34.61 µg.mL⁻¹), THB21 (35.47 µg.mL⁻¹) and THB22 (42.84 µg.mL⁻¹). The extracts from THB10, THB11, THB36, THB62, THB76, THB103, THB118, THB142, THB151 and THB153 showed IC₅₀ values between 50 to 100 µg.mL⁻¹. Among 31 bacterial extracts screened for antiplasmodial activity, THB49, THB50, THB51, THB61, THB74, THB75, THB88, THB89, THB90, THB102, THB104, THB116, THB117, THB130, THB131, THB141 and THB152 showed IC₅₀ values more than 100 µg.mL⁻¹ (Table 2). The screened extracts showed significant antiplasmodial activity ($P < 0.05$) between the concentrations and time of exposure.

The microscopic observation of uninfected erythrocytes added with the ethyl acetate extracts from 31 bacterial isolates and uninfected erythrocytes from the blank column of the 96–well plate showed no morphological differences after 48 h of incubation. The analysis of preliminary biochemical constituents reveals that, the extracts from bacterial isolates have variety of biochemical constituents, namely alkaloids and reducing sugars (Table 3).

4. Discussion

Malaria is a devastating infectious disease and a major cause of morbidity and mortality throughout the world. There are up to 500 million clinical cases and 3 million deaths annually[36]. The drugs currently used to treat malaria come from three families such as quinolines, antifolates and artemisinin derivatives[37]. This major public health problem is aggravated by the widespread diffusion of drug resistant strains of *Plasmodium falciparum*. This increasing resistance to multiple drugs may be the result of defective DNA repair or other pathways responsible for genomic stability[38]. Antiplasmodial compounds from natural resources would be the alternative to overcome the resistance problem. In that, the marine invertebrates play a major role and harbor numerous microorganisms within their tissues[14]. Sponge mesohyl was referred as “micro–environments” providing a broad variety of ecological niches[39].

All the collected *Clathria indica* sponge samples reported to harbour associated bacteria and this might be due to bacteria can contribute up to 40% of the sponge biomass and are probably permanently associated with the host sponge unless they are disturbed by external stress factors[40–44]. The present study also observed variable counts in different months and different seasons. Similarly, Taylor et al. reported that, the substantial variability of microbial communities between different species of sponges although there was little variability of microbial communities within each sponge species[45]. Taylor et al. observed that, the sponge associated microorganisms considerably varied with geographic[14].

In vitro screening for antiplasmodial activity is one of the key tools for identification lead compounds for new antiplasmodials. In this study, the THB23 showed excellent antiplasmodial activity and it could be comparable with the positive control chloroquine. The antiplasmodial nature of sponge associated microbes due to the microbes can sense, adapt and respond to their environment quickly and can compete for defense and survival by the generation of unique secondary metabolites. The marine microorganisms showed potential antiplasmodial activity against *Plasmodium falciparum*[46–48]. According to Rasoanaivo et al. 13%, 32% and 55% of extracts from isolated bacterial isolates were classified as active, weakly active and inactive respectively (Figure 1)[34].

The ethyl acetate extracts of potential strains showed the presence of reducing sugars and alkaloids. The presence of these biochemical constituents in the potential isolates extracts, especially sugars and alkaloids[49,50]. The mode of action could be due to the inhibition of *Plasmodium falciparum* merozoites invasion into the erythrocytes[51] and disruption of *Plasmodium falciparum* rosettes[52] by the carbohydrates; inhibition of *Plasmodium falciparum* fatty acid biosynthesis[53], inhibition of hemozoin biocrystallization by the alkaloids[54]. Alkaloids are nitrogenous compounds derived from many biogenetic precursors and possessing antimalarial activity[55]. El Sayed et al. reported that, the marine sponge associated microbial alkaloids (curcuphenol) have antiplasmodial activity[56]. Numerous alkaloids isolated from marine sources have antimalarial activity[57]. These findings could encourage the sponge associated bacteria derived compounds for the

antiplasmodial drug development. It is concluded that, the ethyl acetate extract of *Clathria indica* associated bacterial isolate THB23 proved as a massive resource to develop potential antiplasmodial drugs. This justifies our continuing research to fractionate the *Clathria indica* associated bacterial extracts to determine the active principles responsible for antiplasmodial activity.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgments

The authors are thankful to the authorities of Alagappa University for providing required facilities and also to Indian Council of Medical Research, New Delhi for financial assistance. The authors are also grateful to Prof. Dr. Hemalatha Balaraman, Molecular Biology and Genetics Unit, Jawaharlal Nehru Centre for Advanced Scientific Research, Bangalore for providing us the parasite culture.

References

- [1] Kim TK, Hewavitharana AK, Shaw PN, Fuerst JA. Discovery of a new source of rifamycin antibiotics in marine sponge actinobacteria by phylogenetic prediction. *Appl Environ Microbiol* 2006; **72**: 2118–25.
- [2] Li ZY, Liu Y. Marine sponge *Craniella australiensis* associated bacterial diversity revelation based on 16S rDNA library and biologically active actinomycetes screening, phylogenetic analysis. *Lett Appl Microbiol* 2006; **43**: 410–416.
- [3] Ravikumar S, Thajuddin N, Suganthi P, Jacob Inbaneson S, Vinodkumar T. Bioactive potential of seagrass bacteria against human bacterial pathogens. *J Env Bio* 2010; **31**: 387–9.
- [4] Prakash Williams G, Babu S, Ravikumar S, Kathiresan K, Arul Prathap S, Chinnapparaj S, et al. Antimicrobial activity of tissue and associated bacteria from benthic sea anemone *Stichodactyla haddoni* against microbial pathogens. *J Environ Biol*. 2007; **28**: 782–93.
- [5] Gandhimathi R, Arunkumar M, Selvin J, Thangavelu T, Sivaramakrishnan S, Kiran GS, et al. Antimicrobial potential of sponge associated marine actinomycetes. *J Med Mycol* 2008; **18**: 16–22.
- [6] Ravikumar S, Jacob Inbaneson S, Uthiraselvam M, Rajini Priya S, Ramu A, Beula Banerjee M. Diversity of endophytic actinomycetes from Karangkadu mangrove ecosystem and its antibacterial potential against bacterial pathogens. *J Pharm Res* 2011; **4**(1): 294–296.
- [7] Ravikumar S, Jacob Inbaneson S, Uthiraselvam M, Kaleeswari R, Ramu A, Beula Banerjee M, et al. Antibacterial Activity of heterotrophic endophytes from Karangkadu mangrove ecosystem, India. *J Pharm Res* 2011; **4**(1): 195–198.
- [8] Kwon HC, Kauffman CA, Jensen PR, Fenical W. Marinomycins A–D, antitumour antibiotics of a new structure class from a marine actino–mycete of the recently discovered genus *Marinispora*. *J Am Chem Soc*. 2006; **128**: 1622–32.
- [9] Adinarayana G, Venkateshan MR, Bapiraju VVSNK, Sujatha P, Premkumar J, Ellaiah P, et al. Cytotoxic compounds from the marine actinobacterium *Streptomyces corchorusii* AUBN1/7. *Rus J Bioorg Chem* 2006; **32**(3): 295–300.
- [10] Kim MJ, Jeong SY, Lee SJ. Isolation, identification and algicidal activity of marine bacteria against *Cochlodinium polykrikoides*. *J Appl Phycol* 2008; **20**: 1069–78.
- [11] Ravikumar S, Krishnakumar S, Jacob Inbaneson S, Gnanadesigan M. Antagonistic activity of marine actinomycetes from Arabian sea coast. *Arc Appl Sci Res* 2010; **2**(6): 273–280.
- [12] Ravikumar S, Jacob Inbaneson S, Sengottuvel R, Ramu A. Assessment of endophytic bacterial diversity among mangrove plants and their antibacterial activity against bacterial pathogens. *Ann Biol Res*. 2010; **1**(4): 240–247.
- [13] Hentschel U, Uscher KM, Taylor MW. Marine sponges as microbial fermenters. *FEMS Microbiol Ecol* 2006; **55**: 167–177.
- [14] Taylor MW, Radax R, Steger D, Wagner M. Sponge-associated microorganisms: evolution, ecology and biotechnological potential. *Microbiol Mol Biol Rev* 2007; **71**: 295–347.
- [15] Lafi FF, Fuerst JA, Fieseler L, Hentschel U. Widespread distribution of Poribacteria in Demospongiae. *Appl Environ Microbiol* 2009; **75**: 5695–5699.
- [17] Hochmuth T, Piel J. Polyketide synthases of bacterial symbionts in sponges – evolution based applications in natural products research. *Phytochem* 2009; **70**: 1841–1849.
- [18] Piel J. Metabolites from symbiotic bacteria. *Nat Prod Rep* 2009; **26**: 338–362.
- [19] Taylor MW, Thacker RW, Hentschel U. Genetics. Evolutionary insights from sponges. *Sci* 2007; **316**: 1854–1855.
- [20] Taylor MW, Hill RT, Piel J, Thacker RW, Hentschel U. Soaking it up: the complex lives of marine sponges and their microbial associates. *ISME J* 2007; **1**: 187–190.
- [21] Webster NS, Cobb RE, Negri AP. Temperature thresholds for bacterial symbiosis with a sponge. *ISME J* 2008; **2**: 830–842.
- [22] Webster NS, Xavier JR, Freckelton M, Motti CA, Cobb R. Shifts in microbial and chemical patterns within the marine sponge *Aplysina aerophoba* during a disease outbreak. *Environ Microbiol* 2008b; **10**: 3366–3376.
- [23] Webster NS, Taylor MW, Behnam F, Luckner S, Rattei T, Whalan S, et al. Deep sequencing reveals exceptional diversity and modes of transmission for bacterial sponge symbionts. *Environ Microbiol* 2010; **12**: 2070–2082.
- [24] Webster NS, Cobb RE, Soo R, Anthony SL, Battershill CN, Whalan et al. Bacterial community dynamics in the marine sponge *Rhopaloeides odorabile* under *in situ* and *ex situ* cultivation. *Mar Biotechnol* 2010; doi:10.1007/s10126-010-9300-4.
- [25] Lee OO, Chui PY, Wong YH, Pawlik JR, Qian PY. Evidence of vertical transmission of bacterial symbionts from adult to embryo in the Caribbean sponge *Svenzea zaei*. *Appl Environ Microbiol* 2011; **75**: 6147–6156.
- [26] Lee OO, Wang Y, Yang J, Lafi FF, Al-Suwailem A, Qian PY.

- Pyrosequencing reveals highly diverse and species specific microbial communities in sponges from the Red sea. *ISME J* 2011; doi:10.1038/ismej.2010.165.
- [27] Kamke J, Taylor MW, Schmitt S. Activity profiles for marine sponge associated bacteria obtained by 16S rRNA gene comparisons. *ISME J* 2010; **4**: 498–508.
- [28] Thoms C, Horn M, Wagner M, Hentschel U, Proksch P. Monitoring microbial diversity and natural product profiles of the sponge *Aplysina cavernicola* following transplantation. *Mar Biol* 2003; **142**: 685–92.
- [29] Kennedy J, Baker P, Piper C, Cotter PD, Walsh M, Mooij MJ et al. Isolation and analysis of bacteria with antimicrobial activities from the marine sponge *Haliclona simulans* collected from Irish waters. *Mar Biotechnol* 2009; **11**: 384–396.
- [30] Blunt JW, Copp BR, Hu WP, Munro MH, Northcote PT, Prinsep MR. Marine natural products. *Nat Prod Rep* 2007; **24**: 31–86.
- [31] Moore GE, Gerner RE, Franklin HA. Cultures of normal human leukocytes. *J Am Med Assoc* 1967; **199**: 519–24.
- [32] Trager W. The cultivation of *Plasmodium falciparum*: applications in basic and applied research in malaria. *Ann Trop Med Parasitol* 1987; **82**: 511–29.
- [33] Ravikumar S, Jacob Inbaneson S, Suganthi P. Seaweeds as a source of lead compounds for the development of new antiparasitic drugs from South East coast of India. *Parasitol Res* 2011; **109**: 47–52.
- [34] Rasoanaivo P, Ratsimamanga Urverg S, Ramanitrasimbola D, Rafatro H, Rakoto Ratsimamanga A. Criblage d'extraits de plantes de Madagascar pour recherche d'activite antipaludique et d'effet potentialisateur de la chloroquine. *J Ethnopharmacol* 1992; **64**: 117–26.
- [35] Waako PJ, Katuura E, Smith P, Folb P. East African medicinal plants as a source of lead compounds for the development of new antimalarial drugs. *Afr J Ecol* 2007; **45**(1): 102–6.
- [36] WHO. World Health Organization malaria reports. World Health Organization, Geneva. 2008.
- [37] Von Seidlein L, Greenwood BM. Mass administration of antimalarial drugs. *Trends Parasitol* 2003; **19**: 452–60.
- [38] Trotta RF, Brown ML, Terrell JC, Geyer JA. Defective DNA repair as a potential mechanism for the rapid development of drug resistance in *Plasmodium falciparum*. *Biochem* 2004; **43**: 4885–91.
- [39] Thiel V, Neulinger SC, Staufenberger T, Schmaljohann R, Imhoff JF. Spatial distribution of sponge-associated bacteria in the Mediterranean sponge *Tethya aurantium*. *FEMS Microbiol Ecol* 2007; **59**: 47–63.
- [40] Lopez-Legentil S, Song B, McMurray SE, Pawlik JR. Bleaching and stress in coral reef ecosystem; hsp70 expression by the giant barrel sponge *Xestospongia muta*. *Mol Ecol* 2008; **17**: 1840–1849.
- [41] Lemoine N, Buell N, Hill A, Hill M. Assessing the utility of sponge microbial symbionts communities as models to study global climate change: a case study with *Halichondria bowerbanki*. In Porifera Research: Biodiversity, innovation and sustainability. Custodio Mr, Lobo Hajdu G, Hajdu E, Muricy G (eds). Rio de Janeiro, Brazil: Serie Livros, Museu Nacional, 2007; pp. 419–425.
- [42] Lopez-Legentil S, Erwin PE, Pawlik JR, Song B. Effects of sponge bleaching on ammonia-oxidizing Archaea: distribution and relative expression of ammonia monooxygenase genes associated with the giant barrel sponge *Xestospongia muta*. *Mol Ecol* 2010; **60**: 561–571.
- [43] Webster NS, Cobb RE, Negri AP. Temperature thresholds for bacterial symbiosis with a sponge. *ISME J* 2008; **2**: 830–842.
- [44] Selvin J, Shanmugha Priya S, Seghal Kiran G, Thangavelu T, Sapna Bai N. Sponge associated marine bacteria as indicators of heavy metal pollution. *Microbiol Res* 2007; **164**: 352–363.
- [45] Taylor MW, Schupp PJ, Dahllof I, Kjelleberg S, Steinberg PD. Host specificity in marine sponge-associated bacteria and potential implications for marine microbial diversity. *Environ Microbiol* 2004; **6**: 121–30.
- [46] McPhail KL, Correa J, Linington RG, Gonzalez J, Ortega-Barria E, Capson TL, Gerwick WH. Antimalarial linear lipopeptides from a Panamanian strain of the marine cyanobacterium *Lyngbya majuscula*. *J Nat Prod* 2007; **70**: 984–8.
- [47] Linington RG, Gonzalez J, Urena LD, Romero LI, Ortega-Barria E, Gerwick WH. Venturamides A and B: Antimalarial constituents of the Panamanian marine cyanobacterium *Oscillatoria* sp. *J Nat Prod* 2007; **70**: 397–401.
- [48] Linington RG, Clark BR, Trimble EE, Almanza A, Urena LD, Kyle DE, et al. Antimalarial peptides from marine cyanobacteria: Isolation and structural elucidation of Gallinamide A. *J Nat Prod*. 2009; **72**(1):14–7.
- [49] Ootoguro K, Ishiyama A, Kobayashi M, Sekiguchi H, Izuhara T, Sunazuka T, et al. *In vitro* and *in vivo* antimalarial activities of a carbohydrate antibiotic, prumycin against drug resistant strains of Plasmodia. *J Antibiot* 2004; **57**(6): 400–2.
- [50] Stierle AC, Cardellina JHI, Singleton FL. A marine Micrococcus produces metabolites ascribed to the sponge *Tedania ignis*. *Experientia* 1988; **44**: 1021.
- [51] Adams Y, Smith SL, Schwartz-Albiez R, Andrews KT. Carrageenans inhibit the *in vitro* growth of *Plasmodium falciparum* and cytoadhesion to CD36. *Parasitol Res*. 2005; **97**: 290–4.
- [52] Rowe A, Berendt AR, Marsh K, Newbold CI. *Plasmodium falciparum*: a family of sulphated glycoconjugates disrupts erythrocyte rosettes. *Exp Parasitol* 1994; **79**: 506–516.
- [53] Tasdemir D, Topaloglu B, Perozzo R, Brun R, O'Neill R, Carballeira NM, et al. Marine natural products from the Turkish sponge *Agelas oroides* that inhibit the enoyl reductases from *Plasmodium falciparum*, *Mycobacterium tuberculosis* and *Escherichia coli*. *Bioorg Med Chem* 2007; **15**: 6834–45.
- [54] Dubar F, Egan TJ, Pradines B, Kuter D, Ncokazi KK, Forge D, et al. The antimalarial ferroquine: role of the metal and intramolecular hydrogen bond in activity and resistance. *ACS Chem Biol* 2010; DOI: 10.1021/cb100322v.
- [55] Saxena S, Pant N, Jain DC, Bhakuni RS. Antimalarial agents from plant sources. *Cur Sci* 2003; **85**(9): 1314–29.
- [56] El Sayed KA, Yousaf M, Hamann MT, Avery MA, Kelly M, Wipf P. Microbial and chemical transformation studies of the bioactive marine sesquiterpenes (S)-(+)-curcuphenol and -curcudiol isolated from a deep reef collection of the Jamaican sponge *Didiscus oxeata*. *J Nat Prod* 2002; **65**(11): 1547–53.
- [57] Peng J, Rao, Karumanchi V, Choo YM, Hamann MT. In modern alkaloids. Fattorusso, E. and O. Tagliatela-Scafati, Eds; Wiley-VCH Publishers, Weinheim, Germany; 2008; p. 189–232.