

Erratum

Erratum to “On the angular distributions of molecular photoelectrons: dipole cross-sections for fixed-in-space and randomly oriented molecules”
[Chem. Phys. Lett. 346 (2001) 341] [☆]

J.C. Arce ^a, J.A. Sheehy ^b, P.W. Langhoff ^{b,c,d,*}, O. Hemmers ^e, H. Wang ^{e,f},
P. Focke ^g, I.A. Sellin ^g, D.W. Lindle ^e

^a Departamento de Química, Universidad del Valle, A.A. 25360 Cali, Colombia

^b Air Force Research Laboratory, AFRL/PRS, Edwards AFB, CA 93524-7680, USA

^c Department of Chemistry, Indiana University, Bloomington, IN 47405-4001, USA

^d San Diego Supercomputer Center, University of California, 9500 Gilman Drive, La Jolla, CA 92093-0505, USA

^e Department of Chemistry, University of Nevada, Las Vegas, NV 89154-4003, USA

^f MAX-Lab, Lund University, P.O. Box 118, 22100 Lund, Sweden

^g Department of Physics, University of Tennessee, Knoxville, TN 37996-1600, USA

Received 16 October 2001

The publisher regrets that the figures appearing in the original publication were inadvertently typeset at reduced magnification; full size Fig. 1 are given on the next page.

[☆] PII of original article S0009-2614(01)00920-4.

* Corresponding author. Fax: +1-858-534-5113.

E-mail address: langhoff@drifter.sdsc.edu (P.W. Langhoff).




Fig. 1. (a) Photoionization anisotropy factor $\beta(h\nu)$ for the K-shell ($1\sigma_g^2, 1\sigma_u^2$) of molecular nitrogen: (●) present experimental values; previous measurements as indicated; (---) Eq. (18). Error bars for the present measurements are smaller than the data symbol shown. (b) Photon energy variations of $\rho(h\nu)$ and of the body-frame angle-averages employed in Eq. (18). The former is inferred from measured photoabsorption anisotropy data [31–33] and the latter from measured body-frame photoelectron angular distributions [6], employing the results of static exchange calculations and high-photon energy limits (arrows) to provide continuous curves, as discussed in the text.