

The postthrombotic syndrome in relation to venous hemodynamics, as measured by means of duplex scanning and strain-gauge plethysmography

José H. Haenen, RVT, Mirian C.H. Janssen, MD, Herman van Langen, PhD, Wim N.J.C. van Asten, PhD, Hub Wollersheim, MD, PhD, Martin A. van 't Hof, PhD, Stefan H. Skotnicki, MD, PhD, and Theo Thien, MD, PhD, Nijmegen, The Netherlands

Purpose: Venous hemodynamics were evaluated in relation to the postthrombotic syndrome (PTS) 7 to 13 years after deep venous thrombosis (DVT).

Methods: The presence of flow, reflux, and compressibility of 1394 vein segments in 82 patients was assessed by means of duplex scanning. The venous outflow resistance was measured by means of strain-gauge plethysmography. The venous hemodynamics were related to the clinical severity of the PTS, characterized by the CEAP (clinical, etiologic, anatomic, pathophysiologic) classification.

Results: In patients with severe clinical symptoms of PTS, the prevalence of reflux was significantly higher. There was no relationship between the severity of the PTS and the noncompressibility or the combination of reflux and noncompressibility or an increased venous resistance. By means of multiple regression analysis with the variables of age, gender, reflux, and venous resistance, age and reflux were shown to be the main contributors to the severity of PTS. Significantly more patients (64%) with severe signs of PTS had a combination of deep and superficial reflux. In each of the traceable vein segments, the mean of the CEAP classification was calculated for the vein segments with and without reflux. In the proximal superficial femoral vein ($P < .001$), distal superficial femoral vein ($P < .05$), and popliteal vein ($P < .05$), a significantly higher mean CEAP classification was found in the veins with reflux, whereas in the distal, long, and short saphenous veins, no such relationship was found.

Conclusion: Most patients with severe PTS had a combination of deep and superficial reflux. Reflux in the deep proximal veins contributes significantly to the PTS. (J Vasc Surg 1999;29:1071-6.)

Deep venous thrombosis (DVT) may cause persistent venous abnormalities, resulting in postthrombotic syndrome (PTS). Venous hypertension caused by a persistent venous obstruction, valve incompetence, or a combination of the two is considered to

be the major cause of PTS.¹⁻⁴ It is unclear whether an obstruction, reflux, or a combination of the two provides the main contribution to PTS. The presence of reflux in the popliteal vein is considered to be important in the pathophysiology of PTS,⁵⁻⁷ but the influence of venous abnormalities in other areas is less clear.

In this study, the venous hemodynamics, in relation to the severity of the PTS, were evaluated in patients 7 to 13 years after DVT. Local venous compressibility, reflux, and the presence of flow were measured by means of color flow duplex scanning, and the venous resistance in the postthrombotic extremity was measured by means of strain-gauge plethysmography. Because duplex scanning provided a local evaluation of the veins, the venous resistance of the entire leg was measured by means of plethysmography.

From the Clinical Vascular Laboratory (Mrs Haenen, Dr van Langen, and Dr van Asten), the Department of Thoracic and Cardiac Surgery (Prof Dr Skotnicki), the Department of General Internal Medicine (Dr Janssen, Dr Wollersheim, and Prof Dr Thien), and the Department of Medical Statistics (Dr van 't Hof), University Hospital Nijmegen.

Reprint requests: J. H. Haenen, University Hospital Nijmegen, Clinical Vascular Laboratory 810, PO Box 9101, 6500 HB Nijmegen, The Netherlands.

Copyright © 1999 by The Society for Vascular Surgery and International Society for Cardiovascular Surgery, North American Chapter.

0741-5214/99/\$8.00 + 0 24/1/96861

Table I. The number of patients in each classification of chronic lower extremity venous disease, the CEAP (clinical, etiologic, anatomic, pathophysiologic) scale

Class	Signs of chronic lower-extremity venous disease	Patients (n)
0	No visible or palpable signs of venous disease	22
1	Telangiectases, reticular veins, malleolar flare	11
2	Varicose veins	11
3	Edema without skin changes	10
4	Skin changes ascribed to venous disease (pigmentation, lipodermatosclerosis)	24
5	Skin changes with healed ulceration	4
6	Skin changes with active ulceration	0

Table II. Characteristics of the patients at follow-up examination for the different CEAP (clinical, etiologic, anatomic, pathophysiologic) classifications

CEAP classification	0 (n = 22)	1 to 3 (n = 32)	4 to 6 (n = 28)
Mean age (years) \pm SD	45 \pm 9	50 \pm 15	64 \pm 14
Mean body mass index (kg/m ²) \pm SD	24 \pm 3	27 \pm 6	25 \pm 4
Men (%)	55	66	82
Prevalence abnormal venous resistance (%)	5	6	21
Reflux			
Not present(%)*	32	22	4
Only deep (%)	41	41	29
Only superficial (%)	4	3	3
Superficial + deep (%)†	23	34	64
Location of thrombi at initial phlebography			
Proximal (%)	4	6	7
Proximal and distal (%)	17	21	24
Distal (%)	6	12	2
Deep (%)	13	28	21
Deep and superficial (%)	13	11	13

* $P < .05$, † $P < .01$ (Chi-square test).

METHODS

The records of all the patients at our institution who had DVT between 1983 and 1989 were reviewed. DVT was diagnosed by means of phlebography. Of the 256 patients, 92 died, 33 were unable to return for reexamination, 26 could not be traced, and 23 refused to participate in the study. The remaining 82 patients were reexamined 7 to 13 years (mean, 10 years) after the initial DVT. The study was approved by the ethics committee of the hospital, and informed consent was obtained from the participants.

All patients had been treated with heparin, administered intravenously for 5 to 10 days during the acute phase, and with oral anticoagulants for at least 12 weeks. All patients had been instructed to wear graduated compression stockings.

A physician, unaware of the findings of the duplex scanning and plethysmography, classified the patients according to the seven-point CEAP scale (clinical, etiologic, anatomic, pathophysiologic, class 0 to 6) of the update of the Ad Hoc Committee on

Reporting Standards in Venous Disease.⁸ For the sake of simplicity, the tables and figures were presented in three categories: (1) patients with no signs of PTS (class 0); (2) patients with mild signs of PTS (classes 1 to 3); and (3) patients with severe signs (classes 4 to 6) of PTS.

The venous resistance in the postthrombotic extremity was measured by means of strain-gauge plethysmography, as described previously.⁹⁻¹¹ The patients were examined in the supine position with pneumatic cuffs around the thighs and strain-gauges around the calves. After inflating the cuffs, the venous volume increased. When the maximum volume was reached, the cuff-pressure was released, resulting in a volume decrease measured by means of plethysmography. The tangent to the slope of the volume decrease was drawn 0.5 seconds after pressure release, and the slope was converted into a venous flow rate. This procedure was performed at five different cuff pressures. The corresponding venous flow rates were plotted against the effective cuff pressures. The slope of the line straight through

the points gave an angle β , and by means of analogy with Ohm's law, the venous resistance could be calculated as $1/\tan \beta$. A venous resistance exceeding 0.8 mm Hg/min was considered abnormal.^{10,11}

Duplex scanning was performed by two experienced vascular technicians using a Toshiba SSA 270A scanner with a 3.75 or 5 MHz probe on the low-flow setting. The vascular technicians scanned the patients without knowledge of the original location of the thrombosis. All veins were examined with the patient in a 45-degree sitting position, with the knees flexed and the feet resting on a footstool. The veins examined were the common femoral vein, the superficial femoral vein (proximal, middle, and distal), the long saphenous vein (proximal, middle, and distal), the popliteal vein, the short saphenous vein, and the posterior and anterior tibial, peroneal, and gastrocnemial veins. In the calf, each artery is accompanied by two veins. The two veins were numbered; number 1 was the most superficial vein, and number 2 was the deeper vein.

In the longitudinal plane, the presence of venous flow and reflux was measured. Proximally, reflux was measured after the Valsalva maneuver in the distal veins by means of distal manual compression with sudden release. Pathological reflux in the proximal veins was defined as a reversed flow duration of more than 1 second; in the distal veins, pathological reflux was defined as a reversed flow duration of more than 0.5 second, as described previously.¹² This previous study showed that our method with the patient in the sitting position is compatible with the method with patients in an upright position and the cuff deflation technique. Superficial or deep reflux was defined as the presence of abnormal venous reflux in at least one vein segment. Combined superficial and deep reflux was defined as the presence of abnormal venous reflux in at least two vein segments. Compressibility was assessed in the transverse plane. A vein was considered noncompressible when it was not totally compressed under gentle pressure of the duplex probe applied to the skin overlying the vein.

The relation between the seven-point CEAP scale and the presence of an abnormal venous resistance and body mass index (BMI) was studied with the chi-square test. The relation between the seven-point CEAP scale and the outcome of the venous evaluation by means of duplex scanning was studied at patient level, adjusted for age with multiple regression. The contribution of age, gender, reflux, and the venous resistance to the seven-point CEAP scale was also studied with multiple regression. The results were considered significant when $P \leq .05$ (two-sided).

RESULTS

The characteristics of the 82 patients are shown in Tables I and II. At the time of admission to the follow-up study, the mean age of the patients was 53 years (range, 21 to 80 years), their mean BMI was 26 kg/m² (SD \pm 5); 56 patients (68%) were men, and 26 (32%) were women. The DVT was located in the right leg in 48 patients (59%) and in the left leg of 34 patients (41%). There was no significant relationship between the seven-point CEAP scale and the BMI.

The venous resistance, as measured by means of strain-gauge plethysmography, was abnormal in only nine patients (11%). The prevalence of an abnormal venous resistance was not significantly higher in patients with severe clinical symptoms of PTS. Significantly more of the patients (64%) with severe signs of PTS had a combination of deep and superficial reflux (Table II).

The duplex findings for all the 1394 vein segments of the patients were related to the clinical findings (Fig 1). Because the classification of PTS increased with age, an adjustment was made for age in the statistical analysis. It was shown by means of regression analysis that only reflux could explain the PTS classification. No relation was found between PTS and noncompressibility or a combination of reflux and noncompressibility. Flow was present in almost all noncompressible vein segments (93%). The nontraceable veins were almost all located distally (95%). These nontraceable veins were not included in further analysis.

Because only reflux was shown to have a significant relationship to the CEAP classification, reflux was evaluated more extensively. In each of the traceable vein segments, the mean of the CEAP classification was calculated for the vein segments with and without reflux (Table III). In the proximal superficial femoral vein ($P < .001$), distal superficial femoral vein ($P < .05$), and popliteal vein ($P < .05$), a significantly higher mean CEAP classification was found in the veins with reflux, whereas in the distal, long, and short saphenous veins, no such significant relationship was found.

After testing the contributions of the factors age, gender, reflux, and the venous resistance to the PTS by means of multiple regression, age ($P < .001$) and reflux ($P < .01$) appeared to be the main contributors to the severity of PTS.

DISCUSSION

It is possible to detect venous abnormalities by means of duplex scanning, sometimes even better than with invasive tests such as phlebography.^{13,14}




Fig 1. The relationship between venous hemodynamics in all segments and the CEAP (clinical, etiologic, anatomic, pathophysiologic) classification.

Table III. The mean of the CEAP (clinical, etiologic, anatomic, pathophysiologic) classification in the traceable veins, with and without reflux

Traceable veins	n	Mean CEAP in veins with reflux	n	Mean CEAP in veins without reflux
Proximal superficial femoral*	40	2.78	42	1.61
Middle superficial femoral	30	2.56	52	1.96
Distal superficial femoral*	19	2.71	58	1.93
Popliteal*	24	2.72	56	1.88
Proximal long saphenous	8	3.03	63	2.26
Middle long saphenous	26	2.29	56	2.13
Distal long saphenous	12	2.56	47	2.32
Short saphenous	15	2.67	62	1.96
Posterior tibial 1	17	2.17	51	2.16
Posterior tibial 2	14	2.62	35	2.27
Anterior tibial 1	25	2.21	51	2.10
Anterior tibial 2	21	2.43	55	2.14
Peroneal 1	5	1.12	61	2.26
Peroneal 2	4	1.08	55	2.23
Gastrocnemial 1	5	1.60	44	2.23
Gastrocnemial 2	7	1.56	37	2.17

* $P < .05$.

Descending phlebography underestimates severe reflux and is usually used as a means of evaluating deep vein reflux,¹³ whereas the presence of superficial reflux seems also to be important.¹⁵⁻¹⁷

In a previous study,¹⁸ the relationship between the initial phlebography and the outcome measured by means of duplex scanning was evaluated in the same patient database (except for the 10 patients with recurrent thrombosis). In the current study, the definition of pathological reflux for the distal veins

was restricted from 1 second to 0.5 second; this definition was derived from another study.¹²

Duplex scanning provides local information on the veins. The consequences of these local abnormalities to the entire leg are uncertain and can be measured with plethysmography.

The prevalence of an abnormal venous resistance was not significantly higher in patients with severe clinical symptoms of PTS. This lack of statistical significance might be caused by the small sample size.

But because, in contrast to other studies,⁴ no relation was found between (reflux and) noncompressibility and the clinical signs of PTS either, another explanation might be that in our patients almost all noncompressible vein segments were partially occluded because flow was present in almost all (93%) noncompressible veins.

Reflux was shown to be the main contributor to the severity of PTS by means of multiple regression analysis, in accordance with other studies.² As clearly shown in Table II, only a minority of the patients with severe signs of PTS had an abnormal venous resistance, whereas most of these patients had reflux.

Other studies showed that in patients with severe signs of PTS the involved incompetent segments were the superficial femoral, popliteal, and the long and short saphenous veins.^{19,20} Although in our study the superficial femoral and popliteal veins showed a statistically significant increased prevalence of reflux with an increasing severeness of PTS, for the long and short saphenous veins this relationship showed a trend without being statistically significant.

This study corroborates the findings of others,²¹ because it indicates that isolated deep reflux in itself does not always cause severe signs of PTS. Perhaps the superficial veins can take over the drainage of venous blood as long as there are sufficient perforating veins. If the perforating veins become insufficient, the venous pressure in the superficial veins might increase and possibly cause an overload of the superficial system, resulting in a gradually developing insufficiency of the superficial veins. When the superficial veins also become insufficient, PTS develops. We did not evaluate the presence of insufficient perforating veins. Other studies²² have shown that duplex scanning misses a substantial number of incompetent perforating veins and therefore is unreliable in these evaluations. The patients in Table II with isolated deep vein reflux and severe signs of PTS might, therefore, have incompetent perforating veins, with an increased venous pressure in the superficial venous system, which is yet not insufficient.

In contrast to other studies,^{17,23,24} no relationship was found between reflux limited to the superficial veins and the PTS classification. This may be caused by the small number of patients with reflux limited only to the superficial vein segments.

The influence of the calf veins on the PTS is still controversial, with some studies reporting a relationship between reflux and PTS in the distal veins^{6,15} and others not reporting this.²⁵ In our study, not only the presence of reflux was analyzed, but also the traceability. Previously, we demonstrated that the

nontraceable veins are probably occluded.¹⁸ No relationship was found between reflux or traceability in the distal calf veins and an increased severity of PTS. Presumably because of the number of distal veins, reflux or an occlusion of one or more distal veins does not seem to contribute significantly to the presence of PTS. The other distal veins can probably drain away enough venous blood to prevent PTS.

Scott et al²⁶ revealed that many of the previously suggested associations with chronic venous insufficiency were in reality caused by the older age of the patients. In the present study, the severity of PTS also increased with age. Therefore, the statistical analysis was adjusted for age. Scott also found that patients with chronic venous insufficiency were men and were obese. The BMI of 21 of our patients was more than 27 kg/m², but these 21 patients were equally distributed among all classes of the CEAP scale. Perhaps our group of patients did not contain enough patients with severe obesity for us to uncover a relationship between obesity and PTS. In Scott's study, the mean BMI (30 kg/m²) was much higher than in our patient population.

A problem with our study is that it is retrospective. This might incorporate a selection of patients, causing a bias in the data, because the patients lost to follow-up could be different than those actually examined. Selection, although present, is minimized in our study because almost 80% of the patients that we invited took part; therefore, not only successfully or unsuccessfully treated patients have been included. Also, patients of different ages were included, with a variation of the location of the thrombi at initial phlebography.

In conclusion, patients with severe PTS most frequently have a combination of deep and superficial reflux. Reflux in the deep proximal veins contributes significantly to PTS. Because not all patients with deep and superficial reflux had severe signs of PTS, mechanisms other than reflux must be involved in the development of PTS.

We thank Adriënne van Maarseveen for her assistance in carrying out duplex examinations.

REFERENCES

1. Franzeck UK, Schalch I, Jager KA, Schneider E, Grimm J, Bollinger A. Prospective 12-year follow-up study of clinical and hemodynamic sequelae after deep vein thrombosis in low-risk patients (Zurich study). *Circulation* 1996;93:74-9.
2. Killewich LA, Martin R, Cramer M, Beach KW, Strandness DE. An objective assessment of the physiologic changes in the postthrombotic syndrome. *Arch Surg* 1985;120:424-6.
3. Janssen MCH, Wollersheim H, van Asten WNJC, de Rooij

- MJM, Novakova IRO, Thien T. The post-thrombotic syndrome: A review. *Phlebology* 1996;11:86-94.
4. Johnson BF, Manzo RA, Bergelin RO, Strandness DE. Relationship between changes in the deep venous system and the development of the postthrombotic syndrome after an acute episode of lower limb deep vein thrombosis: A one- to six-year follow-up. *J Vasc Surg* 1995;21:307-12.
 5. Shull KC, Nicolaides AN, Fernandes J, et al. Significance of popliteal reflux in relation to ambulatory venous pressure and ulceration. *Arch Surg* 1979;114:1304-6.
 6. Johnson BF, Manzo RA, Bergelin RO, Strandness DE. The site of residual abnormalities in the leg veins in long-term follow-up after deep vein thrombosis and their relationship to the development of the post-thrombotic syndrome. *Int Angiol* 1996;15:14-9.
 7. Payne SP, London NJ, Newland CJ, Bell PR, Barrie WW. Investigation and significance of short saphenous vein incompetence. *Ann R Coll Surg Engl* 1993;75:354-7.
 8. Porter JM, Moneta GL. Reporting standards in venous disease: An update. *J Vasc Surg* 1995;21:635-45.
 9. Klein Rouweler BJF, Brakkee AJM, Kuiper JP. Plethysmographic measurement of venous flow resistance and venous capacity in the human leg. Part one: Method. *Phlebology* 1989;4:241-50.
 10. Klein Rouweler BJF, Brakkee AJM, Kuiper JP. Plethysmographic measurement of venous flow resistance and venous capacity in human legs. Part two: Normal values related to age, sex and site of measurement. *Phlebology* 1989;4:251-7.
 11. Janssen MCH, Haenen JH, Asten WNJC, et al. Clinical and haemodynamic sequelae of deep venous thrombosis: Retrospective evaluation after 7-13 years. *Clin Sci* 1997;93:7-12.
 12. Haenen JH, van Langen H, Janssen MCH, et al. Venous duplex scanning of the leg; range, variability and reproducibility. *Clin Sci* 1999;96:271-7.
 13. Baker SR, Burnand KG, Sommerville KM, Thomas ML, Wilson NM, Browse NL. Comparison of venous reflux assessed by duplex scanning and descending phlebography in chronic venous disease. *Lancet* 1993;13:400-3.
 14. Neglen P, Raju S. A comparison between descending phlebography and duplex Doppler investigation in the evaluation of reflux in chronic venous insufficiency: A challenge to phlebography as the "gold standard." *J Vasc Surg* 1992;16:687-93.
 15. Labropoulos N, Leon M, Nicolaides AN, Giannoukas AD, Volteas N, Chan P. Superficial venous insufficiency: Correlation of anatomic extent of reflux with clinical symptoms and signs. *J Vasc Surg* 1994;20:953-8.
 16. van Rij AM, Solomon C, Christie R. Anatomic and physiologic characteristics of venous ulceration. *J Vasc Surg* 1994;20:759-64.
 17. Shami SK, Sarin S, Cheatle TR, Scurr JH, Smith PD. Venous ulcers and the superficial venous system. *J Vasc Surg* 1993;17:487-90.
 18. Haenen JH, Janssen MCH, van Langen H, et al. Duplex ultrasound in the hemodynamic evaluation of the late sequelae of deep venous thrombosis. *J Vasc Surg* 1998;27:472-8.
 19. van Bemmelen PS, Bedford G, Beach K, Strandness DE. Status of the valves in the superficial and deep venous system in chronic venous disease. *Surgery* 1991;109:730-4.
 20. Sakurai T, Matsushita M, Nishikimi N, Nimura Y. Hemodynamic assessment of femoropopliteal venous reflux in patients with primary varicose veins. *J Vasc Surg* 1997;26:260-4.
 21. Hanrahan LM, Araki CT, Rodriguez AA, Kechejian GJ, LaMorte WW, Menzoian JO. Distribution of valvular incompetence in patients with venous stasis ulceration. *J Vasc Surg* 1991;13:805-12.
 22. Pierik EGJM, Toonder IM, van Urk H, Wittens CHA. Validation of duplex ultrasonography in detecting competent and incompetent perforating veins in patients with venous ulceration of the lower leg. *J Vasc Surg* 1997;26:49-52.
 23. Bass A, Chayen D, Weinman EE, Ziss M. Lateral venous ulcer and short saphenous vein insufficiency. *J Vasc Surg* 1997;25:654-7.
 24. Labropoulos N, Leon M, Nicolaides AN, et al. Venous reflux in patients with previous deep venous thrombosis: Correlation with ulceration and other symptoms. *J Vasc Surg* 1994;20:20-6.
 25. Labropoulos N, Delis K, Nicolaides AN, Leon M, Ramaswami G. The role of the distribution and anatomic extent of reflux in the development of signs and symptoms in chronic venous insufficiency. *J Vasc Surg* 1996;23:504-10.
 26. Scott TE, LaMorte WW, Gorin DR, Menzoian JO. Risk factors for chronic venous insufficiency: A dual case-control study. *J Vasc Surg* 1995;22:622-8.

Submitted Sep 3, 1998; accepted Dec 15, 1998.