

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/41848233>

Synthesis of 2 β -substituted-14-epi-previtamin D₃ and testing of its genomic activity

ARTICLE *in* THE JOURNAL OF STEROID BIOCHEMISTRY AND MOLECULAR BIOLOGY · MARCH 2010

Impact Factor: 3.63 · DOI: 10.1016/j.jsbmb.2010.02.035 · Source: PubMed

CITATIONS

6

READS

21

8 AUTHORS, INCLUDING:



[Ken-ichiro Takagi](#)

Teijin Pharma Limited

13 PUBLICATIONS 67 CITATIONS

SEE PROFILE



[Atsushi Kittaka](#)

Teikyo University

164 PUBLICATIONS 1,944 CITATIONS

SEE PROFILE



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at ScienceDirect

Journal of Steroid Biochemistry and Molecular Biology

journal homepage: www.elsevier.com/locate/jsbmb

Synthesis of 2 β -substituted-14-epi-previtamin D₃ and testing of its genomic activity[☆]

Daisuke Sawada^a, Yuya Tsukuda^a, Hiroshi Saito^b, Ken-ichiro Takagi^b, Eiji Ochiai^b, Seiichi Ishizuka^b, Kazuya Takenouchi^b, Atsushi Kittaka^{a,*}

^a Faculty of Pharmaceutical Sciences, Teikyo University, 1091-1, Suwarashi, Sagamiko, Sagami-hara, Kanagawa 229-0195, Japan

^b Teijin Institute for Bio-medical Research, Teijin Pharma Ltd., Tokyo 191-8512, Japan

ARTICLE INFO

Article history:

Received 19 October 2009

Received in revised form 10 February 2010

Accepted 25 February 2010

Keywords:

Vitamin D₃

14-epi-Previtamin D₃

Osteocalcin

Vitamin D receptor

ABSTRACT

2 β -Substituted analogs of 14-epi-previtamin D₃ were synthesized for the first time by the thermal isomerization of the corresponding 14-epi-vitamin D₃ that were available using coupling reaction between the A-ring phosphine oxide derived from a chiral epoxide and CD-ring *cis*-hydrindanone. The VDR binding affinity and transactivation activity of osteocalcin promoter in HOS cells were evaluated, and the new analogs were found to be less active, 0.01–0.18% of VDR binding affinity compared with the natural hormone and EC₅₀ 1.0–9.1 nM for transactivation activity, than 14-epi-previtamin D₃ with 0.5% (VDR) and EC₅₀ 0.46 nM, respectively.

© 2010 Elsevier Ltd. All rights reserved.

1. Introduction

It is well established that vitamin D₃ is present in thermal equilibrium with previtamin D₃ via [1,7]-sigmatropic rearrangement. In this equilibrium, the vitamin D form (**A**) with the 6-*s-trans* triene structure is more stable and dominant than the 6-*cis* isomer of the previtamin D form (**B**) (Scheme 1). The biologically most active metabolite of vitamin D₃, 1 α ,25(OH)₂D₃ (**1**), also contains 5–10% of its previtamin D form, 1 α ,25(OH)₂preD₃ (**pre-1**) at 37 °C in similar equilibrium. The major isomer, the vitamin D form (**A**), has been the focus of therapeutic evaluation rather than the previtamin D form, because previtamin D is easily transformed to vitamin D through thermal equilibrium and is almost impossible to isolate in pure form [1]. While **1** is a ligand of the nuclear vitamin D receptor (VDR), regulates gene transcription, and exhibits various biological responses as a hormone [2], **pre-1** is thought to be a weak ligand of VDR and a poor activator of the above genomic actions [3]; however, **pre-1** has been studied as a responsible compound for rapid responses [4], such as stimulation of intestinal Ca²⁺ transport (transcaltachia), activation of PKC and MAP kinases, and so on, which are called non-genomic actions [5].

Okamura and coworkers reported that the thermal equilibrium ratio between the vitamin D form (**A**) and previtamin D form (**B**) at 80 °C was reversed by epimerizing the CD-ring bridgehead hydrogen of C14 [6]. Briefly, 14-epi-1 α ,25(OH)₂preD₃ (**14-epi-pre-1**) was major and dominant to 14-epi-1 α ,25(OH)₂D₃ (**14-epi-1**), and the former was isolated as a stable single isomer at room temperature. Therefore, we focused on the synthesis of **14-epi-pre-1** analogs with A-ring modification to investigate their more detailed biological properties and potential as therapeutic agents of the previtamin D₃ skeleton.


Previously, we found that 2 α -alkyl and 2 α -(ω -hydroxyalkyl) substitution afforded great enhancements for VDR binding affinity and the subsequent genomic actions [7]. In the preceding paper, we reported the synthesis and biological evaluations of 2 α -substituted **14-epi-pre-1** [8]. Here, we prepared analogs with 2 β -substitution (**14-epi-pre-1a-c**), because 2 β -substitution is known as an important modification for vitamin D derivatization (Scheme 2) [9].

14-epi-pre-1 could be prepared from **14-epi-1** by thermal isomerization; therefore, we synthesized **14-epi-1** analogs as temporary first targets. The **14-epi-1** analogs were divided into two fragments, CD-ring and A-ring, which were coupled by the Roche coupling method [10]. The CD-ring fragment **2** [6,8] is the known compound, which was obtained by epimerization at H14 in Grundmann's ketone derivative derived from vitamin D₃ [11,12]. The A-ring fragments, the phosphine oxides **3a-c**, could be synthesized from dimethyl D-tartrate, and we could introduce various alkyl groups at the 2 β -position by nucleophilic epoxide ring-opening reactions [13].

[☆] Special issue selected article from the 14th Vitamin D Workshop held at Brugge, Belgium on October 4–8, 2009.

* Corresponding author. Tel.: +81 42 685 3713; fax: +81 42 685 3713.

E-mail address: akittaka@pharm.teikyo-u.ac.jp (A. Kittaka).




Scheme 1. Equilibrium between vitamin D₃ and previtamin D₃.


2. Results and discussion

2β-Substituted A-ring fragments (**3a–c**) were prepared from the known epoxide **4** derived from dimethyl D-tartrate (**Scheme 3**) [13,14]. Using the nucleophilic epoxide ring-opening reaction of **4**, three substitutions were introduced as follows: (1) methyl cuprate gave a methyl substitution, (2) an allyl group brought


by Grignard reagent was treated with 9-borabicyclo[3,3,1]nonane (9-BBN), and then with H₂O₂ to afford a hydroxypropyl substitution, and (3) propylene glycol gave a hydroxypropoxy substitution. After their primary hydroxyls were protected as TBDPS ether, **5a–c** were converted into bromide **6a–c** by the known procedure [15]. Methanolysis of both acetyl groups under basic conditions led to epoxide formation, and the resultant hydroxyl group was



Scheme 2. Retrosynthetic analysis of 2β-substituted 14-epi-1α,25(OH)₂preD₃.



Scheme 3. Synthesis of the A-ring fragments. Conditions: (a) for **5a** MeLi, CuI, Et₂O, 98%; (b) for **5b** (i) allylmagnesium chloride, toluene, (ii) 9-BBN, THF, H₂O₂, NaOH, (iii) TBDPSCI, imidazole, DMF, 90% (3 steps); (c) for **5c** (i) propylene glycol, KOtBu, (ii) TBDPSCI, imidazole, DMF, 92% (2 steps); (d) Pd/C, H₂, MeOH; (e) MeC(OMe)₃, PPTS, CH₂Cl₂; (f) AcBr, CH₂Cl₂, 60% for **6a**, 52% for **6b**, 55% for **6c** (3 steps); (g) K₂CO₃, MeOH; (h) BzCl, Et₃N, CH₂Cl₂, 80% for **7a**, 95% for **7b**, 83% for **7c** (2 steps); (i) (trimethylsilyl)acetylene, nBuLi, BF₃·OEt₂, THF; (j) TBSOTf, iPr₂EtN, CH₂Cl₂; (k) K₂CO₃, MeOH, 68% for **8a**, 83% for **8b**, 95% for **8c** (3 steps); (l) SO₃·Py, Et₃N, DMSO, 77% for **9a**, 99% for **9b**, 95% for **9c**; (m) vinylmagnesium chloride, THF, 93% (α/β 46/47) for **10a**, 95% (α/β 37/58) for **10b**, 73% (α/β 15/58) for **10c** (2 steps); (n) TBSOTf, iPr₂EtN, CH₂Cl₂, 100% for **11aα**, 97% for **11aβ**, 99% for **11bβ**, 99% for **11cβ**; (o) nBuLi, (CH₂O)_n, THF; (p) Red-Al, Et₂O, then I₂, THF; (q) Pd(PPh)₄, Et₃N, MeCN; (r) (i) NCS, Me₂S, CH₂Cl₂, (ii) nBuLi, PPh₃, THF, then 30% H₂O₂, 49% for **3aα**, 57% for **3aβ**, 27% for **3bβ**, 28% for **3cβ** (4 steps).




Scheme 4. Determination of the stereochemistry of the 1-hydroxy group of **10a–c**. Conditions: (a) for **10aβ**, TBAF, THF, 100%; (b) for **10bβ**, PPTS, EtOH, 60%; (c) dimethoxypropane, PPTS, DMF, 70% for **13aβ**, 90% for **13bβ**.

transformed into benzyl ester **7a–c**. The addition of (trimethylsilyl)acetylene to the epoxide using *n*BuLi was straightforward, and the generated secondary alcohol was protected as TBS ether, and removal of the terminal TMS group and the benzoyl group gave alkyne **8a–c**. The primary alcohol was oxidized to aldehyde by DMSO and SO₃•pyridine complex (**9a–c**), to which the vinyl group was introduced to give a diastereomixture of alcohol **10a–c**. The stereochemistry of the new hydroxy groups is discussed below (Scheme 4), and both isomers of **10a** and the major isomer of **10b** and **10c** were used for further transformation after column chromatography. The hydroxy group of **10a–c** was protected by the TBS group to obtain 2β-substituted enyne **11a–c**. According to the known procedure, enyne **11a–c** were transformed into phosphine oxide in four steps to give **3a–c**, respectively [8,13]. As above, we were able to prepare four A-ring fragments.

As shown in Scheme 4, the minor diastereomer of **10c** (**10cα**) was converted to the phosphine oxide **3cα** by the same strategy as in Scheme 3, and it was identical to the known compound reported by Hatakeyama et al. [13]. Therefore, the stereochemistry of its 1-hydroxy group (steroidal numbering) was found to be α-configuration, and the major diastereomer of **10c** was determined to have the 1β-hydroxy group (**10cβ**). For determination of the stereochemistry in **10a** and **10b**, the TBS groups of the major diastereomers (**10aβ** and **10bβ**) were removed, and the resultant 1,3-dihydroxy groups of **12aβ** and **12bβ** were converted into acetone **13aβ** and **13bβ**, respectively. NOE analysis is described in Scheme 4, and the stereochemistry of 1,3-dihydroxy groups was determined as *syn*, that is, **10aβ** and **10bβ** had 1β,3β-dihydroxy groups. As above, we found that all of the major diastereomer of **10a–c** had 1β-hydroxy groups.

Using the CD- and A-ring fragments prepared as above, we examined the coupling reaction under basic conditions with *n*BuLi (Scheme 5) [6,10]. Small excess amounts of the A-ring fragment worked well and we obtained the coupled products **14a–c** in moderate yields. At this point, isomerization to the previtamin D form was seldom observed, probably because TBS groups at the A-ring should have steric hindrance to prevent from reaching the transition state for the [1,7]-sigmatropic hydrogen shift between the vitamin D form and the previtamin D form. Then, all silyl groups in **14a–c** were removed in one step with excess TBAF, and most of the deprotected compounds remained in the vitamin D form (**14-epi-1a–c**), and small amounts of the previtamin D form (**14-epi-pre-1a–c**) were produced under these reaction conditions. However, once they were heated at 80 °C in benzene, isomerization was found to proceed smoothly by ¹H NMR observation. After 2 h, a large proportion of the vitamin D form had been converted into the previtamin D form, and the isomerization seemed to reach thermal equilibrium, at which the ratio of the compounds was about 5/95 (vitamin D/previtamin D) based on ¹H NMR studies. Using HPLC, the mixture of both forms was separated, and we were able to obtain **14-epi-pre-1a–c** as pure forms, which were used for further biological studies.

The VDR binding affinity and the osteocalcin promoter trans-activation activity of the new compounds were evaluated using chick intestinal VDR and HOS cells, respectively. The results are summarized in Table 1 in comparison with the natural hormone **1** and 14-epi-1α,25(OH)₂preD₃ (**14-epi-pre-1**), which was synthesized in a similar manner in our laboratory. The new compounds showed lower activity than the natural hormone **1**, and also than



Scheme 5. Coupling reaction and synthesis of 2β-substituted 14-epi-1α,25(OH)₂preD₃.

Table 1

Relative binding affinity for chick intestinal VDR and osteocalcin promoter transactivation activity in HOS cells of 2β-substituted 14-epi-1α,25(OH)₂preD₃.

Compound	VDR ^a	Osteocalcin transactivation activity (EC ₅₀ , nM)
1	100	0.03
14-epi-pre-1	0.5	0.46
14-epi-pre-1aα	0.08	1.34
14-epi-pre-1aβ	0.08	9.12
14-epi-pre-1bβ	0.18	1.01
14-epi-pre-1cβ	0.01	1.24

^a The potency of **1** is normalized to 100.

14-epi-pre-1 regardless of the stereochemistry at the 1-hydroxy group.

3. Conclusion

We synthesized 2β-substituted analogs of **14-epi-1** for the first time and were able to isolate these new analogs (**14-epi-pre-1a-c**) after thermal isomerization at 80 °C. We evaluated their VDR binding affinity and transactivation activity of osteocalcin promoter in HOS cells. It was found that 2β-modified analogs of 14-epi-1α,25-dihydroxyvitamin D₃ were considerably less active than the natural hormone (**1**) and than **14-epi-pre-1**, although 2β-modification of **1** afforded important knowledge to the vitamin D SAR studies.

Acknowledgments

We are grateful to Ms. Junko Shimode and Ms. Ayako Kawaji (Teikyo University) for the spectroscopic measurements. This work

was supported in part by a Grant-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology, Japan (to D.S.) and by a Grant-in-Aid from the Japan Society for the Promotion of Science (to A.K.).

References

- [1] M.L. Curtin, W.H. Okamura, 1α,25-Dihydroxyvitamin D₃: synthesis of the 9,14,19,19,19-pentadeuterio derivatives and a kinetic study of its [1,7]-sigmatropic shift to 1α,25-dihydroxyvitamin D₃, *J. Am. Chem. Soc.* 113 (1991) 6958–6966.
- [2] D. Feldman, J.W. Pike, F.H. Glorieux (Eds.), *Vitamin D*, 2nd ed., Elsevier Academic Press, New York, 2005.
- [3] (a) H.F. DeLuca, Evolution of our understanding of vitamin D, *Nutr. Rev.* 66 (Suppl. 2) (2008) S73–S87;
(b) A.J. Brown, E. Slatopolsky, *Vitamin D analogs: therapeutic applications and mechanisms for selectivity*, *Mol. Aspects Med.* 29 (2008) 433–452.
- [4] (a) M.T. Mizwicki, D. Keidel, C.M. Bula, J.E. Bishop, L.P. Zanello, J.-M. Wurtz, D. Moras, A.W. Norman, Identification of an alternative ligand-binding pocket in the nuclear vitamin D receptor and its functional importance in 1α,25(OH)₂-vitamin D₃ signaling, *PNAS* 101 (2004) 12876–12881;
(b) I. Nemere, M.C. Dormanen, M.W. Hammond, W.H. Okamura, A.W. Norman, Identification of a specific binding protein for 1α,25-dihydroxyvitamin D₃ in basal-lateral membranes of chick intestinal epithelium and relationship to transcaltachia, *J. Biol. Chem.* 269 (1994) 23750–23756.
- [5] A.W. Norman, M.T. Mizwicki, D.P.G. Norman, Steroid-hormone rapid actions, membrane receptors and a conformational ensemble model, *Nat. Rev. Drug Discov.* 3 (2004) 27–41.
- [6] D.F. Maynard, W.G. Trankle, A.W. Norman, W.H. Okamura, 14-Epi stereoisomers of 25-hydroxy- and 1α,25-dihydroxyvitamin D₃: synthesis, isomerization to previtamins, and biological studies, *J. Med. Chem.* 37 (1994) 2387–2393.
- [7] For our review articles, see:
(a) N. Saito, S. Honzawa, A. Kittaka, Recent results on A-ring modification of 1α,25-dihydroxyvitamin D₃: design and synthesis of VDR-agonists and antagonists with high biological activity, *Curr. Top. Med. Chem.* 6 (2006) 1273–1288;
(b) N. Saito, A. Kittaka, Highly potent vitamin D receptor antagonists: design, synthesis, and biological evaluation, *Chembiochem* 7 (2006) 1478–1490;
(c) N. Saito, A. Kittaka, Design and synthesis of highly potent vita-

- min D receptor antagonists based on the structural development of vitamin D₃-26,23-lactone, *J. Synth. Org. Chem. Jpn.* 65 (2007) 947–958.
- [8] D. Sawada, T. Katayama, Y. Tsukuda, N. Saito, M. Takano, H. Saito, K. Takagi, E. Ochiai, S. Ishizuka, K. Takenouchi, A. Kittaka, Synthesis of 2 α -substituted-14-*epi*-previtamin D₃ and its genomic activity, *Bioorg. Med. Chem. Lett.* 19 (2009) 5397–5400.
- [9] (a) N. Tsugawa, K. Nakagawa, M. Kurobe, Y. Ono, N. Kubodera, K. Ozono, T. Okano, *In vitro* biological activities of a series of 2 β -substituted analogues of 1 α ,25-dihydroxyvitamin D₃, *Biol. Pharm. Bull.* 23 (2000) 66–71;
(b) Y. Ono, H. Watanabe, A. Shiraishi, S. Takeda, Y. Higuchi, K. Sato, N. Tsugawa, T. Okano, T. Kobayashi, N. Kubodera, Synthetic studies of vitamin D analogs. XXIV. Synthesis of active vitamin D₃ analogs substituted at 2 β -position and their preventive effects on bone mineral loss in ovariectomized rats, *Chem. Pharm. Bull.* 45 (1997) 1626–1630;
(c) K. Miyamoto, E. Murayama, K. Ochi, H. Watanabe, N. Kubodera, Synthetic studies of vitamin D analogs. XIV. Synthesis and calcium regulation activity of vitamin D₃ analogues bearing a hydroxyalkoxy group at the 2 β -position, *Chem. Pharm. Bull.* 41 (1993) 1111–1113.
- [10] B. Lythgoe, Simonsen lecture. Synthetic approaches to vitamin D and its relatives, *Chem. Soc. Rev.* 9 (1980) 449–475.
- [11] (a) H.H. Inhoffen, G. Quinkert, S. Siegmund, D. Kampe, G.F. Domagk, Vitamin D series. XXI. Hydrindan compounds derived from vitamin D₃, *Chem. Ber.* 90 (1957) 664–673;
(b) J. Kiegiel, P.M. Wovkulich, M.R. Uskokovic, Chemical conversion of vitamin D₃ to its 1,25-dihydroxy metabolite, *Tetrahedron Lett.* 32 (1991) 6057–6060.
- [12] R.R. Sicinski, K.L. Perlman, H.F. DeLuca, Synthesis and biological activity of 2-hydroxy and 2-alkoxy analogs of 1 α ,25-dihydroxy-19-norvitamin D₃, *J. Med. Chem.* 37 (1994) 3730–3738.
- [13] (a) J. Maeyama, H. Hiyamizu, K. Takahashi, J. Ishihara, S. Hatakeyama, N. Kubodera, Two convergent approaches to the synthesis of 1 α ,25-dihydroxy-2 β -(3-hydroxypropoxy)vitamin D₃ (ED-71) by the Lythgoe and the Trost coupling reactions, *Heterocycles* 70 (2006) 295–307;
(b) S. Hatakeyama, H. Irie, T. Shintani, Y. Noguchi, H. Yamada, M. Nishizawa, An efficient route to a key A-ring synthon for 1 α ,25-dihydroxyvitamin D₃ and its analogs, *Tetrahedron* 50 (1994) 13369–13376.
- [14] K.C. Nicolaou, D.P. Papahatjis, D.A. Claremon, R.L. Magolda, R.E. Dolle, Total synthesis of ionophore antibiotic X-14547A, *J. Org. Chem.* 50 (1985) 1440–1456.
- [15] T. Hanazawa, M. Koiwa, G.P.-J. Hareau, F. Sato, Optically active *trans*-4-(*tert*-butyldimethylsiloxy)methyl)-5-(*tert*-butyldimethylsiloxy)-2-cyclohexenone as a useful chiral building block for preparation of substituted cyclohexane rings: synthesis and its highly stereoselective reaction with RCu(CN)Li, *Tetrahedron Lett.* 41 (2000) 2659–2662.