

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/13801066>

# Quenching of 7-chloro-4-nitrobenzo-2-oxa-1,3-diazole-modified Na<sup>+</sup>/K<sup>+</sup>-ATPase reveals a higher accessibility of the low-affinity ATP-binding site

ARTICLE *in* FEBS LETTERS · JANUARY 1998

Impact Factor: 3.17 · DOI: 10.1016/S0014-5793(97)01460-9 · Source: PubMed

---

CITATIONS

10

---

READS

15

3 AUTHORS, INCLUDING:



Evžen Amler

Charles University in Prague

129 PUBLICATIONS 1,179 CITATIONS

SEE PROFILE

# Quenching of 7-chloro-4-nitrobenzo-2-oxa-1,3-diazole-modified $\text{Na}^+/\text{K}^+$ -ATPase reveals a higher accessibility of the low-affinity ATP-binding site

Holger Linnertz<sup>1,a,b</sup>, Petra Urbanova<sup>a</sup>, Evzen Amler<sup>a,\*</sup>

<sup>a</sup>Institute of Physiology, Academy of Sciences of the Czech Republic, Videnska 1083, Cz-142 20 Prague 4, Czech Republic

<sup>b</sup>Institute of Biochemistry and Endocrinology, Justus-Liebig-University Giessen, Frankfurter Str. 100, D-35392 Giessen, Germany

Received 30 September 1997; revised version received 31 October 1997

**Abstract** 7-Chloro-4-nitrobenzo-2-oxa-1,3-diazole (NBD-Cl) labeled  $\text{Na}^+/\text{K}^+$ -ATPase covalently with two different inactivation constants ( $K_i = 2.5 \mu\text{M}$ ;  $K_i' = 10 \mu\text{M}$ ). It apparently modified the two different ATP-binding sites of the enzyme since it decreased the activity of the  $\text{E}_2\text{ATP}$  site, i.e. the  $\text{K}^+$ -activated para-nitrophenylphosphatase activity, in an enzyme whose high-affinity  $\text{E}_1\text{ATP}$  site had been blocked by fluorescein 5'-isothiocyanate (FITC). It also reduced the activity of the  $\text{E}_1\text{ATP}$  site, i.e. the  $\text{Na}^+$ -activated protein phosphorylation, in an enzyme whose low-affinity  $\text{E}_2\text{ATP}$  site had been blocked by  $\text{Co}(\text{NH}_3)_4\text{PO}_4$ . Fluorescence quenching experiments with KI, CsCl and  $\text{MnCl}_2$  of the NBD-Cl-labeled  $\text{Na}^+/\text{K}^+$ -ATPase revealed two differently accessible types of fluorophores depending on the ATP site: The  $\text{E}_2\text{ATP}$  site apparently differs from the  $\text{E}_1\text{ATP}$  site in that it is more open because the fluorophore labeling in the  $\text{E}_2\text{ATP}$  site was sterically better accessible for quenchers.

© 1997 Federation of European Biochemical Societies.

**Key words:**  $\text{Na}^+/\text{K}^+$ -ATPase; NBD-Cl; ATP-binding site; ATP analog; Fluorescence anisotropy; Fluorescence quenching

## 1. Introduction

$\text{Na}^+/\text{K}^+$ -ATPase (EC 3.6.1.37) is an integral membrane protein which transports sodium and potassium ions against an electrochemical potential gradient. The process of cation transport has been described by the so-called Albers-Post model in which a single ATP site per catalytic  $\alpha$  subunit exists in two main conformations which differ in their affinities for ATP: The high-affinity  $\text{E}_1\text{ATP}$  site is associated with  $\text{Na}^+$  export while the low-affinity  $\text{E}_2\text{ATP}$  site is implicated in  $\text{K}^+$  import [1]. This model, however, is inconsistent with the observations of backdoor phosphorylation and of [ $^{86}\text{Rb}$ ] occlusion in an  $\text{E}_1\text{ATP}$  site blocked enzyme [2,3] and with the

phenomena of superphosphorylation from [ $\gamma\text{-}^{32}\text{P}$ ]ATP [4], double labeling with ATP analogs [5] or double phosphorylation from para-nitrophenylphosphate [6]. All these observations, including the observation of a positive cooperative effect of 2'-O-DANS-8- $\text{N}_3$ -ATP in the inactivation of  $\text{Na}^+/\text{K}^+$ -ATPase [7] led to the conclusion that two simultaneously existing ATP sites need to cooperate during ATP-driven  $\text{Na}^+/\text{K}^+$  transport [7,8]. Unfortunately, the available data cannot differentiate between the possibility that two ATP-binding sites exist on a single  $\alpha$  subunit and the possibility that the sodium pump works as a functional dimer ( $\alpha\beta$ )<sub>2</sub> with two cooperating ATP sites. To answer this question, covalent and preferably fluorescent labeling of the  $\text{E}_2\text{ATP}$ -binding site is needed.

To reach this goal, 7-chloro-4-nitrobenzo-1,3-diazole (NBD-Cl) was used to label the ATP sites in  $\text{Na}^+/\text{K}^+$ -ATPase. Cantley et al. hypothesized from their data that two different tyrosine residues within the ATP sites may be modified [9], but Repke's group concluded that ATP may protect the enzyme against modification of sulphhydryl groups [10].

It was of interest to learn whether modification of these two types of amino acid residues could underlie the inactivation of the partial activities of the enzyme, i.e.  $\text{Na}^+$ -dependent phosphorylation at the  $\text{E}_1\text{ATP}$  site [11] and  $\text{K}^+$ -activated phosphatase activity at the  $\text{E}_2\text{ATP}$  site [12,13]. Therefore, we performed several experiments to study the effect of NBD-Cl on the enzyme with specifically blocked  $\text{E}_1\text{ATP}$ - (FITC-pretreated) or  $\text{E}_2\text{ATP}$ -binding site ( $\text{Co}(\text{NH}_3)_4\text{PO}_4$ -pretreated). In addition, NBD-Cl is a suitable label to obtain key information on the accessibility or 'depth' of the two ATP-binding sites and, hence, fluorescence quenching by iodide, cesium and manganese(II) ions of NBD-Cl-labeled  $\text{Na}^+/\text{K}^+$ -ATPase was studied in an enzyme preparation where the  $\text{E}_1\text{ATP}$  site had been protected with FITC or  $\text{Cr}(\text{H}_2\text{O})_4\text{AdoPP}[\text{CH}_2]_P$  and whose  $\text{E}_2\text{ATP}$  site had been protected with  $\text{Co}(\text{NH}_3)_4\text{ATP}$  or  $\text{Co}(\text{NH}_3)_4\text{PO}_4$ . The differences in the accessibility of the NBD-Cl label to all these quenchers is consistent with the assumption that the  $\text{E}_2\text{ATP}$  site is better accessible to quenchers than the  $\text{E}_1\text{ATP}$  site.

## 2. Materials and methods

All chemicals were of the highest available purity and were obtained from Bio-Rad (Munich, Germany), Boehringer-Mannheim (Mannheim, Germany), E. Merck (Darmstadt, Germany) and Molecular Probes (Eugene, OR, USA). Lab-Trol protein standard is a product of Merz and Dade (Munich, Germany). [ $\gamma\text{-}^{32}\text{P}$ ]ATP was from Amersham Buchler (Braunschweig, Germany). Calculation and presentation of data were done with the program GraphPad Prism (GraphPad Software Inc., San Diego, CA, USA).

\*Corresponding author. Fax: +42 (2) 475/2249.  
E-mail: AMLER@BIOMED.CAS.CZ

<sup>1</sup>This work is part of the Ph.D. thesis of H.L.

**Abbreviations:**  $\text{Cr}(\text{H}_2\text{O})_4\text{AdoPP}[\text{CH}_2]_P$ ,  $\beta,\gamma$  bidentate complex of chromium(III)-tetraaqua-adenylyl [ $\beta,\gamma$ -methylene] diphosphonate;  $\text{Co}(\text{NH}_3)_4\text{ATP}$ ,  $\beta,\gamma$  bidentate complex of cobalt(III)-tetramino-adenosine-5'-triphosphate;  $\text{Co}(\text{NH}_3)_4\text{PO}_4$ , tetramine cobalt(III)phosphate; FITC, fluorescein 5'-isothiocyanate; NBD-Cl, 7-chloro-4-nitrobenzo-2-oxa-1,3-diazole, 7-chloro-4-nitro-benzofurazane;  $\text{E}_1\text{ATP}$  site, nucleotide-binding site of  $\text{Na}^+/\text{K}^+$ -ATPase with high affinity for ATP;  $\text{E}_2\text{ATP}$  site, nucleotide-binding site of  $\text{Na}^+/\text{K}^+$ -ATPase with low affinity for ATP

### 2.1. Enzyme and assays

$\text{Na}^+/\text{K}^+$ -ATPase in the range of 20–25 units/mg protein was isolated by modification of Jørgensen's procedure from pig kidney [14] and measured by a coupled spectrometric assay [15]. One enzyme unit (U) is defined as the hydrolysis of 1  $\mu\text{mole}$  ATP per min at  $37^\circ\text{C}$ . Protein was determined by Lowry's method [16] using Lab-Trol as a protein standard. Lab-Trol is a mixture of proteins and enzymes used for the calibration of assays in clinical chemical analysis.

### 2.2. Inactivation of $\text{Na}^+/\text{K}^+$ -ATPase by NBD-Cl

$\text{Na}^+/\text{K}^+$ -ATPase was inactivated with 7-chloro-4-nitrobenzo-2-oxa-1,3-diazole (NBD-Cl) as follows: 1 U of purified  $\text{Na}^+/\text{K}^+$ -ATPase was incubated in a total volume of 1 ml with different concentrations of NBD-Cl in 20 mM Tris/HCl (pH 7.25) and 15 mM NaCl at  $37^\circ\text{C}$ . The time course of inactivation of  $\text{Na}^+/\text{K}^+$ -ATPase was followed by transferring aliquots of 50  $\mu\text{l}$  to the coupled optical test [15]. The initial rates of inactivation were calculated and analyzed according to the method of Piszkiwics et al. [7,17].

### 2.3. Inactivation of $\text{Na}^+/\text{K}^+$ -ATPase by FITC

$\text{Na}^+/\text{K}^+$ -ATPase in a final concentration of 1 U/ml (65 mg/ml) was incubated overnight at  $37^\circ\text{C}$  in a solution containing 20 mM Tris/HCl (pH 7.25) and 10  $\mu\text{M}$  FITC. The control enzyme was treated in the same way but without FITC. The inactivated enzyme (residual activity  $\approx 1\%$ ) was spun down, washed in 20 mM Tris/HCl (pH 7.25) and resuspended in a solution of 20 mM Tris/HCl (pH 7.25), 15 mM NaCl and various concentrations (0–500  $\mu\text{M}$ ) NBD-Cl. After incubation for 1 h at  $37^\circ\text{C}$  the enzyme was spun down, washed twice with 20 mM Tris/HCl and  $\text{K}^+$ -activated para-nitrophenylphosphatase was estimated. For details see [2].

### 2.4. Frontdoor phosphorylation of $\text{Na}^+/\text{K}^+$ -ATPase

$\text{Na}^+/\text{K}^+$ -ATPase in a final concentration of 3 U/ml (195  $\mu\text{g}/\text{ml}$ ) was incubated overnight at  $37^\circ\text{C}$  in a solution containing 20 mM Tris/HCl (pH 7.25) and 1 mM  $\text{Co}(\text{NH}_3)_4\text{PO}_4$  (a control without  $\text{Co}(\text{NH}_3)_4\text{PO}_4$  was run in parallel). The inactivated enzyme was centrifuged at  $100\,000\times g$ , washed in 20 mM Tris/HCl (pH 7.25) and incubated in 20 mM Tris/HCl, 15 mM NaCl with 100  $\mu\text{M}$  or 1 mM NBD-Cl. After different incubation times at  $37^\circ\text{C}$  the enzyme was washed twice with 20 mM Tris/HCl (pH 7.25) and frontdoor phosphorylation was estimated according to Thoenges and Schoner [7]. Background labeling was subtracted using the value of a sample where 1 U of  $\text{Na}^+/\text{K}^+$ -ATPase had been quenched with 250  $\mu\text{l}$  of 25% trichloroacetic acid before phosphorylation from  $[\gamma\text{-}^{32}\text{P}]\text{ATP}$  started. Controls were run in parallel with the native enzyme (no  $\text{Co}(\text{NH}_3)_4\text{PO}_4$  pretreatment) to detect the influence of 1 mM NBD-Cl.

### 2.5. Fluorescence measurements

$\text{Na}^+/\text{K}^+$ -ATPase in a final concentration of 2.5 U/ml (163 mg/ml) was incubated overnight at  $37^\circ\text{C}$  in a solution containing 20 mM Tris/HCl (pH 7.5), 15 mM NaCl and 1 mM  $\text{Co}(\text{NH}_3)_4\text{ATP}$  or 25  $\mu\text{M}$   $\text{Cr}(\text{H}_2\text{O})_4\text{AdoPP}[\text{CH}_2]\text{P}$  (residual activity  $< 5\%$ ). The enzyme was centrifuged at  $100\,000\times g$  and washed with 20 mM Tris/HCl (pH 7.5) and resuspended in 20 mM Tris/HCl (pH 7.5), 15 mM NaCl and 1 mM NBD-Cl (the native samples contained no  $\text{Co}(\text{NH}_3)_4\text{ATP}$  or  $\text{Cr}(\text{H}_2\text{O})_4\text{AdoPP}[\text{CH}_2]\text{P}$ ). After a 1 h incubation at  $37^\circ\text{C}$  the enzyme was centrifuged and washed again. The fluorescence was read at  $\lambda_{\text{ex}}$  475 nm and  $\lambda_{\text{em}}$  520 nm and fluorescence intensity quenching was achieved by adding increasing concentrations (0–800 mM) of the quenchers (KI, CsCl and  $\text{MnCl}_2$ , respectively). A Specol 211 spectro-




Fig. 1. Inactivation of the  $\text{Na}^+/\text{K}^+$ -ATPase activity by NBD-Cl.  $\text{Na}^+/\text{K}^+$ -ATPase was inactivated with NBD-Cl as follows: 1 U of  $\text{Na}^+/\text{K}^+$ -ATPase was incubated in a total volume of 1 ml with (1, 5, 10, 25, 50, 100 and 500 mM) NBD-Cl in 20 mM Tris/HCl (pH 7.25) and 15 mM NaCl at  $37^\circ\text{C}$ . The residual  $\text{Na}^+/\text{K}^+$ -ATPase activity was assayed upon transferring 50  $\mu\text{l}$  aliquots spectrophotometrically [15]. Closed symbols show the residual rate constants of inactivation with 1–25 mM NBD-Cl and the insert with the open symbols the constants of inactivation with 25–500 mM NBD-Cl. Mean values of two experiments are shown.

photometer was used for absorbance measurements. Steady-state fluorescence data were collected in quartz cuvettes in a Perkin-Elmer LS-5 fluorometer equipped with monochromators and analyzed as described in [18]. Two Glan-Thompson polarizers were used for determination of steady-state anisotropy values. All measurements were performed at  $37^\circ\text{C}$ .

To detect the protective effect of  $\text{Co}(\text{NH}_3)_4\text{ATP}$  against NBD-Cl labeling, FITC-pretreated  $\text{Na}^+/\text{K}^+$ -ATPase was incubated at a final concentration of 2 U/ml for 2 h at  $37^\circ\text{C}$  in a solution containing 20 mM Tris/HCl (pH 7.5) and 1 mM  $\text{Co}(\text{NH}_3)_4\text{ATP}$  (control without  $\text{Co}(\text{NH}_3)_4\text{ATP}$ ). After washing the enzyme was incubated in 20 mM Tris/HCl buffer (pH 7.5) for 1 h at  $37^\circ\text{C}$  with increasing concentrations of NBD-Cl (0–500  $\mu\text{M}$ ) and washed again. The fluorescence of these probes was read at  $\lambda_{\text{ex}}$  475 nm and  $\lambda_{\text{em}}$  520 nm. A control without NBD-Cl was run in parallel as background to exclude non-specific effects.

## 3. Results

Inactivation rate constants of  $\text{Na}^+/\text{K}^+$ -ATPase at 0–500  $\mu\text{M}$  NBD-Cl revealed two different  $K_i$  values, viz. 2.5  $\mu\text{M}$  (Fig. 1) and 10  $\mu\text{M}$  (Fig. 1, insert) and implied a two-site modification process. To learn whether NBD-Cl may affect the partial activity of the  $\text{E}_2$ ATP site we studied the inactivation of  $\text{K}^+$ -activated para-nitrophenylphosphatase by this reagent (Table 1) in a FITC-prelabeled  $\text{Na}^+/\text{K}^+$ -ATPase where the  $\text{E}_1$ ATP site was thus blocked [2]. All inactivation effects of

Table 1

Inactivation of  $\text{K}^+$ -dependent para-nitrophenylphosphatase activity in native and FITC pretreated enzyme

NBD-Cl (mM)	Relative velocity in native enzyme (%)	Relative velocity in FITC-pretreated enzyme (%)
0	100	100
20	69.7	67.6
40	33.3	34.5
60	13.8	16.7
100	6.5	6.3
500	0.7	1.3

$\text{Na}^+/\text{K}^+$ -ATPase in a final concentration of 1 U/ml (65 mg/ml) was incubated overnight at  $37^\circ\text{C}$  in a solution containing 20 mM Tris/HCl (pH 7.25) and 10  $\mu\text{M}$  FITC and a control without FITC was run in parallel. The enzyme was then incubated in 20 mM Tris/HCl (pH 7.25), 15 mM NaCl and various concentrations (0–500  $\mu\text{M}$ ) NBD-Cl. After incubation for 1 h at  $37^\circ\text{C}$  remaining  $\text{K}^+$ -activated para-nitrophenylphosphatase was estimated. Mean values of two experiments are shown.




Fig. 2. Protective effect of  $\text{Co}(\text{NH}_3)_4\text{ATP}$  towards NBD-Cl labeling. FITC-pretreated  $\text{Na}^+/\text{K}^+$ -ATPase was incubated for 2 h at  $37^\circ\text{C}$  in a solution containing 20 mM Tris/HCl (pH 7.5) with 1 mM (○) and without (■)  $\text{Co}(\text{NH}_3)_4\text{ATP}$  and then in 20 mM Tris/HCl buffer (pH 7.5) for 1 h at  $37^\circ\text{C}$  with increasing concentrations of NBD-Cl (0–500  $\mu\text{M}$ ). The fluorescence of these probes was read at  $\lambda_{\text{ex}}$  475 nm and  $\lambda_{\text{em}}$  520 nm. A control without NBD-Cl was run in parallel as background. Mean values and standard deviation of four experiments are shown.

NBD-Cl on the partial activities could be prevented by millimolar concentrations of ATP (data not shown). The protective effect against NBD-Cl labeling was even more pronounced when the FITC-inactivated enzyme was additionally treated by the  $\text{E}_2\text{ATP}$  site specific MgATP complex analog  $\text{Co}(\text{NH}_3)_4\text{ATP}$  (Fig. 2). The incorporation of the NBD-Cl label was almost completely blocked by this ATP derivative. Apparently, NBD-Cl reacted with the  $\text{E}_2\text{ATP}$  site in the same way independent of whether the  $\text{E}_1\text{ATP}$  site was blocked by FITC or not. To study the effect of NBD-Cl on the high-affinity ATP-binding site ( $\text{E}_1\text{ATP}$  site) of  $\text{Na}^+/\text{K}^+$ -ATPase, its effect on  $\text{Na}^+$ -dependent phosphorylation from  $[\gamma\text{-}^{32}\text{P}]\text{ATP}$  (frontdoor) in an  $\text{E}_2\text{ATP}$  site blocked enzyme (by  $\text{Co}(\text{NH}_3)_4\text{PO}_4$ ) was measured. As is evident from Table 2, NBD-Cl modified the  $\text{E}_1\text{ATP}$  site in the untreated control enzyme much more efficiently than in the  $\text{E}_2\text{ATP}$  site blocked enzyme. Generally, less NBD-Cl was necessary to modify the  $\text{E}_2\text{ATP}$  site than the  $\text{E}_1\text{ATP}$  site (compare Tables 1 and 2). These experiments clearly showed that NBD-Cl modified both




Fig. 3. Fluorescence analysis of fluorescence parameters of NBD-Cl-labeled  $\text{Na}^+/\text{K}^+$ -ATPase. Stern-Volmer plots of quenching of NBD-Cl-labeled  $\text{Na}^+/\text{K}^+$ -ATPase by  $\text{Mn}^{2+}$ -ions (insert shows quenching by  $\text{Cs}^+$ -ions) bound to enzyme pretreated with 1 mM  $\text{Co}(\text{NH}_3)_4\text{ATP}$  (□, ○) and 25  $\mu\text{M}$   $\text{Cr}(\text{H}_2\text{O})_4\text{AdoPP}[\text{CH}_2]\text{P}$  (■, ●). Mean values of three experiments are shown.

ATP sites and, unlike FITC, was unable to discriminate between these sites.

Besides inactivation, modification of  $\text{Na}^+/\text{K}^+$ -ATPase by NBD-Cl resulted in a spectral shift of the NBD-Cl absorption spectrum from 330 nm to 425 nm and in the appearance of fluorescence with a maximum at 520 nm. Both the spectral shift and the appearance of fluorescence, were a clear indication of NBD-Cl modifications of the enzyme. This was observed when the enzyme was labeled either with or without  $\text{Co}(\text{NH}_3)_4\text{ATP}$  or  $\text{Cr}(\text{H}_2\text{O})_4\text{AdoPP}[\text{CH}_2]\text{P}$ . The former compound is known as a specific inhibitor of the  $\text{E}_2\text{ATP}$  site [2,5,7] while the latter blocks specifically the  $\text{E}_1\text{ATP}$  site [2,3]. The absorption or fluorescence spectra of enzyme-bound NBD-Cl did not differ significantly if the labeling of the enzyme was done in the presence of  $\text{Co}(\text{NH}_3)_4\text{ATP}$  or  $\text{Cr}(\text{H}_2\text{O})_4\text{AdoPP}[\text{CH}_2]\text{P}$  ( $\text{E}_1\text{ATP}$  site or  $\text{E}_2\text{ATP}$  site is blocked). Quenching experiments, however, clearly distinguished between the two types of ATP site blocking. We used both positively and negatively charged quenchers:  $\text{Cs}^+$ ,  $\text{Mn}^{2+}$  and  $\text{I}^-$ . In all cases, the Stern-Volmer plot was not monotonic (Fig. 3) and the modified Stern-Volmer plot revealed a fraction of inaccessible fluorophores.

Table 2  
Effect of NBD-Cl on  $\text{Na}^+$ -dependent frontdoor phosphorylation

Time (min)	Phosphorylation of $\text{Co}(\text{NH}_3)_4\text{PO}_4$ -pretreated enzyme (%) <sup>a</sup>		Phosphorylation of native enzyme (%) <sup>b</sup>
	+0.1 mM NBD-Cl	+1 mM NBD-Cl	+1 mM NBD-Cl
0	100	100	100
5	n.d.	91	37
15	86	81	30
30	n.d.	74	15
45	72	65	6.5
60	68	50	6.0
90	61	n.d.	n.d.
120	58	n.d.	n.d.

$\text{Na}^+/\text{K}^+$ -ATPase (3 U/ml; 195  $\mu\text{g}/\text{ml}$ ) where the  $\text{E}_2\text{ATP}$  site was blocked with 1 mM  $\text{Co}(\text{NH}_3)_4\text{PO}_4$  (see Section 2) was incubated in 20 mM Tris/HCl, 15 mM NaCl and 100  $\mu\text{M}$  or 1 mM NBD-Cl. A native control enzyme was treated in the same way with 1 mM NBD-Cl. Frontdoor phosphorylation from  $[\gamma\text{-}^{32}\text{P}]$  was studied as a function of the exposure time of the enzyme at  $37^\circ\text{C}$  with NBD-Cl. For experimental details see Section 2 and [5,7]. Mean values of three experiments are shown.

<sup>a</sup>Control is  $\text{Co}(\text{NH}_3)_4\text{PO}_4$ -pretreated  $\text{Na}^+/\text{K}^+$ -ATPase without NBD-Cl measured after the same incubation time (100% is 33.3 pmole phosphate/unit).

<sup>b</sup>Control is native  $\text{Na}^+/\text{K}^+$ -ATPase without NBD-Cl measured after the same incubation time (100% is 100 pmole phosphate/unit). n.d., not determined.

Interestingly, only a minute difference, if any, was detected between fluorescence quenching of  $\text{Na}^+/\text{K}^+$ -ATPase labeled with NBD-Cl in the presence of  $\text{Cr}(\text{H}_2\text{O})_4\text{AdoPP}[\text{CH}_2]\text{P}$  ( $\text{E}_1$ ATP site is blocked) and in the absence of any analog. The enzyme pretreated with  $\text{Co}(\text{NH}_3)_4\text{ATP}$  ( $\text{E}_2$ ATP site is blocked), on the other hand, showed different fluorescence properties as compared with the enzyme labeled without ATP analogs. The presence of  $\text{Co}(\text{NH}_3)_4\text{ATP}$  clearly changed the bimolecular quenching constant calculated for low concentrations of the quenchers, i.e. as an initial slope. The obtained Stern-Volmer quenching constants were  $(9.6 \pm 0.8) \cdot 10^{-4}$  for  $\text{Cs}^+$ ,  $(3.3 \pm 0.3) \cdot 10^{-2}$  for  $\text{I}^-$  and  $(2.0 \pm 0.2) \cdot 10^{-3}$  for  $\text{Mn}^{2+}$  in the case of  $\text{Cr}(\text{H}_2\text{O})_4\text{AdoPP}[\text{CH}_2]\text{P}$ -pretreated enzyme. For  $\text{Co}(\text{NH}_3)_4\text{ATP}$ -pretreated  $\text{Na}^+/\text{K}^+$ -ATPase where the low-affinity ATP-binding site was occupied, the bimolecular quenching constants decreased to  $(4.6 \pm 0.4) \cdot 10^{-4}$  for  $\text{Cs}^+$ ,  $(2.3 \pm 0.2) \cdot 10^{-2}$  for  $\text{I}^-$  and  $(1.2 \pm 0.2) \cdot 10^{-3}$  for  $\text{Mn}^{2+}$ . Additionally, we also observed a different profile of anisotropy changes when the fluorescence anisotropy was followed during incubation of the enzyme with NBD-Cl. The anisotropy after a 6 h incubation of the  $\text{Cr}(\text{H}_2\text{O})_4\text{AdoPP}[\text{CH}_2]\text{P}$ -pretreated enzyme with  $100 \mu\text{M}$  NBD-Cl at  $37^\circ\text{C}$  (virtually a total binding) was 0.128 while in the presence of  $\text{Co}(\text{NH}_3)_4\text{ATP}$  the anisotropy value increased to 0.137.

#### 4. Discussion

We confirmed in this study earlier reports that NBD-Cl inhibits  $\text{Na}^+/\text{K}^+$ -ATPase with two different affinities [9,10]. This was evident not only from the kinetics of inactivation with  $K_i$  values of 2.5 and  $10 \mu\text{M}$  (Fig. 1), but also from the fact that NBD-Cl inactivated both the  $\text{K}^+$ -activated para-nitrophenylphosphatase activity in an enzyme where the  $\text{E}_1$ ATP site was blocked by FITC (Table 1) and the  $\text{Na}^+$ -dependent frontdoor phosphorylation in an enzyme with the  $\text{E}_2$ ATP site blocked by  $\text{Co}(\text{NH}_3)_4\text{PO}_4$  (Table 2). Thus, we can conclude that NBD-Cl is a fluorescent probe labeling both the low- and high-affinity ATP-binding sites. To our knowledge, this is the first study describing a covalent fluorescent label with an affinity for both ATP-binding sites. In combination with site-specific MgATP derivatives or fluorescent pseudo ATP analogs, this probe is a potent tool for clarifying the structure and mechanism of the low- and the high-affinity ATP-binding sites on  $\text{Na}^+/\text{K}^+$ -ATPase.

Fluorescence intensity studies of NBD-Cl bound to  $\text{Na}^+/\text{K}^+$ -ATPase in this work revealed two populations of fluorophores (Fig. 3). The bimolecular quenching constants for all three quenchers ( $\text{Cs}^+$ ,  $\text{I}^-$  and  $\text{Mn}^{2+}$ ) were significantly higher for the  $\text{Cr}(\text{H}_2\text{O})_4\text{AdoPP}[\text{CH}_2]\text{P}$ - $\text{Na}^+/\text{K}^+$ -ATPase than for the enzyme treated with  $\text{Co}(\text{NH}_3)_4\text{ATP}$ . Hence, the accessibility of the quenchers to the fluorophore bound to the  $\text{E}_2$ ATP site was higher than to the fluorophore bound to the  $\text{E}_1$ ATP site.

Modification of either of the two ATP sites resulted in a rotational immobilization of the fluorophore which was reflected in a relatively high fluorescence anisotropy. This anisotropy was higher when the  $\text{E}_2$ ATP-binding site was blocked than when the blockage affected the  $\text{E}_1$ ATP-binding site. Consequently, the  $\text{E}_2$ ATP site seems to be sterically more open as compared with the  $\text{E}_1$ ATP site. The fact that the high-affinity binding site ( $\text{E}_1$ ATP site) is sterically poorly accessible was shown previously by FITC labeling [19].

**Acknowledgements:** H.L. thanks for a FEBS short term fellowship. This work was supported by the Fonds der Chemischen Industrie, Frankfurt/M, the Deutsche Forschungsgemeinschaft (Bonn Bad-Godesberg) through Graduiertenkolleg 'Molekularbiologie und Pharmakologie' at the Justus-Liebig-University, Giessen, and grants No. 204/95/0624 of the Grant Agency of the Czech Republic, No. 5011505 of the Grant Agency of Czech Academy of Sciences and No. 95.00923CT04 of CNR. Cooperation between the institutes was supported by a bilateral German-Czech governmental program through 'IWTZ' No. TSR-088-97.

#### References

- [1] Skou, J.C. and Esmann, M. (1992) *J. Bioenerg. Biomembr.* 24, 249–261.
- [2] Linnertz, H., Thönges, D. and Schoner, W. (1995) *Eur. J. Biochem.* 232, 420–424.
- [3] Hamer, E. and Schoner, W. (1993) *Eur. J. Biochem.* 213, 743–748.
- [4] Peluffo, R.D., Garrahan, P.J. and Rega, A.F. (1992) *J. Biol. Chem.* 267, 6596–6601.
- [5] Buxbaum, E. and Schoner, W. (1991) *Eur. J. Biochem.* 195, 407–419.
- [6] Yamayaki, A., Kaya, S., Tsuda, T., Araki, Y., Hayashi, Y. and Taniguchi, K. (1994) *J. Biochem.* 116, 1360–1369.
- [7] Thönges, D. and Schoner, W. (1997) *J. Biol. Chem.* 272, 16315–16321.
- [8] Thönges, D., Linnertz, H. and Schoner, W. (1997) *Ann. N.Y. Acad. Sci.*, in press.
- [9] Cantley, L.C., Gelles, J. and Josephson, L. (1978) *Biochemistry* 17, 418–425.
- [10] Grosse, R., Eckert, K., Malur, J. and Repke, K.R.H. (1978) *Acta Biol. Med. Germ.* 37, 83–96.
- [11] Karlisch, S.J. (1980) *J. Bioenerg. Biomembr.* 12, 111–136.
- [12] Linnertz, H., Lastres Becker, I., Krumscheid, R., Amler, E., Thönges, D. and Schoner, W. (1997) *Physiol. Res.* 46, 345–349.
- [13] Buxbaum, E. and Schoner, W. (1990) *Eur. J. Biochem.* 193, 355–360.
- [14] Jørgensen, P.L. (1975) *Biochim. Biophys. Acta* 401, 399–415.
- [15] Schoner, W., von Illberg, C., Kramer, R. and Seubert, W. (1967) *Eur. J. Biochem.* 1, 334–343.
- [16] Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J. (1951) *J. Biol. Chem.* 193, 2165–2175.
- [17] Piszkievics, D. and Smith, E.L. (1971) *Biochemistry* 10, 4544–4552.
- [18] Amler, E., Abbott, A. and Ball, W.J. (1992) *Biophys. J.* 61, 553–568.
- [19] Abbot, A.J., Amler, E. and Ball, W.J. (1991) *Biochemistry* 30, 1692–1701.