

Non-coding RNAs with essential roles in neurodegenerative disorders



Evgenia Salta, Bart De Strooper

The importance of various classes of regulatory non-protein-coding RNA molecules (ncRNAs) in the normal functioning of the CNS is becoming increasingly evident. ncRNAs are involved in neuronal cell specification and patterning during development, but also in higher cognitive processes, such as structural plasticity and memory formation in the adult brain. We discuss advances in understanding of the function of ncRNAs in the CNS, with a focus on the potential involvement of specific species, such as microRNAs, endogenous small interfering RNAs, long intergenic non-coding RNAs, and natural antisense transcripts, in various neurodegenerative disorders. This emerging field is anticipated to profoundly affect clinical research, diagnosis, and therapy in neurology.

Lancet Neurol 2012;
11: 189–200

Centre for Human Genetics and Leuven Institute for Neurodegenerative Disorders (LIND), University of Leuven, Leuven, Belgium (E Salta PhD, B De Strooper MD); and VIB Centre for the Biology of Disease (VIB11), Leuven, Belgium (E Salta, B De Strooper)

Correspondence to:
Dr Bart De Strooper, VIB Centre for the Biology of Disease (VIB11), Herestraat 49, Box 602, 3000 Leuven, Belgium
Bart.Destrooper@cme.vib-kuleuven.be

Introduction

The idea that the genome exerts its functions only via classical genes and proteins seems more and more a naive oversimplification of a fascinating system of feed-forward and feedback loops that involves various RNA molecules. Indeed, a large part of the genomic code is only transcribed as RNA and is never translated into protein. Only in the past decade has non-protein-coding RNA (ncRNA) been recognised to have important functions. The phenomenon of so-called pervasive transcription^{1–3} provides the transcriptome and the proteome with an incredibly complex layer of regulation and fine-tuning.

ncRNAs are abundantly expressed in mammalian CNS in specific spatiotemporal patterns.^{4–7} MicroRNAs (miRNAs) and other ncRNAs are features of virtually all neuronal activity, including neurogenesis, neuronal patterning, neurotransmission, and synaptic plasticity (figure 1).^{8–14} They maintain cellular homeostasis with time and energy efficiency, which is particularly relevant in the nervous system, where rigorous requirements for continuous signalling and plasticity necessitate the rapid and accurate buffering of gene products along long processes and to distant synapses (figure 1).^{8,10,15} The implications of this knowledge for the understanding of neurological and psychiatric diseases have only begun to be clarified compared with those for other diseases, and possible medical applications in diagnosis and therapy are not yet being sufficiently explored. Notably, the number of scientific publications on the roles of ncRNAs in cancer pathology is around ten times higher than that on their involvement in physiological and pathological processes in the CNS. In cancer, miRNAs, long intergenic non-coding RNAs (lincRNAs), and other long non-coding RNAs (lncRNAs) exert oncogenic or tumour-suppressing effects on tumorigenesis, invasion, angiogenesis, and metastasis.^{16–19}

In this Review we provide an overview of ncRNA biology in the CNS and discuss possible roles in neurological disorders, with a focus on neurodegeneration (webappendix). We draw lessons from pioneering translational studies in cancer and cardiovascular disease and note where more thorough and systematic work is

needed in human tissues and in animal models before findings can be translated into daily clinical practice for neurological disorders.

Biology of ncRNAs

Molecules of ncRNA are roughly classified into two size groups: small (<400 nucleotides) and long (>400 nucleotides). Small ncRNAs comprise well characterised infrastructural RNA, such as ribosomal RNAs, transfer RNAs, and small nuclear/spliceosomal RNAs. Other types of small ncRNAs are regulatory RNAs, such as miRNAs, small interfering RNAs (siRNAs), piwi-interacting RNAs, splice junction-associated RNAs, and small nucleolar RNAs.²⁰ Long ncRNAs can exceed 100 000 nucleotides and cover a wide range of heterogeneous regulatory molecules, such as lincRNAs,²¹ natural antisense transcripts (NATs),²² and non-coding RNA repeats (table, figure 2).³³ In many cases, however, ncRNAs have mixed characteristics and do not clearly fall into any of these categories.

Why 98.8% of the human genome consists of non-coding DNA (ncDNA) has puzzled researchers for many years.³⁹ High-resolution and high-throughput screening technologies, such as microarrays and next-generation sequencing (deep sequencing), have shown that in fact most of this ncDNA encodes ncRNA.^{2,39} Strong evidence, including conservation of promoter sequences, dynamic expression during development and disease, and expression patterns and responsiveness to stimuli that are specific to tissue, cell, and subcellular structures, argues for the functionality of these ncRNAs.^{2,20} Nevertheless, further studies are needed to address the extent to which each of the transcripts is operational and to identify the transcripts that affect specific molecular pathways in a physiological context.

The ncRNA machinery involves RNA–RNA, RNA–DNA, and RNA–protein interactions, and affects transcription, stability, processing, and translation of messenger RNA (mRNA), alternative splicing, and epigenetic events, such as RNA editing and chromatin modification.^{9,10} These regulatory phenomena are effected on neighbouring transcripts (in cis) or at loci distant from their own transcription (in trans) and depend mainly on

See Online for webappendix




Figure 1: Physiological roles of regulatory ncRNAs in the CNS

Examples of ncRNAs involved in the physiology of the CNS and affecting processes ranging from neuronal development to maturation. ncRNAs can be involved in the maintenance of the stem cell status of neuronal stem cells and their differentiation into neuronal, astrocytic, or oligodendroglial precursors, and also in the maintenance, lineage specification, maturation, and differentiation of precursor cells into their respective cell types. Additionally, ncRNAs affect processes such as activation and proliferation of astrocytes, survival, maturation, and homeostasis of neurons, and myelination of mature oligodendrocytes. At the level of the synapse, ncRNAs might regulate local translation, neurite outgrowth, synaptogenesis, neurotransmitter signalling, synaptic plasticity, and long-term potentiation. ncRNA=non-protein-coding RNA. lincRNA=long intergenic non-coding RNA. lncRNA=long non-coding RNA. snRNA=small nuclear RNA. eRNA=enhancer RNA. endo-siRNA=endogenous small interfering RNA. BC1=brain cytoplasmic RNA of 150 nucleotides. BC200=brain cytoplasmic RNA of 200 nucleotides.

the complementarity between the ncRNA sequence and DNA or RNA molecules. In many cases, structural changes in regulatory or structural proteins can also be induced by the secondary structure of particular ncRNAs;² this phenomenon is similar to the mechanism of action of regulatory proteins. Notably, ncRNAs are regulated by the CNS in an activity-dependent manner, which results in rapid changes in their expression.^{10,15}

In view of their broad biology and function, it is unsurprising that the numbers of ncRNAs recognised as contributing to key pathogenic mechanisms are increasing. Below we look at the characteristics and roles of specific ncRNA species in the context of neurological diseases.

miRNAs and neurodegenerative disorders

miRNAs were discovered in *Caenorhabditis elegans* almost 20 years ago,⁴⁰ and they now represent the most extensively studied class of ncRNAs. This species of RNA is particularly important for the fine-tuning of neuronal networks.⁴¹ miRNA genes are transcribed mainly by RNA polymerase II. The resulting long primary transcript can contain several

miRNA stretches, but subsequent processing by a series of enzymes, including nuclear Drosha and cytoplasmic Dicer, results in one or more double-stranded, mature miRNA molecules of around 21 nucleotides in length. One of the strands becomes incorporated into the RNA-induced ribonucleoprotein silencing complex and, after base pairing with the 3'-UTR of the target mRNA, induces most of the time mRNA degradation or repression.^{23,24,42} Of note, translation activation has also been reported.⁴³ Each of the roughly 700 miRNAs identified so far in the human genome can theoretically bind to a few hundred mRNAs.⁴⁴ This feature raises the question of whether all the interactions are equally important; in many reported examples, one particular miRNA has affected one specific mRNA that is responsible for most of the biological effects exerted. For instance, in phenotype rescue experiments the mir-310–313 cluster regulated synaptic strength at the neuromuscular junction mainly via one target, Khc-73,¹¹ whereas activity-dependent dendritic growth, which was mediated by mir-132, was attributed to the translational regulation of the Rho family GTPase-activating protein p250GAP.⁴⁵

A rather crude and general way to investigate the biological relevance of the miRNA machinery in the CNS is to genetically delete the miRNA-processing enzyme Dicer. This enzyme (a classical protein) controls final maturation of precursor miRNA into the tiny operational miRNAs. The full deletion of Dicer in mice is embryonic lethal,⁴⁶ whereas conditional inactivation of Dicer selectively in Purkinje cells or in midbrain dopaminergic, cortical, or hippocampal neurons leads to degeneration of the cells.⁴⁷ Follow-up studies have investigated which miRNAs are specifically involved in these degenerative phenotypes. Changes in miRNAs have been detected in various neurodegenerative disorders (eg, Alzheimer's disease,^{48–54} Parkinson's disease,^{55,56} Huntington's disease,^{57–60} and prion disease^{61,62}), and proteins involved in these diseases can be regulated by miRNAs. Thus, the hypothesis that changes in miRNAs are involved in the neurodegenerative disease process, either causally or as part of positive feedback loops, is gaining ground. Available studies in patients, however, have only provided snapshots of the disease process and have not shown how miRNAs change with disease progression. The observed alterations might also be only epiphenomena of the underlying disease process. In view of the important regulatory roles of miRNAs in brain physiology, the reality is probably that some miRNA changes are part of the disease process and others are due to the loss of cells or other secondary alterations. To gain understanding of both possibilities is crucial and might yield novel diagnostic markers and deeper insight into the neurodegenerative disease process itself.

Alzheimer's disease

The list of protein targets relevant to neurodegenerative diseases is rapidly growing. In Alzheimer's disease, two of the most important proteins are extensively controlled (at least in vitro) by miRNAs: amyloid precursor protein (APP), which carries the amyloid- β peptide that precipitates in the amyloid plaques, and BACE1, which cleaves APP in the generation of amyloid- β peptide (figure 3 and webappendix). Of note, post-mortem analysis of brains from patients with Alzheimer's disease has revealed increased levels of BACE1 but not of its mRNA,^{50,67,68} which indicates post-transcriptional regulation of the protein under pathological conditions. The marked decrease in the expression of miR-29a/b-1 and miR-107 and increased concentrations of BACE1 as Alzheimer's disease progresses suggest a direct functional link between these miRNAs and the protease in the disease process.^{49,50} Several neuronal miRNAs are differentially regulated after exposure to amyloid- β peptide, which suggests that aberrant miRNA function contributes to the elusive toxic effects of amyloid- β peptide in patients with Alzheimer's disease.⁶⁹ The link to tau pathology in this disease remains less clear, but deletion of Dicer in adult mouse brain led to changes in tau phosphorylation and neurodegeneration, which supports a potential link

between dysfunctional miRNA pathways and tangle pathology during the disease course.⁷⁰ In one report, mir-132 was noted to be involved in the abnormal tau splicing pattern in sporadic progressive supranuclear palsy, which is a neurodegenerative tauopathy.⁷¹

Parkinson's disease

Evidence for a direct or indirect role of miRNAs in the pathogenesis of Parkinson's disease is accumulating (figure 4 and webappendix). One of the pathological hallmarks of this disease is fibrillar aggregates (Lewy bodies) in the dopaminergic neurons of the substantia nigra that contain α -synuclein. The expression of α -synuclein is fine-tuned by specific miRNAs (figure 4 and webappendix). Furthermore, a report has implicated the most important gene in Parkinson's disease, *LRRK2*, in the negative regulation of let-7 and mir-184* functions in dopaminergic neurons. The respective targets of these

	Origin	Mechanisms and functions
Short		
miRNAs ^{23,24}	Sense, intergenic or intronic	Incorporate into RISC (miRISC), base pair to 3'-UTR of mRNA targets, and mainly induce translational repression or deadenylation and degradation
endo-siRNAs ^{24,25}	Sense or antisense, intergenic or exonic	Incorporate into RISC (siRISC), base pair to mRNA target, and induce degradation or heterochromatin formation
piRNAs ²⁶	Sense or antisense, intergenic	Epigenetic and possible translational control via complementarity with DNA or RNA sequences
PASRs ¹	Sense, intergenic (promoter region)	Unknown
TASRs ¹	Antisense, intergenic (3'-UTR end of genes)	Unknown
snoRNAs ²⁷	Sense, intergenic or intronic	Pre-RNA processing or nucleoside modification (2'-O-ribose methylation and pseudouridylation) of other RNA molecules
snRNAs ²⁸	Sense, intergenic or intronic	pre-mRNA splicing
tiRNAs ^{29,30}	Sense or antisense, intergenic (5'-UTR transcription initiation sites)	Possibly promotes transcription via epigenetic regulation
spliRNAs ³⁰	Sense, exonic (splice donor site)	Possible epigenetic regulation
Long		
lincRNAs ³¹	Sense or antisense, intergenic	Epigenetic regulation
NATs ³²	Antisense, exonic or intronic	mRNA transcription, splicing, stability, and translation, epigenetic modifications, and precursors of endo-siRNAs
RNA expansion repeats ³³	Sense or antisense, exonic or intronic	Epigenetic regulation and RNA toxic effects
ENORs ³⁴	Sense or antisense, intergenic	Transcriptional regulation, genomic imprinting, precursors of other short and long ncRNAs
eRNAs ³⁵	Sense, intergenic (enhancer region)	Transcriptional regulation
PALRs ¹	Sense, overlap with promoter and first exon of genes	Unknown
miRNAs=micro RNAs. RISC=RNA-induced silencing complex. mRNA=messenger RNA. endo-si RNAs=endogenous small interfering RNAs. piRNAs=piwi-interacting RNAs. PASRs=promoter-associated short RNAs. TASRs=termini-associated short RNAs. snoRNAs=small nucleolar RNAs. snRNAs=small nuclear RNAs. tiRNAs=transcription initiation RNAs. spliRNAs=splice junction-associated RNAs. lincRNAs=long intergenic non-coding RNAs. NATs=natural antisense transcripts. ENORs=expressed non-coding regions. eRNAs=enhancer RNAs. PALRs=promoter-associated long RNAs.		
Table: Non-protein-coding RNA species		




Figure 2: Complexity of transcription patterns of ncRNAs

The two strands of genomic DNA are displayed together with the basic building blocks of a gene. ncRNAs can be transcribed by one of three different RNA polymerases (RNA polymerase I, II, or III) from intergenic (DNA sequences between two genes) and intragenic (intronic or mostly unstable exonic transcripts) genomic regions. Transcription occurs in patterns nested in and overlapping with protein-coding transcripts but also in regions where no classical genes are found (even heterochromatin sequences²). Both sense and antisense orientations of the genome are used.^{2,8,9,36,37} Some ncRNAs undergo further modifications similar to authentic mRNAs, including polyadenylation (conferring stability)³⁸ or 5' end nucleotide capping (which provides stability but is also important for cellular transport and localisation of the transcripts).³⁶ Notably, sometimes ncRNAs undergo alternative splicing.⁹ Clearly, these RNA molecules are processed and regulated in ways as complex as those that affect proteins. ncRNA=non-protein-coding RNA. endo-siRNA=endogenous small interfering RNA. piRNA=piwi-interacting RNA. snRNA=small nuclear RNA. snoRNA=small nucleolar RNA. miRNA=microRNA. sdRNA=sno-derived RNA. lincRNA=long intergenic non-coding RNA. eRNA=enhancer RNA. PALR=promoter-associated long RNA. PASR=promoter-associated short RNA. tiRNA=transcription initiation RNA. spliRNA=splice junction-associated RNA. mRNA=messenger RNA. NAT=natural antisense transcript. TASR=termini-associated short RNA.

miRNAs, E2F1 and DP, are in turn overexpressed, which results in defective cell division and eventually cell death.⁷⁵ These observations need confirmation in mammalian brain to definitively establish a role for LRRK2 in this pathway. Finally, mir-133b, a dopaminergic neuron-specific miRNA that is downregulated in Parkinson's disease, forms a negative feedback loop with the transcription factor PITX3, which in the CNS is expressed exclusively in midbrain dopaminergic neurons.⁵⁵ Although a direct role of mir-133b in the loss of dopaminergic neurons is not supported by this report, its downregulation might counteract a neuroprotective mechanism.

Other neurodegenerative disorders

The pathogenic mechanism in Huntington's disease is not completely understood. Trinucleotide expansion in the gene that encodes huntingtin is, however, known to impair the cytoplasmic binding of this protein to REST/NRSF, which represses transcription of neuronal genes in non-neuronal cells. This effect could partly explain the increased nuclear translocation of REST/NRSF and the consequent aberrant decrease in neuronal gene expression in the cortex of patients with Huntington's disease.^{58,60} Several miRNAs are regulated by REST/CoREST.⁶⁰ miR-9 and

miR-9* are thought to be part of a double negative feedback loop, as they are regulated by the REST repressor complex but they also target REST and CoREST expression.⁵⁷ Multiple studies of miRNAs in mouse and human brain tissue have provided further links to neurodegenerative diseases, including spinocerebellar ataxia type 1,^{76,77} amyotrophic lateral sclerosis,⁷⁸ frontotemporal dementia,⁷⁹ prion diseases,^{61,62} spinal muscular atrophy,⁸⁰ and myotonic dystrophy type I.⁸¹ Collectively, these data clearly support pivotal functions for miRNAs in neurodegeneration via interaction with and regulation of crucial players in the pertinent pathways or through their cardinal roles in regulating the survival and function of neurons, which affect mitochondrial function,^{82,83} apoptosis,⁵² inflammation,⁸⁴ and synaptic plasticity.⁸⁵

Single-nucleotide polymorphisms

Few studies have investigated the importance of single-nucleotide polymorphisms (SNPs) identified in genome-wide association studies for ncRNA biology. Many SNPs reside in the non-coding parts of the genome. Somewhat incorrectly, the increased disease risk associated with an identified SNP is frequently interpreted to imply roles for protein-encoding genes near the polymorphism.

Of the thousands of SNPs that have been characterised, many occur at experimentally validated target sites of miRNAs, but the variation in actual miRNA sequences is low.^{86–89} Most of those SNPs were predicted to create novel miRNA-binding sites. Potentially, such polymorphisms might contribute to the phenotypic differences between individuals.^{86,89} Cancer research is at the forefront in the analysis of this phenomenon,⁹⁰ but the study of these SNPs in neurodegeneration has only just started. For example, bioinformatic analysis has shown that seven genes associated with the risk of Alzheimer's disease bear SNPs in their 3'-UTRs at miRNA-binding sites, which suggests a possible role for variable miRNA regulation of these genes in the course of the disease.⁹¹ Additionally, aberrant regulation of APP expression by miRNAs owing to the introduction of SNPs in the 3'-UTR has been related to the pathophysiology of Alzheimer's disease.⁹² A polymorphism in an miRNA-binding site in *GRN* was proposed to result in decreased progranulin concentrations, which might in turn

represent a risk factor in frontotemporal dementia and Alzheimer's disease.^{79,93} A 3'-UTR SNP associated with Parkinson's disease and translational control by miR-433, however, did not correlate with a specific population of patients with this disease.⁹⁴

Future challenges

An increase is expected in the number of functional studies in mouse models that directly explore the effects on disease processes of loss of specific miRNAs in neurons or glial cells. The generation of a complete mouse embryonic stem cell library with all miRNA genes conditionally targeted⁹⁵ makes this task highly feasible. Major challenges are the expansion of the number of patients assessed and stratification of the data by disease stage, brain area, and quality of material used. Single-cell analysis will also become very important in the identification of cell-specific changes. The ability to detect alterations in miRNAs in the brain by analysis of cerebrospinal fluid, or even better the analysis of plasma,




Figure 3: Regulatory ncRNAs in Alzheimer's disease

Some of the paradigms of ncRNA regulation in the pathogenesis of Alzheimer's disease that involve microRNAs, natural antisense transcripts, and other ncRNAs are shown. More specifically, APP mRNA is regulated by miR-101,^{63,64} miR-106a, miR-520c,⁶⁵ miR-124,⁵³ and let-7⁶⁶ and BACE1 mRNA is regulated by miR-298,⁶⁷ miR-107,⁴⁹ and miR-29a/b-1.⁵⁰ BACE1-AS reacts to deposition of amyloid-β peptide and in turn binds to and stabilises BACE1 mRNA, resulting in increased BACE1 expression and APP processing. BC1 and BC200 reside in synapses, where they are involved in activity-dependent translational regulation. Finally, 17A, an antisense small ncRNA, affects the alternative splicing of the GABABR2 receptor, which leads to aberrant GABA_B signalling and increased production of amyloid-β peptide. APP= amyloid precursor protein. ncRNA=non-protein-coding RNA. BC1=brain cytoplasmic RNA of 150 nucleotides. BC200=brain cytoplasmic RNA of 200 nucleotides. BACE1-AS=BACE1 antisense transcript. mRNA=messenger RNA.




Figure 4: Regulatory ncRNAs in Parkinson's disease

Examples of ncRNAs possibly directly or indirectly involved in the dopaminergic phenotype of Parkinson's disease. The expression of α -synuclein is fine-tuned by specific microRNAs (eg, mir-7,^{72,73} mir-433,⁷⁴ and mir-153⁷⁵). Under pathogenic conditions LRRK2 mediates a microRNA-dependent transcriptional regulation mechanism in dopaminergic neurons, whereas mir-133b is involved in a negative feedback loop that results in perturbed transcriptional control. Finally, PINK1-AS and PINK1 mRNA levels are co-regulated during mitochondrial biogenesis. ncRNAs=non-protein-coding RNAs. mRNA=messenger RNA.

would remarkably improve the diagnostic toolbox for Alzheimer's disease, Parkinson's disease, Huntington's disease, and prion and other neurological diseases.

Other small regulatory ncRNAs

The number of small regulatory ncRNAs identified in the CNS is rapidly growing. siRNAs that occur naturally in plants have been used in neurology research to manipulate expression of genes, for instance, in the context of antisense therapies. Endogenous siRNAs have also been noted in mammalian cells. They apparently originate from transposable elements, cis-natural antisense transcripts, mammalian pseudogenes, and long foldback transcripts called hairpin RNAs.²⁵ In contrast to miRNAs, siRNAs require perfect sequence complementarity with their targets. Furthermore, they are derived from long, double-stranded RNAs and their maturation process is independent of Drosha. One study revealed with deep sequencing that expression of many endogenous siRNA-like RNAs strikingly changes after hippocampus-dependent learning.⁹⁶ Some of these short RNAs align to important neuronal proteins, such as BACE1 and hSIRT3 and, therefore, they are proposed to be functionally involved in learning-associated neuronal processes. Further experimental data are needed, however, to confirm the role of endogenous siRNAs in cognition and memory.

Certain small nucleolar RNAs identified in mouse brain have been implicated in learning and memory,^{27,97} and small nuclear RNAs in the spliceosome, which is responsible for the splicing of precursor mRNA in eukaryotes, seem to have additional functions.²⁸ The assembly of small nuclear RNA–ribonucleoprotein complexes is directed by the survival of motor neuron protein (SMN), whereas deficiency of SMN leads to spinal muscular atrophy with characteristic motor neuron degradation.⁹⁸ Regulation of alternative splicing in the brain by small ncRNAs has been described,⁹⁹ as has altered expression of piwi-interacting RNAs in response to induced ischaemia in rat brains.¹⁰⁰

Some newly identified small RNA species include promoter-associated short RNAs and termini-associated short RNAs, which originate, respectively, from regions at the 5' or 3' end of genes,¹ tiny RNAs associated with transcription initiation sites,²⁹ and RNAs associated with splice junctions.³⁰ Many of these small ncRNAs seem to originate from longer sense or antisense transcripts, which indicates a complex system of cis-regulatory and trans-regulatory interplay.¹⁰¹ Tiling arrays (high-density microarrays consisting of oligonucleotide probes that cover the whole genome of a given species) have revealed that almost half of the protein-coding genes in the human genome are flanked by promoter-associated short RNAs and termini-associated short RNAs (22–200 nucleotides long), some of which are expressed at the same level as their overlapping genes. This finding supports claims of biological relevance,¹ but their functions in the CNS have not yet been investigated.

Two small ncRNAs of particular note are a brain cytoplasmic RNA of 200 nucleotides (BC200) and one of 150 nucleotides (BC1; figure 3). They are transcribed by RNA polymerase III and are specifically expressed in primate and rodent brains, respectively.^{102,103} These two ncRNAs are actively transported to dendrites, where they participate in activity-dependent translational regulation during synaptogenesis, and in control of the excitation–repression equilibrium in the synaptodendritic compartment, which promotes neuronal plasticity.^{103,104} Upregulation and cellular mislocalisation of BC200 have been seen in regions of disease-affected brain in patients with Alzheimer's disease. Thus, BC200 seems to have a reactive or even causative role in neurodegeneration related to Alzheimer's disease, possibly by perturbed translational control at the synapse.¹⁰⁵

Finally, a unique, small, double-stranded, non-coding RNA has been identified that interacts with and converts the REST/NRSF complex from repressor to activator. This action triggers the expression of neuron-specific genes in adult hippocampal stem cells and directs them towards a neuronal identity.¹⁰⁶

lincRNAs

Despite their abundance in the genome, only a handful of lincRNAs have so far been functionally characterised. This species of ncRNA is transcribed by RNA polymerase II from genomic loci located between two genes, and varies in length from 2000 to 20000 nucleotides.³¹ lincRNAs display limited sequence conservation across species, which raises the question of whether they represent merely transcriptional noise.^{107,108} However, in a study that used a chromatin-immunoprecipitation sequencing approach (ultra-high-throughput sequencing used initially to detect transcription factors associated with transcriptionally active DNA regions), more than 1000 evolutionarily conserved lincRNAs were predicted that might counteract the function of transcription factors and, therefore, repress transcription.³¹

Another chromatin-immunoprecipitation sequencing study identified a brain-specific cluster of lincRNAs that could be implicated in brain ageing, hippocampal development, oligodendrocyte myelination, synaptic transmission, and signalling pathways involving the transcription factor CREB, GABA, and G-protein-coupled receptors.³¹ Experimental evidence from a transcriptomics analysis in mice reveals a role for AK044422, a lincRNA abundantly expressed in brain tissue, in lineage specification of GABAergic neurons.¹⁰⁹ Of note is the fact that AK044422 RNA additionally serves as an miRNA precursor. A follow-up study identified almost 3300 lincRNAs and suggested that a common function is binding to and guidance of chromatin remodelling complexes (enzymatic complexes that catalyse chromatin modifications and lead to transcriptional silencing or activation), such as the polycomb repressive complex 2 and CoREST (a primary cofactor of REST), which has been implicated in Huntington's disease.^{110–112}

NATs

NATs are RNA polymerase II transcripts that originate from the opposite strand of annotated, protein-coding sense mRNAs. Many of them have mRNA-like properties, such as 5' capping, 3' polyadenylation, and alternative splicing. Various studies have suggested that more than 70% of human and mouse genes undergo such antisense transcription.³² NATs might also encode proteins, although most represent non-coding transcripts. The mechanisms of action of NATs involve both cis and trans interactions with complementary sense transcripts,^{113,114} and affect a wide variety of cellular processes, including transcription, splicing, stability, and translation of mRNA, epigenetic modifications (leading to an increase or decrease in the levels of the sense mRNA transcript), and generation of endogenous siRNAs.^{32,115}

Many NATs have been specifically detected in the CNS and have been postulated to regulate important neuronal processes, such as stem cell renewal and proliferation, stress responses,¹¹⁶ cortical neuron specification and migration,¹¹⁷ oligodendrocyte differentiation,¹¹⁸ myelination,¹¹⁹ synaptic transmission, plasticity,^{120,121} and long-term memory formation.¹²² BACE1 antisense transcript, a conserved non-coding transcript, becomes upregulated when cells are exposed to the amyloid- β peptide (figure 3).¹²³ It interacts with BACE1 transcript and seems to stabilise this mRNA, which leads to increased BACE1 expression and a further increase in amyloid- β production through a positive feed-forward loop.¹²³ BACE1 and BACE1 antisense transcripts show similar expression patterns and their levels are raised in brain samples from patients with Alzheimer's disease and in *APP* transgenic mice.¹²³ BACE1 antisense transcript is thought to act by masking the binding site of mir-485-5p on BACE1 mRNA; miRNAs repress translation of their target mRNA.¹²⁴

Another amyloid-responsive NAT is the neuronal NAT-Rad18, which is upregulated after exposure of rat

cortical neurons to amyloid- β peptide and leads to post-transcriptional downregulation of the DNA repair protein Rad18.¹²⁵ This finding indicates that RNA control machinery is involved in the DNA damage repair system in Alzheimer's disease. Additionally, a NAT against apolipoprotein E might be particularly relevant to this disease: levels of both apolipoprotein E and apolipoprotein E-AS1 increase in response to CNS injury, and the latter might regulate expression of the former.¹²⁶

One of the genetic forms of Parkinson's disease is caused by mutations in the mitochondrial gene *PINK1*.¹²⁷ Antisense transcripts of the *PINK1* locus have been identified in human beings and mice and are concordantly regulated with *PINK1* mRNA during mitochondrial biogenesis in vivo.^{128,129} In Huntington's disease, a NAT called HAR1 targets REST/NRSF and has been implicated in the pathogenesis of the disease, as greatly decreased levels have been seen in the striatum of patients.¹³⁰

Collectively, these data suggest that NATs are important for the fine-tuning of gene expression. In view of their abundance in the CNS, these transcripts are expected to influence almost all important signalling pathways in neurobiology, and identification of many more naturally occurring antisense transcripts with regulatory roles in a variety of neurodegenerative disorders is anticipated.

Other long regulatory ncRNAs

New classes and subclasses of lincRNAs with distinct and overlapping genomic and functional features are frequently being reported. For instance, long expressed non-coding regions (also known as macroRNAs or ENORs) constitute a class of very long sense or antisense, mainly nuclear, ncRNAs that are specifically enriched in the CNS and function through antisense transcriptional regulation and genomic imprinting. They serve as precursors of other lincRNAs, miRNAs, and small nucleolar RNAs.³⁴

Enhancer RNAs represent a separate class of lincRNAs and are transcribed by RNA polymerase II from activity-regulated enhancers (DNA sequences that can activate gene transcription from remote positions) that are frequently associated with the transcription of neuronal genes.^{13,35,131} A brain-specific enhancer RNA, termed *Evf2*, is transcribed from an intergenic enhancer locus and acts through recruitment of positive (DLX family) and negative (MECP-2) transcription factors to the enhancer region, thereby increasing the transcription of the downstream gene. Of note, this first lincRNA knockout mouse model showed that *Evf2* might act both in cis and in trans to control the transcription of the homoeodomain transcription factors *Dlx-5* and *Dlx-6* and also of the enzyme *Gad1* (necessary for the conversion of glutamate to GABA) in the developing mouse brain. This effect ensures normal development of GABAergic inhibitory interneurons in the hippocampus, and proper synaptic activity in the embryonic and adult brain.¹³²

Promoter-associated long RNAs are non-coding RNA transcripts that originate from the same genomic loci as

Search strategy and selection criteria

We searched PubMed with the search terms “non coding RNA”, “non coding RNA and neurodegeneration”, “non coding RNA and disease”, and “non coding RNA and central nervous system” for papers published from 1987 to November, 2011. We also manually searched the reference lists of published work to identify further relevant references. Only papers published in English were reviewed. The final reference list was generated on the basis of originality and relevance to the broad scope of this Review.

promoter-associated short RNAs and overlap with the 5' end and the first exons and introns of certain genes.¹

An RNA toxic gain-of-function mechanism involving a variety of non-coding sense, antisense, intronic, or exonic RNA repeats has been implicated in several nucleotide repeat expansion disorders. These include different spinocerebellar ataxias,^{133–136} myotonic dystrophies,^{137,138} Huntington's disease-like 2,¹³⁹ and fragile X tremor ataxia.¹⁴⁰

Many more lncRNA species have been detected in specific regions of the CNS. These include Gomafu¹⁴¹ (an mRNA-like, brain-specific repetitive ncRNA that is preferentially expressed in rat brain and further gives rise to a small nucleolar RNA),^{142,143} Ntab,¹⁴⁴ Sox2dot, an isoform of a non-coding Sox2-overlapping transcript that is enriched in regions of adult neurogenesis,¹⁴⁵ and metastasis-associated lung adenocarcinoma transcript 1 (a highly abundant neuronal ncRNA that has been implicated in synapse formation).¹⁴⁶ The latter lncRNA was identified as a binding target of TAR DNA-binding protein 43, a nuclear protein that forms inclusion bodies in frontotemporal lobar degeneration and amyotrophic lateral sclerosis.^{147,148}

Conclusions and future directions

Although still in its early days, the study of ncRNAs in neurological disease, and in neurodegeneration in particular, is clearly a research area of great potential. We expect that further in-depth study of the specific patterns and expression profiles of these ncRNAs in different diseases will ultimately result in the identification of new diagnostic markers. The initial expression-profiling studies of miRNAs have shown the potential of this approach. The available information, however, should be interpreted with caution. Differences in brain sampling area, variability between individuals, insufficient clinical stratification,⁴⁷ and, in many cases, inherent bias related to the platforms used to detect the RNA molecules, could introduce obscure errors that make comparisons difficult. Moreover, to what extent the altered expression of RNA in the brain will be reflected in alterations that can be measured in biological fluids, such as CSF or plasma, remains unclear. miRNAs, for instance, have been detected in serum and plasma,¹⁴⁹ but their origins remain controversial. One

possibility is secretion from cells via microvesicles and exosomes (membranous vesicles released from the cell surface that contribute to cell-to-cell communication) in peripheral blood or CSF.^{150–154} Some studies have validated changes in miRNA concentrations in body fluids in the context of cancer and cardiovascular disorders,^{155,156} which suggests that further work in neurological diseases might be fruitful.

Further study of ncRNA biology might eventually result in novel therapeutic applications. So-called RNA-based drugs, which are synthetic RNA molecules generated to mimic the mechanisms of action of endogenous ncRNAs—typically siRNAs or miRNAs—have been developed for many diseases.¹⁵⁷ The best developed and most studied approach is RNA interference, which involves synthetic double-stranded siRNAs, short hairpin RNAs, and artificial miRNA precursors that are mainly expressed by viral vectors and intended to suppress the translation of specific disease-related molecules.¹⁵⁷ Antisense molecules that antagonise miRNAs, such as antagomirs, locked nucleic acids, and morpholinos, have generally been chemically modified to improve specificity and plasma half-life in vivo.¹⁵⁸ Of note, the first miRNA-based therapeutic agent, a mir-122 locked nucleic acid antisense drug, is currently being tested in a phase 2 clinical trial in chronic hepatitis C patients.¹⁵⁹ While RNA-interference technology might become useful for the treatment of malignant and cardiovascular diseases,^{160,161} neurobiology has its own particular issues, the most important being the blood–brain barrier.¹⁶² Advances in siRNA delivery into the brain have, however, been reported.¹⁶³ Finally, issues concerning toxic effects in the CNS and off-target effects have to be seriously considered.^{164–166}

An exponential increase in knowledge with regard to the role of ncRNAs in the CNS seems likely. We predict that this research will rapidly translate into increasingly precise diagnostic applications for neurology clinics. The prospects for novel treatments are realistic, but new and more efficient brain-targeting strategies are required. The versatility and straightforward nature of RNA chemistry, however, makes rapid translation into therapeutic strategies a reasonable hope.

Contributors

ES and BDS jointly did literature searches, prepared the figures and tables, and wrote and revised the paper.

Conflicts of interest

We declare that we have no conflicts of interest.

Acknowledgments

Funding was received from the European Research Council, Fund for Scientific Research, Flanders, KU Leuven, the VIB, Methusalem grant from KU Leuven and the Flemish government, the Foundation for Alzheimer Research (SAO/FRMA), and the Interuniversity Attraction Poles Program (IAP P6/43). BDS is the Arthur Bax and Anna Vanluffelen chair for Alzheimer's disease.

References

- 1 Kapranov P, Cheng J, Dike S, et al. RNA maps reveal new RNA classes and a possible function for pervasive transcription. *Science* 2007; **316**: 1484–88.

- 2 Berretta J, Morillon A. Pervasive transcription constitutes a new level of eukaryotic genome regulation. *EMBO Rep* 2009; **10**: 973–82.
- 3 Brosius J. Waste not, want not—transcript excess in multicellular eukaryotes. *Trends Genet* 2005; **21**: 287–88.
- 4 Mehler MF, Mattick JS. Non-coding RNAs in the nervous system. *J Physiol* 2006; **575**: 333–41.
- 5 Mercer TR, Dinger ME, Sunkin SM, Mehler MF, Mattick JS. Specific expression of long noncoding RNAs in the mouse brain. *Proc Natl Acad Sci USA* 2008; **105**: 716–21.
- 6 Kapsimali M, Kloosterman WP, de Bruijn E, Rosa F, Plasterk RH, Wilson SW. MicroRNAs show a wide diversity of expression profiles in the developing and mature central nervous system. *Genome Biol* 2007; **8**: R173.
- 7 Bak M, Silahatoglu A, Moller M, et al. MicroRNA expression in the adult mouse central nervous system. *RNA* 2008; **14**: 432–44.
- 8 Cao X, Yeo G, Muotri AR, Kuwabara T, Gage FH. Noncoding RNAs in the mammalian central nervous system. *Annu Rev Neurosci* 2006; **29**: 77–103.
- 9 Mehler MF, Mattick JS. Noncoding RNAs and RNA editing in brain development, functional diversification, and neurological disease. *Physiol Rev* 2007; **87**: 799–823.
- 10 Mercer TR, Dinger ME, Mariani J, Kosik KS, Mehler MF, Mattick JS. Noncoding RNAs in long-term memory formation. *Neuroscientist* 2008; **14**: 434–45.
- 11 Tsurudome K, Tsang K, Liao EH, et al. The *Drosophila* miR-310 cluster negatively regulates synaptic strength at the neuromuscular junction. *Neuron* 2010; **68**: 879–93.
- 12 Gao FB. Context-dependent functions of specific microRNAs in neuronal development. *Neural Dev* 2010; **5**: 25.
- 13 Qureshi IA, Mehler MF. Non-coding RNA networks underlying cognitive disorders across the lifespan. *Trends Mol Med* 2011; **17**: 337–46.
- 14 Siegel G, Saba R, Schrat G. microRNAs in neurons: manifold regulatory roles at the synapse. *Curr Opin Genet Dev* 2011; **21**: 491–97.
- 15 St Laurent G 3rd, Faghihi MA, Wahlestedt C. Non-coding RNA transcripts: sensors of neuronal stress, modulators of synaptic plasticity, and agents of change in the onset of Alzheimer's disease. *Neurosci Lett* 2009; **466**: 81–88.
- 16 Huarte M, Rinn JL. Large non-coding RNAs: missing links in cancer? *Hum Mol Genet* Oct 2010; **19**: R152–61.
- 17 Iorio MV, Croce CM. MicroRNAs in cancer: small molecules with a huge impact. *J Clin Oncol* 2009; **27**: 5848–56.
- 18 Gibb EA, Brown CJ, Lam WL. The functional role of long non-coding RNA in human carcinomas. *Mol Cancer* 2011; **10**: 38.
- 19 Tsai MC, Spitale RC, Chang HY. Long intergenic noncoding RNAs: new links in cancer progression. *Cancer Res* 2011; **71**: 3–7.
- 20 Mattick JS. The central role of RNA in human development and cognition. *FEBS Lett* 2011; **585**: 1600–16.
- 21 Qureshi IA, Mattick JS, Mehler MF. Long non-coding RNAs in nervous system function and disease. *Brain Res* 2010; **1338**: 20–35.
- 22 Werner A. Natural antisense transcripts. *RNA Biol* 2005; **2**: 53–62.
- 23 Krol J, Loedige I, Filipowicz W. The widespread regulation of microRNA biogenesis, function and decay. *Nat Rev Genet* 2010; **11**: 597–610.
- 24 Carthew RW, Sontheimer EJ. Origins and mechanisms of miRNAs and siRNAs. *Cell* 2009; **136**: 642–55.
- 25 Okamura K, Lai EC. Endogenous small interfering RNAs in animals. *Nat Rev Mol Cell Biol* 2008; **9**: 673–78.
- 26 Thomson T, Lin H. The biogenesis and function of PIWI proteins and piRNAs: progress and prospect. *Annu Rev Cell Dev Biol* 2009; **25**: 355–76.
- 27 Dieci G, Preti M, Montanini B. Eukaryotic snoRNAs: a paradigm for gene expression flexibility. *Genomics* 2009; **94**: 83–88.
- 28 Valadkhan S. snRNAs as the catalysts of pre-mRNA splicing. *Curr Opin Chem Biol* 2005; **9**: 603–08.
- 29 Taft RJ, Glazov EA, Cloonan N, et al. Tiny RNAs associated with transcription start sites in animals. *Nat Genet* 2009; **41**: 572–78.
- 30 Taft RJ, Simons C, Nahkuri S, et al. Nuclear-localized tiny RNAs are associated with transcription initiation and splice sites in metazoans. *Nat Struct Mol Biol* 2010; **17**: 1030–34.
- 31 Guttman M, Amit I, Garber M, et al. Chromatin signature reveals over a thousand highly conserved large non-coding RNAs in mammals. *Nature* 2009; **458**: 223–27.
- 32 Werner A, Sayer JA. Naturally occurring antisense RNA: function and mechanisms of action. *Curr Opin Nephrol Hypertens* 2009; **18**: 343–49.
- 33 Nakamori M, Thornton C. Epigenetic changes and non-coding expanded repeats. *Neurobiol Dis* 2010; **39**: 21–27.
- 34 Furuno M, Pang KC, Ninomiya N, et al. Clusters of internally primed transcripts reveal novel long noncoding RNAs. *PLoS Genet* 2006; **2**: e37.
- 35 Kim TK, Hemberg M, Gray JM, et al. Widespread transcription at neuronal activity-regulated enhancers. *Nature* 2010; **465**: 182–87.
- 36 Carninci P. Constructing the landscape of the mammalian transcriptome. *J Exp Biol* 2007; **210**: 1497–506.
- 37 Mattick JS. The central role of RNA in the genetic programming of complex organisms. *An Acad Bras Cienc* 2010; **82**: 933–39.
- 38 Cheng J, Kapranov P, Drenkow J, et al. Transcriptional maps of 10 human chromosomes at 5-nucleotide resolution. *Science* 2005; **308**: 1149–54.
- 39 Lindberg J, Lundeberg J. The plasticity of the mammalian transcriptome. *Genomics* 2010; **95**: 1–6.
- 40 Lee RC, Feinbaum RL, Ambros V. The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell* 1993; **75**: 843–54.
- 41 Saba R, Schrat GM. MicroRNAs in neuronal development, function and dysfunction. *Brain Res* 2010; **1338**: 3–13.
- 42 Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 2004; **116**: 281–97.
- 43 Vasudevan S, Tong Y, Steitz JA. Switching from repression to activation: microRNAs can up-regulate translation. *Science* 2007; **318**: 1931–34.
- 44 Christensen M, Schrat GM. microRNA involvement in developmental and functional aspects of the nervous system and in neurological diseases. *Neurosci Lett* 2009; **466**: 55–62.
- 45 Wayman GA, Davare M, Ando H, et al. An activity-regulated microRNA controls dendritic plasticity by down-regulating p250GAP. *Proc Natl Acad Sci USA* 2008; **105**: 9093–98.
- 46 Bernstein E, Kim SY, Carmell MA, et al. Dicer is essential for mouse development. *Nat Genet* 2003; **35**: 215–17.
- 47 Hebert SS, De Strooper B. Alterations of the microRNA network cause neurodegenerative disease. *Trends Neurosci* 2009; **32**: 199–206.
- 48 Lukiw WJ. Micro-RNA speciation in fetal, adult and Alzheimer's disease hippocampus. *Neuroreport* 2007; **18**: 297–300.
- 49 Wang WX, Rajeev BW, Stromberg AJ, et al. The expression of microRNA miR-107 decreases early in Alzheimer's disease and may accelerate disease progression through regulation of β -site amyloid precursor protein-cleaving enzyme 1. *J Neurosci* 2008; **28**: 1213–23.
- 50 Hebert SS, Horre K, Nicolai L, et al. Loss of microRNA cluster miR-29a/b-1 in sporadic Alzheimer's disease correlates with increased BACE1/ β -secretase expression. *Proc Natl Acad Sci USA* 2008; **105**: 6415–20.
- 51 Wang H, Liu J, Zong Y, et al. miR-106b aberrantly expressed in a double transgenic mouse model for Alzheimer's disease targets TGF- β type II receptor. *Brain Res* 2010; **1357**: 166–74.
- 52 Wang X, Liu P, Zhu H, et al. miR-34a, a microRNA up-regulated in a double transgenic mouse model of Alzheimer's disease, inhibits bcl2 translation. *Brain Res Bull* 2009; **80**: 268–73.
- 53 Smith P, Al Hashimi A, Girard J, Delay C, Hebert SS. *In vivo* regulation of amyloid precursor protein neuronal splicing by microRNAs. *J Neurochem* 2011; **116**: 240–47.
- 54 Sethi P, Lukiw WJ. Micro-RNA abundance and stability in human brain: specific alterations in Alzheimer's disease temporal lobe neocortex. *Neurosci Lett* 2009; **459**: 100–04.
- 55 Kim J, Inoue K, Ishii J, et al. A MicroRNA feedback circuit in midbrain dopamine neurons. *Science* 2007; **317**: 1220–24.
- 56 Miñones-Moyano E, Porta S, Escaramis G, et al. MicroRNA profiling of Parkinson's disease brains identifies early downregulation of miR-34b/c which modulate mitochondrial function. *Hum Mol Genet* 2011; **20**: 3067–78.
- 57 Packer AN, Xing Y, Harper SQ, Jones L, Davidson BL. The bifunctional microRNA miR-9/miR-9* regulates REST and CoREST and is downregulated in Huntington's disease. *J Neurosci* 2008; **28**: 14341–46.

- 58 Marti E, Pantano L, Banez-Coronel M, et al. A myriad of miRNA variants in control and Huntington's disease brain regions detected by massively parallel sequencing. *Nucleic Acids Res* 2010; **38**: 7219–35.
- 59 Johnson R, Zuccato C, Belyaev ND, Guest DJ, Cattaneo E, Buckley NJ. A microRNA-based gene dysregulation pathway in Huntington's disease. *Neurobiol Dis* 2008; **29**: 438–45.
- 60 Lee ST, Chu K, Im WS, et al. Altered microRNA regulation in Huntington's disease models. *Exp Neurol* 2011; **227**: 172–79.
- 61 Saba R, Goodman CD, Huzarewich RL, Robertson C, Booth SA. A miRNA signature of prion induced neurodegeneration. *PLoS One* 2008; **3**: e3652.
- 62 Montag J, Hitt R, Opitz L, Schulz-Schaeffer WJ, Hunsmann G, Motzkus D. Upregulation of miRNA hsa-miR-342-3p in experimental and idiopathic prion disease. *Mol Neurodegener* 2009; **4**: 36.
- 63 Long JM, Lahiri DK. MicroRNA-101 downregulates Alzheimer's amyloid- β precursor protein levels in human cell cultures and is differentially expressed. *Biochem Biophys Res Commun* 2011; **404**: 889–95.
- 64 Vilardo E, Barbato C, Ciotti M, Cogoni C, Ruberti F. MicroRNA-101 regulates amyloid precursor protein expression in hippocampal neurons. *J Biol Chem* 2010; **285**: 18344–51.
- 65 Patel N, Hoang D, Miller N, et al. MicroRNAs can regulate human APP levels. *Mol Neurodegener* 2008; **3**: 10.
- 66 Niwa R, Zhou F, Li C, Slack FJ. The expression of the Alzheimer's amyloid precursor protein-like gene is regulated by developmental timing microRNAs and their targets in *Caenorhabditis elegans*. *Dev Biol* 2008; **315**: 418–25.
- 67 Boissonneault V, Plante I, Rivest S, Provost P. MicroRNA-298 and microRNA-328 regulate expression of mouse β -amyloid precursor protein-converting enzyme 1. *J Biol Chem* 2009; **284**: 1971–81.
- 68 Yang LB, Lindholm K, Yan R, et al. Elevated beta-secretase expression and enzymatic activity detected in sporadic Alzheimer disease. *Nat Med* 2003; **9**: 3–4.
- 69 Schonrock N, Ke YD, Humphreys D, et al. Neuronal microRNA deregulation in response to Alzheimer's disease amyloid- β . *PLoS One* 2010; **5**: e11070.
- 70 Hebert SS, Papadopoulou AS, Smith P, et al. Genetic ablation of *Dicer* in adult forebrain neurons results in abnormal tau hyperphosphorylation and neurodegeneration. *Hum Mol Genet* 2010; **19**: 3959–69.
- 71 Smith PY, Delay C, Girard J, et al. MicroRNA-132 loss is associated with tau exon 10 inclusion in progressive supranuclear palsy. *Hum Mol Genet* 2011; **20**: 4016–24.
- 72 Junn E, Lee KW, Jeong BS, Chan TW, Im JY, Mouradian MM. Repression of α -synuclein expression and toxicity by microRNA-7. *Proc Natl Acad Sci USA* 2009; **106**: 1305–07.
- 73 Doxakis E. Post-transcriptional regulation of alpha-synuclein expression by mir-7 and mir-153. *J Biol Chem* 2010; **285**: 12726–34.
- 74 Wang G, van der Walt JM, Mayhew G, et al. Variation in the miRNA-433 binding site of FGF20 confers risk for Parkinson disease by overexpression of alpha-synuclein. *Am J Hum Genet* 2008; **82**: 283–89.
- 75 Gehrke S, Imai Y, Sokol N, Lu B. Pathogenic LRRK2 negatively regulates microRNA-mediated translational repression. *Nature* 2010; **466**: 637–41.
- 76 Bilen J, Liu N, Burnett BG, Pittman RN, Bonini NM. MicroRNA pathways modulate polyglutamine-induced neurodegeneration. *Mol Cell* 2006; **24**: 157–63.
- 77 Lee Y, Samaco RC, Gatchel JR, Thaller C, Orr HT, Zoghbi HY. miR-19, miR-101 and miR-130 co-regulate ATXN1 levels to potentially modulate SCA1 pathogenesis. *Nat Neurosci* 2008; **11**: 1137–39.
- 78 Williams AH, Valdez G, Moresi V, et al. MicroRNA-206 delays ALS progression and promotes regeneration of neuromuscular synapses in mice. *Science* 2009; **326**: 1549–54.
- 79 Rademakers R, Eriksen JL, Baker M, et al. Common variation in the miR-659 binding-site of GRN is a major risk factor for TDP43-positive frontotemporal dementia. *Hum Mol Genet* 2008; **17**: 3631–42.
- 80 Haramati S, Chapnik E, Sztainberg Y, et al. miRNA malfunction causes spinal motor neuron disease. *Proc Natl Acad Sci USA* 2010; **107**: 13111–16.
- 81 Perbellini R, Greco S, Sarra-Ferraris G, et al. Dysregulation and cellular mislocalization of specific miRNAs in myotonic dystrophy type 1. *Neuromuscul Disord* 2011; **21**: 81–88.
- 82 Aschrafi A, Schwlechter AD, Mameza MG, Natera-Naranjo O, Gioio AE, Kaplan BB. MicroRNA-338 regulates local cytochrome c oxidase IV mRNA levels and oxidative phosphorylation in the axons of sympathetic neurons. *J Neurosci* 2008; **28**: 12581–90.
- 83 Ouyang YB, Lu Y, Yue S, Giffard RG. miR-181 targets multiple Bcl-2 family members and influences apoptosis and mitochondrial function in astrocytes. *Mitochondrion* 2011; published online Sept 17. DOI:10.1016/j.mito.2011.09.001.
- 84 Lukiw WJ, Zhao Y, Cui JG. An NF- κ B-sensitive micro RNA-146a-mediated inflammatory circuit in Alzheimer disease and in stressed human brain cells. *J Biol Chem* 2008; **283**: 31315–22.
- 85 Kocerha J, Faghihi MA, Lopez-Toledano MA, et al. MicroRNA-219 modulates NMDA receptor-mediated neurobehavioral dysfunction. *Proc Natl Acad Sci USA* 2009; **106**: 3507–12.
- 86 Glinsky GV. SNP-guided microRNA maps (MirMaps) of 16 common human disorders identify a clinically accessible therapy reversing transcriptional aberrations of nuclear import and inflammasome pathways. *Cell Cycle* 2008; **7**: 3564–76.
- 87 Saunders MA, Liang H, Li WH. Human polymorphism at microRNAs and microRNA target sites. *Proc Natl Acad Sci USA* 2007; **104**: 3300–05.
- 88 Richardson K, Lai CQ, Parnell LD, Lee YC, Ordovas JM. A genome-wide survey for SNPs altering microRNA seed sites identifies functional candidates in GWAS. *BMC Genomics* 2011; **12**: 504.
- 89 Mishra PJ, Bertino JR. MicroRNA polymorphisms: the future of pharmacogenomics, molecular epidemiology and individualized medicine. *Pharmacogenomics* 2009; **10**: 399–416.
- 90 Zhang L, Liu Y, Song F, et al. Functional SNP in the microRNA-367 binding site in the 3' UTR of the calcium channel ryanodine receptor gene 3 (RYS3) affects breast cancer risk and calcification. *Proc Natl Acad Sci USA* 2011; **108**: 13653–58.
- 91 Mallick B, Ghosh Z. A complex crosstalk between polymorphic microRNA target sites and AD prognosis. *RNA Biol* 2011; **8**: 665–73.
- 92 Delay C, Calon F, Mathews P, Hebert SS. Alzheimer-specific variants in the 3'UTR of amyloid precursor protein affect microRNA function. *Mol Neurodegener* 2011; **6**: 70.
- 93 Dickson DW, Baker M, Rademakers R. Common variant in GRN is a genetic risk factor for hippocampal sclerosis in the elderly. *Neurodegener Dis* 2010; **7**: 170–74.
- 94 de Mena L, Cardo LF, Coto E, et al. FGF20 rs12720208 SNP and microRNA-433 variation: no association with Parkinson's disease in Spanish patients. *Neurosci Lett* 2010; **479**: 22–25.
- 95 Prosser HM, Koike-Yusa H, Cooper JD, Law FC, Bradley A. A resource of vectors and ES cells for targeted deletion of microRNAs in mice. *Nat Biotechnol* 2011; **29**: 840–45.
- 96 Smalheiser NR, Lugli G, Thimmapuram J, Cook EH, Larson J. Endogenous siRNAs and noncoding RNA-derived small RNAs are expressed in adult mouse hippocampus and are up-regulated in olfactory discrimination training. *RNA* 2011; **17**: 166–81.
- 97 Matera AG, Terns RM, Terns MP. Non-coding RNAs: lessons from the small nuclear and small nucleolar RNAs. *Nat Rev Mol Cell Biol* 2007; **8**: 209–20.
- 98 Zhang Z, Lotti F, Dittmar K, et al. SMN deficiency causes tissue-specific perturbations in the repertoire of snRNAs and widespread defects in splicing. *Cell* 2008; **133**: 585–600.
- 99 Massone S, Vassallo I, Fiorino G, et al. 17A, a novel non-coding RNA, regulates GABA B alternative splicing and signaling in response to inflammatory stimuli and in Alzheimer disease. *Neurobiol Dis* 2011; **41**: 308–17.
- 100 Dharap A, Nakka VP, Vemuganti R. Altered expression of PIWI RNA in the rat brain after transient focal ischemia. *Stroke* 2011; **42**: 1105–09.
- 101 St Laurent G 3rd, Wahlestedt C. Noncoding RNAs: couplers of analog and digital information in nervous system function? *Trends Neurosci* 2007; **30**: 612–21.
- 102 DeChiara TM, Brosius J. Neural BC1 RNA: cDNA clones reveal nonrepetitive sequence content. *Proc Natl Acad Sci USA* 1987; **84**: 2624–28.

- 103 Tiedge H, Chen W, Brosius J. Primary structure, neural-specific expression, and dendritic location of human BC200 RNA. *J Neurosci* 1993; **13**: 2382–90.
- 104 Zhong J, Chuang SC, Bianchi R, et al. Regulatory BC1 RNA and the fragile X mental retardation protein: convergent functionality in brain. *PLoS One* 2010; **5**: e15509.
- 105 Mus E, Hof PR, Tiedge H. Dendritic BC200 RNA in aging and in Alzheimer's disease. *Proc Natl Acad Sci USA* 2007; **104**: 10679–84.
- 106 Kuwabara T, Hsieh J, Nakashima K, Taira K, Gage FH. A small modulatory dsRNA specifies the fate of adult neural stem cells. *Cell* 2004; **116**: 779–93.
- 107 Pang KC, Frith MC, Mattick JS. Rapid evolution of noncoding RNAs: lack of conservation does not mean lack of function. *Trends Genet* 2006; **22**: 1–5.
- 108 Ponting CP, Oliver PL, Reik W. Evolution and functions of long noncoding RNAs. *Cell* 2009; **136**: 629–41.
- 109 Mercer TR, Qureshi IA, Gokhan S, et al. Long noncoding RNAs in neuronal-glial fate specification and oligodendrocyte lineage maturation. *BMC Neurosci* 2010; **11**: 14.
- 110 Khalil AM, Guttman M, Huarte M, et al. Many human large intergenic noncoding RNAs associate with chromatin-modifying complexes and affect gene expression. *Proc Natl Acad Sci USA* 2009; **106**: 11667–72.
- 111 Zuccato C, Tartari M, Crotti A, et al. Huntingtin interacts with REST/NRSF to modulate the transcription of NRSE-controlled neuronal genes. *Nat Genet* 2003; **35**: 76–83.
- 112 Buckley NJ, Johnson R, Zuccato C, Bithell A, Cattaneo E. The role of REST in transcriptional and epigenetic dysregulation in Huntington's disease. *Neurobiol Dis* 2010; **39**: 28–39.
- 113 Osato N, Suzuki Y, Ikeo K, Gojobori T. Transcriptional interferences in cis natural antisense transcripts of humans and mice. *Genetics* 2007; **176**: 1299–306.
- 114 Li JT, Zhang Y, Kong L, Liu QR, Wei L. Trans-natural antisense transcripts including noncoding RNAs in 10 species: implications for expression regulation. *Nucleic Acids Res* 2008; **36**: 4833–44.
- 115 Faghihi MA, Wahlestedt C. Regulatory roles of natural antisense transcripts. *Nat Rev Mol Cell Biol* 2009; **10**: 637–43.
- 116 Korneev S, O'Shea M. Natural antisense RNAs in the nervous system. *Rev Neurosci* 2005; **16**: 213–22.
- 117 Pollard KS, Salama SR, Lambert N, et al. An RNA gene expressed during cortical development evolved rapidly in humans. *Nature* 2006; **443**: 167–72.
- 118 Tochitani S, Hayashizaki Y. Nkx2.2 antisense RNA overexpression enhanced oligodendrocytic differentiation. *Biochem Biophys Res Commun* 2008; **372**: 691–96.
- 119 Okano H, Aruga J, Nakagawa T, Shiota C, Mikoshiba K. Myelin basic protein gene and the function of antisense RNA in its repression in myelin-deficient mutant mouse. *J Neurochem* 1991; **56**: 560–67.
- 120 Ling KH, Hewitt CA, Beissbarth T, et al. Spatiotemporal regulation of multiple overlapping sense and novel natural antisense transcripts at the Nrgn and Camk2n1 gene loci during mouse cerebral corticogenesis. *Cereb Cortex* 2011; **21**: 683–97.
- 121 Pruunsild P, Kazantseva A, Aid T, Palm K, Timmusk T. Dissecting the human BDNF locus: bidirectional transcription, complex splicing, and multiple promoters. *Genomics* 2007; **90**: 397–406.
- 122 Korneev SA, Straub V, Kemenes I, et al. Timed and targeted differential regulation of nitric oxide synthase (NOS) and anti-NOS genes by reward conditioning leading to long-term memory formation. *J Neurosci* 2005; **25**: 1188–92.
- 123 Faghihi MA, Modarresi F, Khalil AM, et al. Expression of a noncoding RNA is elevated in Alzheimer's disease and drives rapid feed-forward regulation of beta-secretase. *Nat Med* 2008; **14**: 723–30.
- 124 Faghihi MA, Zhang M, Huang J, et al. Evidence for natural antisense transcript-mediated inhibition of microRNA function. *Genome Biol* 2010; **11**: R56.
- 125 Parenti R, Paratore S, Torrisi A, Cavallaro S. A natural antisense transcript against Rad18, specifically expressed in neurons and upregulated during β -amyloid-induced apoptosis. *Eur J Neurosci* 2007; **26**: 2444–57.
- 126 Seitz A, Gourevitch D, Zhang XM, et al. Sense and antisense transcripts of the apolipoprotein E gene in normal and ApoE knockout mice, their expression after spinal cord injury and corresponding human transcripts. *Hum Mol Genet* 2005; **14**: 2661–70.
- 127 Morais VA, Verstreken P, Roethig A, et al. Parkinson's disease mutations in PINK1 result in decreased Complex I activity and deficient synaptic function. *EMBO Mol Med* 2009; **1**: 99–111.
- 128 Scheele C, Petrovic N, Faghihi MA, et al. The human PINK1 locus is regulated *in vivo* by a non-coding natural antisense RNA during modulation of mitochondrial function. *BMC Genomics* 2007; **8**: 74.
- 129 Chiba M, Kiyosawa H, Hiraiwa N, Ohkohchi N, Yasue H. Existence of Pink1 antisense RNAs in mouse and their localization. *Cytogenet Genome Res* 2009; **126**: 259–70.
- 130 Johnson R, Richter N, Jauch R, et al. The human accelerated region 1 noncoding RNA is repressed by REST in Huntington's disease. *Physiol Genomics* 2010; published online Feb 23. DOI:10.1152/physiolgenomics.00019.2010.
- 131 Orom UA, Derrien T, Beringer M, et al. Long noncoding RNAs with enhancer-like function in human cells. *Cell* 2010; **143**: 46–58.
- 132 Bond AM, Vangompel MJ, Sametsky EA, et al. Balanced gene regulation by an embryonic brain ncRNA is critical for adult hippocampal GABA circuitry. *Nat Neurosci* 2009; **12**: 1020–27.
- 133 Sopher BL, Ladd PD, Pineda VV, et al. CTCF regulates ataxin-7 expression through promotion of a convergently transcribed, antisense noncoding RNA. *Neuron* 2011; **70**: 1071–84.
- 134 Koob MD, Moseley ML, Schut LJ, et al. An untranslated CTG expansion causes a novel form of spinocerebellar ataxia (SCA8). *Nat Genet* 1999; **21**: 379–84.
- 135 Moseley ML, Zu T, Ikeda Y, et al. Bidirectional expression of CUG and CAG expansion transcripts and intranuclear polyglutamine inclusions in spinocerebellar ataxia type 8. *Nat Genet* 2006; **38**: 758–69.
- 136 Matsuura T, Yamagata T, Burgess DL, et al. Large expansion of the ATTCT pentanucleotide repeat in spinocerebellar ataxia type 10. *Nat Genet* 2000; **26**: 191–94.
- 137 Brook JD, McCurrach ME, Harley HG, et al. Molecular basis of myotonic dystrophy: expansion of a trinucleotide (CTG) repeat at the 3' end of a transcript encoding a protein kinase family member. *Cell* 1992; **69**: 385.
- 138 Liquori CL, Ricker K, Moseley ML, et al. Myotonic dystrophy type 2 caused by a CCTG expansion in intron 1 of ZNF9. *Science* 2001; **293**: 864–67.
- 139 Holmes SE, O'Hearn E, Rosenblatt A, et al. A repeat expansion in the gene encoding junctophilin-3 is associated with Huntington disease-like 2. *Nat Genet* 2001; **29**: 377–78.
- 140 Hagerman PJ, Hagerman RJ. The fragile-X premutation: a maturing perspective. *Am J Hum Genet* 2004; **74**: 805–16.
- 141 Sone M, Hayashi T, Tarui H, Agata K, Takeichi M, Nakagawa S. The mRNA-like noncoding RNA Gomafu constitutes a novel nuclear domain in a subset of neurons. *J Cell Sci* 2007; **120**: 2498–506.
- 142 Komine Y, Tanaka NK, Yano R, et al. A novel type of non-coding RNA expressed in the rat brain. *Brain Res Mol Brain Res* 1999; **66**: 1–13.
- 143 Cavaille J, Vitali P, Basyuk E, Huttenhofer A, Bachellerie JP. A novel brain-specific box C/D small nucleolar RNA processed from tandemly repeated introns of a noncoding RNA gene in rats. *J Biol Chem* 2001; **276**: 26374–83.
- 144 French PJ, Bliss TV, O'Connor V. Ntab, a novel non-coding RNA abundantly expressed in rat brain. *Neuroscience* 2001; **108**: 207–15.
- 145 Amaral PP, Neyt C, Wilkins SJ, et al. Complex architecture and regulated expression of the Sox2ot locus during vertebrate development. *RNA* 2009; **15**: 2013–27.
- 146 Bernard D, Prasanth KV, Tripathi V, et al. A long nuclear-retained non-coding RNA regulates synaptogenesis by modulating gene expression. *EMBO J* 2010; **29**: 3082–93.
- 147 Tollervey JR, Curk T, Rogelj B, et al. Characterizing the RNA targets and position-dependent splicing regulation by TDP-43. *Nat Neurosci* 2011; **14**: 452–58.
- 148 Polymenidou M, Lagier-Tourenne C, Hutt KR, et al. Long pre-mRNA depletion and RNA missplicing contribute to neuronal vulnerability from loss of TDP-43. *Nat Neurosci* 2011; **14**: 459–68.
- 149 Mitchell PS, Parkin RK, Kroh EM, et al. Circulating microRNAs as stable blood-based markers for cancer detection. *Proc Natl Acad Sci USA* 2008; **105**: 10513–18.
- 150 Hunter MP, Ismail N, Zhang X, et al. Detection of microRNA expression in human peripheral blood microvesicles. *PLoS One* 2008; **3**: e3694.

- 151 Baraniskin A, Kuhnhen J, Schlegel U, et al. Identification of microRNAs in the cerebrospinal fluid as marker for primary diffuse large B-cell lymphoma of the central nervous system. *Blood* 2011; **117**: 3140–46.
- 152 Kosaka N, Iguchi H, Ochiya T. Circulating microRNA in body fluid: a new potential biomarker for cancer diagnosis and prognosis. *Cancer Sci* 2010; **101**: 2087–92.
- 153 Arroyo JD, Chevillet JR, Kroh EM, et al. Argonaute2 complexes carry a population of circulating microRNAs independent of vesicles in human plasma. *Proc Natl Acad Sci USA* 2011; **108**: 5003–08.
- 154 Cogswell JP, Ward J, Taylor IA, et al. Identification of miRNA changes in Alzheimer's disease brain and CSF yields putative biomarkers and insights into disease pathways. *J Alzheimers Dis* 2008; **14**: 27–41.
- 155 Schwarzenbach H, Hoon DS, Pantel K. Cell-free nucleic acids as biomarkers in cancer patients. *Nat Rev Cancer* 2011; **11**: 426–37.
- 156 Edwards JK, Pasqualini R, Arap W, Calin GA. MicroRNAs and ultraconserved genes as diagnostic markers and therapeutic targets in cancer and cardiovascular diseases. *J Cardiovasc Transl Res* 2010; **3**: 271–79.
- 157 Cooper TA, Wan L, Dreyfuss G. RNA and disease. *Cell* 2009; **136**: 777–93.
- 158 Weinberg MS, Wood MJ. Short non-coding RNA biology and neurodegenerative disorders: novel disease targets and therapeutics. *Hum Mol Genet* 2009; **18**: R27–39.
- 159 Lanford RE, Hildebrandt-Eriksen ES, Petri A, et al. Therapeutic silencing of microRNA-122 in primates with chronic hepatitis C virus infection. *Science* 2010; **327**: 198–201.
- 160 Wang Z, Rao DD, Senzer N, Nemunaitis J. RNA interference and cancer therapy. *Pharm Res* 2011; published online Oct 19. DOI:10.1007/s11095-011-0604-5.
- 161 Poller W, Fechner H. Development of novel cardiovascular therapeutics from small regulatory RNA molecules—an outline of key requirements. *Curr Pharm Des* 2010; **16**: 2252–68.
- 162 Pecot CV, Calin GA, Coleman RL, Lopez-Berestein G, Sood AK. RNA interference in the clinic: challenges and future directions. *Nat Rev Cancer* 2011; **11**: 59–67.
- 163 Kumar P, Wu H, McBride JL, et al. Transvascular delivery of small interfering RNA to the central nervous system. *Nature* 2007; **448**: 39–43.
- 164 Ehlert EM, Eggers R, Niclou SP, Verhaagen J. Cellular toxicity following application of adeno-associated viral vector-mediated RNA interference in the nervous system. *BMC Neurosci* 2010; **11**: 20.
- 165 Martin JN, Wolken N, Brown T, Dauer WT, Ehrlich ME, Gonzalez-Alegre P. Lethal toxicity caused by expression of shRNA in the mouse striatum: implications for therapeutic design. *Gene Ther* 2011; **18**: 666–73.
- 166 Jackson AL, Linsley PS. Recognizing and avoiding siRNA off-target effects for target identification and therapeutic application. *Nat Rev Drug Discov* 2010; **9**: 57–67.