

*Biochemistry*. Author manuscript; available in PMC 2009 January 9.

Published in final edited form as:

Biochemistry. 2006 March 7; 45(9): 3069-3076. doi:10.1021/bi0518141.

# Mini-αB-crystallin:

A functional element of  $\alpha B$ -crystallin with chaperone-like activity

**Jaya Bhattacharyya**, **E.G. Padmanabha Udupa**, **Jing Wang**, and **K. Krishna Sharma**\*

From the Departments of Ophthalmology and Biochemistry, University of Missouri, Columbia, MO 65212, USA

# **Abstract**

 $\alpha$ -Crystallin is a member of the family of small heat shock proteins (sHSP) and is composed of two subunits,  $\alpha$ A-crystallin and  $\alpha$ B-crystallin, which exhibit molecular chaperone-like properties. In a previous study we found that residues 70-88 in  $\alpha$ A-crystallin can function like a molecular chaperone by preventing the aggregation and precipitation of denaturing substrate proteins. In this study we show that the complementary sequence in  $\alpha$ B-crystallin, residues 73-92

(DRFSVNLDVKHFSPEELKVK), is the functional chaperone site of  $\alpha B$ -crystallin. Like the mini- $\alpha A$ -crystallin chaperone, the mini- $\alpha B$ -crystallin chaperone interacts with bis-ANS and also possesses significant  $\beta$ -sheet and random coil structure. Deletion of four residues (DRFS) from the N-terminus or deletion of C-terminus LKVK residues from the 73-92 peptide abolishes the chaperone-like activity against denaturing alcohol dehydrogenase. However, removal of DRFS or HFSPEELKVK is necessary to completely abolish the anti-aggregation property of the peptide in insulin reduction assay. Substitution of Asp at a site corresponding to D80 in  $\alpha B$ -crystallin, with D-Asp or  $\beta$ -Asp results in a significant loss of chaperone-like activity. Kynurenine modification of His in the peptide abolishes the anti-aggregation property of the mini-chaperone. These data suggest that the 73-92 region in  $\alpha B$ -crystallin is one of the substrate binding sites during chaperone activity.

 $\alpha$ -Crystallin, the most abundant protein in the lens belongs to the group of small heat shock proteins (sHSPs).  $\alpha$ -Crystallin exists as a polydisperse aggregate with an average molecular mass of 800 kDa and is composed of  $\alpha$ A- and  $\alpha$ B-subunits, each with a molecular mass of 20 kDa each (1,2). The sequences of the subunits of  $\alpha$ -crystallin have high sequence homology between them (1) and are highly conserved among species (3). Although  $\alpha$ A-crystallin is primarily expressed in the lens,  $\alpha$ B-crystallin is found in other tissues as well (4-7). Extensive studies have not yet delineated the quaternary structure of  $\alpha$ -crystallin or its subunits. Available data suggest that the  $\alpha$ -crystallin subunits have high amounts of  $\beta$ -sheet structure. Like all sHSPs,  $\alpha$ -crystallin displays chaperone-like activity by preventing the aggregation of substrate proteins (8-11). This property has been implicated in the maintenance of lens transparency. Both A and B subunits in  $\alpha$ -crystallin display chaperone-like activity independent of each other and as heteroaggregates (12).

Because the chaperone-like function of  $\alpha$ -crystallin may have an integral role in lens transparency, identification of the critical residues required for the chaperone-like function of  $\alpha$ -crystallin subunits has been the focus of numerous studies. The C-terminal region of  $\alpha$ -crystallin (13,14), Asp<sup>69</sup> of  $\alpha$ -crystallin (15), Phe<sup>71</sup> of  $\alpha$ -crystallin (16), the most hydrophobic N-terminal phenylalanine-rich region in  $\alpha$ B-crystallin (17) and Arg<sup>120</sup> of  $\alpha$ B-crystallin (18) have all been suggested as necessary for chaperone-like activity.

<sup>\*</sup>To whom correspondence should be addressed: Tel: 573-882-8478, Fax: 573-884-4100 e-mail: sharmak@health.missouri.edu

Hydrophobic sites in  $\alpha$ -crystallin have been hypothesized to be responsible for the chaperone-like activity (19,20), as in the case of molecular chaperone GroEL (21). We have previously determined that residues 50-54 and 79-99 in  $\alpha$ A-crystallin and residues 75-103 in  $\alpha$ B-crystallin are hydrophobic sites recognized by the environment-sensitive probe 1,1'-bi(4-anilino)n-naphthalene sulfonic acid (bis-ANS) (22). Because we found that prior binding of bis-ANS to  $\alpha$ -crystallin diminishes chaperone-like activity (23), we concluded that hydrophobic sites participate in  $\alpha$ -crystallin chaperone-like activity. A similar observation has been made for isolated  $\alpha$ A- and  $\alpha$ B-crystallins (24). We have also shown that residues 57-69 and residues 93-107 in  $\alpha$ B-crystallin are involved in binding with ADH (12). In addition, we have determined that a 19-amino-acid residue from  $\alpha$ A-crystallin (residues 70-88, mini- $\alpha$ A-crystallin), which contains bis-ANS-binding sequences, possesses an anti-aggregating property similar to that of  $\alpha$ A-crystallin (25).

Here we report that  $\alpha B$ -crystallin residues 73-92 can independently prevent the aggregation of denatured substrate proteins, similar to the action of native  $\alpha B$ -crystallin. We have coined the term "mini- $\alpha B$ -crystallin" to describe these specific residues. In this study, we describe the smallest peptide sequence in  $\alpha B$ -crystallin that shows an anti-aggregation property and the critical residues required for such chaperone-like function.

#### **EXPERIMENTAL PROCEDURES**

#### **Materials**

 $\gamma$ -Crystallin was isolated from young bovine lenses and purified, as described previously (8).  $\alpha$ B-Crystallin was purified according to the procedure described elsewhere (12). All peptides used in this study were synthesized in the peptide synthesis core facility at the University of Missouri. The synthetic peptides were purified by reversed-phase HPLC and their masses were confirmed by mass spectrometry prior to use. Biuret method was used to estimate the peptide concentrations used in various assays. Bis-ANS was obtained from Molecular Probes, Inc. (Junction City, OR). A stock solution of bis-ANS was prepared in 95% ethanol, and the concentration was determined by absorbance at 385 nm using an extinction coefficient,  $\epsilon_{385}$  = 16,790 cm<sup>-1</sup> M<sup>-1</sup> (26). Insulin and yeast ADH were purchased from Sigma and Biozyme Laboratories, USA, respectively. All other chemicals were of the highest grade commercially available.

## Assay of chaperone-like activity

The capacity of the different peptides to prevent DTT-induced insulin aggregation (27), oxidation-induced  $\gamma$ -crystallin aggregation (28) and heat-induced ADH aggregation (12) was determined. Insulin and  $\gamma$ -crystallin assays were performed at 37°C, whereas ADH assay was conducted at 48°C. All chaperone assays were carried out in 1 ml reaction volumes that contained 50 mM sodium phosphate buffer, pH 7.2 and 0.1 M NaCl. Aggregation of denaturing proteins was monitored by measuring the apparent absorption at 360 nm for all the substrate proteins as a function of time in a Shimadzu spectrophotometer equipped with a multi-cell transporter and a Peltier temperature regulator.

#### **CD Spectroscopy of peptides**

The far-UV CD spectra of purified peptides were recorded at 25°C over a range of 190-250 nm on an Aviv 62DS spectropolarimeter using 0.1 mg/ml concentration of the peptide in 10 mM phosphate buffer, pH 7.2. A quartz cell of 0.1-cm path length was used and five spectra for each sample were taken and averaged.

# **Bis-ANS** binding studies

Fluorescence of bis-ANS bound to peptides was measured using a Jasco FP750 spectrofluorimeter. bis-ANS solution,  $16~\mu moles$ , was added to 0.1~mg/ml peptide solution in 0.05~M phosphate buffer, pH 7.2. The mixture was thoroughly mixed and then incubated for 15~minutes. Fluorescence emission spectra were then recorded at 400-600~nm using an excitation wavelength of 390~nm. The excitation and emission slits were set at 5~nm.

## Modification of mini-αB-crystallin with kynurenine

Purified peptide was incubated with kynurenine (1:5 ratio of peptide to kynurenine) at 37°C for 6 days. The modified peptide was purified by HPLC fitted with a Vydac C18 column and 0 to 80% acetonitrile gradient containing 0.1% trifluoroacetic acid. The kynurenine modification of the peptide was confirmed by mass spectrometric analysis. The chaperone-like activity of the modified peptide was measured using insulin reduction assay.

# Size exclusion chromatography and HPLC analysis

Biogel P-30 gel filtration chromatography was performed to isolate the complex formed between denaturing ADH and mini- $\alpha$ B-crystallin, as described previously for mini- $\alpha$ A-crystallin and ADH studies (25). Following gel filtration, the complex was treated with urea and filtered through a 10 kDa filter, and the filtrate was analyzed in a C18 column (25). Sephadex G-75 chromatography was used to determine retropeptide aggregation.

#### RESULTS AND DISCUSSION

## Chaperone-site and concept of mini-chaperone

The A and B subunits of α-crystallin contain an α-crystallin domain, an N-terminal domain and a C-terminal extension. Figure 1 shows the amino acid residues contributing to these domains in αB-crystallin. The hydrophobic sites on this highly conserved protein have been implicated in playing a role in the chaperone-like activity of αB-crystallin (2,19,20,23, 29-31). Bis-ANS-binding and cross-linking studies have shown that hydrophobic sites in  $\alpha$ crystallin subunits are involved in binding of substrate proteins to αA- and αB-crystallin during chaperone action (22). In a separate experiment, the 83-90 region in αB-crystallin was identified as another hydrophobic site-specific probe 1,5-AZNS-binding site (22). The 1,5-AZNS-labeled peptide sequence was sandwiched between the two bis-ANS-binding sequences in αB-crystallin. The structural differences between bis-ANS and 1,5-AZNS may contribute to the difference in the site of photoincorporation of these two probes to αB-crystallin. In a related study we observed that one (i.e., FSVNLDV) of the two regions in αB-crystallin identified as bis-ANS-binding sequences also contributes to the mellitin-binding region (12). These studies, taken together, suggest that the hydrophobic mellitin-binding site in αBcrystallin subunit is part of the chaperone site. Additional insight into the role of hydrophobic sites in chaperone activity of  $\alpha$ -crystallin subunits came from our novel finding that a peptide consisting a hydrophobic sequence in  $\alpha$ A-crystallin, which also includes the residues from substrate binding site, can mimic the anti-aggregation properties of the parent protein (25). We previously reported that a hydrophobic melittin-binding sequence,

KFVIFLDVKHFSPEDLTVK, in  $\alpha A$ -crystallin (residues 70-88) by itself displays chaperone-like anti-aggregation activity (25). Because the A and B subunits of  $\alpha$ -crystallins have significant sequence homology, we hypothesized that an  $\alpha B$ -crystallin peptide that shows significant sequence homology to  $\alpha A$ -crystallin chaperone-like peptide will also demonstrate chaperone-like activity. Therefore, we investigated whether residues 73-92 of  $\alpha B$ -crystallin (DRFSVNLDVKHFSPEELKVK; mini- $\alpha B$ -crystallin) prevent the aggregation of denatured substrate proteins. This region, (i.e., residues 73-92) is known to be a highly conserved sequence in  $\alpha B$ -crystallin of various species.

## Chaperone-like activity of mini-αB-crystallin

Figure 2 shows the anti-aggregating property of mini-αB-crystallin against DTT-induced aggregation of insulin at 37°C (Figure 2A), oxidation-induced aggregation of γ-crystallin at 37°C (Figure 2B) and thermal aggregation of ADH at 48°C (Fig 2C). As Fig 2 shows, only a portion of  $\alpha$ B-crystallin, represented as mini- $\alpha$ B-crystallin, is sufficient to prevent the aggregation of proteins denatured by heat or chemical methods. Anti-aggregation activity of the peptide was found to be concentration dependent. The data show that 40 µM of the peptide suppresses aggregation of 2.7 μM of ADH by 80%, comparable to the activity of mini-αAcrystallin reported previously (25). The formation of an ADH-mini-αB-crystallin complex was confirmed by gel filtration and HPLC analysis of the reaction mixture (data not shown). The mini-αB-crystallin was found to suppress insulin B chain aggregation very efficiently. A 1:1 molar ratio of mini-αB-chaperone and insulin nearly completely suppressed aggregationinduced light scattering. However, the mini- $\alpha B$ -peptide activity was lower than native  $\alpha B$ crystallin activity when ADH or  $\gamma$ -crystallin were used as substrates and resembled the pattern of mini-αA-crystallin (25). The αA-crystallin was 8-fold more effective in suppressing aggregation-induced light scattering than the mini-αA-crystallin on molar basis when ADH was used as the substrate (25). The different efficiency of native  $\alpha A$ - and  $\alpha B$ -crystallin toward different proteins has been previously observed by us (32) and others (33,34). Therefore, it can be concluded from these observations that the different chaperone-like activity of crystallins reported previously applies to mini-αB-crystallin as well.

Suppression of heat-induced aggregation of ADH by mini- $\alpha B$ -crystallin was tested by adding the peptide after the start of aggregation (Figure 3). The addition of mini- $\alpha B$ -crystallin after 20 and 40 minutes of heat-induced ADH aggregation significantly impeded further aggregation. The aggregated proteins were not solubilized, as in the case of  $\alpha_s$ -casein (35). Therefore, this property of mini- $\alpha B$ -crystallin is consistent with that of other molecular chaperones: the chaperones bind to aggregation-prone proteins and do not solubilize preformed aggregates (36).

After the anti-aggregation property of mini- $\alpha$ B-crystallin was confirmed, we investigated the ability of a retro-sequence of mini- $\alpha$ B-crystallin (KVKLEEPSFHVDLNVSFRD) to suppress heat-induced aggregation of ADH. The retro-sequence was found to not possess antiaggregation activity (Figure 4). Gel chromatograpy on Sephadex G-75 (profile not shown) did not show any evidence of retro-peptide aggregation, suggesting that aggregation of the peptide is not the reason for the loss of activity. Further, the retro-peptide did not form light scattering aggregates during the assay (Figure 4). A similar result was obtained with mini- $\alpha$ A-crystallin and its retropeptide (27). Therefore, the amino-acid sequence, rather than the composition in mini- $\alpha$ B-crystallin, appears to determine its chaperone-like activity.

## Effect of truncation or substitution on the chaperone-like activity of mini-αB-crystallin

Since the mini-chaperone sequence is highly conserved in several species, we prepared several peptides that are truncated forms of the active peptide chaperone to determine the minimum sequence required for chaperone-like activity. Table 1 shows the relative anti-aggregation activity of truncated forms of mini- $\alpha B$ -crystallin against DTT-induced insulin aggregation at 37°C and against ADH aggregation at 48°C. The data show that deletion of two residues - DR from the amino terminal end and VK from the C-terminal of the peptide - partially reduced chaperone-like activity of the peptide against both the substrates. Deletion of 8 residues from the C-terminal of the peptide yielded a peptide that showed significant protection against insulin aggregation but no ability to suppress aggregation of denaturing ADH. On the other hand, deletion of 4 residues (DRFS) from the N-terminal end of the peptide completely abolished the anti-aggregation property of the peptide against both substrates, but removal of DR residues had no significant effect. Furthermore, substitution of F with A in N-terminal

region with A resulted in complete loss of activity. These results suggest that, like the N-terminal region F of mini- $\alpha$ A-crystallin (25), the F in the terminal region of mini- $\alpha$ B-crystallin may be essential for chaperone-like activity of the peptide. The critical role of F of mini- $\alpha$ A-crystallin (corresponding to F71 of  $\alpha$ A-crystallin) was confirmed by a site-directed mutagenesis study in which the mutant  $\alpha$ A-crystallin was found to not display chaperone activity at physiological temperatures (16).

#### Structural features of native and substituted mini-αB-crystallin

The emission maximum of bis-ANS, a hydrophobic site-specific probe, shifts to a lower wavelength when it interacts with proteins or peptides that have significant hydrophobicity (26). Binding of bis-ANS to the peptide chaperone resulted in a shift in the emission maximum of the fluorophore, to 499 nm from 520 nm in aqueous media (data not shown). The far-UV CD spectra of the peptide in buffer showed significant amounts of β-sheet and random coil structure (Fig. 5). Compared with mini-αA-crystallin (25,27), mini-αB-crystallin demonstrated a slightly greater, non-compact conformation, which may be due to its random coil content being greater than its  $\beta$ -sheet content. The conformation and affinity of mini- $\alpha$ B-crystallin and its substituted forms to hydrophobic probe were determined by far-UV CD spectra and bis-ANS binding studies (Figures 5 and Table I). The studies were performed to identify any correlation between the peptide hydrophobicity and chaperone-like activity. When compared to mini- $\alpha$ B-crystallin (Peptide 1), the peptides-3 to 6 (Table I) showed a decrease in the  $\beta$ -sheet content and an increase in the random coil content (Figure 5). Other truncated or Ala-substituted mini- $\alpha$ B-peptides were also mostly in random-coiled form, with negligible  $\beta$ -sheet content, and did not bind bis-ANS (data not shown). All mutated or truncated peptides, with the exception of peptide 5, were found to bind significantly less bis-ANS than mini-αB-crystallin (Table I). The binding of bis-ANS to peptide 5 was comparable to that of mini-αB-crystallin (Peptide 1).

Previously we reported that mini-αA-crystallin regains most of its secondary structure upon refolding from its heat-denatured conformation and that the chaperone-like activity of the native and heat-treated mini-αA-crystallin was not significantly different (27). To investigate the effect of heat treatment on mini-αB-crystallin a known amount of the peptide was heated to 60°C in assay buffer. After 30 min at 60°C, the peptide solution was cooled to room temperature and subjected to CD spectroscopy. Compared to the far-UV CD spectrum of native mini-αB-crystallin, the spectrum of mini-αB-crystallin after thermal unfolding and refolding showed a more random-coiled structure (trace 2 in Figure 5). Surprisingly, the chaperone-like activity of the refolded peptide was comparable to that of the native peptide. The activities of the native and heat-treated peptides were comparable during DTT-induced insulin reduction assay (Figure 6A) and temperature-induced ADH denaturation assay (Figure 6B). Similar data were obtained when citratesynthase was used in place of ADH (not shown). Therefore, it appears that although the secondary structure of refolded mini-αB-crystallin is more disrupted than that of its native state, the structural differences do not affect the chaperone-like activity of the peptide. In this aspect the peptide behaves like a major random-coiled protein  $\alpha$ -casein, which has been reported to possess anti-aggregation properties against denatured substrate proteins (35). Therefore, a combination of hydrophobicity and ionic-interactions stemming from the amphipathic nature of the peptide seems to determine the activity. This conclusion is also based on the observations that a) removal of charged K-x-K residues from the C-terminus of the peptide resulted in the loss of activity, b) substitution of D with A (peptide 5) reduced the chaperone-like activity and c) substitution of hydrophobic F with less hydrophobic A at the N-terminal region of the peptides abolished the chaperone-like activity.

## Effect of substitution with D-Asp or β-Asp on chaperone-like activity

Studies of the effects of substitutions at various locations in the peptide chaperone revealed that the substitution of hydrophillic amino acid residues D80 and H83 with Ala preserved most of the chaperone-like activity of mini- $\alpha$ B-crystallin (Table 1). However, when substitution was with D-Asp or  $\beta$ -Asp at a position corresponding to D80 in  $\alpha$ B-crystallin, the peptide lost its ability to suppress aggregation of denaturing ADH (Figure 7) and both forms of the peptide had also lost the  $\beta$ -sheet content when examined by CD spectroscopy (data not shown). A modest (about 10%) increase in the ADH aggregation in presence of D-Asp peptides was observed. This effect is most likely due to the interaction of D-Asp-peptide with denaturing ADH. In the past we have observed that modified crystallin peptides increase the aggregation of denaturing ADH (37). The presence of D-Asp or  $\beta$ -Asp likely alters the peptide structure in such a way that it becomes less favorable for the interaction. The presence of D-Asp or  $\beta$ -Asp in peptides is known to contribute to the structural change (38,39).

## Effect of modification of His with kynurenine on mini-αB-crystallin chaperone-like activity

The post-translational modification of  $\alpha$ -crystallin subunits has been attributed to the loss of chaperone-like activity (40). Truscot et al (41) reported that His residues in αB-crystallin are susceptible to modification by the tryptophan metabolite kynurenine. Because H83 of αBcrystallin is at the chaperone site, we investigated whether the His present in mini-αB-crystallin chaperone is susceptible to modification by kynurenine and whether such a modification leads to a loss in chaperone-like activity of the peptide. As shown in Fig. 8, the modification of miniαB-crystallin chaperone by kynurenine resulted in more than an 80% loss of chaperone-like activity. Therefore, the age-related modification of  $\alpha B$ -crystallin by kynurenine likely leads to diminished chaperone-like activity. Furthermore, the Lys in αB-crystallin is also susceptible to glycation. We have previously shown that glycation of α-crystallin decreases ANS binding and reduces chaperone-like activity of  $\alpha$ -crystallin (22). Others have shown that Lys in  $\alpha$ Bcrystallin chaperone site is susceptible to modification (42). Since Lys 92 is the C-terminal residue on mini-αB-crystallin chaperone, glycation likely affects the chaperone site and interferes with the binding of denaturing proteins. While we did not measure the chaperonelike activity of kynurenine-modified, glycated \alpha B-crystallin, these two modifications are likely to reduce significantly the chaperone-like activity of the protein. On the basis of data from this study, the loss of chaperone activity we observed in an earlier study (22) can be attributed to the structural changes at the chaperone site. Deamidation of a residue, N78, at the chaperone site has been shown to diminish the ability of the protein to suppress aggregation of denaturing proteins (43).

## **CONCLUDING REMARKS**

Both subunits of the sHSP  $\alpha$ -crystallin display chaperone-like activity. Previous studies have shown that the hydrophobic region in the  $\alpha$ -crystallin domain of these proteins is the binding site for denaturing proteins (22). Site-directed mutagenesis study and the demonstration of chaperone-like activity by a 19mer peptide, DFVIFLDVKHFSPEDLTVK (71-88 residues of  $\alpha$ A-crystallin), have confirmed that in  $\alpha$ A-crystallin the beginning part of  $\alpha$ -crystallin domain is the chaperone site (16,24). The present study documents that the corresponding sequence in  $\alpha$ B-crystallin, DRFSVNLDVKHFSPEELKVK, is a chaperone site as well. During a pin array study, the DRFSVNLDVK sequence (residues 73-82 in  $\alpha$ B-crystallin) was also identified as one of the chaperone sites (44). In a related study, the same authors reported that residues 75-82 in  $\alpha$ B-crystallin are involved in subunit interaction (45). However, the mechanism of function of the same region as chaperone site and subunit interaction site are yet to be determined. The peptide array study (44) also identified 9-20, 43-58, 113-120, 131-138, 141-148 and 157-164 sequences in  $\alpha$ B-crystallin as target protein binding sites. The same study also reported that synthetic peptides 73-85 and 131-141 inhibited thermal aggregation of  $\beta$ H-crystallin, ADH

and CS, with 73-85 sequence showing the maximal activity. The peptides identified after pin array study were less efficient in suppressing the aggregation of ADH compared to the activity of  $\alpha B$ -peptide 73-92 (this study) or the  $\alpha A$ -peptide that represented the chaperone site 70-88, as reported previously (25). It is yet to be determined whether the peptides corresponding to the remainder of the binding sites determined by pin array studies function independently as mini-chaperones. Earlier we reported that peptide TSLSPFYLRPPSFLRA (42-57 sequence in  $\alpha B$ -crystallin), identified as  $\alpha A$ -crystallin interacting region and contains the chaperone interactive domain reported by Ghosh et. al (44) has no chaperone-like activity. Likewise, several other peptides representing other regions of  $\alpha B$ -crystallin showed no chaperone-like activity (data not shown) during in vitro aggregation assays. We hypothesize that a number of mutations affecting the chaperone-like activity of  $\alpha A$ - or  $\alpha B$ -crystallins impact the structure of the mutant protein in such a way that the chaperone site 70-88 in  $\alpha A$ - or 73-92 in  $\alpha B$ -crystallin is masked partially or fully. Further studies are required to prove this hypotheses.

Mini- $\alpha$ A- and mini- $\alpha$ B-crystallins are the shortest sequences in sHSPs capable of displaying maximal anti-aggregation activity. In a previous study we found that a 21-mer peptide (residues 40-60) of yeast ADH also shows anti-aggregation activity (46). However, the yeast ADH peptide was less active than mini- $\alpha$ A- or mini- $\alpha$ B-crystallins. The demonstration of diminished activity following modification of the chaperone peptide with kynurenine may explain in part the decrease in the chaperone activity of  $\alpha$ -crystallin isolated from the aged lenses.

## **ACKNOWLEDGEMENTS**

We thank Dr. B.J. Ortwerth for helpful discussions on this project and Kathleen Connolly for technical support. We thank Sharon Morey for help with preparation of the manuscript.

This work was supported by National Institute of health Grants EY 11981 and EY 14795, to K.K.S., a grant-in-aid from Research to Prevent Blindness (RPB) to the department and a Lew Wasserman Merit Award (to K.K.S. from RPB). Portions of this work were presented at the annual meeting of Association for Research in Vision and Ophthalmology in 2002 and 2003.

#### ABBREVIATIONS

sHSPS, small heat shock proteins; bis-ANS, 1,1'-bi(4-anilino) napthalene-5,5'-disulphonic acid; ADH, alcohol dehydrogenase; 1,5-AZNS, 1-azidonapthalene 5-sulphonate..

#### REFERENCES

- Ingolia TD, Craig EA. Four small drosophila heat shock proteins are related to each other and to mammalian α-crystallin. Proc. Natl. Acad. Sc.i U S A 1982;79:2360–2364.
- 2. Groenen PJTA, Merck KB, de Jong WW, Bloemendal H. Structure and modification of the junior chaperone α-crystallin. Eur. J Biochem 1994;225:1–19. [PubMed: 7925426]
- 3. Sax CM, Piatygorsky J. Expression of the alpha-crystallin/small heat-shock protein/molecular chaperone. Adv. Enzymol. Related Areas. Mol. Biol 1994;69:155–201.
- 4. Bhat SP, Nagineni CN. αB-subunit of lens specific protein α-crystallin is present in other ocular and non-ocular tissues. Biochem. Biophys. Res. Commun 1989;158:319–325. [PubMed: 2912453]
- 5. Iwaki T, Kume-Iwaki T, Liem RKH, Goldman JE. αB-crystallin is accumulated in non-lenticular tissues and accumulates in Alexander's disease brain. Cell 1989;57:71–78. [PubMed: 2539261]
- 6. Kato K, Shinohara H, Kurobe N, Goto S, Inaguma Y, Ohshima K. Immunoreactive  $\alpha A$ -crystallin in rat non-lenticular tissues detected with sensitive immunoassay method. Biochim. Biophys. Acta 1991;1080:173–180. [PubMed: 1932094]
- 7. Srinivasan AN, Nagineni CN, Bhat SP. Alpha A-crystallin is expressed in non-ocular tissues. J. Biol. Chem 1992;267:23337–23341. [PubMed: 1429679]
- 8. Horwitz J. A-crystallin can function as a molecular chaperone. Proc. Natl. Acad. Sci. U S A 1992;89:10449–10453. [PubMed: 1438232]

 Merck KB, Groenen PJTA, Voorter CEM, de Haard-Hoekman WA, Horwitz J, Bloemendal H, de Jong WW. Structural and functional similarities of bovine alpha-crystallin and mouse small heat-shock protein. A family of chaperones. J. Biol. Chem 1993;268:1046–1052. [PubMed: 8093449]

- 10. Rao PV, Huang Q-I, Horwitz J, Zigler JS Jr. Evidence that alpha-crystallin prevents non-specific protein aggregation in the intact eye lens. Biochim. Biophys. Acta 1995;1245:439–447. [PubMed: 8541324]
- 11. Farahbakhsh ZT, Huang QL, Ding LL, Altenbach C, Steinhoff HJ, Horwitz J, Hubbell WL. Interaction of alpha crystallin with spin-labeled peptides. Biochemistry 1995;34:509–516. [PubMed: 7819243]
- Sharma KK, Kaur H, Kester K. Functional elements in molecular chaperone alpha crystalline: Identification of binding sites in αB-crystallin. Biochem. Biophys. Res. Commun 1997;239:217–222. [PubMed: 9345298]
- Takemoto L, Emmons T, Horwitz J. The C-terminal region of alpha-crystallin: involvement in protection against heat-induced denaturation. Biochem. J 1993;294:435–438. [PubMed: 8373358]
- 14. Andley UP, Mathur S, Griest TA, Petrash JM. Cloning, expression and chaperone-like activity of human alpha-crystallin. J. Biol. Chem 1996;271:31973–31980. [PubMed: 8943244]
- 15. Smulders RH, Merck KB, Aendekerk J, Horwitz J, Takemoto L, Slingsby C, Bloemendal H, De Jong WW. The mutation Asp69Ser affects the chaperone-like activity of alpha A-crystallin. Eur. J. Biochem 1995;232:834–838. [PubMed: 7588723]
- 16. Santhoshkumar P, Sharma KK. Phe71 is essential for chaperone-like function in αA-crystallin. J. Biol. Chem 2001;276:47094–47099. [PubMed: 11598124]
- 17. Plater ML, Goode D, Crabbe MJ. Effects of site-directed mutations on the chaperone-like activity of alpha B-crystallin. J. Biol. Chem 1996;271:28558–28566. [PubMed: 8910485]
- 18. Bova MP, Yaron O, Huang Q, Ding L, Haley DA, Stewart PL, Horwitz J. Mutation R120G in αB-crystallin, which is linked to a desmin-related myopathy, results in an irregular structure and defective chaperone-like function. Proc. Natl. Acad. Sci. U S A 1999;96:6137–6142. [PubMed: 10339554]
- Raman B, Rao CM. Chaperone-like activity and quaternary structure of alpha-crystallin. J. Biol. Chem 1994;269:27264–27268. [PubMed: 7961635]
- Das KP, Surewicz WK. Temperature-induced exposure of hydrophobic surfaces and its effect on the chaperone activity of alpha-crystallin. FEBS Lett 1995;369:321–325. [PubMed: 7649280]
- Seale JW, Martinez JL, Horowitz PM. Photoincorporation of 4,4'-Bis(1-anilino-8-naphthalenesulfonic acid) into the apical domain of GroEL: Specific information from a nonspecific probe. Biochemistry 1995;34:7443–7449. [PubMed: 7779787]
- 22. Sharma KK, Kumar GS, Murphy AS, Kester K. Identification of 1,1'Bi(4-anilino)naphthalene-5,5'-disulfonic acid binding sequences in α-crystallin. J. Biol. Chem 1998;273:15474–15478. [PubMed: 9624133]
- 23. Sharma KK, Kaur H, Kumar GS, Kester K. Interaction of 1,1'Bi(4-anilino)naphthalene-5,5'-disulfonic acid with  $\alpha$ -crystallin. J. Biol. Chem 1998;273:8965–8970. [PubMed: 9535881]
- Kumar MS, Kapoor M, Sinha S, Reddy GB. Insights into hydrophobicity and the chaperone-like function of alpha A- and alpha B-crystallins: an isothermal titration calorimetric study. J Biol. Chem 2005;280:21726–21730. [PubMed: 15817465]
- 25. Sharma KK, Kumar RS, Kumar GS, Quinn PT. Synthesis and characterization of a peptide identified as functional element in  $\alpha$ A-crystallin. J. Biol. Chem 2000;275:3767–3771. [PubMed: 10660525]
- 26. Sudhakar K, Fay PJ. Exposed Hydrophobic Sites in Factor VIII and Isolated Subunits. J. Biol. Chem 1996;271:23015–23021. [PubMed: 8798489]
- 27. Bhattacharyya J, Sharma KK. Conformational specificity of mini-alpha A-crystallin as a molecular chaperone. J. Pept. Res 2001;57:428–434. [PubMed: 11350603]
- 28. Kumar RS, Sharma KK. Synthetic  $\alpha$ A-crystallin functional element has chaperone-like activity toward oxidized  $\gamma$ -crystallin. J. Pept. Res 2000;56:157–164. [PubMed: 11007272]
- 29. Raman B, Rao CM. Chaperone-like activity and temperature-induced structural changes of alphacrystallin. J. Biol. Chem 1997;272:23559–23564. [PubMed: 9295293]
- 30. Das BK, Liang JJ. Detection and characterization of alpha-crystallin intermediate with maximal chaperone-like activity. Biochem. Biophys. Res. Commun 1997;236:370–374. [PubMed: 9240443]

31. Smulders RH, de Jong WW. The hydrophobic probe 4,4'-bis(1-anilino-8-naphthalene sulfonic acid) is specifically photoincorporated into the N-terminal domain of alpha B-crystallin. FEBS Lett 1997;409:101–104. [PubMed: 9199512]

- 32. Santhoshkumar P, Sharma KK. Analysis of alpha crystallin chaperone function using restriction enzymes and citrate synthase. Mol. Vision 2001;7:172–177.
- 33. Horwitz J. Alpha crystallin. Exp. Eye Res 2003;76:145–153. [PubMed: 12565801]
- 34. Kumar MS, Reddy PY, Kumar PA, Surolia I, Reddy GB. Effect of dicarbonyl-induced browning on alpha-crystallin chaperone-like activity: physiological significance and caveats of in vitro aggregation assays. Biochem. J 2004;379:273–282. [PubMed: 14711370]
- 35. Bhattacharyya J, Das KP. Molecular chaperone-like properties of an unfolded protein,  $\alpha_s$ -Casein. J. Biol. Chem 1999;274:15505–15509. [PubMed: 10336443]
- 36. Fenton WA, Kashi Y, Furtak K, Horwich AL. Residues in chaperonin GroEL required for polypeptide binding and release. Nature 1994;371:614–619. [PubMed: 7935796]
- 37. Padmanabha Udupa EG, Sharma KK. Effect of Oxidized βB3-Crystallin Peptide on Lens β<sub>L</sub>-Crystallin: Interaction with βB2-Crystallin. Invest. Ophthalmol. Vis. Sci 2005;46:2514–2521. [PubMed: 15980243]
- 38. Vyas SB, Duffy LK. Stabilization of secondary structure of Alzheimer beta-protein by aluminum(III) ions and D-Asp substitutions. Biochem. Biophys. Res. Commun 1995;206:718–723. [PubMed: 7826392]
- 39. Mizuno A, Fujii N, Akaboshi M, Momose Y, Shimo-oka T. Changes of secondary structure detected by laser Raman spectroscopy in model peptides of human alpha A-crystallin due to substitution of D-aspartyl residues for L-aspartyl residues. Japanese J. Ophthalmol 2000;44:354–359.
- 40. Bloemendal H, deJong W, Jaenicke R, Lubsen NH, Slingsby C, Tardieu A. Ageing and vision: structure, stability and function of lens crystallins. Prog. Biophys. and Mol. Biol 2004;86:407–485. [PubMed: 15302206]
- 41. Aquilina JA, Truscott RJ. Kynurenine binds to the peptide binding region of the chaperone alphaB-crystallin. Biochem. Biophys. Res. Commun 2001;285:1107–1113. [PubMed: 11478768]
- 42. Ortwerth BJ, Slight SH, Prabhakaram M, Sun Y, Smith JB. Site-specific glycation of lens crystallins by ascorbic acid. Biochim. Biophys. Acta 1992;1117:207–215. [PubMed: 1525182]
- 43. Gupta R, Srivastava OP. Effect of deamidation of Asparagine 146 on functional and structural properties of human lens  $\alpha B$ -crystallin. Invest. Ophthalmol. Vis. Sci 2004;45:206–214. [PubMed: 14691175]
- 44. Ghosh JG, Estrada MR, Clark JI. Interactive domains for chaperone activity in the small heat shock protein, human αB-crystallin. Biochemistry 2005;44:14854–14869. [PubMed: 16274233]
- 45. Gosh JG, Clark JI. Insights into the domains required for dimerization and assembly of human  $\alpha$ B-crystallin. Protein Science 2005;14:684–694. [PubMed: 15722445]
- 46. Bhattacharyya J, Santhoshkumar P, Sharma KK. A peptide sequence YSGVCHTDLHAWHGDWPLPVK (40-60) in yeast alcohol dehydrogenase prevents the aggregation of denatured substrate proteins. Biochem. Biophys. Res. Commun 2003;307:1–7. [PubMed: 12849973]

Bhattacharyya et al.

Page 10

# FIGURE 1.

<--C-terminal extension-->

 $\alpha B$ -crystallin sequence showing the mini- $\alpha B$ -crystallin chaperone site. The bis-ANS-binding region is underscored and the mini-chaperone sequence is shown in bold. Mellitin-binding region is within the brackets. 1,5-AZNS binding sequence is shown in italics. The ligand binding sites are taken from Ref 22.




FIGURE 2.

Aggregation of denaturing proteins in the presence of mini- $\alpha$ B-crystallin. Light scattering at 360 nm of 0.4 mg/ml insulin and 20 mM DTT (A), 0.25 mg/ml  $\gamma$ -crystallin (B) and 0.3 mg/ml ADH (C) in the absence and presence of mini- $\alpha$ B-crystallin. Insulin and  $\gamma$ -crystallin assays were done at 37°C, whereas ADH assay was performed at 48°C. 2A: diamond, insulin; square, insulin+ 0.05 mg/ml mini- $\alpha$ B-crystallin; triangle, insulin+ 0.1 mg/ml mini- $\alpha$ B-crystallin. 2B: diamond,  $\gamma$ -crystallin; square,  $\gamma$ -crystallin + 0.05 mg/ml mini- $\alpha$ B-crystallin; triangle,  $\gamma$ -crystallin + 0.1 mg/ml mini- $\alpha$ B-crystallin; diamond, ADH+ 0.1 mg/ml mini- $\alpha$ B-crystallin.




FIGURE 3. Effect of the addition of mini- $\alpha$ B-crystallin on ADH aggregation. The aggregation of 0.3 mg of ADH at 48°C in 50 mM phosphate buffer containing 0.1 M NaCl was measured in the presence and absence of mini- $\alpha$ B-crystallin. Mini- $\alpha$ B-crystallin, 0.1 mg, was added 20 min and 40 min after initiation of heat-induced aggregation (shown by arrows) and the assay was continued for 65 min. (Open triangle, ADH; filled square and open square, ADH + mini- $\alpha$ B-crystallin added.)




FIGURE 4. ADH aggregation in the presence and absence of retro-mini- $\alpha B$ -crystallins. The aggregation of 0.3 mg of ADH at 48°C in 50 mM phosphate buffer containing 0.1 M NaCl was measured in the absence and presence of 0.1 mg of mini- $\alpha B$ -crystallin and retro-mini- $\alpha B$ -crystallin. (filled circle, ADH; filled square, ADH+ mini- $\alpha B$ -crystallin; open square, ADH+ retro-mini- $\alpha B$ -crystallin; open circle, retro-mini- $\alpha B$ -crystallin.)




FIGURE 5. Far-UV CD spectra of native, mutant and heat-treated mini- $\alpha B$ -crystallins. The sequences of peptides 1 and 3-6 are shown in Table 1 and the experimental conditions are described under methods. Trace 2 is the spectra for peptide 1 after heat treatment.




FIGURE 6. Effect of heat treatment on chaperone-like activity of mini-αB-crystallin. The peptide was heated to 60°C in assay buffer. After 30 min at 60°C the peptide solution was cooled to room temperature and the residual chaperone-like activity was measured. A) Light scattering was

measured at 360 nm for 0.4 mg/ml insulin and 20 mM DTT at 37°C in the absence (diamond) and presence of native 0.1 mg/ml mini- $\alpha B$ -crystallin (triangle) and 0.1 mg/ml refolded mini- $\alpha B$ -crystallin (square) after thermal unfolding. B) Light scattering at 360 nm of 0.4 mg/ml ADH in the absence (diamond) and presence of 0.1 mg of native (open square) and heat-treated (circle) mini- $\alpha B$ -crystallin at 48°C.




FIGURE 7. Effect of D-Asp or β-Asp substitution on chaperone-like activity of mini- $\alpha$ B-crystallin. DTT-induced aggregation of 0.3 mg of insulin in presence and absence 0.1 mg of substituted and native mini- $\alpha$ B-crystallin was measured at 37°C in 1 ml of 50 mM phosphate buffer, pH 7.0. (square, insulin; circle, insulin+ β-Asp containing mini- $\alpha$ B-crystallin; diamond, insulin+ D-Asp containing mini- $\alpha$ B-crystallin; triangle, insulin+ mini- $\alpha$ B-crystallin.)




FIGURE 8. Effect of His modification in mini- $\alpha$ B-crystallin by kynurenine on chaperone-like activity. DTT-induced aggregation of 0.3 mg of insulin in presence and absence 0.1 mg of kynurenine-modified mini- $\alpha$ B-crystallin was measured at 37°C in 1 ml of 50 mM phosphate buffer, pH 7.0. (triangle, insulin; square, Insulin + kynurenine modified mini- $\alpha$ B-crystallin; diamond, insulin + 0.1 mg mini- $\alpha$ B-crystallin.)

Table 1 Effect of truncation and substitution on anti-aggregation property of mini-\$\alpha\$B-crystallin

Peptides	Percent protection		Dia ANG hindina
	ADH assay	Insulin assay	Bis-ANS binding
DRFSVNLDVKHFSPEELKVK (Peptide1)	84	86	32 ± 2
DRFSVNLDVKHFSPEELK (Peptide 3)	64	70	$11 \pm 0.5$
DRFSVNLDVKHFSPEE	0	52	ND
DRFSVNLDVKHFSP	0	63	ND
DRFSVNLDVKHF	0	64	ND
DRFSVNLDVK	0	0	ND
FSVNLDVKHFSPEELKVK (Peptide 4)	70	74	$8 \pm 0.4$
VNLDVKHFSPEELKVK	0	0	ND
FSVNLAVKHFSPEELKVK (Peptide 5)	24	67	$27 \pm 1.5$
FSVNLDVKAFSPEELKVK (Peptide 6)	69	72	$9 \pm 0.5$
DRASVNLDVKHFSPEELKVK (Peptide 7)	0	ND	ND

Relative activity of the truncated forms of mini- $\alpha B$ -crystallin against ADH aggregation at 48°C and DTT-induced insulin aggregation at 37°C was measured as described under methods. The fluorescence intensity of Bis-ANS bound to peptides was recorded, as described under methods.

ND: not determined